Notably, only the IES-R was evaluated against a “gold standard” PTSD assessment (i.e., the Clinician-Administered PTSD Scale) in this patient population (4). In contrast, the PTSS-14 was evaluated against the Post-traumatic Stress Diagnostic Scale (PDS), which is another patient-reported outcome measure (5). Hence, the measurement properties of the PTSS-14 have not been as rigorously evaluated as the IES-R, and thus the internal validity of the PTSS-14 in acute respiratory failure survivors remains uncertain. Moreover, although the PTSS-14 was administered at three points (4–14 days, 2 months, and 3 months post-ICU discharge) in the validation study, the PDS was administered only at 3 months post-ICU discharge (5). Despite this, the area under the receiver operating characteristics curve for the PTSS-14 was calculated for all three points, with the authors recommending a PTSS-14 threshold of 45 based on the highest area under the receiver operating characteristics curve from the 2-month assessment (5).

This recommendation may be problematic, given that the PTSS-14 and PDS were not administered at the same point. Notably, Sjoding and colleagues only used the sensitivity/specificity (86%/97%) of the PTSS-14 (vs. PDS) at the 2-month assessment. However, at the 3-month assessment, which is when the PDS and PTSS-14 were both administered in the validation study, the PTSS-14 sensitivity/specificity (100%/84%) were very similar to those of the IES-R at the recommended threshold of 1.6 (100%/85%). Had Sjoding and colleagues used the 3-month sensitivity/specificity estimates for the PTSS-14, their results and conclusions would have changed, given very little difference in the measurement properties of the PTSS-14 versus IES-R.

Several other aspects of the PTSS-14 and its validation study are noteworthy, given a goal of measuring PTSD symptoms. First, the validation study was conducted in the United Kingdom and did not include any patients from ethnic minorities (5), potentially limiting its external validity. In addition, of the 14 questions included in the PTSS-14, only two (14%) focus on the mandatory *Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition*, PTSD symptom criterion of intrusion/re-experiencing, whereas two (14%) focus on symptoms not relevant for PTSD (i.e., mood swings and muscle tension). Hence, there may be limitations in the ability of the PTSS-14 for measuring PTSD symptoms.

Finally, as opposed to the IES-R (which does prompt patients to respond to its questions specifically with respect to their ICU experience), the PTSS-14 does not anchor patient respondents to any specific traumatic event when answering each item,

which could give rise to PTSD symptoms being reported from a pre-ICU event, as recently highlighted in the critical care literature (6).

Hence, we believe that these issues regarding the PTSS-14, and its evaluation by Sjoding and colleagues, are valuable considerations when reviewing their recent publication.

Author disclosures are available with the text of this letter at www.atsjournals.org.

Ann M. Parker, M.D.
Sina Nikayin, M.D.
O. Joseph Bienvenu, M.D., Ph.D.
Dale M. Needham, M.D., Ph.D.
Johns Hopkins University School of Medicine
Baltimore, Maryland

ORCID ID: 0000-0001-9024-038X (S.N.).

References

- Sjoding MW, Schoenfeld DA, Brown SM, Hough CL, Yealy DM, Moss M, Angus DC, Iwashyna TJ; NHLBI Prevention and Early Treatment of Acute Lung Injury (PETAL) Network. Power calculations to select instruments for clinical trial secondary endpoints: a case study of instrument selection for post-traumatic stress symptoms in subjects with acute respiratory distress syndrome. *Ann Am Thorac Soc* 2017;14:110–117.
- Hodgson CL, Turnbull AE, Iwashyna TJ, Parker A, Davis W, Bingham CO III, Watts NR, Finfer S, Needham DM. Core domains in evaluating patient outcomes after acute respiratory failure: international multidisciplinary clinician consultation. *Phys Ther* 2017;97:168–174.
- Turnbull AE, Rabiee A, Davis WE, Nasser MF, Venna VR, Lolitha R, Hopkins RO, Bienvenu OJ, Robinson KA, Needham DM. Outcome measurement in ICU survivorship research from 1970 to 2013: a scoping review of 425 publications. *Crit Care Med* 2016;44:1267–1277.
- Bienvenu OJ, Williams JB, Yang A, Hopkins RO, Needham DM. Posttraumatic stress disorder in survivors of acute lung injury: evaluating the Impact of Event Scale-Revised. *Chest* 2013;144:24–31.
- Twigg E, Humphris G, Jones C, Bramwell R, Griffiths RD. Use of a screening questionnaire for post-traumatic stress disorder (PTSD) on a sample of UK ICU patients. *Acta Anaesthesiol Scand* 2008;52:202–208.
- Patel MB, Jackson JC, Morandi A, Girard TD, Hughes CG, Thompson JL, Kiehl AL, Elstad MR, Wasserstein ML, Goodman RB, et al. Incidence and risk factors for intensive care unit-related post-traumatic stress disorder in veterans and civilians. *Am J Respir Crit Care Med* 2016;193:1373–1381.

Copyright © 2017 by the American Thoracic Society

Reply

We thank Dr. Parker and colleagues for their comments regarding the validation study of the Posttraumatic Stress Symptoms-14 instrument we cited in our manuscript examining the power of

instruments to measure differences in clinical outcomes. We agree there are many issues to consider when choosing a specific instrument to measure an outcome for a clinical trial. Issues stressed in their letter included the importance of considering how rigorously the validation of the instrument was performed in similar populations.

Our manuscript examined how the power of two instruments can be compared for their ability to detect statistically significant differences in outcome rates between trial arms. Our results show that in general, specificity is more important than sensitivity for statistical power to detect a difference between the arms when the outcome has a low prevalence, a result potentially counterintuitive to clinicians.

Supported by the National Institute of Health, National Heart Lung, and Blood Institute U01 Grants HL123009-01, HL123010-01, HL123004-01, HL123022-01, HL122989-01, HL123008-01, HL123027-01, HL123020-01, HL123018-01, HL123031-01, HL123033-01, HL122998-01, and HL123023-01 and a grant to M.W.S. from the National Heart Lung, and Blood Institute (T32HL007749).

Author Contributions: All authors were involved in the conceptualization, drafting, revisions, and final approval of this manuscript.

Our results also show that the reverse becomes true as the prevalence of the outcome exceeds 50%, where sensitivity is likely more important.

As we state in our article (1), using this approach to compare two instruments requires an investigator to make some assumptions about the performance of each instrument in the study population. Typically, information about an instrument's sensitivity and specificity must be obtained from previous validation studies. Therefore, the points made by Dr. Parker and colleagues are important when considering whether to apply results from a particular validation study to a given design application. Our key contention was not that any specific instrument was superior for detecting a true difference in a specific outcome between the two arms in a trial. Instead, we contend that the approach we outline allows clinical trialists to quantitatively assess the trade-offs across a range of plausible instrument performance. We hope that these points are complementary to the important points made by Dr. Parker and colleagues.

Author disclosures are available with the text of this letter at www.atsjournals.org.

Michael W. Sjoding, M.D., M.S.
University of Michigan
Ann Arbor, Michigan

Samuel M. Brown, M.D., M.S.
University of Utah School of Medicine
Salt Lake City, Utah

Marc Moss, M.D.
University of Colorado
Denver, Colorado

Derek C. Angus, M.D., M.P.H.
University of Pittsburgh
Pittsburgh, Pennsylvania

Theodore J. Iwashyna, M.D., Ph.D.
University of Michigan
Ann Arbor, Michigan

and

Veterans Affairs Center for Clinical Management Research
Ann Arbor, Michigan

Reference

- 1 Sjoding MW, Schoenfeld DA, Brown SM, Hough CL, Yealy DM, Moss M, Angus DC, Iwashyna TJ; NHLBI Prevention and Early Treatment of Acute Lung Injury (PETAL) Network. Power calculations to select instruments for clinical trial secondary endpoints: a case study of instrument selection for post-traumatic stress symptoms in subjects with acute respiratory distress syndrome. *Ann Am Thorac Soc* 2017;14:110–117.

Copyright © 2017 by the American Thoracic Society