Published in final edited form as:

J Racial Ethn Health Disparities. 2015 June ; 2(2): 176–183. doi:10.1007/s40615-014-0061-z.

The Relationship Between Education and Prostate-Specific Antigen Testing Among Urban African American Medicare Beneficiaries

Mohammad Khalid Hararah,

University of Wisconsin Madison School of Medicine and Public Health, Madison, WI, USA

Craig Evan Pollack,

Department of General Internal Medicine, Johns Hopkins School of Medicine, Baltimore, MD, USA

Mary A. Garza,

Department of Behavioral and Community Health, School of Public Health, University of Maryland, College Park, MD, USA

Hsin-Chieh Yeh,

Department of General Internal Medicine, Johns Hopkins School of Medicine, Baltimore, MD, USA

Diane Markakis,

Department of Epidemiology, Johns Hopkins Bloomberg School of Public Health, Baltimore, MD, USA

Darcy F. Phelan-Emrick,

Department of Epidemiology, Johns Hopkins Bloomberg School of Public Health, Baltimore, MD, USA

Jennifer Wenzel,

Department of Acute and Chronic Care, Johns Hopkins School of Nursing, Baltimore, MD, USA

Gary R. Shapiro,

Health Partners Cancer Program and Institute for Education and Research, Minneapolis, MN, USA, Institute for Advanced Studies in Aging (IASIA), Falls Church, VA, USA

Lee Bone,

Department of Health Behavior and Society, Johns Hopkins Bloomberg School of Public Health, Baltimore, MD, USA

Correspondence to: Mohammad Khalid Hararah; Jean G. Ford.

Conflict of interest

Mohammad Khalid Hararah, Craig Evan Pollack, Mary A. Garza, Hsin-Chieh Yeh, Diane Markakis, Darcy F. Phelan-Emrick, Jennifer Wenzel, Gary R. Shapiro, Lee Bone, Lawrence Johnson, and Jean G. Ford all declare that they have no conflict of interest.

Informed consent

All procedures followed were in accordance with the ethical standards of the responsible committee on human experimentation (institutional and national) and with the Helsinki Declaration of 1975, as revised in 2000 (5). Informed consent was obtained from all patients for being included in the study.

Lawrence Johnson, and

Park West Health Systems, Baltimore, MD, USA

Jean G. Ford

Department of Medicine, The Brooklyn Hospital Center, Brooklyn, NY, USA

Abstract

Purpose—We examined the association between socioeconomic status (SES) and prostate-specific antigen (PSA) cancer screening among older African American men.

Methods—We analyzed baseline data from a sample of 485 community-dwelling African American men who participated in the Cancer Prevention and Treatment Demonstration Trial. The outcome was receipt of PSA screening within the past year. SES was measured using income and educational attainment. Sequential multivariate logistic regression models were performed to study whether health care access, patient–provider relationship, and cancer fatalism mediated the relationship between SES and PSA screening.

Results—Higher educational attainment was significantly associated with higher odds of PSA screening in the past year (odds ratio (OR) 2.08 for college graduate compared to less than high school graduate, 95 % confidence interval (CI) 1.03–4.24); income was not. Health care access and patient–provider communication did not alter the relationship between education and screening; however, beliefs regarding cancer fatalism partially mediated the observed relationship.

Conclusion—Rates of prostate cancer screening among African American men vary by level of educational attainment; beliefs concerning cancer fatalism help explain this gradient. Understanding the determinants of cancer fatalism is a critical next step in building interventions that seek to ensure equitable access to prostate cancer screening.

Keywords

Prostate-specific antigen; Prostate cancer; Cancer screening; African American; Education; Socioeconomic status

Purpose

Socioeconomic status (SES), especially one's level of education and income, is an important determinant of the health care that one receives. Men with lower socioeconomic status, typically as indicated by lower levels of education and income, are less likely to receive prostate-specific antigen (PSA) screening and active treatment for their prostate cancer than those with higher socioeconomic status [1–7]. Less well studied, however, is examining how socioeconomic status affects health *within* racial/ethnic groups [8].

Examining the contribution of socioeconomic status to rates of PSA screening among African American men is especially important given the high burden of prostate cancer morbidity and mortality in this group [9]. While guidelines no longer recommending routine PSA screening for average-risk white men [10], few African American men were included in the large screening trials [11, 12]. With continuing uncertainty regarding the optimal

approach for PSA screening among African American men, informed decision-making is frequently recommended [13, 14].

In this context, we sought to first examine the association between socioeconomic status and PSA screening among older African American men living in Baltimore and, second, explore potential mechanisms that may mediate any observed association. We specifically examined whether lower access to health care, difficulties with doctor–patient communication, and higher beliefs around cancer fatalism may help explain socioeconomic differences in PSA screening [2, 4–6, 15, 16]. Each of these factors have been previously associated with rates of PSA screening among men and are more likely to be found among men with lower socioeconomic status [2, 4, 3].

Methods

This analysis is based on data from the baseline interview of 485 Baltimore City, community-dwelling African American men, age 65–75 years participating in the Cancer Prevention and Treatment Demonstration Project (CPTD). CPTD was funded by the Centers for Medicare & Medicaid Services (CMS) between April 2006 and December 2010. The Johns Hopkins University Institutional Review Board approved the study.

Study Participants

Study participants were recruited using two strategies. First, between October 2006 and June 2008, individuals were recruited using the Medicare membership database as well as from clinical (Johns Hopkins affiliated clinics and federally qualified health centers) and community-based settings, such as senior centers, apartment buildings, and community events. During this period, 744 individuals (27 % male, 73 % female) were recruited into the study. Second, in July 2008, enhanced population-based recruitment was initiated wherein participants were recruited through Medicare enrollment rosters provided by CMS. During this second phase of recruitment, which continued through March 2010, 1,849 individuals were enrolled in the study (26.5 % male, 73.5 % female).

African American residents of Baltimore City were eligible to participate in the CPTD study if they were aged 65 years or older, enrolled in Medicare Parts A and B, and had either no known history of cancer or cancer in remission for five years or longer. Study exclusion criteria included inability to provide informed consent and residence in an institutionalized setting, including a chronic care facility. Patients who were not enrolled in fee-for-service Medicare were excluded as the larger trial planned to obtain insurance claims data on patients. A total of 685 men completed the in-person, baseline interview. We excluded 7 men who did answer the main outcome questions on PSA screening and 193 men who were over 75 years and older as PSA screening was not routine recommended for this age group at the time when the study was conducted [17].

Variables

PSA Screening—In a baseline interview, participants were asked “Have you ever had a PSA test?” and for those who responded affirmatively, “How long ago was it since you have your last PSA?” Screening in the past year was our primary outcome.

Socioeconomic Status—Socioeconomic status was measured using educational attainment (categorized as less than high school, high school, some college, and college graduate) and household income measured by increments of \$10,000 (categorized as less than \$10,000, \$10,000 to \$30,000, and greater than \$30,000 based on the sample distribution).

Demographic Characteristics—Marital status (married/living with partner vs single/separated/divorced), family history of prostate cancer in a first-degree relative, and self-reported health status (excellent/very good/good vs fair/poor) were included.

Health Care Access—We examined three measures of access to health care: (1) whether they have a usual place for health care, (2) whether there is a particular doctor, nurse practitioner, or physician's assistant they usually see, and (3) whether respondents have Medicaid and/or Medigap enrollment.

Patient–Provider Relationship—Participants were asked how often the following statements reflected their experiences with the health care system: “Health care professional listens to you carefully,” “Health care professional explains things in way you can understand,” “Health care professional shows you respect,” and “Health care professional spends enough time with you.” Responses were dichotomized into always/usually versus sometimes/never.

Cancer Fatalism—Cancer susceptibility and beliefs were assessed by how much a participant agreed or disagreed with the following statements: “There is nothing you can do to lower your chance of getting cancer” and “It seems everything causes cancer.” [18] Responses were categorized as “strongly agree/agree,” “no opinion”, and “disagree/strongly disagree.”

Statistical Analyses

Exploratory data analysis was used to determine the variability and distribution of the data, followed by bivariate analyses. Bivariate analyses with chi-square tests or two-sided Fisher's exact test were performed with each independent variable and PSA within the past year. We then examined the independent association between income and education and the main outcome (PSA testing within the past year) in logistic regression analyses that adjusted for marital status, family history, and self-reported health status. To test whether health care access, patient–provider relationship, and beliefs regarding cancer fatalism individually mediated the observed relationship between socioeconomic status and PSA testing, we followed the approach of Baron and Kenny in which we (1) tested whether each potential mediator was associated with PSA screening in unadjusted analyses, (2) tested whether each potential mediator was associated with socioeconomic status, and (3) examined whether the association between socioeconomic status and PSA screening was attenuated after adjustment for the potential mediator [19]. Because we did not observe a significant association between income and PSA screening, we focused on the relationship between potential mediators and education only in the tests of the association between mediators and

socioeconomic status. Analyses were conducted using STATA version 11.2 for Windows (StataCorp LP, College Station 148 TX).

Results

Demographic information, patient–provider relationship, beliefs around cancer fatalism, and health access characteristics are presented in Table 1. Among the 485 African American men between 65 and 75 years old, 56 % of the sample reported having a PSA screening within the past year, and 81 % reported having a PSA screening during their lifetime. With regard to education, 36 % of the sample had less than a high school education and 13 % had earned a bachelor’s degree or greater.

In bivariate analyses, participants who had higher levels of education and those with higher incomes were significantly more likely to report PSA screening in the past year. With regard to potential mediators, men who reported screening in the past year were more likely to have a usual place of health care and a particular health care professional they usually see, report that their health care professional explained things understandably and showed respect for what they had to say, and disagree with the statement that there is nothing that you can do to lower your chance of prostate cancer and that everything causes cancer.

In Table 2, the association between education and the potential mediators is reported. We focus on education because, as discussed later, income was not independently associated with PSA screening in adjusted analyses. Higher educational attainment is significantly associated with Medigap supplemental enrollment, while no other health access variables had an association with education. Higher education was not significantly associated with measures of the patient–provider relationship. Higher beliefs in cancer fatalism were found among men with lower levels of education (p value <0.01 for each statement).

Results of the unadjusted and adjusted logistic models are presented in Table 3. In unadjusted models, we find similar results as the bivariate analyses in Table 1, where men who reported screening in the past year were at higher odds of having a bachelor’s degree or higher, having an income of greater than or equal to \$30,000, being married or living with a partner, and having better reported health. Again, we find that men who reported screening in the past year were at higher odds of having a usual place of care and a particular health care professional they usually see, and reporting that their health care professional explained things understandably and showed respect for what they had to say. Notably, we also see that men who agreed with the statements that there is nothing that you can do to lower your chance of prostate cancer and that everything causes cancer were at lower odds of having been screened in the prior year (odds ratio (OR) 0.56, 95 % confidence interval (CI) 0.36–0.86, and OR 0.58, 95 % CI 0.40–0.84), respectively.

Adjusted model I includes both income and education along with demographic factors. Only men with a bachelor’s degree or higher were significantly associated with having a PSA screening within the past year (odds ratio (OR) 2.08, 95 % confidence interval (CI) 1.03–4.24) compared to less than high school diploma. Income level was not significantly

associated with screening. Marital status remains significantly associated with PSA screening (OR 1.91, 95 % CI 1.27–2.88).

In model II, we added measures of health care access to model I. Individuals who reported having a usual place of care had approximately 28 times higher odds of having a PSA test within the past year though with wide confidence intervals indicating instability of the estimate (OR 27.89, 95 % CI 3.59–216.64). Reporting supplemental insurance was not significantly correlated with having PSA testing. The addition of health access variables did not appear to alter the relationship between education and PSA screening. Model III adds patient–provider communication variables to model I. None of the communication variables were associated with PSA screening status within the past year, and the inclusion of these variables also did not alter the point estimate between education and PSA screening status. In model IV, we added variables measuring fatalism about cancer to model I. The inclusion of these items attenuated the point estimate between educational attainment and PSA testing in the negative direction, making it no longer statistically significant. Individuals who reported having a bachelor’s degree or higher had approximately 1.83 times higher odds of being screened in the previous year than those without a high school diploma (OR 1.83, 95 % CI 0.89–3.77).

Discussion

Among older African American men living in Baltimore, we found that education was positively significantly associated with reported prostate cancer screening by PSA testing. Beliefs concerning cancer fatalism (i.e., that “there is nothing you can do to lower cancer risk” and “everything causes cancer”) were more common among men with lower socioeconomic status and, after the inclusion of these beliefs in a multivariable model, education was no longer significantly associated with PSA screening. The results present important evidence on socioeconomic differences in cancer screening within racial/ethnic group as well as potential factors that may be targeted in addressing these differences.

Although PSA screening remains controversial, most guidelines recommend informed decision-making for African American men [10, 13]. Patients with lower socioeconomic status are often less likely to engage in shared decision-making [20]. In previous studies, patients with fatalistic beliefs about cancer are at decreased likelihood of engaging in preventative behaviors [21]. In our current study, we found that those with higher beliefs in cancer fatalism were less likely to undergo screening, although the results did not remain significant in our fully adjusted model. It is possible that addressing cancer fatalism may be an important way to reduce socioeconomic disparities within racial/ethnic groups.

Studies using short education programs or brief media interventions targeting cancer fatalism have been successful in reducing fatalistic beliefs about cancer survivorship; this suggests that cancer fatalism is modifiable [22, 23]. Future research should be aimed at sources of cancer fatalism and assess the impact of interventions to reduce cancer fatalism. If fatalistic beliefs about cancer prevention are largely attributable to information overload or are culturally related, health educators might target these areas. Health care workers and

educators could develop simpler cancer prevention messages that can be widely disseminated and understood by less educated individuals.

Annual income was not the major determinant of screening status after adjusting for education and other known confounders. The finding that varying income levels are insufficient to explain socioeconomic differentials of cancer screening is consistent with previous research on PSA screening and other health service use, particularly in relation to countries with universal health care (in comparison to the Medicare) [24, 8]. Moreover, focusing on urban African American Medicare beneficiaries in Baltimore City may lead to less heterogeneity in income distribution and have limited our ability to see significant associations, especially for those at the higher end of the income distribution [25]. There also may be other markers of socioeconomic status such as wealth, which may be particularly meaningful among older adults when yearly income may not be stable [26, 27].

Although access to care did not mediate the role of education, it was a strong predictor for PSA screening in its own right. Men who have a usual place of care were 28 times more likely to have had a PSA within the past year, albeit with wide confidence intervals. Increasing access to care may be an important step in increasing rates of screening among urban, African American Medicare beneficiaries, regardless of level of education.

Limitations

Our findings contain several limitations that should be considered in the interpretation of the results. First, we focus on African American Medicare beneficiaries in Baltimore City. We do not have comparison data for the rates of PSA screening among African Americans in Baltimore based solely on a population-based sample. Moreover, annual screening rates were slightly higher in this study than other previous national estimates that described frequency in screening [28, 5, 3]. Thus, our results may not be broadly generalizable. Second, questions on cancer fatalism were not specific to prostate cancer. Similarly, patient-provider communication was measured using general self-reported communication items, which may not capture the full extent of the patient-provider relationship and were not specific to prostate cancer screening. Third, the outcome measure of having undergone prostate cancer screening within the past year was self-reported and subject to recall bias [29, 30]. Fourth, consistent with previous studies, our measure of income included a high degree of missing values. Incomes were reported to be low, which is consistent with Baltimore City's relative deprivation compared to the remainder of the state [25]. The lower range of incomes likely limited our ability to detect significant associations between income and PSA, especially among men with higher incomes. In addition, with a small sample size, our ability to detect significant associations is diminished. Finally, this study occurs in the context of uncertainty over the optimal use of PSA screening among African American men [13, 10]. Related to this, while it is possible that educational differences in the observed rates of PSA screening reflect differences in informed choices, such explanations based on patient preferences are often misleading [31]. Patients with lower socioeconomic status have been found to have lower levels of informed decision-making [15].

Conclusion

The results demonstrate that urban, African American men with lower levels of education status tend to have lower rates of PSA screening. These differences may, at least in part, be explained by cancer fatalism. Understanding the determinants of cancer fatalism and why it is more prominent among low-income men is a critical next step in building interventions that seek to ensure equitable access to prostate cancer screening.

Acknowledgments

We acknowledge the Community Advisory Committee for their contributions to this demonstration project and allowing me to work with them.

Funding/support

This study was supported by Centers for Medicare and Medicaid Services (Grant #1A0CMS300066 and U54CA153710). Additionally, Dr. Mary A. Garza was supported, in part, through her Mentored Research Scientist Development Award to Promote Diversity (K01CA140358). Dr. Pollack's salary is supported by the National Cancer Institute (NCI) and Office of Behavioral and Social Sciences (K07 CA151910). Dr. Wenzel was supported, in part, by her American Cancer Society Mentored Research Scholar Grant (117902-MRSGT-09-152-01-CPPB).

References

1. Hammond WP, Matthews D, Mohottige D, Agyemang A, Corbie-Smith G. Masculinity, medical mistrust, and preventive health services delays among community-dwelling African-American men. *J Gen Intern Med*. 2010; 25(12):1300–8. DOI: 10.1007/s11606-010-1481-z [PubMed: 20714819]
2. Woods VD, Montgomery SB, Herring RP, Gardner RW, Stokols D. Social ecological predictors of prostate-specific antigen blood test and digital rectal examination in black American men. *J Natl Med Assoc*. 2006; 98(4):492–504. [PubMed: 16623061]
3. Haque R, Van Den Eeden SK, Jacobsen SJ, Caan B, Avila CC, Slezak J, et al. Correlates of prostate-specific antigen testing in a large multiethnic cohort. *Am J Manage Care*. 2009; 15(11):793–9.
4. Winterich JA, Grzywacz JG, Quandt SA, Clark PE, Miller DP, Acuna J, et al. Men's knowledge and beliefs about prostate cancer: education, race, and screening status. *Ethn Dis*. 2009; 19(2):199–203. [PubMed: 19537233]
5. Ross LE, Taylor YJ, Richardson LC, Howard DL. Patterns in prostate-specific antigen test use and digital rectal examinations in the Behavioral Risk Factor Surveillance System, 2002–2006. *J Natl Med Assoc*. 2009; 101(4):316–24. [PubMed: 19397221]
6. Gilligan T, Wang PS, Levin R, Kantoff PW, Avorn J. Racial differences in screening for prostate cancer in the elderly. *Arch Intern Med*. 2004; 164(17):1858–64. DOI: 10.1001/archinte.164.17.1858 [PubMed: 15451760]
7. Ford ME, Vernon SW, Havstad SL, Thomas SA, Davis SD. Factors influencing behavioral intention regarding prostate cancer screening among older African-American men. *J Natl Med Assoc*. 2006; 98(4):505–14. [PubMed: 16623062]
8. Braveman PA, Cubbin C, Egerter S, Williams DR, Pamuk E. Socioeconomic disparities in health in the United States: what the patterns tell us. *Am J Public Health*. 2010; 100(Suppl 1):S186–96. DOI: 10.2105/AJPH.2009.166082 [PubMed: 20147693]
9. Cancer Facts & Figures 2011. 2011
10. Moyer VA, on behalf of the USPSTF. Screening for prostate cancer: U.S. Preventive Services Task Force recommendation statement. *Ann Intern Med*. 2012; 157(2):120–34. DOI: 10.7326/0003-4819-157-2-201207170-00459 [PubMed: 22801674]
11. Andriole GL, Crawford ED, Grubb RL 3rd, Buys SS, Chia D, Church TR, et al. Mortality results from a randomized prostate-cancer screening trial. *N Engl J Med*. 2009; 360(13):1310–9. DOI: 10.1056/NEJMoa0810696 [PubMed: 19297565]

12. Schroder FH, Hugosson J, Roobol MJ, Tammela TL, Ciatto S, Nelen V, et al. Screening and prostate-cancer mortality in a randomized European study. *N Engl J Med*. 2009; 360(13):1320–8. DOI: 10.1056/NEJMoa0810084 [PubMed: 19297566]
13. Carter HB, Albertsen PC, Barry MJ, Etzioni R, Freedland SJ, Greene KL, et al. Early detection of prostate cancer: AUA Guideline. *J Urol*. 2013; 190(2):419–26. DOI: 10.1016/j.juro.2013.04.119 [PubMed: 23659877]
14. Wolf AM, Wender RC, Etzioni RB, Thompson IM, D'Amico AV, Volk RJ, Brooks DD, Dash C, Guessous I, Andrews K, DeSantis C, Smith RA. American Cancer Society guideline for the early detection of prostate cancer: update 2010. *CA Cancer J Clin*. 60(2):70–98. DOI: 10.3322/caac.20066 [PubMed: 20200110]
15. Bowen DJ, Hannon PA, Harris JR, Martin DP. Prostate cancer screening and informed decision-making: provider and patient perspectives. *Prostate Cancer Prostatic Dis*. 2011; 14(2):155–61. DOI: 10.1038/pcan.2010.55 [PubMed: 21243007]
16. Gilligan T. Social disparities and prostate cancer: mapping the gaps in our knowledge. *Cancer Causes Control*. 2005; 16(1):45–53. DOI: 10.1007/s10552-004-1291-x [PubMed: 15750857]
17. Screening for prostate cancer: U.S. Preventive Services Task Force recommendation statement. *Ann Intern Med*. 2008; 149(3):185–191. [PubMed: 18678845]
18. Befort CA, Nazir N, Engelman K, Choi W. Fatalistic cancer beliefs and information sources among rural and urban adults in the USA. *J Cancer Educ: Off J Am Assoc Cancer Educ*. 2013; doi: 10.1007/s13187-013-0496-7
19. Baron RM, Kenny DA. The moderator-mediator variable distinction in social psychological research: conceptual, strategic, and statistical considerations. *J Pers Soc Psychol*. 1986; 51(6): 1173–82. [PubMed: 3806354]
20. Williams RM, Zincke NL, Turner RO, Davis JL, Davis KM, Schwartz MD, et al. Prostate cancer screening and shared decision-making preferences among African-American members of the Prince Hall Masons. *Psychooncology*. 2008; 17(10):1006–13. DOI: 10.1002/pon.1318 [PubMed: 18203246]
21. Niederdeppe J, Levy AG. Fatalistic beliefs about cancer prevention and three prevention behaviors. *Cancer Epidemiol Biomarkers Prev*. 2007; 16(5):998–1003. DOI: 10.1158/1055-9965.EPI-06-0608 [PubMed: 17507628]
22. Powe BD, Weinrich S. An intervention to decrease cancer fatalism among rural elders. *Oncol Nurs Forum*. 1999; 26(3):583–8. [PubMed: 10214599]
23. Morgan PD, Fogel J, Tyler ID, Jones JR. Culturally targeted educational intervention to increase colorectal health awareness among African Americans. *J Health Care Poor Underserved*. 2010; 21(3 Suppl):132–47. DOI: 10.1353/hpu.0.0357
24. Katz SJ, Hofer TP. Socioeconomic disparities in preventive care persist despite universal coverage. Breast and cervical cancer screening in Ontario and the United States. *JAMA*. 1994; 272(7):530–4. [PubMed: 8046807]
25. United States Census Bureau: State & County QuickFacts. 2014. <http://quickfacts.census.gov/qfd/states/24/2404000.html>. Accessed July 12, 2014
26. Pollack CE, Chideya S, Cubbin C, Williams B, Dekker M, Braveman P. Should health studies measure wealth? A systematic review. *Am J Prev Med*. 2007; 33(3):250–64. DOI: 10.1016/j.amepre.2007.04.033 [PubMed: 17826585]
27. Braveman PA, Cubbin C, Egerter S, Chideya S, Marchi KS, Metzler M, et al. Socioeconomic status in health research: one size does not fit all. *JAMA*. 2005; 294(22):2879–88. DOI: 10.1001/jama.294.22.2879 [PubMed: 16352796]
28. Han PK, Kobrin S, Breen N, Joseph DA, Li J, Frosch DL, et al. National evidence on the use of shared decision making in prostate-specific antigen screening. *Ann Fam Med*. 2013; 11(4):306–14. DOI: 10.1370/afm.1539 [PubMed: 23835816]
29. Jordan TR, Price JH, King KA, Masyk T, Bedell AW. The validity of male patients' self-reports regarding prostate cancer screening. *Prev Med*. 1999; 28(3):297–303. DOI: 10.1006/pmed.1998.0430 [PubMed: 10072749]

30. Chan EC, Vernon SW, Ahn C, Greisinger A. Do men know that they have had a prostate-specific antigen test? Accuracy of self-reports of testing at 2 sites. *Am J Public Health*. 2004; 94(8):1336–8. [PubMed: 15284039]
31. Armstrong K, Hughes-Halbert C, Asch DA. Patient preferences can be misleading as explanations for racial disparities in health care. *Arch Intern Med*. 2006; 166(9):950–4. DOI: 10.1001/archinte.166.9.950 [PubMed: 16682567]

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 1Baseline characteristics by PSA screened status within the past year ($n=485$)

	Not screened		Screened		Total		p value	
	N	%	N	%	N	%		
Socioeconomic Status								
Income								
<\$10,000	42	(19.7 %)	29	(10.7 %)	71	(14.6 %)	<0.01	
\$10,000–\$30,000	77	(36.2 %)	89	(32.7 %)	166	(34.2 %)		
\$30,000	55	(25.8 %)	121	(44.5 %)	176	(36.3 %)		
Unknown	39	(18.3 %)	33	(12.1 %)	72	(14.9 %)	<0.01	
Education								
<High school	87	(41.6 %)	83	(30.9 %)	170	(35.6 %)		
High school	57	(27.3 %)	73	(27.1 %)	130	(27.2 %)		
Some college	49	(23.4 %)	65	(24.2 %)	114	(23.9 %)		
Bachelors degree	16	(7.7 %)	48	(17.8 %)	64	(13.4 %)	<0.01	
Demographic characteristics								
Marital status								
Married/lives with partner	87	(41.6 %)	166	(62.4 %)	253	(53.3 %)		
Widowed/divorced/lives alone	122	(58.4 %)	100	(37.6 %)	222	(46.7 %)		
Family Hx of PCa								
No	185	(86.9 %)	230	(85.5 %)	415	(86.1 %)		
Yes	15	(7.0 %)	28	(10.4 %)	43	(8.9 %)		
Self-reported health								
Excellent/very good/good	156	(73.2 %)	226	(83.4 %)	382	(78.9 %)		
Fair/poor	57	(26.8 %)	45	(16.6 %)	102	(21.1 %)		
Health care access								
Usual place for health care?								
No	21	(10.0 %)	1	(0.4 %)	22	(4.6 %)		
Yes	189	(90.0 %)	268	(99.6 %)	457	(95.4 %)		
Particular doctor/NP/PA usually see?								
No	47	(22.1 %)	22	(8.1 %)	69	(14.2 %)		

	Not screened		Screened		Total		p value
	N	%	N	%	N	%	
Yes	166	(77.9 %)	250	(91.9 %)	416	(85.8 %)	
Enrolled in Medicaid/Supplement?							0.17
No	111	(53.9 %)	128	(47.6 %)	239	(50.3 %)	
Yes	95	(46.1 %)	141	(52.4 %)	236	(49.7 %)	
Enrolled in Medicaid, i.e., dual?							0.54
No	178	(86.4 %)	234	(88.3 %)	412	(84.5 %)	
Yes	28	(13.6 %)	31	(11.7 %)	59	(12.5 %)	
Provider–patient relationship							
Doctor listens to you carefully?							0.02
Sometimes/never	22	(10.6 %)	13	(4.8 %)	35	(7.3 %)	
Usually/always	186	(89.4 %)	257	(95.2 %)	443	(92.7 %)	
Doctor explains things in way you understand?							0.01
Sometimes/never	25	(12.0 %)	15	(5.6 %)	40	(8.4 %)	
Usually/always	183	(88.0 %)	255	(94.4 %)	438	(91.6 %)	
Doctor show respect for what you had to say?							<0.01
Sometimes/never	20	(9.7 %)	7	(2.6 %)	27	(5.7 %)	
Usually/always	187	(90.3 %)	263	(97.4 %)	450	(94.3 %)	
Doctor spends enough time with you?							<0.01
Sometimes/never	28	(13.5 %)	16	(6.0 %)	44	(9.2 %)	
Usually/always	180	(86.5 %)	253	(94.1 %)	433	(90.8 %)	
Cancer fatalism							
There is nothing you can do lower chance of getting cancer							0.02
Disagree	140	(65.7 %)	208	(76.5 %)	348	(71.8 %)	
Agree	60	(28.2 %)	50	(18.4 %)	110	(22.7 %)	
It seems everything causes cancer							0.01
Disagree	93	(43.7 %)	156	(57.4 %)	249	(51.3 %)	
Agree	111	(52.1 %)	108	(39.7 %)	219	(45.2 %)	

Table 2

Baseline characteristics by educational level ($n=478$)

	Education Level						<i>p</i> value		
	<High school		High school		Some college			Bachelor's	
	<i>N</i>	%	<i>N</i>	%	<i>N</i>	%		<i>N</i>	%
Total (<i>n</i> =478)	250	(100 %)	181	(100 %)	141	(100 %)	95	(100 %)	
PSA screened <i>within past year</i>	83	(30.9 %)	73	(27.1 %)	65	(24.2 %)	48	(17.8 %)	<0.01
Health care access									
Usual place for health care	162	(35.5 %)	124	(27.2 %)	107	(23.5 %)	63	(13.8 %)	0.58
Usual Doctor/NP/PA	146	(35.2 %)	114	(27.5 %)	100	(24.1 %)	55	(13.3 %)	0.95
Enrolled in Medigap/Supplement	66	(28.1 %)	64	(27.2 %)	60	(25.5 %)	45	(19.2 %)	<0.01
Provider–patient relationship									
Doctor listens to you carefully <i>usually/always</i>	157	(35.8 %)	117	(26.7 %)	103	(23.5 %)	62	(14.1 %)	0.48
Doctor explains things in way you understand <i>usually/always</i>	149	(34.3 %)	120	(27.7 %)	105	(24.2 %)	60	(13.8 %)	0.27
Doctor shows respect for what you had to say <i>usually/always</i>	158	(35.4 %)	120	(26.9 %)	105	(23.5 %)	63	(14.1 %)	0.45
Doctor spends enough time with you <i>usually/always</i>	148	(34.5 %)	118	(27.5 %)	103	(24.0 %)	60	(14.0 %)	0.05
Cancer fatalism									
Agrees there is nothing you can do lower chance of getting cancer	53	(50.0 %)	23	(21.7 %)	21	(19.8 %)	9	(8.5 %)	<0.01
Agrees, it seems everything causes cancer	94	(43.3 %)	64	(29.5 %)	40	(18.4 %)	19	(8.8 %)	<0.01

 $N=478$, because 7 persons did not answer the educational attainment question

Table 3

Factors associated with PSA screening within the past year

	Unadjusted OR (95 % CI)	Model I	Model II	Model III	Model IV
Socioeconomic status					
Income					
<\$10,000	1.00	1.00	1.00	1.00	1.00
\$10,000–30,000	1.67 (0.95, 2.93)	1.30 (0.71, 2.35)	1.45 (0.78, 2.70)	1.25 (0.68, 2.30)	1.33 (0.73, 2.42)
\$30,000	3.18 (1.80, 5.64) ^a	1.68 (0.86, 3.28)	1.88 (0.93, 3.79)	1.61 (0.81, 3.17)	1.66 (0.84, 3.26)
Unknown	1.22 (0.63, 2.37)	0.81 (0.39, 1.70)	1.04 (0.48, 2.24)	0.89 (0.42, 1.88)	0.85 (0.40, 1.79)
Education					
<High school	1.00	1.00	1.00	1.00	1.00
High school	1.34 (0.85, 2.12)	1.09 (0.67, 1.77)	1.14 (0.69, 1.88)	1.10 (0.67, 1.82)	1.01 (0.61, 1.65)
Some college	1.39 (0.86, 2.24)	1.23 (0.73, 2.05)	1.35 (0.79, 2.31)	1.29 (0.76, 2.19)	1.09 (0.65, 1.86)
Bachelors degree	3.14 (1.66, 5.96) ^a	2.08 (1.03, 4.24) ^b	2.11 (1.03, 4.37) ^b	2.07 (1.01, 4.21) ^b	1.83 (0.89, 3.77) ^c
Demographics					
Married/lives with partner	2.32 (1.61, 3.37) ^a	1.91 (1.27, 2.88) ^a	1.76 (1.14, 2.70) ^a	1.71 (1.12, 2.60) ^b	1.94 (1.28, 2.94) ^a
Family history of PCa	1.50 (0.78, 2.89)	1.37 (0.68, 2.79)	1.54 (0.73, 3.26)	1.48 (0.71, 3.06)	1.43 (0.69, 2.92)
Fair/poor self-reported Health	0.54 (0.35, 0.85) ^a	0.69 (0.43, 1.11)	0.65 (0.40, 1.06) ^c	0.69 (0.43, 1.11)	0.73 (0.45, 1.18)
Health care access					
Usual source of care	29.78 (3.97, 223.29) ^a		27.89 (3.59, 216.64) ^a		
Usual Doctor/NP/PA	3.22 (1.87, 5.54) ^a	–	–		
Medigap/Supplement insurance	1.29 (0.89, 1.85)		0.73 (0.50, 1.11)		
Enrolled in Medicaid	0.84 (0.49, 1.46)	–			
Provider–patient relationship					
Doctor listens to you carefully <i>usually/always</i>	2.34 (1.15, 4.76) ^b		–	1.09 (0.36, 3.34)	
Doctor explains things in way you understand <i>usually/always</i>	2.32 (1.19, 4.53) ^b		–	1.04 (0.35, 3.06)	
Doctor show respect for what you have to say <i>usually/always</i>	4.01 (1.67, 9.70) ^a		–	1.65 (0.32, 8.45)	
Doctor spends enough time with you <i>usually/always</i>	2.46 (1.30, 4.70) ^a		–	1.58 (0.61, 4.13)	
Cancer fatalism					
Agrees there is nothing you can do lower chance of getting cancer	0.56 (0.36, 0.86) ^a		–	–	0.64 (0.40, 1.04) ^c

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

	Unadjusted OR (95 % CI) ^a	Model I	Model II	Model III	Model IV
Agrees that everything causes cancer	0.58 (0.40, 0.84) ^a		–	–	0.68 (0.45, 1.02) ^c

OR odds ratio, CI confidence interval

- ^a *p* value 0.01
- ^b *p* value 0.05
- ^c *p* value 0.10