Published in final edited form as:

Glob Public Health. 2018 August ; 13(8): 1126–1143. doi:10.1080/17441692.2016.1251603.

Explaining Ethnic Disparities in Preterm Birth in Argentina and Ecuador

George L. Wehby, MPH, PhD¹, Mariela Pawluk, PhD², Kwame A. Nyarko, PhD³, and Jorge S. López-Camelo, PhD^{2,4}

¹University of Iowa, Departments of Health Management and Policy, Economics, and Preventive & Community Dentistry, and Public Policy Center, Iowa City, IA, & National Bureau of Economic Research, Cambridge, MA, United States of America

²Center of Medical Education and Clinical Investigation/Centro de Educación Médica e Investigación Clínica (CEMIC), Buenos Aires, Argentina

³University of Iowa, Department of Health Management and Policy, Iowa City, IA, and University of Tennessee, Health Sciences Center, Memphis, TN, United States of America

⁴Latin American Collaborative Study of Congenital Anomalies/Estudio Colaborativo Latinoamericano de Malformaciones Congénitas (ECLAMC), Buenos Aires, Argentina

Abstract

Background—Little is understood about racial/ethnic disparities in infant health in South America. We quantified the extent to which the disparity in preterm birth rate (PTB; < 37 gestational weeks) between infants of Native only ancestry and those of European only ancestry in Argentina and Ecuador are explained by household socioeconomic, demographic, healthcare use, and geographic location indicators.

Methods—The samples included 5199 infants born between 2000 and 2011 from Argentina and 1579 infants born between 2001 and 2011 from Ecuador. An Oaxaca-Blinder type decomposition model adapted to binary outcomes was estimated to explain the disparity in PTB risk across groups of variables and specific variables.

Results—Maternal use of prenatal care services significantly explained the PTB disparity, by nearly 57% and 30% in Argentina and Ecuador, respectively. Household socioeconomic status explained an additional 26% of the PTB disparity in Argentina.

Conclusions—Differences in maternal use of prenatal care may partly explain ethnic disparities in PTB in Argentina and Ecuador. Improving access to prenatal care may reduce ethnic disparities in PTB risk in these countries.

Corresponding author: George L. Wehby, Ph.D., Associate Professor, University of Iowa, Research Associate, National Bureau of Economic Research, Departments of Health Management and Policy, Economics, and Preventive & Community Dentistry, and Public Policy Center, University of Iowa, 145 N. Riverside Dr., 100 College of Public Health Bldg., Room N248, Iowa City, Iowa 52242-2007, Phone: 319-384-3814, Fax: 319-384-4371, george-wehby@uiowa.edu.

Disclosures: The authors have no conflicts of interest with this work.

Keywords

Child health; preterm birth; racial disparities; ethnic disparities; health inequalities

INTRODUCTION

Socioeconomic disparities in infant health outcomes have been reported in various contexts in several South American countries including Argentina, Brazil, Colombia, Ecuador, Chile and Uruguay (Antunes et al. 2006; Hertel-Fernandez et al. 2007; Jewell 2014; Ortiz et al. 2013; Wehby and McCarthy, 2013; Wehby et al. 2015). Of all relevant socioeconomic indicators, race and ethnicity have been the least investigated factors in South American populations, although some evidence points to extensive disparities. Wehby et al. (2012) reported ethnic disparities in early neurodevelopment in a pooled sample from five South American countries. Similarly, Nyarko et al. (2013a) found large disparities in low birth weight (LBW; <2500 grams) and preterm birth (PTB; <37 weeks) rates between infants of European and African ancestry in Brazil. Most recently, Wehby et al. (2015) reported disparities in LBW and PTB by ethnic ancestry in 7 out of 8 South American countries with more prominent disparities for PTB. Other studies have also reported racial/ethnic disparities in household factors relevant for infant health such as child health insurance coverage in Argentina, Brazil, and Ecuador (Wehby et al. 2011) or in prenatal care use and quality in Brazil (Matijasevich et al. 2009; Victora et al. 2010).

Racial and ethnic disparities in child health are of great concern for policymakers since they broadly affect large population groups and because early deficits in health do not only reduce wellbeing during childhood but are also carried forward and expressed in greater health risks and lower socioeconomic achievement during adulthood. Furthermore, these disparities are thought to be highly driven by social and economic mechanisms that can be effectively targeted by policies and interventions once identified. Despite extensive research on mechanisms underlying racial/ethnic disparities in health in developed settings such as the USA (e.g., Lhila and Long, 2012), direct research on this question specifically in South American settings is sparse. One study has directly explored contributors to racial disparities in LBW and PTB in Brazil using an Oaxaca-Blinder type decomposition model, and found number of prenatal visits and geographic location to be the most prominent factors in explaining these disparities (Nyarko et al, 2013a). No other work has comprehensively investigated and quantified the pathways that explain racial/ethnic disparities in infant health in South American settings. As mentioned above, the underlying mechanisms for racial/ethnic disparities may vary between countries. For example, Woodhouse et al. (2014) found that the effects of prenatal care on birth weight vary extensively between several South American countries, suggesting that the extent to which prenatal care explains racial/ethnic disparities is not necessarily generalizable across populations.

In this study, we examined the extent to which various household characteristics including socioeconomic, demographic, and healthcare indicators as well as geographic location explain disparities in PTB (< 37 gestational weeks) by ethnic ancestry in Argentina and Ecuador, which were recently shown to have large disparities in this outcome (Wehby et al.

2015). In our dataset (described below), these two countries had the largest disparities in PTB between infants of Native only ancestry and those of European only ancestry, and had sufficient samples for addressing this question. PTB is a major adverse birth outcome that is associated with increased infant mortality and risk of neurodevelopmental complications later in childhood (Blencowe et al. 2012; Kent et al. 2012; Morse et al. 2009). Prior work has focused on explaining PTB disparities in Brazil using the same dataset (Nyarko et al, 2013a). No other empirical published work directly investigates and explains ethnic disparities in PTB in Argentina and Ecuador.

METHODS

Data and Sample

Data were obtained from the Latin American Collaborative Study of Congenital Malformations (ECLAMC), a consortium involving a network of hospitals in South America that is focused on epidemiological surveillance for birth defects (Castilla & Orioli 2004). ECLAMC is based on a voluntary collaboration with a group of health professionals (mostly pediatricians) who monitor the occurrence of birth defects among newborns in their hospitals. Affiliated professionals enroll into ECLAMC infants born alive with birth defects as well as a group of infants born in the same hospital without congenital anomalies and matched one-to-one to the infants with birth defects by sex and birth date. ECLAMC professionals collect data on enrolled newborns about infant health and demographics, household demographic and socioeconomic indicators, and maternal health conditions and use of prenatal care services through interviews with the mothers before infants are discharged from the hospital after birth and through abstraction of medical records. All infants are recruited following the same enrollment procedures and data are systematically collected using the same questionnaires across all affiliated hospitals. ECLAMC professionals received training in data collection and study procedures. This dataset has been employed in numerous studies of infant health in South America (Wehby et al. 2010; Nyarko et al. 2013b; Wehby et al. 2014; Woodhouse et al. 2014).

We included infants born in Argentina between 2000 and 2011 and in Ecuador between 2001 and 2011 and enrolled into ECLAMC; 2001 is the first year of ECLAMC data collection in Ecuador. Since birth defects may influence fetal growth and gestational age, our analytical sample only included infants without birth defects. We excluded infants with recorded gestational ages outside of the range 19.5 to 46.5 weeks to reduce data errors, a standard practice in this literature (Nyarko et al. 2013b). We also only included singleton births, which were nearly 99% of the sample. The main analytical sample with complete data on all study variables included 5119 infants born in 32 hospitals in 19 cities in Argentina and 1579 infants born in 8 hospitals in 7 cities in Ecuador.

Ethnic Ancestry

We measured ethnic variation in ancestry following multiple previous studies using ECLAMC data (Nyarko et al. 2013a; Wehby et al. 2014; Wehby et al. 2012; Wehby et al. 2011; Wehby et al. 2010; Woodhouse et al. 2014). This measure of ethnic ancestry was innovated by ECLAMC in order to systematically capture ethnic variation across multiple

South American countries that are highly ethnically admixed. Mothers were interviewed by ECLAMC professionals about all the ancestries of the child through both maternal and paternal lineages across as many family generations as the mother was able to recall (e.g., child's parents/grandparents/ great grandparents/great great-grandparents, etc.). Ancestry was systematically coded into unique or mixed ancestry groups.

We focus on the two most frequent groups among unique (i.e. non-mixed) ancestries in our study samples, which were Native only and European only. Native only ancestry meant that the mother only knew about specific child ancestors who were born in Latin America as far back as she could remember. In other words, the mother did not know about any ancestors of the child who were born outside of Latin America. As measured, Native only ancestry does not necessarily mean indigenous ancestry. Indeed for the majority of children, Native only ancestry does not indicate that the child is only of indigenous ancestry but rather that all the child ancestors as far back as the mother could recall (generally up to great great-grandparents) were born in Latin America. Similarly, European only ancestry indicated that the mother recalled specific ancestors of the child who were born in Europe for both parental lineages, and there were no known ancestors from other continents such as Africa or Asia.

It is important to note that many of the children reported to be of Native ancestry only are likely to have had some European ancestry at one point in time. However, in most of those cases, the European ancestors likely migrated to Latin America much earlier than in cases that report specific European ancestors. Therefore, reporting Native only ancestry indicates in most cases having some level of early admixture between indigenous and European ancestries, in contrast to reporting European only ancestry, which in most cases indicates recent European migration [generally up to great (or great-great) grandparent].

We excluded mixed Native and European ancestry because compared to European only ancestry, there was no significant difference in PTB rates in Argentina and only a marginally significant difference in Ecuador. (Wehby et al, 2015). Non-European and non-Native ancestries such as African or Asian ancestry were not commonly reported for these two countries (~4% in Argentina and ~8% in Ecuador). Besides being the only two non-mixed ancestries with sufficient sample size for our analysis, there is a large disparity in PTB rate between infants of Native only ancestry and European only ancestry (Wehby et al. 2015), with a much higher rate among infants of Native only ancestry. Only a very small group had missing data on ancestry (~2% in Argentina and ~1% in Ecuador). These observations were excluded from the analysis.

Analytical Framework

We modeled PTB risk as a function of conceptually relevant demographic and socioeconomic indicators, maternal health, obstetric history, and healthcare use, and geographic effects. These factors have been shown to be relevant for birth outcomes in South American populations (Wehby et al. 2010; Woodhouse et al. 2014) and may also vary by ancestry and potentially explain disparities in PTB (Nyarko et al. 2013a). We focused first on broader constructs defined below which we evaluated for their potential in explaining the observed disparities as follows:

$$PTB = f(\text{Ancestry, Years, Maternal Healthcare Use, Maternal Health and Obstetric History, Demographics, Household Socioeconomics Status, Geographic Location}),$$

(1)

where *Ancestry* was an indicator for native only ancestry versus European only ancestry and *Years* included fixed effects for year of birth to account for any time trends in PTB.

Maternal Healthcare Use—Maternal use of healthcare services during pregnancy was measured by number of prenatal care visits throughout pregnancy and by indicators for receiving immunizations in each trimester. Previous studies have suggested a role for prenatal care in reducing PTB risk (Wehby et al. 2009b). In order to account for potential non-linear effects of prenatal care visits, we included binary (0/1) variables for numbers of prenatal care visits including separate indicators for no prenatal visits, 1–2 visits, 3–5 visits, 6–8 visits, and 9 or more visits (which was used as the reference category). We did not include waiting time before initiating prenatal care as it has been shown in previous studies to be highly confounded in this population, in that mothers who are more likely to deliver preterm were more likely to initiate prenatal care earlier (Wehby et al. 2009b; Wehby et al. 2009c). Immunizations were included as indicators of the intensity/quality of prenatal care.

Maternal Health and Obstetric History—Maternal health has direct effects on pregnancy complications and illnesses during pregnancy; acute and chronic illness indicators have been shown to be associated with higher PTB risk in ECLAMC birth samples from Argentina and Brazil (Wehby et al. 2009a; Wehby et al. 2009b). Maternal health was measured using an indicator for any acute illnesses during pregnancy such as flu or urinary tract infections and another indicator for any chronic illnesses such as diabetes or hypertension. These were based on maternal self-reports of illnesses during pregnancy. Maternal obstetric history included an indicator for history of conception difficulty and number of previous pregnancies, also self-reported by the mother. Maternal obstetric history captures maternal experience and health knowledge related to pregnancy and fetal health.

Demographics—Household demographic characteristics included indicators for infant sex, maternal age (grouped into six categories to capture non-linear effects: 19 years or younger, 20–24 years, 25–29 years, 30–34 years, 35–39 years, and 40 years or older), and paternal age (grouped into three categories: 24 years or younger, 25–39 years, and 40 years or older). These factors may capture parental preferences and experiences that are relevant for pregnancy progress and fetal health.

Household Socioeconomic Status—Household socioeconomic status was measured by indicators for maternal and paternal education and occupational activity coded into multiple categories. Socioeconomic status may affect PTB in several ways such as by influencing maternal nutrition, psychosocial status, exposure to environmental hazards, health information, and benefit from prenatal care, and has been shown to be associated with

PTB risk in South American countries such as Argentina and Brazil (Wehby et al. 2009a; Wehby et al. 2009b). Maternal occupational activity also captures maternal time costs and occupational exposures.

Geographic Location—We also measured geographic location by including indicators (fixed effects) for the city where the child was born. Geographic differences in health outcomes, socioeconomic conditions, and environmental risks are commonly reported and may play an important role in ethnic disparities due to the residential clustering and segregation by ethnicity. Residential segregation can adversely affect pregnancy and infant health outcomes by reducing access to quality health care, social capital, nutritional resources, and economic opportunities.

PTB Disparity Decomposition

In order to quantify the contributions of household characteristics and geographic effects to the PTB disparities, we employed a model that extends the Oaxaca-Blinder decomposition model to non-linear models with binary outcomes (Fairlie 2005; Oaxaca 1973). The model allows for estimating the extent to which differences in multiple characteristics between two population groups explain their difference in an outcome in a multivariate framework. The explanatory variables can be evaluated in terms of their individual contribution to explaining the outcome disparity or grouped into categories that are each evaluated as one group. This model has been successfully applied in previous studies including work explaining ethnic disparities in child health insurance coverage in several South American countries (Wehby et al. 2011), and in birth outcomes in Brazil (Nyarko et al. 2013b), and racial disparities in birth weight (Lhila and Long, 2012) and child dental health in the US (Guarnizo-Herreno & Wehby 2012).

In the case of linear models, a standard decomposition of outcome differences between two population groups essentially replaces the means of explanatory (right-hand-side) variables, one at a time, for one group by the values of the other group, and re-estimates the change in outcome difference between the two groups after each replacement. The outcome disparity between two groups that is explained by a certain variable is simply the change in their outcome difference (at the outcome means) after replacing one group's mean of that specific explanatory variable by the other. While this approach can be applied for a binary outcome by using a linear probability model, it is suboptimal for many reasons such as predicting out of range (0–1) probabilities and ignoring non-linear effects of covariates on probability. Therefore, we focus on the non-linear extension of this model developed specifically for binary outcomes (Fairlie 2005).

We first estimated equation (1) listed above using logistic regression. We clustered the standard errors at the hospital of birth to account for any non-independence of observations born within the same hospital. From this logistic regression, we predicted PTB probability for each child. Next within each ancestral group, each child was ranked by their predicted PTB probability in order to match observations between the two groups by their rank of PTB probability. Since the groups differed in size, a subgroup from the larger or “majority” ancestral group (based on sample size and not rates of the outcome) equal in size to the

smaller or “minority” group was randomly selected for this matching. The children from the “majority” subgroup, rank-ordered by their probability, were then matched one-to-one with those in the “minority” group based on their probability rank.

After this matching and for each of the explanatory variables in equation (1), one variable at a time, the value of the variable was replaced for each observation in the “minority” group by the value of its matched observation from the “majority” subgroup. The average of the changes in the probabilities of PTB across observations in the “minority” group (predicted from equation 1) after substituting their values on a certain variable with those from the matched “majority” subgroup represents the disparity in PTB risk that is explained by the difference in values of that specific variable between the “minority” and “majority” groups. This was repeated for each explanatory variable holding the other variables fixed as follows:

$$D_k = \frac{1}{N^M} \sum_{i=1}^{N^M} F \left(a_0 + \sum_{j=1}^{k-1} \beta_j X_{ij}^M + \beta_k X_{ik}^O + \sum_{j=k+1}^K \beta_j X_{ij}^O \right) - F \left(a_0 + \sum_{j=1}^{k-1} \beta_j X_{ij}^M + \beta_k X_{ik}^M + \sum_{j=k+1}^K \beta_j X_{ij}^O \right). \quad (2)$$

In equation (2), M represents the minority group and O represents the “majority” subgroup; N^M is the number of observations in the “minority” group; j indicates the order of the variables from 1 to K for substituting their values in the “minority” group by those of “majority” subgroup; X represents the explanatory variables with X^M representing variables taking their values from the “minority” group and X^O representing variables taking their values from the “majority” subgroup; and F is the logit cumulative density function. The disparity in PTB risk (D) explained by the k^{th} variable from a total of K variables (or categories of variables) ordered from 1 to K was estimated by averaging the change in PTB probabilities when replacing the values of k for observations in the “minority” group by those of their matched observations from the randomly selected “majority” subgroup. This equation is sequentially estimated for each explanatory variable in the order from 1 to K (shown above for the k^{th} variable). The values of variables ordered 1 through $k-1$ are from the “minority” group, while the values of variables ordered $k+1$ to K are based on the “majority” subgroup. The total explained disparity is the sum of the proportions of PTB disparity accounted for by each explanatory variable individually. Alternatively, the total explained disparity may be estimated by simultaneously replacing the values of all explanatory variables in the “minority” group by those in the matched “majority” subgroup.

Instead of evaluating individual variables, the model can be estimated for groupings/ categories of variables, one group at a time. In that case, the values of all variables in a category are simultaneously replaced in the “minority” group with those from the “majority” subgroup. We consider this model to be more powerful for understanding the main constructs that explain these disparities since variables within the same category may be related to each other and therefore not necessarily individually statistically significant in explaining the disparity, but still jointly relevant. We first estimated this model for the five categories of explanatory variables described above: *Maternal Healthcare Use*, *Maternal*

Health and Obstetric History, Demographics, Household Socioeconomics Status, and Geographic Location. Next, we estimated the model for the variables themselves irrespective of the categories in order to identify the most prominent ones within each category for explaining the disparity.

In order to ensure that the results are not specific to one randomly selected subgroup from the “majority” sample for matching, we randomly selected 2000 subgroups from the “majority” sample (each equal in size to the “minority” group), estimated equation (2) for each “majority” subgroup, and averaged the results across the 2000 subgroups. These averaged results approximate those from matching the entire “majority” sample to the “minority” sample. Furthermore, with each randomly selected “majority” subgroup, the ordering of the variable (or category) j was randomly assigned to ensure that the results are not dependent on an arbitrary order. The results therefore approximate the average estimates from all possible variable (or category) orderings and “majority” subgroups.

RESULTS

Sample Description

Table 1 shows descriptive statistics of all model variables by child’s ancestry and country. The two countries were similar in their PTB rates by ethnic ancestry. The PTB rate was 10.4% among children of Native ancestry alone and 7.4% among those of European ancestry alone in Argentina, compared to 12.7% and 7.0% in Ecuador.

PTB Disparity Decomposition

The results from the decomposition model for the five evaluated categories of variables (*Maternal Healthcare Use, Maternal Health and Obstetric History, Demographics, Household Socioeconomics Status, and Geographic Location*) are in Table 2 (the associations of the explanatory variables with PTB risk from the logistic regression in equation 1 are in Supplementary Table S1 for Argentina and Supplementary Table S2 for Ecuador). For each country, we show the total difference in PTB rate between child of Native only and European only ancestry, the explained difference based on categories of variables that were significant at least at $p < 0.05$, and the contribution of each category of variables to explaining the difference, whether significant or not.

Maternal healthcare use significantly ($p < 0.05$) explained the PTB disparity by about 57% in Argentina and 30% Ecuador (Table 2). In Argentina, *household socioeconomic status* explained an additional 26% of the PTB disparity. None of the other categories of variables were significant in explaining this disparity. We describe further the results for each country below.

Argentina

Children of European only ancestry had lower PTB rate than those of Native only ancestry by about 3 percentage-points (Table 1). Of the five evaluated variable categories, only maternal *healthcare use* and *household socioeconomic status* significantly explained the PTB rate disparity between these two ethnic groups, by nearly 1.7 percentage-points and 0.8

percentage-points, respectively, of the total 3 percentage-point difference in PTB rate (Table 2). Table 3 shows the contributions of specific variables to the PTB disparity on their own when variables were not grouped in categories. Indicators for no prenatal visits, using 1–2 visits, and using 3–5 visits (all relative to using 9 or more visits as the reference category) were significant. Among these, use of 3–5 visits explained the most - nearly 1 percentage-point of the PTB disparity. This result is driven by both the decline in PTB risk with greater use of prenatal visits (Supplementary Table S1) and the fact that prenatal care use was noticeably lower among mothers of Native only ancestry compared to those of European only ancestry (Table 1). For example, almost 9% of mothers of Native only ancestry had 2 or fewer prenatal visits throughout the pregnancy compared to 3% of those of European only ancestry. Similarly, nearly 21% of Native only mothers had 9 or more visits compared to 36% of mothers of European only ancestry. Immunizations did not significantly explain the PTB disparity; in fact, the results suggest that the disparity would have been larger if the third trimester vaccination rate among Native only mothers were as low as that among European only mothers.

Among the *household socioeconomic status* indicators, two variables were individually significant (Table 3): mothers completing secondary school and fathers who have a high occupational status (executive, owner, or boss). Each explained close to 0.5 percentage-points of the PTB disparity. There was a stark difference in socioeconomic status by ancestry, with a greater proportion of lower socioeconomic status households among Native only mothers (Table 1). For example, almost 50% of Native only mothers did not attend high school, whereas nearly 77% of European only mothers attended high school or had higher education. Similarly, less than 4% of Native only mothers attended or graduated from University, compared to 25% of European only mothers. Such differences were also observed in father's education and in parental occupational status. For example, only 2% of Native only fathers had a high occupational status, compared to 17% of European only fathers. None of the other variables significantly explained the PTB disparity when included on their own.

Ecuador

Children of European only ancestry had lower PTB rate than those of Native only ancestry by nearly 6 percentage-points (Table 1). Similar to Argentina, differences in *maternal healthcare use* emerged as an important factor, explaining up to 1.7 percentage-points of the total PTB disparity (Table 2). None of the other categories significantly explained this disparity. Table 3 shows the contributions of specific variables to the PTB disparity irrespective of variable categorization. Not receiving prenatal care or only receiving 1–2 prenatal visits (versus 9 visits) significantly explained the disparity, collectively by nearly 1 percentage-point. Similar to Argentina, prenatal care was associated with lower PTB risk in Ecuador (Supplementary Table S3). Also, prenatal care use was lower among Native only mothers than those of European only ancestry (Table 1). For instance, nearly 8% of European only mothers received 2 or fewer visits (including no prenatal care), compared to 2% of European only mothers. Receiving 9 or more visits was also more frequent among European only mothers (15% versus 9%). None of the other healthcare service use measures significantly explained the disparity.

Three other demographic and socioeconomic variables significantly explained the PTB disparity in Ecuador. The higher proportion of very young Native only mothers (18% versus 5%) explained 1 percentage-point of the disparity. Similarly, the lower proportion of Native only mothers with high occupational status (2% versus 10%) explained nearly 1 percentage-point. Finally, the lower proportion of Native only fathers with complete secondary school (28% versus 36%) explained 0.3 percentage-points.

DISCUSSION

We were able to explain an important fraction of the PTB disparity between infants and Native only ancestry and those of European only ancestry in samples from Argentina and Ecuador. Maternal use of prenatal care services emerges as the most important category of variables from all the evaluated household characteristics, explaining up to 57% and 30% of the PTB disparity in Argentina and Ecuador, respectively. In both countries, an increase in prenatal visits was associated with lower PTB risk, consistent with previous studies (e.g., Woodhouse et al. 2014). There were stark socioeconomic differences by ancestry, with a greater proportion of Native only children born into lower socioeconomic status households than European only children in both countries. In Argentina, parental education and occupational status significantly explained as a group nearly 26% of the PTB disparity. In Ecuador, these variables were not significant as a group, but differences in proportions of high maternal occupations and fathers with complete secondary school education significantly explained the disparity. Native only mothers in Ecuador were much more likely to be very young than European only mothers, which also significantly explained the disparity. The other evaluated variables did not appear to play a key role in explaining the PTB disparity. Interestingly, city of birth did not emerge as a key factor in explaining these disparities, unlike for Brazil where geographic differences in infant health and race/ethnicity distribution were much more prominent (Nyarko et al, 2013a). These findings indicate that mechanisms underlying racial/ethnic disparities in health may differ between South American countries and that study findings are not necessarily generalizable across populations.

Given that prenatal care is a relatively low cost intervention, policies that increase access to prenatal care such as by mandating coverage of prenatal care services in private insurance plans or providing these services in public programs may help in reducing this ethnic disparity in PTB. Increasing prenatal care utilization may also be associated with an improvement in additional perinatal outcomes such fetal growth and birth weight (Coria-Soto et al. 1996; Ickovics et al. 2003; Kogan & Alexander 1998; Krueger & Scholl 2000). The World Health Organization recommends a minimum of 5 prenatal visits during pregnancy (Villar et al. 2001). However, a sizeable fraction of Native only women in the study samples obtained fewer than 5 visits including 27% in Argentina and 23% in Ecuador.

These gaps in prenatal care use from the minimum recommended levels illustrate important opportunities for developing maternal health policies to improve access to prenatal care and potentially reduce ethnic disparities in PTB. Inadequate prenatal care use has been shown to be strongly linked to low socioeconomic status in South American settings including in Argentina (Wehby et al. 2009b; Wehby et al. 2009c). As noted above, Native only mothers

are more likely to be of lower socioeconomic status than European only mothers. Therefore, comprehensive policies that improve the socioeconomic environment of poor women of child bearing age in the study countries are needed. Such efforts may ultimately increase prenatal care utilization in this group.

Our study is unique in analyzing rich infant-level data that was collected and analyzed using similar methods across two countries. Measuring race and ethnicity in South American populations is challenging because of the high racial and ethnic admixture and varying perceptions of race and ethnicity between countries. Race and ethnicity are not routinely measured for research or administrative purposes and are vaguely characterized in many South American countries. This does not however mean that there are no perceptions of racial or ethnic variation in the population. The ancestry measure that we employ is innovative and flexible allowing for defining and comparing ethnic groups of sufficient sample sizes. However, a weakness of this measure is dependence on the mother's recall of the child's lineage, which may introduce error and bias that vary with maternal education or socioeconomic status, which even though we adjust for, may not be sufficient for completely addressing this issue. The lack of data on the exact recall window and the uncertainty about the child's ancestry and the likely variation of these across mothers suggest measurement error in our ancestry measure. This may result in underestimating the PTB disparities by ethnic ancestry. However, the fact that we observe significant disparities and that we are able to explain an important fraction of these disparities suggests that our results may be even more pronounced without such potential measurement error in ancestry. In other words, this measurement error is unlikely to bias the effects of the explanatory variables upward (i.e. towards explaining more of the disparity) but rather downward (i.e. towards explaining less of the disparity). Another limitation of the ancestry measure is its inability to separate in many cases between indigenous ancestry and early European migrations which could be both reported in the "Native" group as discussed above; the extent of admixture in this group may vary between the two countries; for e.g., the Native only group in Ecuador likely includes more indigenous ancestry than the Native only group in Argentina. Clearly however, cases in the Native only group represent a different ancestry overall from those in the European only group.

A few other limitations are worth discussing. We did not have measures on several conceptually relevant variables such as household income, detailed data on all maternal health problems, etiology of preterm birth (medically indicated versus spontaneous preterm labor), maternal health behaviors (e.g., nutrition and stress) and psychosocial status during pregnancy, and neighborhood economic and social indicators and were, therefore, unable to evaluate these potential mechanisms. Even though we find no effects of the city of birth on explaining disparities, neighborhood characteristics within each city may still be relevant. This limitation may have contributed to not explaining a greater fraction of the disparity in Argentina, but it may also be relevant for explaining the differences in use of prenatal services in Ecuador.

A related limitation is that some of the household variables we analyze such as prenatal care use and maternal health and obstetric history are potentially endogenous (i.e. correlated with unobservable confounders) and may have biased effects, which would also bias how much

they can explain of the PTB disparities. This is especially important for prenatal care which emerged as the key variable for explaining the PTB disparities. On the one hand, there is the possibility of confounders such as smoking and nutrition that are likely to be correlated with prenatal care and birth outcomes in the same direction. Such confounders may result in overestimating the effects of prenatal care on PTB and on explaining the ethnic disparity. On the other hand, other unobserved confounders such as pregnancy complications or history of complications in prior pregnancy (e.g. PTB status of prior births) may result in an opposite bias due to adverse selection with women who have greater health risks obtaining more prenatal care. Self-selection into prenatal care based on unobservable confounders has generally been shown in several previous studies (including in studies of South American populations) to result in underestimating the benefits from prenatal care (e.g., Wehby et al. 2009b; Wehby et al. 2009c). If so, prenatal care differences could explain even more of the PTB disparity when accounting for the bias from unobservable confounders than in the current model.

These opposite sources of confounding and prior evidence on bias resulting in underestimation of prenatal care effects suggest that it is unlikely that this limitation substantially affects the main implications of our findings. It is impossible however to fully gauge the extent of the bias in this specific context without adequate data on such confounders or exogenous variation in prenatal care to estimate causal effects. ECLAMC began collecting data on smoking during pregnancy, a potential confounder of prenatal care, in recent years. There is however a concern about underreporting of smoking especially for countries such as Ecuador, where 2–3% of the mothers report smoking during pregnancy. In Argentina, nearly 27% of the mothers report smoking during pregnancy.

To examine whether controlling for smoking modified prenatal care effects, we re-estimated the multivariate logistic regression for PTB described above with and without controlling for smoking for births in 2004 through 2011 in Argentina. We found no difference in prenatal care effects on PTB between the two models (Supplementary Table S5). Smoking did not have a significant association with PTB. Also, the extent of the ethnic gap in preterm birth explained by maternal use of health services (nearly 2 percentage-points) did not change based on whether smoking was controlled for or not.

To examine the extent of confounding from maternal health characteristics, we added to the above sensitivity model specific indicators for the most common illnesses reported in the analysis. We did not include indicators for specific illnesses in the main model because most of these illnesses are too infrequent to be individually analyzed in a meaningful way. In this additional check, we added indicators for anemia, flu, urinary tract infections, hypertension and/or pre-eclampsia (pre-eclampsia too rare to analyze alone), and diabetes. Adding these variables resulted in no changes in the effects of prenatal care (Supplementary Table S5), or in the extent to which maternal use of services explained of the ethnic PTB disparity (nearly 2 percentage points). Of these indicators, only the indicator for hypertension and/or pre-eclampsia was significantly associated with an increase in preterm birth risk (OR=2.3; 95% CI: 1.1–4.8). Furthermore, these indicators and the other measures of maternal health and obstetric history as a group continued not to explain the ethnic PTB disparity. Notwithstanding the potential measurement error in smoking and these health indicators,

these sensitivity checks provide assurance for the key results of our main analysis. Like smoking, chronic illnesses were too rare to meaningfully analyze as separate conditions (consistent with the lower rate of any chronic illness compared to Argentina). Controlling for the flu, and urinary tract infections, the two most common acute illnesses, had no impact on the effects of prenatal care or how much use of healthcare services explained the ethnic disparity in PTB.

Another potential caveat is excluding observations from the analytical sample due to missing data. It is unlikely that there are systematic biases in missing data that would bias the conditional relationships between the explanatory variables and PTB or how much they explain of the ethnic disparity in PTB. Imputing missing data introduces the risk of inflating measurement error which may outweigh any benefit gained from a greater sample size. As a sensitivity check however, we imputed the variables with the highest rate of missing data including PTB, number of prior pregnancies, maternal occupational status, and father's characteristics (age, education, and occupational activity). We also imputed prenatal visits given its importance in explaining the disparities that we observed. For imputation, we employed multivariate models (with different functional forms depending on the distribution of the imputed variables) that included ethnic ancestry, maternal age, maternal education, year and hospital of birth fixed effects as predictors. When imputing PTB and prenatal visits, we also included birth weight as a predictor. These imputations increased the sample size by 1692 observations for Argentina and by 480 observations for Ecuador. We did not find any appreciable difference in the results; the main result continues to be that use of prenatal care is the key factor in explaining the ethnic disparity in PTB in these samples and has consistent effects on PTB.

Our samples were recruited from selective hospitals and may not necessarily be representative of the entire population. Therefore, it is uncertain whether our results are nationally generalizable. Representativeness varies between the two countries because of differences in how many cities are represented in each sample (19 cities in 10 provinces and federal capital district in Argentina, and 7 cities in 5 provinces in Ecuador; list of cities and provinces and total ECLAMC births by city are included in Supplementary Tables S4 and S5). While generalizability is less certain at the country level, it is rather high at the city level. Many ECLAMC hospitals are considered large community hospitals that together cover a relatively large percentage of births in their respective cities. Combined, the total number of births in ECLAMC hospitals in the covered cities represents about 11% and 13.1 % of the total births in these cities in Argentina and Ecuador, respectively, based on 2010 census data for Argentina and 2013 census data in Ecuador. Our analytical sample is obviously a small fraction of the total number of births in these hospitals; nonetheless, this indicates that the ECLAMC hospitals from which our sample is selected are major providers of maternity care in their communities. Furthermore, as shown in Table 1, the samples have extensive socioeconomic and demographic diversity, which is expected to enhance the generalizability of the results. Also, ancestry was not related to any inclusion/exclusion criteria for enrolling these samples, which make them particularly useful for studying ethnic disparities in the study countries as they are not limited by ancestry biases in selection. Finally, the overwhelming majority of births in these countries occur in healthcare facilities; therefore, there is no real bias from excluding at-home births (Woodhouse et al., 2014). In

other words, even though our results may not necessarily be generalizable to the entire birth population in each country, they are likely generalizable to a meaningful proportion of the birth population.

We are aware of no nationally representative child-level data from these countries with data on ethnic ancestry that provide the level of detail on household characteristics that are available in ECLAMC data and that we can employ in this analysis. Therefore, this dataset presents a unique resource for understanding these disparities until adequate nationally representative datasets become available. The lack of such national datasets substantially hinders research on social, economic, and healthcare determinants of infant health in the study countries. National datasets could build on existing birth record registries by expanding the collection of data to cover additional key demographic, socioeconomic, and health variables such as the ones utilized in the current study. These datasets should also include sufficient detail on maternal health (e.g., indicators of specific health problems) and infant health (e.g., etiology of preterm birth, fetal growth indicators) to increase the capacity of future maternal and child health research in these countries.

CONCLUSION

We find that differences in use of prenatal care services explain an important fraction of ethnic/racial disparities in PTB in Argentina and Ecuador. Because prenatal care is a relatively low cost intervention, public policies that increase access to prenatal care may bring important population health benefits by reducing ethnic disparities in PTB these two countries.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

The study was supported by grant 1R03TW008110A2 from the National Institutes of Health / Fogarty International Center, USA. The authors thank all health professionals and coordinators of the Latin American Collaborative Study of Congenital Anomalies (ECLAMC) for their efforts in data collection.

References

- Antunes JLF, Peres MA, Mello TRdC, Waldman EA. Multilevel assessment of determinants of dental caries experience in Brazil. *Community Dentistry and Oral Epidemiology*. 2006; 34:146–152. [PubMed: 16515679]
- Blencowe H, Cousens S, Oestergaard MZ, et al. National, regional, and worldwide estimates of preterm birth rates in the year 2010 with time trends since 1990 for selected countries: a systematic analysis and implications. *The Lancet*. 2012; 379:2162–2172.
- Castilla EE, Orioli IM. ECLAMC: the Latin-American collaborative study of congenital malformations. *Community Genet*. 2004; 7:76–94. [PubMed: 15539822]
- Coria-Soto IL, Bobadilla JL, Notzon F. The effectiveness of antenatal care in preventing intrauterine growth retardation and low birth weight due to preterm delivery. *International Journal for Quality in Health Care*. 1996; 8:13–20. [PubMed: 8680812]
- Fairlie RW. An extension of the Blinder-Oaxaca decomposition technique to logit and probit models. *Journal of Economic and Social Measurement*. 2005; 30:305–316.

- Guarnizo-Herreno CC, Wehby GL. Explaining Racial/Ethnic Disparities in Children's Dental Health: A Decomposition Analysis. *Am J Public Health*. 2012; 102:859–66. [PubMed: 22420801]
- Hertel-Fernandez AW, Giusti AE, Sotelo JM. The Chilean infant mortality decline: improvement for whom? Socioeconomic and geographic inequalities in infant mortality, 1990–2005. *Bulletin of the World Health Organization*. 2007; 85:798–804. [PubMed: 18038062]
- Ickovics JR, Kershaw TS, Westdahl C, et al. Group prenatal care and preterm birth weight: results from a matched cohort study at public clinics. *Obstetrics & Gynecology*. 2003; 102:1051–1057. [PubMed: 14672486]
- Jewell RT. Infant Mortality in Uruguay: The Effect of Socioeconomic Status on Survival. *The Journal of developing areas*. 2014:48.
- Kent AL, Wright IMR, Abdel-Latif ME. Wales tNS Group ACTNICUA. Mortality and Adverse Neurologic Outcomes Are Greater in Preterm Male Infants. *Pediatrics*. 2012; 129:124–131. [PubMed: 22184652]
- Kogan M, Alexander G. Social and behavioral factors in preterm birth. *Prenatal and Neonatal Medicine*. 1998; 3:29–31.
- Krueger PM, Scholl TO. Adequacy of prenatal care and pregnancy outcome. *JAOA: Journal of the American Osteopathic Association*. 2000; 100:485–492. [PubMed: 10979253]
- Lhila A, Long S. What is driving the black-white difference in low birthweight in the US? *Health Economics*. 2012; 21:301–15. [PubMed: 21294220]
- Matijasevich A, Santos IS, Silveira MF, et al. Inequities in maternal postnatal visits among public and private patients: 2004 Pelotas cohort study. *BMC Public Health*. 2009; 9:335. [PubMed: 19751521]
- Morse SB, Zheng H, Tang Y, Roth J. Early School-Age Outcomes of Late Preterm Infants. *Pediatrics*. 2009; 123:e622–e629. [PubMed: 19336353]
- Nyarko KA, Lopez-Camelo J, Castilla EE, Wehby GL. Explaining Racial Disparities in Infant Health in Brazil. *American Journal of Public Health*. 2013a; 103:1675–1684. [PubMed: 23409894]
- Nyarko KA, Lopez-Camelo J, Castilla EE, Wehby GL. Does the Relationship between Prenatal Care and Birth Weight Vary by Oral Clefts? Evidence Using South American and United States Samples. *The Journal of pediatrics*. 2013b; 162:42–49. e1. [PubMed: 22835882]
- Oaxaca R. Male-Female wage differentials in urban labor markets. *International Economic Review*. 1973; 14:693–709.
- Ortiz J, Van Camp J, Wijaya S, Donoso S, Huybregts L. Determinants of child malnutrition in rural and urban Ecuadorian highlands. *Public Health Nutrition*. 2013; 17:2122–2130. [PubMed: 24073991]
- Victora CG, Matijasevich A, Silveira M, et al. Socio-economic and ethnic group inequities in antenatal care quality in the public and private sector in Brazil. *Health Policy Plan*. 2010; 25:253–61. [PubMed: 20123940]
- Villar J, Ba'aqeel H, Piaggio G, et al. WHO antenatal care randomised trial for the evaluation of a new model of routine antenatal care. *The Lancet*. 2001; 357:1551–1564.
- Wehby G. Child health insurance and early preventive care in three South American countries. *Health Policy Plan*. 2013; 28:328–38. [PubMed: 22791558]
- Wehby G, Gili JA, Pawluk M, Castilla EE, López-Camelo JS. Disparities in Birth Weight and Gestational Age by Ethnic Ancestry in South American countries. *Int J Public Health*. 2015; 60(3): 343–51. [PubMed: 25542227]
- Wehby G, McCarthy AM, Castilla EE, Murray JC. 2012The Impact of Household Investments on Early Child Neurodevelopment and on Racial and Socioeconomic Developmental Gaps - Evidence from South America. *Forum for Health Economics & Policy*. :14.
- Wehby G, Murray JC, Castilla EE, Lopez-Camelo JS, Ohsfeldt RL. Prenatal care effectiveness and utilization in Brazil. *Health Policy Plan*. 2009a; 24:175–88. [PubMed: 19282483]
- Wehby G, Murray JC, McCarthy AM, Castilla EE. Racial Gaps in Child Health Insurance Coverage in Four South American Countries: The Role of Wealth, Human Capital, and Other Household Characteristics. *Health Services Research*. 2011; 46:2119–2138. [PubMed: 21210797]
- Wehby GL, Castilla EE, Lopez-Camelo J. The impact of altitude on infant health in South America. *Economics & Human Biology*. 2010; 8:197–211. [PubMed: 20594925]

- Wehby GL, McCarthy AM. Economic gradients in early child neurodevelopment: A multi-country study. *Social Science & Medicine*. 2013; 78:86–95. [PubMed: 23273409]
- Wehby GL, Murray JC, Castilla EE, Lopez-Camelo JS, Ohsfeldt RL. Prenatal care demand and its effects on birth outcomes by birth defect status in Argentina. *Economics & Human Biology*. 2009b; 7:84–95. [PubMed: 19059012]
- Wehby GL, Murray JC, Castilla EE, Lopez-Camelo JS, Ohsfeldt RL. Quantile effects of prenatal care utilization on birth weight in Argentina. *Health Economics*. 2009c; 18:1307–1321. [PubMed: 19142894]
- Wehby GL, Nyarko KA, Lopez-Camelo JS, Ohsfeldt RL. Fetal health shocks and early inequalities in health capital accumulation. *Health Economics*. 2014; 23:69–92. [PubMed: 23339079]
- Woodhouse C, Lopez Camelo J, Wehby GL. A comparative analysis of prenatal care and fetal growth in eight South American countries. *PLoS One*. 2014; 9:e91292. [PubMed: 24625630]

Table 1

Description of Study Variables by Child's Ancestry and Country

	Argentina		Ecuador	
	Native only [N=3883] % or mean (SD)	European only [N=1236] % or mean (SD)	Native only [N=660] % or mean (SD)	European only [N=919] % or mean (SD)
Preterm Birth	10.4	7.4	12.7	7.0
Maternal Healthcare				
No prenatal visits	2.3	1.0	1.7	0.3
1 to 2 prenatal visits	6.6	2.3	6.4	1.7
3 to 5 prenatal visits	30.9	19.8	27.9	17.5
6 to 8 prenatal visits	39.4	40.9	54.7	65.8
9 or more visits	20.7	36.1	9.4	14.6
Vaccination 1 st trimester	14.8	8.1	27.9	18.7
Vaccination 2 nd trimester	58.2	58.9	50.5	54.7
Vaccination 3 rd trimester	56.1	43.5	28.9	36.0
Maternal Health and Obstetric History				
Any acute illness				
Yes	38.5	38.9	29.2	36.3
No	61.5	61.1	70.8	63.7
Any chronic illness				
Yes	15.0	14.8	1.8	3.4
No	85.0	85.2	98.2	96.6
Conception difficulty				
Yes	5.9	4.3	3.8	3.0
No	94.1	95.7	96.2	97.0
Number of prior pregnancies	1.8 (2.0)	1.4 (1.8)	1.0 (1.4)	0.8 (1.1)
Demographics				
Female infants	46.1	47.2	40.9	47.3
Male infants	53.9	52.8	59.1	52.7
Maternal age <20	21.5	13.8	18.0	5.2
Maternal age 20–24	32.9	24.4	33.6	24.9
Maternal age 25–29	22.7	27.4	22.6	35.1
Maternal age 30–34	13.5	20.0	16.5	20.6
Maternal age 35–39	6.7	10.7	6.5	11.0
Maternal age 40	2.5	3.7	2.7	3.2
Paternal age 24	36.5	25.4	36.4	20.2
Paternal age 25–39	54.4	63.2	56.4	69.9
Paternal age 40	9.1	11.4	7.3	9.9

	Argentina		Ecuador	
	Native only [N=3883] % or mean (SD)	European only [N=1236] % or mean (SD)	Native only [N=660] % or mean (SD)	European only [N=919] % or mean (SD)
<i>Household Socioeconomic Status</i>				
<i>Maternal education^I</i>				
Primary school incomplete	13.4	6.0	10.8	2.2
Primary school complete	36.1	17.1	31.7	9.2
Secondary school incomplete	30.6	25.7	28.2	18.6
Secondary school complete	16.1	26.1	19.2	32.0
University incomplete	2.7	11.5	5.6	18.9
University complete	1.0	13.6	4.6	19.0
<i>Maternal occupation</i>				
Stay-at-home or unemployed	82.8	65.1	47.3	21.0
Unskilled blue collar	8.4	5.3	22.4	10.0
Skilled blue collar	3.4	2.7	10.0	7.1
Independent worker	1.3	4.0	1.4	2.4
Clerk	3.5	11.6	17.0	49.9
Executive, owner, or boss	0.6	11.3	2.0	9.6
<i>Paternal education^I</i>				
Primary school incomplete	14.0	6.1	7.0	1.2
Primary school complete	40.7	23.7	30.9	8.5
Secondary school incomplete	24.3	23.5	23.3	14.1
Secondary school complete	18.3	26.5	27.7	36.1
University incomplete	2.0	8.7	6.1	19.4
University complete	0.9	11.5	5.0	20.7
<i>Paternal occupation</i>				
Stay-at-home or unemployed	10.7	8.2	6.7	1.6
Unskilled blue collar	35.1	19.5	32.1	8.8
Skilled blue collar	22.9	15.6	21.2	13.8
Independent worker	11.9	17.3	5.2	6.1
Clerk	17.5	22.7	32.1	60.2
Executive, owner, or boss	1.9	16.7	2.7	9.5

Notes: The table shows % of categorical variables and means (standard deviations) for continuous variables. The sample distribution across the city of birth is shown in Supplementary Tables S4 and S5 online.

^I“Primary school incomplete” refers to parents who did not graduate from primary school; it also includes a small group of illiterate parents as well as literate parents without formal schooling. “Primary school complete” refers to parents who graduated from primary school but did not have higher education. “Secondary school incomplete” refers to parents who attended but did not graduate from secondary school. “Secondary school complete” refers to parents who graduated from secondary school but did not attend university. “University incomplete” refers to parents who attended but did not graduate from university. “University complete” refers to parents who graduated from university.

Decomposition of Disparities in Preterm Birth Rate between Children of Native only Ancestry and Those of European Only Ancestry across Categories of Variables

Table 2

	Argentina	Ecuador
	Native only vs European only [N=5119]	Native only vs European only [N=1579]
Difference in PTB rate	-0.0296	-0.0576
Explained difference ¹	-0.0245	-0.0170
% explained	82.8	29.5
% unexplained	17.2	70.5
Variable Categories		
Maternal healthcare use	-0.0168 *** (0.0040)	-0.0170 ** (0.0066)
Maternal health and obstetric history	0.0015 (0.0010)	0.0017 (0.0020)
Demographics	-0.0007 (0.0015)	-0.0084 (0.0053)
Household socioeconomic status	-0.0077 ** (0.0037)	-0.0148 (0.0184)
Geographic Location	-0.0011 (0.0035)	-0.0185 (0.0157)

Notes: The Table shows the total difference in PTB rate between children of Native only and European only ancestry, the explained difference based on categories of variables that were significant at least at p<0.1, and the contribution of each category of variables to explaining the difference, whether significant or not (standard errors in parentheses); "explained" disparities that are of the opposite sign as the total disparity (e.g., maternal health and obstetric history) indicate that the PTB disparity would have been greater if the distribution of the variables in those categories were the same between children of Native only ancestry and those of European only ancestry. The models include year of birth fixed effects.

** indicates p<0.05;

*** indicates p<0.01.

¹ Explained difference is based on the categories that significantly explained the disparity (at p<0.05), which includes maternal healthcare use for both countries, and household socioeconomic status for Argentina (contributions of insignificant categories not included).

Table 3

Decomposition of The Disparity in Preterm Birth Rate by Child's Ancestry across Variables without Grouping Variables in Categories in Decomposition

	Argentina [N=5119]	Ecuador [N=1579]
No prenatal visits	−0.0051 *** (0.0014)	−0.0026 ** (0.001)
1 to 2 prenatal visits	−0.0087 *** (0.0025)	−0.0066 *** (0.002)
3 to 5 prenatal visits	−0.0110 *** (0.0039)	−0.0051 (0.0032)
6 to 8 prenatal visits	0.0046 * (0.0027)	0.0083 ** (0.0036)
9 or more visits	Reference	Reference
Vaccination 1 st trimester	0.0007 (0.002)	−0.0071 (0.0046)
Vaccination 2 nd trimester	−0.0000 (0.0003)	−0.0011 (0.0012)
Vaccination 3 rd trimester	0.0028 ** (0.0012)	−0.0028 (0.0025)
Any acute illness	0.0003 (0.0004)	0.0013 (0.0017)
Any chronic illness	0.0000 (0.0001)	−0.0011 ** (0.0004)
Conception difficulty	−0.0000 (0.0003)	0.0001 (0.0006)
Number of prior pregnancies	0.0012 (0.0008)	0.0012 (0.0022)
Female infants	−0.0002 (0.0002)	−0.0008 (0.0005)
Maternal age <20	0.0003 (0.001)	−0.0112 *** (0.0041)
Maternal age 20–24	−0.0002 (0.0005)	−0.0008 (0.0008)
Maternal age 25–29	Reference	Reference
Maternal age 30–34	0.0004 (0.0006)	0.0046 * (0.0026)
Maternal age 35–39	0.0014 (0.0009)	0.0002 (0.0006)
Maternal age 40	0.0005 (0.0006)	−0.0000 (0.0003)
Paternal age 24	−0.0015 (0.0011)	−0.0007 (0.0022)
Paternal age 25–39	Reference	Reference
Paternal age 40	−0.0000 (0.0003)	−0.0000 (0.0003)
<i>Maternal education</i>		
Primary school incomplete	Reference	Reference
Primary school complete	0.0028 (0.0022)	−0.0005 (0.0085)
Secondary school incomplete	0.0000 (0.0005)	0.0006 (0.0023)
Secondary school complete	−0.0047 *** (0.0015)	−0.0045 (0.0046)
Attended university ¹	−0.0041 (0.0051)	−0.0053 (0.0098)
<i>Maternal occupation</i>		
Stay-at-home or unemployed	Reference	Reference
Unskilled blue collar	0.0005 (0.0004)	0.0055 (0.0045)
Skilled blue collar	0.0000 (0.0001)	0.0003 (0.001)
Independent worker	−0.0003 (0.0007)	−0.0003 (0.0005)
Clerk	0.0002 (0.001)	−0.0009 (0.0176)
Executive, owner, or boss	−0.0008 (0.0025)	−0.0076 ** (0.0039)
<i>Paternal education</i>		

	Argentina [N=5119]	Ecuador [N=1579]
Primary school incomplete	Reference	Reference
Primary school complete	0.0015(0.0015)	0.0070(0.0054)
Secondary school incomplete	−0.0001(0.0004)	0.0037(0.0028)
Secondary school complete	−0.0002(0.0014)	−0.0034 ^{**} (0.0017)
Attended university ^I	−0.0002(0.0031)	−0.0051(0.0053)
<i>Paternal occupation</i>		
Stay-at-home or unemployed	Reference	Reference
Unskilled blue collar	0.0021(0.0014)	0.0043(0.006)
Skilled blue collar	0.0011(0.0008)	0.0023(0.0016)
Independent worker	−0.0017(0.0012)	−0.0001(0.0003)
Clerk	0.0002(0.0007)	−0.0128(0.0098)
Executive, owner, or boss	−0.0041 ^{**} (0.0018)	0.0020(0.0029)

Notes: The Table shows the extent to which each variable contributes to the total difference in PTB rate between children of Native only and European only ancestry, regardless of significance (standard errors in parentheses). Infants of European only ancestry have lower PTB rates than those of Native only ancestry by 0.0296 in Argentina and 0.0576 in Ecuador. Contributions of variables that explain this disparity are negatively signed. Contributions of variables that are of the opposite (positive) sign to this disparity (e.g., obtaining 6–8 versus 9 or more visits) indicate that the PTB disparity would have been greater if the distributions of those variables (their proportions or means) were the same between the two ancestry groups. Detailed results for the city of birth are not shown for brevity but are available from the authors upon request. The models include year of birth fixed effects.

* indicates $p < 0.1$;

** indicates $p < 0.05$;

*** indicates $p < 0.01$

^I Attended university includes both parents who attended university but did not graduate as well as university graduates.