Research Article

Evaluating Treatment and Generalization Patterns of Two Theoretically Motivated Sentence Comprehension Therapies

Carrie A. Des Roches,^a Sofia Vallila-Rohter,^{a,b} Sarah Villard,^a
Yorghos Tripodis,^c David Caplan,^{a,d} and Swathi Kiran^a

Purpose: The current study examined treatment outcomes and generalization patterns following 2 sentence comprehension therapies: object manipulation (OM) and sentence-to-picture matching (SPM). Findings were interpreted within the framework of specific deficit and resource reduction accounts, which were extended in order to examine the nature of generalization following treatment of sentence comprehension deficits in aphasia.

Method: Forty-eight individuals with aphasia were enrolled in 1 of 8 potential treatment assignments that varied by task (OM, SPM), complexity of trained sentences (complex, simple), and syntactic movement (noun phrase, *wh*-movement). Comprehension of trained and untrained sentences was probed before and after

treatment using stimuli that differed from the treatment stimuli.

Results: Linear mixed-model analyses demonstrated that, although both OM and SPM treatments were effective, OM resulted in greater improvement than SPM. Analyses of covariance revealed main effects of complexity in generalization; generalization from complex to simple linguistically related sentences was observed both across task and across movement.

Conclusions: Results are consistent with the complexity account of treatment efficacy, as generalization effects were consistently observed from complex to simpler structures. Furthermore, results provide support for resource reduction accounts that suggest that generalization can extend across linguistic boundaries, such as across movement type.

For many years, research has demonstrated the presence of comprehension deficits in aphasia that become particularly apparent when the thematic roles of a sentence can plausibly be interchanged (Caplan, Baker, & Dehaut, 1985; Caplan, Waters, DeDe, Michaud, & Reddy, 2007; Caramazza & Zurif, 1976; Grodzinsky, 2000). When thematic roles are semantically reversible, as in the sentence “The boy pushed the girl,” thematic role assignment is critical: Listeners must understand who is doing the action (the agent) and who is receiving the action (the theme) in order to correctly interpret such sentences.

Thus, the presence of comprehension deficits of reversible sentences in aphasia has led to the hypothesis that impairments are rooted in a difficulty using syntactic structure to interpret the meanings of sentences (Caplan et al., 1985).

Theoretical perspectives that have been proposed to explain the nature of these comprehension deficits in persons with aphasia (PWA) tend to fall into two categories: specific deficit accounts and resource reduction accounts. Specific deficit models argue that sentence comprehension deficits in aphasia stem from a specific impairment in syntactic interpretation. The observation that PWA show impaired comprehension of certain sentence types with shared syntactic structures (Grodzinsky, 2000; Grodzinsky, Piñango, Zurif, & Draí, 1999) has led to hypotheses that deficits are rooted in a lost ability to execute particular syntactic computations or operations. For example, Grodzinsky’s trace deletion hypothesis (Grodzinsky, 1986, 1995a, 1995b, 2000), which is based on Chomsky’s (1995) model of syntactic structure, posits that PWA have lost the ability to relate syntactic constituents of a sentence to their original syntactic position through the coindexation of “traces.” Likewise, mapping

^aSpeech and Hearing Sciences, Boston University Sargent College, MA

^bDepartment of Communication Sciences and Disorders, MGH-Institute of Health Professions, Boston, MA

^cDepartment of Biostatistics, Boston University, MA

^dNeuropsychology Laboratory, Massachusetts General Hospital, Boston

Correspondence to Swathi Kiran: kirans@bu.edu

Editor: Anastasia Raymer

Associate Editor: Gayle DeDe

Received September 14, 2015

Revision received March 4, 2016

Accepted May 2, 2016

DOI: 10.1044/2016_AJSLP-15-0134

Disclosure: The authors have declared that no competing interests existed at the time of publication.

deficit accounts suggest that impairments arise as a result of faulty mapping of thematic role information and syntax, leading to a reduced integration of sentence form and sentence meaning (Linebarger, 1995; Linebarger, Schwartz, & Saffran, 1983; Schwartz, Linebarger, Saffran, & Pate, 1987). It is important to note that these models attribute comprehension deficits to syntactic structure and interpretation.

An alternative account of sentence comprehension deficits in aphasia is the resource reduction theory (Caplan, DeDe, & Michaud, 2006; Caplan & Waters, 2006; Caplan et al., 2007; DeDe & Caplan, 2006; Haarmann, Just, & Carpenter, 1997). This theory maintains that the difficulty extracting the semantic meaning of sentences in PWA is due to limitations in their ability to perform parsing and interpretative operations (Caplan et al., 2007). In other words, PWA may not have deficits specific to syntactic parsing operations but may instead have general deficits in processing resources. Reduced abilities in domains such as verbal working memory, phonological store, and short-term semantic memory, for example, may constrain an individual's ability to perform the computations necessary for correct interpretation of syntactic structure. Resource reduction models predict that comprehension deficits will surface across multiple sentence types, particularly across sentences that contain a similar number of operations and consequently, pose similar processing demands. It is interesting to note that specific deficit accounts and resource reduction accounts make different predictions about the sentences that will be affected by deficits in aphasia. By extension, therefore, specific deficit and resource reduction accounts also make different predictions about mechanisms of therapy and about how treatment-related improvements may generalize to untrained sentence types and contexts. The current study is motivated in part by what observed patterns of treatment-induced generalization may be able to tell us about accounts of sentence comprehension deficits in aphasia.

Sentence comprehension therapies for PWA have been developed and evaluated on the basis of factors related to specific deficit and resource reduction hypotheses, three of which are central to the current study: syntax, complexity, and task. First, a number of therapies have been developed that are based on principles of syntax. Mapping therapy, for example, is designed to repair impaired mapping between sentence constituents and thematic roles (Byng, 1988; Byng, Nickels, & Black, 1994; Haendiges, Berndt, & Mitchum, 1996; Jones, 1986; Le Dorze, Jacob, & Coderre, 1991; Marshall, Pring, & Chiat, 1993; Mitchum, Haendiges, & Berndt, 1995; Nickels, Byng, & Black, 1991; Rochon, Laird, Bose, & Scofield, 2005; Schwartz, Saffran, Fink, Myers, & Martin, 1994; for a review, see Fink, 2001). Likewise, the treatment of underlying forms (TUF; Thompson & Shapiro, 2005) guides PWA through the processes of relearning specific syntactic operations such as reordering constituents of sentences that involve noun phrase (NP) movement or *wh*-movement. Mapping therapies and TUF have been influenced by specific deficit accounts that propose that syntactic structure is at the source of comprehension deficits. These therapies

lead to improved comprehension and often to improved sentence production.

The second factor relevant to sentence comprehension treatment is complexity. Therapy-related improvements have been seen on trained sentences as well as on untrained sentences, particularly following TUF, and examinations of these patterns of progress and generalization have revealed that complexity may play an important role. One of the goals of TUF is to promote generalization from trained to untrained sentences; therefore, researchers systematically probe progress on syntactically related and unrelated untrained sentence structures. It has been found that generalization following TUF is greatest to linguistically related sentences (Ballard & Thompson, 1999; Jacobs & Thompson, 2000; Thompson & Shapiro, 2005; Thompson, Shapiro, & Roberts, 1993; Thompson, Shapiro, Tait, Jacobs, & Schneider, 1996; Thompson et al., 1997), lending support for specific deficit hypotheses that suggest that specific syntactic operations lie at the root of deficits. Furthermore, generalization occurs frequently from complex to simple structures but less frequently from simple to complex structures (Thompson, Ballard, & Shapiro, 1998; Thompson & Shapiro, 2005, 2007; Thompson, Shapiro, Kiran, & Sobecks, 2003; Thompson et al., 1993, 1997). Together, these observations have led to the establishment of the complexity account of treatment efficacy (CATE), which states that training complex sentences leads to generalization to less complex sentences with shared underlying linguistic properties (Ballard & Thompson, 1999; Jacobs & Thompson, 2000; Thompson & Shapiro, 1995, 2005, 2007; Thompson et al., 1997, 1998, 2003). These generalization results can be interpreted within specific deficit accounts. Specific deficit models attribute sentence comprehension deficits to impaired representations; one would therefore predict that treatment-related gains in comprehension would generalize to sentence structures that contain similar representations (e.g., simple sentences that share linguistic properties of trained sentences). Likewise, resource reduction accounts justify generalization to simple structures that pose similar or fewer resource demands (Kiran et al., 2012) but provide no explanation for the specificity of generalization to linguistically related structures.

The final component examined in treatment studies and evaluations of generalization (and critical to the current study) is the task used to train sentence comprehension. Kiran et al. (2012) recently evaluated two sentence comprehension treatments, comparing outcomes following two different task manipulations. In their study, Kiran et al. (2012) enrolled 15 PWA in one of two pilot therapies adapted from assessment measures: sentence-to-picture matching (SPM) and object manipulation (OM). Treatment was founded on principles of syntactic processing and comprehension. In both protocols, participants were explicitly trained to assign thematic roles in a sentence and were probed on their interpretation of canonical and noncanonical sentences. Individuals were trained on simple or complex sentences that involved NP or *wh*-movement. Both OM and SPM treatments resulted in overall gains. Overall effect sizes were similar across treatment methods, yet more

participants were found to improve following OM treatment than following SPM treatment. Researchers posited that the enactment component of OM may have differently engaged resources than SPM, leading to improved progress with therapy. Preliminary generalization analyses demonstrated the presence of generalization to untrained structures and merit further study.

Therefore, the focus of the current study is to extend the findings of Kiran et al. (2012) by further examining the treatment efficacy of both OM and SPM sentence comprehension therapy and by systematically examining how patterns of generalization may be affected by syntax, complexity, and task. The current study enrolled a large group of participants in treatment and varied syntax (NP or *wh*-movement), complexity (simple or complex), and task (SPM or OM) across participants' treatment. The aim of the study was to systematically examine the nature and extent of generalization, probing whether generalization occurs across task (e.g., OM to SPM), across different levels of complexity (e.g., complex *wh*-movement sentences to simple *wh*-movement sentences), and across syntax, which in this study corresponds to two types of syntactic movement and is referred to as *movement* (e.g., NP movement to *wh*-movement sentences; see Figure 1 for a schematic of target generalization patterns). On the basis of the findings of Kiran et al. (2012), it is predicted that treatment will result in overall gains and that treatment outcomes will be superior following OM relative to SPM treatment, indicating that task type influences the effectiveness of sentence comprehension treatment. Generalization results will be examined in the context of specific deficit and resource reduction accounts. Specific deficit models would predict generalization across task to sentences with the same movement (e.g., OM *wh*-movement to SPM *wh*-movement) because syntactic mechanisms are thought to drive change. Generalization would also be expected across complexity—that is, from complex to simple sentences with the same movement (e.g., complex *wh*-movement to simple *wh*-movement sentences)—but not across movement (e.g., *wh*-movement to NP movement sentences). Resource reduction accounts might also predict generalization across tasks (e.g., OM to SPM) and across complexity in the setting of similar or reduced resource demands. In contrast to specific deficit models, resource reduction accounts could also explain generalization across movement (e.g., *wh*-movement to NP movement sentences), as these models posit that comprehension is not constrained by the syntactic structure of the sentences but rather by the resources required to complete the syntactic operations.

Method

Participants

Forty-eight participants (30 males, 18 females) with aphasia secondary to stroke or traumatic brain injury were recruited from hospitals and aphasia centers in the greater Boston area. The average age of participants was 59.6 years ($SD = 13.7$ years; range = 26–86 years), and the average

number of months postonset upon enrollment in the study was 47.7 months ($SD = 57.6$ months; range = 6–278 months; see online Supplemental Material S1 for individual demographic information). Forty-one participants were premorbidly right-handed and seven (P8, P13, P17, P18, P19, P30, and P48) were premorbidly left-handed. All participants had vision and hearing that were near normal or corrected to near normal. Thirty-seven participants identified themselves as White, nine as Black, and one as Asian; one participant declined to provide information about race.

Participants completed a screening battery prior to treatment to determine eligibility and to establish baseline language and cognitive abilities.¹ To be eligible for therapy, participants had to demonstrate adequate comprehension of the nouns and verbs that composed treatment sentences. To this end, participants completed two screeners in which they were shown either a pair of nouns (noun screener) or a pair of verbs (verb screener) and were instructed to indicate the picture that corresponded to an auditorily presented target. The average accuracy score for enrolled participants on noun screeners was 94.1% ($SD = 6.8\%$). The average accuracy for verb screeners was 85.0% ($SD = 11.6\%$). Participants also completed OM and SPM screeners to assess comprehension of active sentences, *wh*-movement sentences, and NP movement sentences (see the Stimuli section for additional details). Screener accuracy on comprehension of active sentences was on average 77.2% ($SD = 21.9\%$). Screener accuracy on comprehension of noncanonical sentences was 47.1% ($SD = 21.2\%$); see online Supplemental Material S1 for individual scores. This difference in comprehension between canonical and noncanonical sentences was expected as treatment was targeted toward the impaired noncanonical sentence forms (NP and *wh*-movement).

Our cognitive–linguistic baseline assessment battery included the following assessments: Western Aphasia Battery–Revised (WAB-R; Kertesz, 2007), Cognitive–Linguistic Quick Test (Helm-Estabrooks, 2001), Northwestern Assessment of Verbs and Sentences (Thompson, 2011), Bells Test (Gauthier, Dehaut, & Joanne, 1989), the Picture Arrangement task ($M = 36.0\%$, $SD = 24.4\%$) from the Wechsler Adult Intelligence Scale (Wechsler, 1997), and several different subtests of the Psycholinguistic Assessment of Language (PAL; Waters & Caplan, 1995): PAL 3 examined auditory lexical decision ($M = 85.9\%$, $SD = 10.7\%$), PAL 5 examined auditory attribute verification ($M = 81.0\%$, $SD = 14.6\%$), PAL 8 examined picture homophone matching ($M = 70.6\%$, $SD = 19.5\%$), PAL 10 examined affixed auditory lexical decision ($M = 80.9\%$, $SD = 12.2\%$), PAL 11 examined affixed auditory word-to-picture matching ($M = 77.4\%$, $SD = 18.1\%$), and PAL 12 examined auditory relatedness ($M = 62.3\%$, $SD = 14.7\%$). The subtests of PAL were used to establish a basic level of auditory processing skills, which were needed to perform the therapy tasks, whereas the remaining tests in the

¹In select cases, participants began treatment after completing eligibility screeners but before fully completing the cognitive–linguistic battery.

Figure 1. Schematic depicting observed generalization patterns. We examined generalization (a) across task, within complexity, and within movement; (b) within task, across complexity, and within movement; and (c) within task, within complexity, and across movement. NP = noun phrase.

baseline assessment battery were used to establish a baseline level of aphasia severity, cognitive abilities, and sentence comprehension and production. With the exception of the Bells Test, the cognitive–linguistic assessments were not used for inclusion criterion; scores on the Bells Test were used to rule out spatial neglect ($M = 92.4\%$, $SD = 15.9\%$). A detailed examination of cognitive–linguistic scores and their influence on the treatment outcomes is outside the scope of the current article. Several scores from the baseline assessments (WAB-R Aphasia Quotient, Cognitive-Linguistic Quick Test Composite Severity, and Northwestern Assessment of Verbs and Sentences Sentence Comprehension Test canonical and noncanonical sentence scores) have been included in online Supplemental Material S1 to provide information on each participant’s overall severity.

Stimuli

Stimuli were created for the screener battery, monitoring battery, and treatment materials. Noun stimuli for all experimental phases consisted of black and white drawings depicting people, vehicles, and objects. Half of the stimulus nouns were animate and half were inanimate. Noun and verb animacy and lexical frequency were balanced across batteries. Auditory stimuli for screening and monitoring phases of the experiment consisted of recordings of a male voice reading each sentence aloud in North American English using a natural rate of speech and neutral tone. All screening and monitoring items were presented using E-Prime software (Schneider & Zuccolotto, 2002). In treatment, stimulus sentences were spoken by the session clinician. All stimulus sentences were also printed on strips of cardstock for treatment.

For SPM screeners, monitors, and treatment, an entire sentence (e.g., “The aunt was kissed by the man”) was depicted in a single drawing and presented to the participant on a computer screen (during screeners and monitors) or large piece of cardstock (during treatment). Foil pictures (i.e., with thematic roles reversed) were also used during both testing and treatment for SPM. Responses were indicated via button press. The visual stimuli for OM in all phases consisted of nouns depicted as paper dolls. On each trial, dolls were placed on a table in random order and participants indicated the agent and theme of the sentence by

manipulating the dolls. Clinicians recorded responses on a score sheet and coded them as correct or incorrect.

Screening Battery

Participants completed an OM and SPM screener, assessing their comprehension of multiple sentence types. Screeners included four noncanonical sentence types targeted in treatment: object relative (OR), object cleft (OC), passive (PA), and unaccusative (UNACC). Screeners also included two control sentence types—sentential complement (3NP) and OR with embedded subject modified by a prepositional phrase (ORCNP)—as well as active sentences, reflexive sentences, pronoun-as-object sentences, object control, subject control, and NP raising. Participants were assessed on their comprehension of five to 10 items of each sentence type, for a total of 110 items per screener. OM and SPM screeners contained the same sentences presented in a different order. Screener averages for active sentences and noncanonical sentences for each participant are included in online Supplemental Material S1.

Monitoring Batteries

Monitoring batteries included items of all sentence types targeted in treatment as well as two control sentence structures. Two versions of OM and SPM monitoring batteries were created that monitored participants on their comprehension of 15 items of target sentence types OR, OC, and PA and of 10 items of target sentence type UNACC (due to the more limited number of pure UNACC verbs). Monitors also included 10 items each of 3NP and ORCNP control sentence types; 3NP sentences controlled for length (matched the longest target sentence, OR, on length), and ORCNP sentences controlled for complexity (contained an increased level of complexity relative to OR). Each monitor contained 75 items. The two versions of the monitor lists contained the same nouns and verbs but with agent and theme reversed. In addition, the order of sentences in each list was reversed between administrations.

Treatment Materials

Participants were trained on one of four sentence types in treatment: OR, OC, PA, or UNACC. Sentence

types were selected to control for syntactic movement (NP or *wh*-movement) and complexity (simple or complex). OR and OC sentences, for example, both involve *wh*-movement. In addition, OR sentences are more complex than OC sentences (see Table 1 for possible treatment structures and sample sentences). Likewise, PA sentences are more complex than UNACC sentences. Participants were trained on 15 OR, OC, PA, or UNACC sentences. As an alternation in the treatment design midway through the study, the majority of participants (29) also saw 15 canonical sentences containing the same number of nouns as their target sentence type in treatment: participants trained on OR sentences saw subject relative sentences, participants trained on OC sentences saw subject cleft sentences, participants trained on PA sentences saw active sentences, and participants trained on UNACC sentences saw unergative sentences. The inclusion of these sentences was designed to encourage participants to think about thematic role assignment and discourage the development of heuristic strategies that were based on the ordering of constituents of target sentence types. This modification was introduced after the first 19 participants were enrolled, and the additional sentences were not part of monitoring probes.

Experimental Design

The experimental design and procedures for this study were largely the same as those used in Kiran et al. (2012). In brief, a single-subject multiple-baseline design was used (Thompson, 2006). After completing screeners, all eligible participants completed three OM and SPM pretreatment baseline monitoring batteries.² Participants were then assigned to one of eight potential treatment assignments (OM or SPM) targeting one of four sentence structures (OR, OC, PA, or UNACC; see Table 1 and Figure 1). Treatment assignment was counterbalanced and participants were pseudorandomly assigned to task and structure combinations, ensuring that they were assigned to a target task/structure on which they had scored below a 73% average during pretreatment monitors. Table 2 provides demographic information of the participants by task of their treatment assignment.

Treatment was administered twice a week for 2 hr each time. Both SPM and OM monitors were administered to all participants throughout the treatment; participants were administered one monitor each week, alternating between the two tasks. Treatment continued for 10 weeks or until a participant exceeded 80% accuracy on their trained task/structure on two consecutive monitors. At the conclusion of treatment, participants completed three posttreatment OM and SPM monitoring baselines to establish the effect size of treatment.

²Two participants were given a fourth and additional pretreatment baseline to ensure baseline stability. One participant completed three pretreatment baselines and then was absent for several weeks. Upon return, he received two additional baselines to ensure stability prior to initiating treatment. For the purpose of this study, the two baselines that immediately preceded treatment were analyzed.

Treatment Protocol

The two sentence comprehension treatment methods explored in this project were SPM and OM. Treatments were the same as those included in Kiran et al. (2012). SPM and OM treatments were matched on the number of steps and basic procedures. Both treatments trained participants to map the nouns in each sentence onto the thematic roles of the verbs. Individuals were trained to think about and identify who was “doing” (agent) and who was “receiving” (theme) each action of a target sentence. In SPM treatment, participants heard a sentence spoken aloud as they viewed a target and foil picture that depicted the meaning of the sentence. In OM treatment, participants heard a sentence spoken aloud and were asked to manipulate dolls to indicate the agent and theme of the sentence. The steps in both treatments were as follows: (a) The pictures (SPM) or dolls (OM) were placed on the table in front of the participant; (b) the clinician read the sentence aloud and asked the participant to either choose the picture that matched the meaning of the sentence (SPM) or use the dolls to perform the action in the sentence (OM); (c) the participant either chose a picture (SPM) or attempted to enact the action (OM) and was given feedback; (d) the clinician produced a sentence strip with the target sentence; (e) the clinician read the sentence aloud and explained its meaning; (f) the clinician trained the agent of the sentence, and the participant was asked three times to indicate who was doing the action; (g) the clinician trained the theme of the sentence, and the participant was asked three times to indicate who was receiving the action; (h) the clinician trained the entire sentence again; and (i) the clinician repeated steps a through c.

Reliability

Because responses for the SPM screeners and monitors were automatically recorded and scored by the E-Prime software, reliability of scoring of the dependent measure was performed only for the OM task. Scoring of 100% of OM screener and monitor responses was checked by the first and third authors. Given the magnitude of treatment sessions (942), research assistants watched videotapes of 47 (5%) OM and SPM treatment sessions to ensure treatment reliability. Agreement between the treatment protocol and the observed implementation of treatment was found to be 96% for this sample.

Data Analysis

Linear mixed-effects models were used to examine the effectiveness of the treatment. Accuracy measured as proportion of success for pre- and posttreatment baseline data was transformed using the log-odds transformation. Seven mixed models were then performed on the log odds to calculate the odds ratio of the probability of a change in performance on trained structures relative to control structures (3NP and ORCNP). Fixed effects included the treated task, trained sentence type, and time point (e.g., pre- vs. posttreatment baseline); participant was a random effect.

Table 1. Possible treatment assignments and control sentences.

Task	Movement	Complexity	Sentence type	Example sentence
Possible treatment assignments				
OM	<i>wh</i> -movement	Complex	OR	The man who the woman fed lowered the boy.
		Simple	OC	It was the husband who the girl greeted.
	NP movement	Complex	PA	The train was hit by the tank.
		Simple	UNACC	The missile was exploding next to the tank.
SPM	<i>wh</i> -movement	Complex	OR	
		Simple	OC	
	NP movement	Complex	PA	
		Simple	UNACC	
Control sentences (OM and SPM)			ORCNP	The son who the mother of the girl poked tickled the man.
			3NP	The man said that the boy kissed the girl.

Note. OM = object manipulation; OR = object relative; OC = object cleft; NP = noun phrase; PA = passive; UNACC = unaccusative; SPM = sentence-to-picture matching; ORCNP = object relative with a complex noun phrase; 3NP = three-noun phrase.

Cohen's *d* effect sizes (ES) were also calculated for each participant for all eight monitored sentences. To calculate ES, the average pretreatment monitor scores were subtracted from the average posttreatment monitor scores, and the result was divided by the standard deviation of pretreatment monitors (Beeson & Robey, 2006). The advantage of using ES in addition to percentage accuracy was that outcome measures could be normalized across different structures, which allowed for comparisons across studies as well as across treatment assignments.

A major focus of this experiment was to examine generalization patterns and, more specifically, to examine how task, complexity, and movement affected generalization. Therefore, three analyses of covariance (ANCOVAs) were completed. For each of these analyses, ES of the untrained structure of interest was the dependent measure, with the initial baseline monitor score of the untrained structure included as a covariate to account for participants' initial accuracy on the examined structure. These ANCOVAs examined the specific patterns of generalization comparing the factors outlined in Figure 1: (a) across task (across task, within complexity, within movement), (b) across complexity (within task, across complexity, within movement), and (c) across movement (within task, within complexity, across movement). Post hoc one-sample comparisons were examined for all adjusted means from the three ANCOVAs to determine which comparisons resulted in significant generalization. All analyses were completed in Statistical Analysis Software (SAS Institute Inc., Cary, NC) and Statistica software (StataCorp, College Station, TX).

Results

Of the 48 participants, 25 were enrolled in OM treatment and 23 were enrolled in SPM treatment (see Figure 2 for each participant's treatment assignment). A *t* test was conducted to ensure that the severity of deficits was balanced across participants enrolled in OM and SPM

treatments.³ A two-tailed independent-samples *t* test comparing the overall sentence comprehension severity levels of participants receiving OM treatment to those of participants receiving SPM treatment was found to be nonsignificant, $t(46) = 1.70$, $p = .22$, indicating comparable initial levels of sentence comprehension between the two groups. Another two-tailed independent-samples *t* test compared WAB-R Aphasia Quotient scores of participants receiving OM treatment to those of participants receiving SPM treatment, which was also nonsignificant, $t(45) = 1.10$, $p = .83$, indicating comparable severity of aphasia between the two groups.

Because the treatment protocol was altered midway in the study to include canonical sentences as part of the protocol, a two-tailed independent-samples *t* test was conducted to compare results of participants enrolled before and after this modification was implemented. The comparison of trained structure ES between these two groups was found to be nonsignificant, $t(46) = 1.64$, $p = .27$. Therefore, results were collapsed across participants for the remainder of analyses.

Effects of Therapy

Seven linear mixed-effects models evaluating improvement on trained structures relative to untrained structures using the log odds of accuracy on pre- and posttreatment monitor scores were performed for each contrast described below. The first mixed model demonstrated that participants trained on the OM task were more likely to improve on their trained structure relative to the OM control structures⁴ by a factor of 6.42 ($p < .01$; see Table 3). Likewise, the second mixed model demonstrated that participants trained on the SPM task were more likely to improve on

³One participant, P29, did not complete an OM pretreatment screener due to time constraints.

⁴When comparing trained structures to control structures, the average of the two within-task control structures (3NP and ORCNP) were compared to the average of the four different trained sentence structures.

Table 2. Participant demographics by task of treatment assignment.

Treatment task		Age (years)	MPO	WAB-R AQ	Active screener average (%)	Noncanonical sentence screener average (%)
OM	Average	56.52	47.48	69.60	78.80	49.75
	Range	26–82	6–260	10.20–99.90	30.00–100.00	26.25–83.75
SPM	Average	63.00	47.91	68.25	75.87	44.89
	Range	29–86	6–278	16.70–97.40	40.00–100.00	27.50–80.00

Note. MPO = months postonset; WAB-R AQ = Western Aphasia Battery–Revised Aphasia Quotient; OM = object manipulation; SPM = sentence-to-picture matching.

their trained structure relative to SPM control structures by a factor of 1.93 ($p = .05$). In a comparison of the probability of OM trained structures improving relative to the probability of SPM trained structures improving, collapsed across structure, it was found that participants who were trained in the OM task were more likely to improve on their trained structure relative to the participants who were trained in the SPM task by a factor of 3.32 ($p < .05$). Analyses conducted across each possible treatment structure (OR, OC, PA, UNACC), collapsing across task, showed that all sentence types except for UNACC led to significantly greater log odds probability of improvement on trained structures relative to control structure, with OR showing the greatest improvement and OC showing the least improvement (OR odds = 7.01, $p < .01$; OC odds = 2.51, $p = .05$; PA odds = 3.64, $p = .01$; UNACC odds = 3.06, $p = .11$).

To evaluate treatment results using ES (the outcome measure used in many treatment studies and included in generalization analyses below), an independent-samples t test was conducted that compared ES on trained structures for OM participants and SPM participants (see Figure 2 for individual ES results). Consistent with results from linear mixed-effects models, results were significant, $t(46) = 6.01$, $p < .01$. The mean ES for participants receiving OM treatment was 5.07, which was greater relative to that for participants receiving SPM treatment (2.14).

Generalization

A series of ANCOVAs were performed to examine generalization patterns and specifically how task, complexity, and movement affected generalization. See Figure 3

Figure 2. Effect size (ES) for trained structures. ^aParticipant did not receive the treatment modification. OM = object manipulation; OR = object relative; OC = object cleft; PA = passive; UNACC = unaccusative; SPM = sentence-to-picture matching.

Table 3. Mixed-model analysis results.

Comparison	df	Odds ratio	SE	p value	CI
OM treatment vs. control	4593	6.42**	1.79	<.01	2.67–15.42
SPM treatment vs. control	4593	1.93*	1.36	.05	0.99–3.78
OM vs. SPM treatment	4593	3.32*	1.04	.02	1.20–9.18
OR trained vs. control	4593	7.01**	1.75	<.01	2.97–16.53
OC trained vs. control	4593	2.51*	1.89	.05	1.00–6.33
PA trained vs. control	4593	3.64*	1.99	.01	1.37–9.65
UNACC trained vs. control	4593	3.06	2.79	.11	0.78–12.03

Note. CI = confidence interval; OM = object manipulation; SPM = sentence-to-picture matching; OR = object relative; OC = object cleft; PA = passive; UNACC = unaccusative.

* $p < .05$. ** $p < .01$.

for individual ES results presented within the framework of generalization analyses.

Across-Task, Within-Complexity, Within-Movement Generalization

In the first ANCOVA, we examined the effects of three parameters of treatment assignment (task treated, complexity of the trained structure, and movement of the trained structure) on across-task patterns of generalization. Therefore, the dependent variable for this analysis

was each participant's ES on their trained sentence on monitors of the opposite task. The initial baseline score in the untrained task served as the covariate. For example, if a participant was trained on OR sentences in the OM task, the dependent variable would be the participant's ES on OR sentences in the SPM task. The covariate would be the initial baseline score on OR sentences in the SPM task. Results showed a significant main effect of complexity, $F(1, 39) = 5.14$, $p = .03$, where participants who were trained on complex structures showed a greater adjusted average ES

Figure 3. Effect size (ES) for structures examined in generalization for participants trained on each sentence structure: (a) participants trained on OM OR and SPM OR, (b) participants trained on OM OC and SPM OC, (c) participants trained on OM PA and SPM PA, and (d) participants trained on OM UNACC and SPM UNACC. OM = object manipulation; OR = object relative; SPM = sentence-to-picture matching; OC = object cleft; PA = passive; UNACC = unaccusative; NP = noun phrase. ^aParticipant did not receive the treatment modification. This figure represents raw ES and does not account for the covariates.

(adjusted for the effect of the covariate) on their trained structure in the opposite task (3.60) compared with participants who were trained on simple structures (1.31; see Figure 4). In other words, participants trained on OR and PA sentences showed greater generalization across task than participants trained on OC and UNACC sentences—the relatively less complex sentence structures. There was no significant main effect of task between participants receiving OM treatment and participants receiving SPM treatment. Likewise, there was no significant main effect of movement between participants trained on NP movement sentences and participants trained on *wh*-movement sentences. There were no significant interactions.

The nonsignificant differences of generalization for the main effects of task and movement raise the question of whether generalization was even present. Thus, post hoc one-sample comparisons were conducted examining the adjusted ES on the main effect of task (OM or SPM) on the ANCOVA and showed that participants treated in the SPM task showed significant generalization to their trained structures in the OM task, $F(1, 23) = 8.19, p < .01$. In contrast, participants treated in the OM task showed no generalization effects to their trained structures in the SPM task, $F(1, 25) = 3.57, p = .07$. Next, post hoc tests further examined the main effect of complexity and showed that only participants trained on complex structures showed significant generalization on their trained structure in the opposite task, $F(1, 25) = 21.96, p < .01$. Likewise,

post hoc tests examined the main effect of movement and showed that only participants trained on *wh*-movement structures showed significant generalization effects on their trained structure in the opposite task, $F(1, 26) = 11.55, p < .01$.

Within-Task, Across-Complexity, Within-Movement Generalization

In the second ANCOVA, we examined the effects of three parameters of treatment assignment (task treated, complexity of the trained structure, and movement of the trained structure) on across-complexity patterns of generalization. The dependent variable for this analysis was each participant's ES in their trained task monitor on the sentence with the same movement but opposite relative complexity. For example, if a participant was trained on OR (complex *wh*-movement) sentences in the OM task, the dependent variable would be the participant's ES on OC (simple *wh*-movement) sentences in the same OM task. The covariate would be the initial baseline score on OC sentences in the OM task. Results showed a significant main effect of complexity, $F(1, 39) = 9.46, p < .01$, where participants who were trained on complex structures showed a greater adjusted average ES on the across-complexity structure (1.61) than participants who were trained on simple structures (-0.33 ; see Figure 5). In other words, the generalization from OR and PA sentences to structures with the same syntactic movement but lower relative complexity

Figure 4. Analysis of covariance results for across-task, within-complexity, within-movement structures, comparing adjusted average effect size (ES; adjusted for baseline performance on measured structure) by parameters of treatment assignment. For example, if a participant was trained on object relative sentences in the OM task, the dependent variable would be the participant's ES on object relative sentences in the SPM task. OM = object manipulation; SPM = sentence-to-picture matching; WH = *wh*-movement sentences; NP = noun phrase movement sentences. Asterisk indicate significant effects.

Figure 5. Analysis of covariance results for within-task, across-complexity, within-movement structures, comparing adjusted average effect size (ES; adjusted for baseline performance on measured structure) by parameters of treatment assignment. For example, if a participant was trained on object relative (complex *wh*-movement) sentences in the OM task, the dependent variable would be the participant's ES on object cleft (simple *wh*-movement) sentences in the same OM task. OM = object manipulation; SPM = sentence-to-picture matching; WH = *wh*-movement sentences; NP = noun phrase movement sentences. Asterisk indicate significant effects.

was greater than the generalization from OC and UNACC sentences to structures with the same syntactic movement but relatively higher complexity. There was no significant main effect of task between participants receiving OM treatment and participants receiving SPM treatment. Likewise, there was no significant main effect of movement between participants trained on NP movement sentences and participants trained on *wh*-movement sentences.

There was a significant interaction between the trained task and complexity on the across-complexity structures, $F(1, 39) = 4.77, p = .04$, where participants who were trained on simple sentences in the OM task showed a negative adjusted average ES (-1.16), whereas participants trained on all other combinations had a positive adjusted average ES (complex sentences in the OM task = 1.77 ; simple sentences in the SPM task = 0.44 ; complex sentences in the SPM task = 1.42). There were no other significant interactions.

Post hoc one-sample comparisons examining generalization in the main effect of task on the ANCOVA showed that participants treated in the SPM task showed significant generalization to their across-complexity structure, $F(1, 23) = 4.61, p = .04$. In contrast, the post hoc analysis revealed that participants treated in the OM task showed no generalization effects to their across-complexity structure, $F(1, 25) = 0.05, p = .82$. Next, post hoc tests further examined the main effect of complexity and showed that only participants trained on complex structures showed significant generalization to their across-complexity structure, $F(1, 25) = 11.59, p < .01$. Post hoc tests examining the main effect of movement showed that participants trained on *wh*-movement

sentences did not show significant generalization to their across-complexity structure. In addition, participants trained on NP movement sentences did not show significant generalization to their across-complexity structure.

The ANCOVA showed a significant interaction effect between task and complexity, where participants trained on simple sentences in the OM task showed a negative adjusted average ES and all other combinations showed a positive effect. Therefore, post hoc comparisons allowed the examination of which of these contrasts showed significant generalization. Three of the four contrasts (i.e., simple sentences in the OM task, complex sentences in the OM task, and complex sentences in the SPM task) showed significant generalization. Unique among these three, participants trained on simple structures in the OM task showed significant negative generalization to the more complex structures in the OM task, $F(1, 11) = 5.96, p = .02$. In other words, participants trained on UNACC and OC in the OM task had decreased ES on PA and OR sentences, respectively. However, participants trained on simple structures in the SPM task showed no significant generalization effect to the more complex structures in the SPM task, $F(1, 12) = 1.04, p = .31$. In addition, participants trained on complex structures in the OM task showed significant generalization on the more simple structures in the OM task, $F(1, 14) = 8.23, p < .01$, and participants trained on complex structures in the SPM task showed significant generalization to the more simple structures in the SPM task, $F(1, 11) = 4.20, p = .05$. Of note, there was a larger magnitude of generalization following OM treatment of complex structures than following SPM treatment of complex structures.

Within-Task, Within-Complexity, Across-Movement Generalization

In the third ANCOVA, we examined the effects of three parameters of treatment assignment (task treated, complexity of the trained structure, and movement of the trained structure) on across-movement patterns of generalization. The dependent variable for this analysis was each participant's ES in their trained task monitor on the sentence of the opposite movement but similar relative level of complexity. For example, if a participant was trained on OR (complex *wh*-movement) sentences in the OM task, the dependent variable would be the participant's ES on PA (complex NP movement) sentences in the OM task. The covariate would be the initial baseline score on PA sentences in the OM task. Results showed a significant main effect of complexity, $F(1, 39) = 7.81, p < .01$, where participants who were trained on complex structures showed a greater adjusted average ES on the structure of opposite movement and matched relative complexity (0.66) than participants trained on simple structures (-0.38 ; see Figure 6). In other words, participants trained on OR and PA sentences showed greater generalization across movement than participants trained on OC and UNACC sentences. There was no significant main effect of task between participants receiving OM treatment and participants receiving SPM treatment. There was no significant main effect of movement between participants trained on NP movement sentences and participants trained on *wh*-movement sentences.

There was a significant interaction between the trained task and movement on the across-movement sentences, $F(1, 39) = 7.16, p = .01$, where participants who were trained on NP movement sentences in the OM task had a negative adjusted average ES (-1.13), whereas participants trained on all other combinations had a positive adjusted average ES (*wh*-movement in the OM task = 0.51; NP movement in the SPM task = 0.44; *wh*-movement in the SPM task = 0.70). There were no other significant interactions.

Post hoc one-sample comparisons examining generalization in the main effect of task on the ANCOVA showed that participants treated in the SPM task showed significant generalization to their across-movement structure, $F(1, 23) = 5.09, p = .03$. In contrast, the post hoc analysis revealed that participants treated in the OM task showed no generalization effects to their across-movement structure, $F(1, 25) = 2.44, p = .13$. Next, post hoc tests examined the main effect of complexity and showed that only participants trained on complex structures showed significant generalization to their across-movement structure, $F(1, 25) = 6.39, p = .02$. Post hoc tests examining the main effect of movement showed that participants trained on *wh*-movement sentences did not show significant generalization to their across-movement structure. In addition, participants trained on NP movement sentences did not show significant generalization to their across-movement structure.

The ANCOVA showed a significant interaction effect between task and movement, where participants

Figure 6. Analysis of covariance results for within-task, within-complexity, across-movement structures, comparing adjusted average effect size (ES; adjusted for baseline performance on measured structure) by parameters of treatment assignment. For example, if a participant was trained on object relative (complex *wh*-movement) sentences in the OM task, the dependent variable would be the participant's ES on passive (complex NP movement) sentences in the OM task. OM = object manipulation; SPM = sentence-to-picture matching; WH = *wh*-movement sentences; NP = noun phrase movement sentences. Asterisk indicate significant effects.

trained on NP movement sentences in the OM task showed a negative adjusted average ES and all other combinations showed a positive effect. Therefore, post hoc comparisons allowed the examination of which of these contrasts showed significant generalization. Participants trained on NP movement structures in the OM task showed a significant negative generalization effect to their across-movement structure, $F(1, 11) = 7.26, p = .01$. The remaining three contrasts (*wh*-movement in the OM task, NP movement in the SPM task, and *wh*-movement in the SPM task) do not show significant generalization to their across-movement structure.

Discussion

The current study was designed to evaluate outcomes of two sentence comprehension therapies—OM and SPM—closely examining patterns of generalization following therapy. Overall, results show that participants improved on their respective trained tasks and structures following both treatments. Results provide support for the rationale that therapy targeted at improving thematic role mapping has a positive effect on sentence comprehension in PWA.

One goal of the study was to directly compare the SPM and OM therapies. Consistent with results from Kiran et al. (2012), these results demonstrate a greater magnitude of treatment effects following OM relative to SPM therapy. There may be important differences in the linguistic and/or cognitive processes targeted by each task that emerge when tasks are implemented in the context of treatment. Research has proposed that the enactment aspect of the OM treatment may lead to improved outcomes due to the relation between language, perceptual, and motor systems (see Kiran et al., 2012; Rizzolatti & Arbib, 1998). Kiran et al. (2012) additionally proposed that the enactment component of OM tasks makes them more demanding than SPM tasks, leading to deeper processing and greater improvements in therapy. The OM task requires formulating a plan for demonstrating comprehension (Hamburger & Crain, 1984; Shankweiler, Crain, Gorrell, & Tuller, 1989)—a task demand that interacts with online syntactic processing and is potentially consequential to progress in therapy (Caplan et al., 2006, 2007). In general, the results of this study are consistent with those of mapping therapy (Schwartz et al., 1994), which also focuses on comprehension. Both of the current treatment protocols focused on thematic role assignment of the verb, and the SPM treatment in particular was quite similar to mapping therapy. However, the enactment portion of the OM treatment was different from mapping therapy and perhaps placed more cognitive demands.

Another major goal of this study was to examine generalization to untrained sentence structures, which was performed using three ANCOVAs each examining the factors of task, complexity, and movement across three scenarios (Figure 1): (a) across task (e.g., generalization from OR in the OM task to OR in the SPM task), (b) across complexity (e.g., generalization from OR in the OM task

to OC in the OM task), and (c) across movement (e.g., generalization from OR in the OM task to PA in the OM task). First, in every analysis, a main effect of complexity was observed and no main effects of either task or movement were observed, underscoring the importance of complexity in generalization patterns. Across the ANCOVAs, more generalization was observed from relatively complex sentence structures (OR and PA) than from relatively simpler sentence structures (OC and UNACC). The pattern of greater instances of generalization from complex to simple structures than from simple to complex structures is consistent with prior treatment research examining the effects of complexity in syntactic (Thompson, 2001, 2006; Thompson et al., 1993, 1997, 1998, 2003) and semantic (Kiran, 2007, 2008; Kiran, Sandberg, & Sebastian, 2011; Kiran & Thompson, 2003) therapies. Therefore, results add to the body of work in support of the CATE, which proposes that treatment targeting complex structures enhances generalization to relatively less complex structures.

Second, although the main effect of task was not significant in any of the ANCOVAs, which indicated no differences between the tasks in generalization patterns, post hoc analyses examining task in all three ANCOVAs demonstrated that participants trained on the SPM task showed a significant generalization effect, whereas participants trained on the OM task did not show significant generalization. These results of greater generalization for SPM than OM were in contrast to the treatment effects where OM had greater effects than SPM. We return to this point later in the discussion.

Third, although the main effect of movement was not significant in any of the ANCOVAs, post hoc analyses examining movement in the first ANCOVA (which examined across-task generalization) showed that participants trained on *wh*-movement sentences showed significant generalization to their trained structure in the opposite task (i.e., OC in the OM task to OC in the SPM task, OC in the SPM task to OC in the OM task, OR in the OM task to OR in the SPM task, and OR in the SPM task to OR in the OM task), whereas participants trained on NP movement sentences did not show significant generalization to their trained structure in the opposite task. The main effect of movement was not significant in the remaining two ANCOVAs. These results indicate that the generalization across task may have different patterns than the generalization seen across structures. We also address this observation later in the discussion.

The nonsignificant main effects on the ANCOVAs, significant interactions on the ANCOVAs, and significant post hoc comparison results require further explanation. Recall that the post hoc results for the main effect of task in each of the ANCOVAs showed that training structures in the SPM task resulted in significant generalization, whereas training structures in the OM task did not. However, these analyses were collapsed across structures, and the interaction effect in ANCOVA across complexity and across movement are important in disentangling the effect of task and other factors. The significant interaction between task

and complexity in the second ANCOVA showed a clear and strong generalization effect of complex structures generalizing to simple structures (i.e., PA to UNACC and OR to OC) for participants receiving OM treatment. As noted, this effect was greater than for participants trained on complex structures receiving SPM treatment. In the same analysis, participants trained on simple structures in the OM task demonstrated a significant negative generalization to complex structures (i.e., UNACC to PA and OC to OR), whereas participants trained on simple structures in the SPM task showed no significant generalization. Therefore, it appears that the stronger positive generalization effect of SPM in the ANCOVA may have been amplified by the negative effects of training simple structures in the OM task. Taken together with the results of treatment, OM appears to be an effective treatment task in terms of both promoting treatment gains on a trained structure and promoting generalization from complex structures to more simple structures. It is possible that when the enactment component of OM treatment is limited to simple structures, it does not promote deep levels of processing and/or patients fall into maladaptive perceptual-motor patterns. In contrast, when trained on complex structures, the OM task led to generalization of simpler structures within task.

The third ANCOVA showed a significant interaction between task and movement such that participants trained on NP movement sentences in the OM task were significantly different than the other three contrasts. Post hoc analyses confirmed that only participants trained on NP movement sentences in the OM task showed a significant negative generalization, whereas the other three contrasts (*wh*-movement in the OM task, NP movement in the SPM task, and *wh*-movement in the SPM task) did not show significant generalization. Similar to the interaction results in the second ANCOVA, it appears that the negative effects of training NP movement sentences in the OM task may have overshadowed the other effects of OM generalization. Together, these interaction effect results partially explain why the SPM task appeared to show a greater generalization effect than the OM task when examining the main effects of task in the across-structure generalizations.

Post hoc comparisons also revealed an interesting effect of movement in generalization. Follow-up on the first ANCOVA, which examined across-task generalization, showed that training *wh*-movement sentences led to significant changes in *wh*-movement sentence performance in the opposite task, whereas training on NP movement sentences did not. Together with the interaction effect of task and movement in the third ANCOVA, these results bring to light a potentially important difference to consider in the relative structure of NP and *wh*-movement sentences. NP movement sentences targeted in this study (UNACC and PA) are relatively less complex than the *wh*-movement sentences (OC and OR) that comprise two clauses or two sets of thematic roles, respectively. Thus, the increased complexity of *wh*-movement sentences may have led to increased generalization. Note that OC sentences are simpler relative to OR sentences and that analyses of generalization across

complexity in the current study were constrained within movement for this reason.

Examined within the framework of models of sentence comprehension deficits in aphasia, generalization patterns are most consistent with resource reduction accounts. Recall that under specific deficit models, syntactic processes are thought to drive impairment. Therefore, treatment effects and generalization would be expected only to sentence structures with shared syntactic properties, which in this study translates to sentence structures with similar movement. The observed generalization across movement in the third ANCOVA, therefore, is not consistent with predictions of specific deficit models. In contrast, under resource reduction accounts, comprehension deficits are not circumscribed by the syntactic structure of sentences but instead are considered to be related to general deficits in processing resources, such as verbal working memory, phonological store, and short-term semantic memory, which are relevant across multiple syntactic structures. The generalization observed in the current study from complex *wh*-movement sentences to complex NP movement sentences (i.e., OR in the OM task to PA in the OM task and OR in the SPM task to PA in the SPM task) is consistent with patterns of generalization predicted to extend to a variety of sentence structures that require fewer resources to process than trained sentences.

The across-movement generalization observed in the current study has not always been found in other treatment studies. In particular, as noted in the introduction, studies of the TUF approach (Ballard & Thompson, 1999; Jacobs & Thompson, 2000; Thompson & Shapiro, 2005; Thompson et al., 1993, 1996, 1997) frequently observed greater generalization to sentences with the same type of movement than to sentences with a different type of movement. Furthermore, it is important to highlight that the focus of the treatments in the current study was on the identification of thematic role in comprehension, which addresses movement only implicitly. Specifically training movement within a syntactic structure might influence the generalization differently than if movement was alluded to implicitly, as was done in the current study's protocol. Therefore, once the individual successfully learns to identify the thematic roles in their trained structure, that mechanism may be applied across sentences that include NP and *wh*-movement. Further, studies that have demonstrated strong within-structure generalization have generally been production treatments (i.e., TUF), whereas the current study focused only on comprehension. The specific nature, demands, and context of therapy are likely significant contributors to generalization and hence may lead to different patterns of generalization.

Overall, these findings provide further evidence of the efficacy of thematic role training on improving sentence comprehension for PWA. Results highlight that OM and SPM treatments are both effective but may function differently; the exact nature of this difference is not fully understood. Although the OM treatment was found to lead to greater treatment gains and to more generalization from complex structures to simple structures, the SPM treatment led to more consistent patterns of generalization. As discussed,

these findings provide support for resource reduction models of sentence comprehension deficits and highlight the importance of considering principles of therapy and levels of processing when interpreting results. Impairment factors and linguistic variables, though out of the scope of the current study, likely each contribute to generalization and merit further analysis.

Although this study examines a substantial number of participants, future studies should continue to systematically examine generalization patterns within theoretically motivated frameworks. Generalization and treatment outcomes should also be examined within a broader context of additional outcome measures, tasks, and contexts. It will also be important to examine the relationship between treatment outcomes and factors such as age, age at onset of injury, severity of aphasia, executive function skills, attention, memory, and learning ability, particularly at the individual participant level.

Acknowledgments

This research was supported by National Institute on Deafness and Other Communication Disorders Grant 4R33DC010461. The authors thank the participants and their families and caregivers. They also thank Gloria Waters and Myrna Schwartz for their contributions and Chaleece Sandberg, Elsa Ascenso, Jason Lucas, Nadia Lonsdale, and Balaji Rangarathnam for their assistance in data collection. In addition, the authors thank Lauren Eiges, Melody Lo, Nicole Woody, Rasika Murali, Sophie Barry, and Zach Smith for their assistance at various stages of the project.

References

- Ballard, K. J., & Thompson, C. K. (1999). Treatment and generalization of complex sentence production in agrammatism. *Journal of Speech, Language, and Hearing Research, 42*, 690–707. doi:10.1044/jslhr.4203.690
- Beeson, P. M., & Robey, R. R. (2006). Evaluating single-subject treatment research: Lessons learned from the aphasia literature. *Neuropsychology Review, 16*, 161–169. doi:10.1007/s11065-006-9013-7
- Byng, S. (1988). Sentence processing deficits: Theory and therapy. *Cognitive Neuropsychology, 5*, 629–676. doi:10.1080/02643298808253277
- Byng, S., Nickels, L., & Black, M. (1994). Replicating therapy for mapping deficits in agrammatism: Remapping the deficit? *Aphasiology, 8*, 315–341. doi:10.1080/02687039408248663
- Caplan, D., Baker, C., & Dehaut, F. (1985). Syntactic determinants of sentence comprehension in aphasia. *Cognition, 21*, 117–175. doi:10.1016/0010-0277(85)90048-4
- Caplan, D., DeDe, G., & Michaud, J. (2006). Task-independent and task-specific syntactic deficits in aphasic comprehension. *Aphasiology, 20*, 893–920. doi:10.1080/02687030600739273
- Caplan, D., & Waters, G. S. (2006). Comprehension disorders in aphasia: The case of sentences that require syntactic analysis. In M. Traxler (Ed.), *The handbook of psycholinguistics* (pp. 939–967). London, United Kingdom: OUP.
- Caplan, D., Waters, G. S., DeDe, G., Michaud, J., & Reddy, A. (2007). A study of syntactic processing in aphasia I: Behavioral (psycholinguistic) aspects. *Brain and Language, 101*, 103–150. doi:10.1016/j.bandl.2006.06.225
- Caramazza, A., & Zurif, E. B. (1976). Dissociation of algorithmic and heuristic processes in language comprehension: Evidence from aphasia. *Brain and Language, 3*, 572–582. doi:10.1016/0093-934X(76)90048-1
- Chomsky, N. (1995). *The minimalist program* (Vol. 1765). Cambridge, MA: MIT Press.
- DeDe, G., & Caplan, D. (2006). Factor analysis of aphasic syntactic comprehension disorders. *Aphasiology, 20*, 123–135. doi:10.1080/02687030500472413
- Fink, R. B. (2001). Mapping treatment: An approach to treating sentence level impairments in agrammatism. *SIG 2 Perspectives on Neurophysiology and Neurogenic Speech and Language Disorders, 11*, 14–23. doi:10.1044/nnsld11.3.14
- Gauthier, L., Dehaut, F., & Joanette, Y. (1989). The Bells test: A quantitative and qualitative test for visual neglect. *International Journal of Clinical Neuropsychology, 11*, 49–54.
- Grodzinsky, Y. (1986). Language deficits and the theory of syntax. *Brain and Language, 27*, 135–159. doi:10.1016/0093-934X(86)90009-X
- Grodzinsky, Y. (1995a). A restrictive theory of agrammatic comprehension. *Brain and Language, 50*, 27–51. doi:10.1006/brln.1995.1039
- Grodzinsky, Y. (1995b). Trace-deletion, theta-roles, and cognitive strategies. *Brain and Language, 51*, 469–497. doi:10.1006/brln.1995.1072
- Grodzinsky, Y. (2000). The neurology of syntax: Language use without Broca's area. *Behavioral and Brain Sciences, 23*, 1–21. doi:10.1017/S0140525X00002399
- Grodzinsky, Y., Piñango, M. M., Zurif, E., & Drai, D. (1999). The critical role of group studies in neuropsychology: Comprehension regularities in Broca's aphasia. *Brain and Language, 67*, 134–147. doi:10.1006/brln.1999.2050
- Haarmann, H. J., Just, M. A., & Carpenter, P. A. (1997). Aphasic sentence comprehension as a resource deficit: A computational approach. *Brain and Language, 59*, 76–120. doi:10.1006/brln.1997.1814
- Haendiges, A. N., Berndt, R. S., & Mitchum, C. C. (1996). Assessing the elements contributing to a “mapping” deficit: A targeted treatment study. *Brain and Language, 52*, 276–302. doi:10.1006/brln.1996.0011
- Hamburger, H., & Crain, S. (1984). Acquisition of cognitive compiling. *Cognition, 17*, 85–136. doi:10.1016/0010-0277(84)90015-5
- Helm-Estabrooks, N. (2001). *Cognitive Linguistic Quick Test: Examiner's manual*. San Antonio, TX: The Psychological Corporation.
- Jacobs, B. J., & Thompson, C. K. (2000). Cross-modal generalization effects of training noncanonical sentence comprehension and production in agrammatic aphasia. *Journal of Speech, Language, and Hearing Research, 43*, 5–20. doi:10.1044/jslhr.4301.05
- Jones, E. V. (1986). Building the foundations for sentence production in a non-fluent aphasic. *International Journal of Language & Communication Disorders, 21*, 63–82. doi:10.3109/13682828609018544
- Kertesz, A. (2007). *Western Aphasia Battery—Revised*. San Antonio, TX: The Psychological Corporation.
- Kiran, S. (2007). Complexity in the treatment of naming deficits. *American Journal of Speech-Language Pathology, 16*, 18–29. doi:10.1044/1058-0360(2007/004)
- Kiran, S. (2008). Typicality of inanimate category exemplars in aphasia treatment: Further evidence for semantic complexity. *Journal of Speech, Language, and Hearing Research, 51*, 1550–1568. doi:10.1044/1092-4388(2008/07-0038)

- Kiran, S., Caplan, D., Sandberg, C., Levy, J., Berardino, A., Ascenso, E., . . . Tripodis, Y. (2012). Development of a theoretically based treatment for sentence comprehension deficits in individuals with aphasia. *American Journal of Speech-Language Pathology*, 21, S88–S102. doi:10.1044/1058-0360(2012/11-0106)
- Kiran, S., Sandberg, C., & Sebastian, R. (2011). Treatment of category generation and retrieval in aphasia: Effect of typicality of category items. *Journal of Speech, Language, and Hearing Research*, 54, 1101–1117. doi:10.1044/1092-4388(2010/10-0117)
- Kiran, S., & Thompson, C. K. (2003). The role of semantic complexity in treatment of naming deficits: Training semantic categories in fluent aphasia by controlling exemplar typicality. *Journal of Speech, Language, and Hearing Research*, 46, 608–622. doi:10.1044/1092-4388(2003/048)
- Le Dorze, G., Jacob, A., & Coderre, L. (1991). Aphasia rehabilitation with a case of agrammatism: A partial replication. *Aphasiology*, 5, 63–85. doi:10.1080/02687039108248520
- Linebarger, M. C. (1995). Agrammatism as evidence about grammar. *Brain and Language*, 50, 52–91. doi:10.1006/brln.1995.1040
- Linebarger, M. C., Schwartz, M. F., & Saffran, E. M. (1983). Sensitivity to grammatical structure in so-called agrammatic aphasics. *Cognition*, 13, 361–392. doi:10.1016/0010-0277(83)90015-X
- Marshall, J., Pring, T., & Chiat, S. (1993). Sentence processing therapy: Working at the level of the event. *Aphasiology*, 7, 177–199. doi:10.1080/02687039308249505
- Mitchum, C. C., Haendiges, A. N., & Berndt, R. S. (1995). Treatment of thematic mapping in sentence comprehension: Implications for normal processing. *Cognitive Neuropsychology*, 12, 503–547. doi:10.1080/02643299508252006
- Nickels, L., Byng, S., & Black, M. (1991). Sentence processing deficits: A replication of therapy. *International Journal of Language & Communication Disorders*, 26, 175–199. doi:10.3109/13682829109012002
- Rochon, E., Laird, L., Bose, A., & Scofield, J. (2005). Mapping therapy for sentence production impairments in nonfluent aphasia. *Neuropsychological Rehabilitation*, 15, 1–36. doi:10.1080/09602010343000327
- Rizzolatti, G., & Arbib, M. A. (1998). Language within our grasp. *Trends in Neurosciences*, 21, 188–194. doi:10.1016/S0166-2236(98)01260-0
- Schneider, W., & Zuccolotto, A. (2002). *E-Prime user's guide*. Pittsburgh, PA: Psychology Software Tools Inc.
- Schwartz, M. F., Linebarger, M. C., Saffran, E. M., & Pate, D. S. (1987). Syntactic transparency and sentence interpretation in aphasia. *Language and Cognitive Processes*, 2, 85–113. doi:10.1080/01690968708406352
- Schwartz, M. F., Saffran, E. M., Fink, R. B., Myers, J. L., & Martin, N. (1994). Mapping therapy: A treatment programme for agrammatism. *Aphasiology*, 8, 19–54. doi:10.1080/02687039408248639
- Shankweiler, D., Crain, S., Gorrell, P., & Tuller, B. (1989). Reception of language in Broca's aphasia. *Language and Cognitive Processes*, 4, 1–33. doi:10.1080/01690968908406355
- Thompson, C. K. (2001). Treatment of underlying forms: A linguistic specific approach for sentence production deficits in agrammatic aphasia. In R. Chapey (Ed.), *Language intervention strategies in adult aphasia* (4th ed., pp. 605–628). Baltimore, MD: Williams & Wilkins.
- Thompson, C. K. (2006). Single subject controlled experiments in aphasia: The science and the state of the science. *Journal of Communication Disorders*, 39, 266–291. doi:10.1016/j.jcomdis.2006.02.003
- Thompson, C. K. (2011). *Northwestern Assessment of Verbs and Sentences*. Unpublished manuscript.
- Thompson, C. K., Ballard, K. J., & Shapiro, L. P. (1998). The role of syntactic complexity in training wh-movement structures in agrammatic aphasia: Optimal order for promoting generalization. *Journal of the International Neuropsychological Society*, 4, 661–674.
- Thompson, C. K., & Shapiro, L. P. (1995). Training sentence production in agrammatism: Implications for normal and disordered language. *Brain and Language*, 50, 201–224. doi:10.1006/brln.1995.1045
- Thompson, C. K., & Shapiro, L. P. (2005). Treating agrammatic aphasia within a linguistic framework: Treatment of underlying forms. *Aphasiology*, 19, 1021–1036. doi:10.1080/02687030544000227
- Thompson, C. K., & Shapiro, L. P. (2007). Complexity in treatment of syntactic deficits. *American Journal of Speech-Language Pathology*, 16, 30–42. doi:10.1044/1058-0360(2007/005)
- Thompson, C. K., Shapiro, L. P., Ballard, K. J., Jacobs, B. J., Schneider, S. S., & Tait, M. E. (1997). Training and generalized production of wh- and NP-movement structures in agrammatic aphasia. *Journal of Speech, Language, and Hearing Research*, 40, 228–244. doi:10.1044/jslhr.4002.228
- Thompson, C. K., Shapiro, L. P., Kiran, S., & Sobecks, J. (2003). The role of syntactic complexity in treatment of sentence deficits in agrammatic aphasia: The complexity account of treatment efficacy (CATE). *Journal of Speech, Language, and Hearing Research*, 46, 591–607. doi:10.1044/1092-4388(2003/047)
- Thompson, C. K., Shapiro, L. P., & Roberts, M. M. (1993). Treatment of sentence production deficits in aphasia: A linguistic-specific approach to wh-interrogative training and generalization. *Aphasiology*, 7, 111–133. doi:10.1080/02687039308249501
- Thompson, C. K., Shapiro, L. P., Tait, M. E., Jacobs, B. J., & Schneider, S. L. (1996). Training Wh-question production in agrammatic aphasia: Analysis of argument and adjunct movement. *Brain and Language*, 52, 175–228. doi:10.1006/brln.1996.0009
- Waters, G. S., & Caplan, D. (1995, September). *The Psycholinguistic Assessment of Language Battery: Preliminary data*. Paper presented at the meeting of the British Aphasiology Society, York, England.
- Wechsler, D. (1997). *WAIS-III administration and scoring manual*. San Antonio, TX: The Psychological Corporation.