Published in final edited form as:

Drug Alcohol Depend. 2016 December 01; 169: 199–200. doi:10.1016/j.drugalcdep.2016.10.034.

Erratum to “Intra-individual changes in Stroop-related activations linked to cigarette abstinence in adolescent tobacco smokers: Preliminary findings” [*Drug Alcohol Depend.* 167 (2016) 182–189]

Sarah W. Yip^{a,b,*}, Iris M. Balodis^{b,c}, Kathleen M. Carroll^b, Suchitra Krishnan-Sarin^{b,1}, and Marc N. Potenza^{a,b,d,e,1}

^aThe National Center on Addiction and Substance Abuse, Department of Psychiatry, Yale University School of Medicine, New Haven, CT, USA

^bDivision of Substance Abuse, Department of Psychiatry, Yale University School of Medicine, New Haven, CT, USA

^cDepartment of Psychiatry and Behavioural Neurosciences, Peter Borris Centre for Addictions Research, McMaster University, Hamilton, ON, Canada

^dDepartment of Neurobiology, Yale University School of Medicine, New Haven, CT, USA

^eYale Child Study Center, Yale University School of Medicine, New Haven, CT, USA

The publisher regrets that Figs. 1 and 2 (A and B) of this article were incorrectly printed in black and white. The color figures are reproduced below for reference.

The publisher would like to apologize for any inconvenience caused.

*Corresponding author at: 1 Church Street, 7th Floor, Suite 731, New Haven, CT 06510-3330, USA. sarah.yip@yale.edu (S.W. Yip).

¹Denotes that senior authorship is shared by Drs. Potenza and Krishnan-Sarin.

Fig. 1. Main effects of Stroop task performance (incongruent versus congruent trials) on neural responses among adolescent smokers at pre-and post treatment. Whole-brain findings, cluster-level corrected at $pFWE < 0.05$ (voxel-level- $p < 0.001$). L = left; R = right.

A. Association between change in anterior cingulate activity and percent days abstinent during smoking cessation treatment

B. Association between change in insula activity and maximum durations of consecutive abstinence during smoking cessation treatment

Fig. 2.

Associations between changes in Stroop-related BOLD response (post-treatment versus baseline) and abstinence during smoking cessation treatment. Left: Individual clusters from uncorrected ($p < 0.01$) whole-brain analyses are shown for illustrative purposes. Right: Extracted signal from peak activations identified in small-volume-correction (SVC) regression analyses are plotted (10 mm sphere) to illustrate associations with abstinence measures. L = left; R = right; μ = BOLD signal change.