In brief

Italy to ease smoking

restrictions: The newly appointed Italian health minister, Francesco Storace, has declared that he will not continue what he described as his predecessor's "excessive" measures in favour of healthier lifestyles, in particular against smoking. He is a former smoker and is planning to make it easier for bars and restaurants to have smoking areas.

Eighteen new polio cases in

Yemen: The recently developed monovalent oral polio vaccine type 1 (mOPV1) will be used to halt an outbreak of polio in Yemen, now totalling 22 cases. It enables a precisely tailored immunological response to the type 1 poliovirus involved. Yemen had been polio-free since disease surveillance began in 1996.

Asian influenza samples

destroyed: 99.6% of the samples of the 1957 H2N2 Asian flu strain inadvertently sent to more than 3000 laboratories worldwide have been recovered and destroyed or are scheduled for destruction, according to the World Health Organization and US officials. The US Centers for Disease Control has formed a committee to investigate how the incident occurred.

Cannabis users have 10-fold increased risk of car crashes:

A New Zealand survey of 571 drivers of cars involved in crashes, in which at least one occupant was admitted to hospital or killed, showed those who habitually used cannabis were 9.5 times more likely to be involved in crashes than non-users (*Addiction* 2005;100:605-11).

Heart disease burden shifting to developing world:

An analysis of cardiovascular disease risk factors and economic indicators for more than 100 countries found that body mass index and cholesterol increased rapidly with national income before flattening and declining. Body mass index and cholesterol also increased with the proportion of people living in cities, indicating that developing countries were facing a major increase in cardiovascular disease (*PLoS Med* 2005;2:e133).

Aggressive treatment of low grade prostate cancer is unnecessary

Susan Mayor *London*

Aggressive treatment of localised low grade prostate cancer is inappropriate, a study has found. The long term study shows a very small risk of progression over 20 years of follow-up in patients treated conservatively (*JAMA* 2005;293:2095-101).

Peter Albertsen and colleagues analysed data from the Connecticut tumour registry, supplemented by hospital record and histology review, for 767 men aged 55 to 74 years diagnosed as having clinically localised prostate cancer between 1 January 1971 and 31 December 1984. All patients were treated conservatively with either observation or immediate or delayed androgen withdrawal therapy, with an average observation period of 24 years.

The results showed that

men with low grade prostate cancer (a Gleason score of 2-4) had a small risk of prostate cancer progression, even after 20 years of management by observation or androgen withdrawal therapy alone. Their risk of dying from prostate cancer was only 6 (95% confidence interval 2 to 11) deaths per 1000 person years.

In contrast, men with poorly differentiated disease (Gleason scores of 7-10) had a high risk of death from prostate cancer within 10 years of diagnosis—121 (90 to 156) deaths per 1000 person years—only three men in this group were alive after 20 years.

Men with moderate grade disease (Gleason scores of 5-6) had an intermediate cumulative risk of prostate cancer progression after 20 years of follow-up.

Overall, prostate cancer mortality in the first 15 years of follow-up was 33 (28 to 38) per 1000 person years and after 15 years of follow-up was 18 (10-29) per 1000 person years. The mortalities for these two follow-up periods were not statistically different after adjusting for differences in tumour histology (ratio 1.1; 95% confidence interval 0.6 to 1.9).

Dr Albertsen, professor and chief of urology at the University of Connecticut Health Center, Farmington, United States, said, "Extended follow-up of our competing risk analysis suggests that prostate cancer progression rates do not increase after 15 years of follow-up.

"Men with low grade prostate cancer have only a small risk of prostate cancer progression... These results do not support aggressive treatment of localised low grade prostate cancer."

He said that, in fact, the data had probably overestimated progression in men treated by observation or androgen withdrawal therapy alone. □

High cholesterol may increase risk of testicular cancer

Roger Dobson *Abergavenny*

High cholesterol concentrations may increase the risk of testicular cancer, a study has shown. The study investigated the relationship between risk of testicular cancer and blood cholesterol. Researchers found that men with the highest cholesterol concentrations had a cancer risk 4.5 times that for the men with the lowest concentrations.

"The highly significant positive association between serum cholesterol and testicular cancer risk found in this population-based cohort study suggests that an elevated concentration of serum cholesterol is a risk factor for testicular cancer," says a report of the study in the *British Journal of Cancer* that was published online (ahead of print publication) on 12 April (www.bjccancer.com, doi: 10.1038/sj.bjc.6602539).

Ann-Britt Wiréhn and colleagues from Karolinska University Hospital and Linköping

Testicular cancer, which has struck cyclist Lance Armstrong (above), may be connected with high cholesterol concentrations

University, Sweden, used data from a 25 year follow-up study of 44 864 men recruited between 1963 and 1965 as part of a mass screening health trial. Among the measures taken at the time was non-fasting serum cholesterol.

The data were matched with the Swedish cancer registry, the Swedish cause of death registry, and records of cancer registra-

tion and death between 1958 and 1987. The national registration system in Sweden using personal code numbers for all residents makes it possible to link registries. Men with any reported cancer before the start of the survey were excluded.

Among the 44 864 men were 21 cases of testicular cancer in the follow-up period. A positive correlation between serum cholesterol concentration and incidence of testicular cancer was found and the estimated hazard ratios calculated for three ranges of cholesterol concentration—below 5.7, 5.7-6.9, and greater than 7.0 mmol/l. The hazard ratios for the middle and highest serum cholesterol categories compared with the lowest were 1.3 (95% confidence interval 0.3 to 5.1) and 4.5 (1.3 to 16.2).

The testes are the commonest site of cancer in men aged 15-44 in Sweden as well as in many other countries. The incidence has increased in recent decades, and a number of possible risk factors have been suggested, including non-descended testes at birth and fat intake.

"To our knowledge, the relation between testicular cancer risk and serum cholesterol has not been evaluated," say the authors. □