Editors' Note: In response to "Incident parkinsonism in older adults without Parkinson disease," Mahlkecht et al. share their own experience of analyzing risk factors for parkinsonism in a separate population. They confirm and expand the study findings, providing additional single markers for prodromal Parkinson disease (PD). Commenting on the same article, Drs. Bohnen and Muller note that factors that contribute to parkinsonism in persons without PD are also likely to contribute to parkinsonism in persons with PD and that physicians need to take age-related pathology and medical comorbidities into consideration when treating gait and balance impairments in PD. Buchman et al., authors of the study, agree with both comments, stress the long subclinical phase in patients with PD, and ask physicians to consider coexisting conditions that may cause parkinsonism.

—Chafic Karam, MD, and Robert C. Griggs, MD

LETTER RE: INCIDENT PARKINSONISM IN OLDER ADULTS WITHOUT PARKINSON DISEASE

Philipp Mahlkecht, Stefan Kiechl, Johann Willeit, Werner Poewe, Klaus Seppi, Innsbruck, Austria:

We read with interest the article by Buchman et al.,¹ which reported that depressive symptoms, neuroticism, urinary incontinence, sleep complaints, and chronic health conditions were associated with incident parkinsonism (48.2% over 5 years) in the elderly. We previously reported on incidence and risk factors of mild parkinsonian signs, a similar concept to possible parkinsonism,¹ in the cohort of the prospective population-based Bruneck Study of 2005.² We now present an analysis of risk factors for parkinsonism according to the definition used by Buchman et al. in this population. Of 393 participants without parkinsonism at baseline undergoing 5-year follow-up, a markedly lower proportion of 16.0% (95% confidence interval [CI] 12.7%–20.0%) developed incident parkinsonism (excluding participants developing secondary parkinsonism). A logistic regression analysis adjusted for age and sex revealed that, out of single markers for prodromal Parkinson disease as defined per movement disorder society research criteria,³ assessed in the Bruneck study,⁴ substantia nigra hyperechogenicity on transcranial sonography (odds ratio 1.99; 95% CI

1.10–3.62, $p = 0.024$), probable REM sleep behavior disorder (6.88; 95% CI 1.10–43.10, $p = 0.039$), olfactory loss (1.96; 95% CI 1.01–3.81, $p = 0.047$), urinary dysfunction (5.15; 95% CI 1.65–16.14, $p = 0.005$), and depression/anxiety (2.85; 95% CI 1.28–6.36, $p = 0.011$) were significantly associated with incident parkinsonism, thus confirming and expanding the findings of Buchman et al.

1. Buchman AS, Leurgans SE, Yu L, et al. Incident parkinsonism in older adults without Parkinson disease. *Neurology* 2016;87:1036–1044.
2. Mahlkecht P, Kiechl S, Stockner H, et al. Predictors for incident mild parkinsonian signs: a prospective population-based study. *Parkinsonism Relat Disord* 2015;21:321–324.
3. Berg D, Postuma RB, Adler CH, et al. MDS research criteria for prodromal Parkinson's disease. *Mov Disord* 2015;30:1600–1611.
4. Mahlkecht P, Gasperi A, Willeit P, et al. Prodromal Parkinson's disease as defined per MDS research criteria in the general elderly community. *Mov Disord* 2016;31:1405–1408.

© 2017 American Academy of Neurology

AUTHOR RESPONSE: INCIDENT PARKINSONISM IN OLDER ADULTS WITHOUT PARKINSON DISEASE

Aron S. Buchman, David A. Bennett, Chicago:

We thank Mahlkecht et al. for the interest in our study, which reported on risk factors for incident parkinsonism in community-dwelling older adults without Parkinson disease (PD).¹ Mahlkecht et al. confirmed and extended some of our findings, growing the literature that demonstrates that PD, like other chronic diseases of aging, has a long subclinical phase in which brain pathology develops without clinical signs or symptoms, followed by mild signs and symptoms of insufficient severity to warrant a diagnosis of PD, with PD representing a late manifestation.² Overall, this conceptual model is similar to the one currently used for Alzheimer disease (AD).³ However, parkinsonian signs are much more common than subclinical PD pathology and are associated with diverse brain pathologies.^{4,5} Thus, like AD dementia, the clinical phenotype of parkinsonism in older adults may be caused by diverse etiologies. A means for identifying the subset of persons with parkinsonian signs who have PD pathology and are therefore likely

to be at a higher risk for progressing to PD is urgently needed.

1. Buchman AS, Leurgans SE, Yu L, et al. Incident parkinsonism in older adults without Parkinson disease. *Neurology* 2016;87:1036–1044.
2. Berg D, Postuma RB, Bloem B, et al. Time to redefine PD? Introductory statement of the MDS Task Force on the definition of Parkinson's disease. *Mov Disord* 2014;29:454–462.
3. Sperling RA, Aisen PS, Beckett LA, et al. Toward defining the preclinical stages of Alzheimer's disease: recommendations from the National Institute on Aging–Alzheimer's Association workgroups on diagnostic guidelines for Alzheimer's disease. *Alzheimers Dement* 2011;7:280–292.
4. Buchman AS, Shulman JM, Nag S, et al. Nigral pathology and parkinsonian signs in elders without Parkinson disease. *Ann Neurol* 2012;71:258–266.
5. Buchman AS, Wilson RS, Shulman JM, et al. Parkinsonism in older adults and its association with adverse health outcomes and neuropathology. *J Gerontol A Biol Sci Med Sci* 2016;71:549–556.

© 2017 American Academy of Neurology

LETTER RE: INCIDENT PARKINSONISM IN OLDER ADULTS WITHOUT PARKINSON DISEASE

Nicolaas I. Bohnen, Martijn Muller, Ann Arbor, MI:

We read with interest the article by Buchman et al.,¹ which reported on the frequency of parkinsonian signs over a 5-year period in older adults without Parkinson disease (PD). The authors showed that 48.2% of older adults who had no or only 1 parkinsonian sign at baseline had 2 or more parkinsonian signs at 5-year follow-up. The observed increase in gait impairment increasing from 19.9% at baseline to 87.4% at follow-up was particularly intriguing. Although the development of parkinsonian signs over time in older adults may signify the effect of age-associated nigrostriatal dopaminergic losses,² we agree with Buchman et al. that gait disturbances in the elderly, in particular, may also be explained by vascular etiology, frailty, medical comorbidities, side effects of medications, or other systemic factors in the absence of prototypical Lewy body parkinsonism.³ The findings of this study are also relevant for patients with PD, as age-related pathophysiology and medical comorbidities may contribute additively to motor impairments (gait and balance disturbances in particular).⁴ Physicians need to take age-related pathology and medical comorbidities into consideration when treating gait and balance impairments in PD.

1. Buchman AS, Leurgans SE, Yu L, et al. Incident parkinsonism in older adults without Parkinson disease. *Neurology* 2016;87:1036–1044.
2. Bohnen NI, Muller ML, Kuwabara H, et al. Age-associated striatal dopaminergic denervation and falls in community-dwelling subjects. *J Rehabil Res Dev* 2009;46:1045–1052.
3. van der Holst HM, van Uden IW, Tuladhar AM, et al. Cerebral small vessel disease and incident parkinsonism: the RUN DMC study. *Neurology* 2015;85:1569–1577.
4. Williams-Gray CH, Mason SL, Evans JR, et al. The CamPaIGN study of Parkinson's disease: 10-year outlook in an incident population-based cohort. *J Neurol Neurosurg Psychiatry* 2013;84:1258–1264.

© 2017 American Academy of Neurology

AUTHOR RESPONSE: INCIDENT PARKINSONISM IN OLDER ADULTS WITHOUT PARKINSON DISEASE

Aron S. Buchman, David A. Bennett, Chicago:

We thank Drs. Bohnen and Muller for their interest in our study on the high incidence of parkinsonism in community-dwelling older adults without PD.¹ They note, and we agree, that many factors other than PD pathology contribute to parkinsonian signs in older persons.^{2,3} Further, Drs. Bohnen and Muller make an excellent point regarding the implications of our findings for aging persons with PD; namely, the factors that contribute to parkinsonism in persons without PD are also likely to contribute to parkinsonism in persons with PD. This may have implications for treatment response and progression of PD. The situation is similar to that of Alzheimer disease (AD). In older persons, mixed pathologies are the most common cause of AD dementia and the coexisting non-AD pathology contributes to both the detection of AD and a more rapid rate of progression.^{4,5} We agree that physicians need to consider coexisting conditions that make an ever-larger contribution to many diseases with age.

1. Buchman AS, Leurgans SE, Yu L, et al. Incident parkinsonism in older adults without Parkinson disease. *Neurology* 2016;87:1036–1044.
2. Buchman AS, Shulman JM, Nag S, et al. Nigral pathology and parkinsonian signs in elders without Parkinson disease. *Ann Neurol* 2012;71:258–266.
3. Buchman AS, Leurgans SE, Nag S, Bennett DA, Schneider JA. Cerebrovascular disease pathology and parkinsonian signs in old age. *Stroke* 2011;42:3183–3189.
4. James BD, Bennett DA, Boyle PA, Leurgans S, Schneider JA. Dementia from Alzheimer disease and mixed pathologies in the oldest old. *JAMA* 2012;307:1798–1800.
5. Schneider JA, Arvanitakis Z, Leurgans SE, Bennett DA. The neuropathology of probable Alzheimer disease and mild cognitive impairment. *Ann Neurol* 2009;66:200–208.

© 2017 American Academy of Neurology

Author disclosures are available upon request (journal@neurology.org).