Differential expression of myocardial heat shock proteins in rats acutely exposed to fluoride

Lakshmikanthan Panneerselvam¹ · Azhwar Raghunath¹ · Ekambaram Perumal¹

Received: 6 January 2017 / Revised: 28 March 2017 / Accepted: 16 April 2017 / Published online: 27 April 2017
© Cell Stress Society International 2017

Abstract Acute fluoride (F^-) toxicity is known to cause severe cardiac complications and leads to sudden heart failure. Previously, we reported that increased myocardial oxidative damage, apoptosis, altered cytoskeleton and AMPK signaling proteins associated with energy deprivation in acute F^- induced cardiac dysfunction. The present study was aimed to decipher the status of myocardial heat shock proteins (Hsps-Hsp27, Hsp32, Hsp40, Hsp60, Hsp70, Hsp90) and heat shock transcription factor 1 (Hsf1) in acute F^- -intoxicated rats. In order to study the expression of myocardial Hsps, male Wistar rats were treated with single oral doses of 45 and 90 mg/kg F^- for 24 h. The expression levels of myocardial Hsps were determined using RT-PCR, western blotting, and immunohistochemical studies. Acute F^- -intoxicated rats showed elevated levels of both the transcripts and protein expression of Hsf1, Hsp27, Hsp32, Hsp60, and Hsp70 when compared to control. In addition, the expression levels of Hsp40 and Hsp90 were significantly declined in a dose-dependent fashion in F^- -treated animals. Our result suggests that differential expression of Hsps in the rat myocardium could serve as a balance between pro-survival and death signal during acute F^- -induced heart failure.

Keywords Acute fluoride toxicity · Cardiac dysfunction · Hsps · Hsf1

Abbreviations

Hsps	Heat shock proteins
Hsf1	Heat shock factor 1
DAB	3,3'-Diaminobenzidine
TMB	3,3',5,5'-Tetramethylbenzidine

Introduction

Acute F^- exposure always posed a high risk to human health during occupational and accidental exposure of hydrofluoric acid, ammonium bifluoride, or sodium fluoride (Lech 2011). Apart from the groundwater contamination of F^- , over fluoridation of public water system plays a major concern in many parts of the world (Gessner et al. 1994; Penman et al. 1997). Acute F^- exposure mainly emerges due to the ingestion of toxic amounts of F^- -containing agents from workplaces, home, laboratories, hospitals, and industries which leads to systemic toxicity (EHC 2002; ATSDR 2003).

Acute F^- exposure severely impairs the cardiovascular system and increases the risk of sudden cardiac failure. Previously, it was well studied that electrolyte imbalances, increased intracellular calcium-, hypocalcemia-, and hyperkalemia-mediated ventricular arrhythmias were the decisive events in the cardiac pathophysiology of acute F^- poisoning in both human case studies (Kumar 2009; Takase et al. 2004; Vohra et al. 2008) and experimental models (Su et al. 2003). In addition, F^- alters the mitochondrial membrane potential in H9c2 cells and inhibits oxidative phosphorylation in rat cardiac mitochondria (Pacauskiene et al. 2009; Yan et al. 2015). Our recent findings showed that F^- exposure causes

Lakshmikanthan Panneerselvam and Azhwar Raghunath contributed equally to this work.

Electronic supplementary material The online version of this article (doi:10.1007/s12192-017-0801-1) contains supplementary material, which is available to authorized users.

✉ Ekambaram Perumal
ekas2009@buc.edu.in

¹ Molecular Toxicology Laboratory, Department of Biotechnology, Bharathiar University, Coimbatore, Tamil Nadu 641 046, India

cardiac dysfunction through (i) increased myocardial oxidants and inhibition of antioxidant machinery, (ii) induction of apoptotic and necrotic stimulus, and (iii) impaired myocardial cytoskeletons and depletion of ATP level leading to the activation of AMPK signaling molecules in rat heart (Panneerselvam et al. 2015, 2017).

Heat shock proteins (Hsps) are the endogenous stress response proteins that confer cytoprotection against xenobiotic insults (Kalmar and Greensmith 2009). Hsps are categorized into subfamilies: HSPB1-Hsp27, Hsp32, DNAJA4-Hsp40, HSPD1-Hsp60, HSPA1B-Hsp70, and HSPC1-Hsp90 according to their molecular weight (16–110 kDa) and their localization in the various subcellular organelles. Among the diverse classes of Hsps-Hsp60, 70, and 90 exhibit molecular chaperone properties which maintain the balance between synthesis and degradation of protein folding and trafficking (Hart et al. 2011). In general, the Hsps expression is either constitutive (HSPB1, HSPD1) or inducible (HSPA1B, Hsp32) and both (HSPA1B, HSPC1). The induction of Hsps is based on the nature of the toxic stimulus (Kalmar and Greensmith 2009). The selection of these Hsps in the present study was mainly based on their role in various xenobiotic-induced cardiotoxic and cardiovascular disease models (Lin et al. 2007; Snoeckx et al. 2001). The expression profile of these Hsps is mainly used to monitor the cardiac pathophysiological consequences upon xenobiotic insults. Earlier literatures showed that single acute arsenic exposure induces Hsp70 expression in the heart, kidney, and liver of rabbits (Brown and Rush 1984). Besides, upregulation of Hsf1 mediated overexpression of Hsp27, and Hsp90 attenuates H₂O₂-induced cardiac failure (Yu et al. 2012). In addition, exercise preconditioning upregulates cardiac Hsf1 and Hsp70 expression in pressure overload rat model (Xu et al. 2015). Our previous study showed that F[−]-induced upregulation of Hsp60 and downregulation of Hsp27 are the indicators for the detection of oxidative stress and apoptosis, in rat liver (Kanagaraj et al. 2015). Further, iron oxide nanoparticle exposure causes alterations in Hsp60, Hsp70, and Hsp90 expressions in mice male accessory glands—seminal vesicles and prostate glands (Sundarraj et al. 2017). However, the status of Hsps expression in acute F[−]-induced cardiotoxicity is yet to be studied. Hence, we made an attempt to explore the molecular events leading to induction of myocardial Hsps in acute F[−] toxicity in male rats.

Materials and methods

Chemicals

Sodium fluoride (Loba Chemie, Mumbai, India), antibodies of Hsp40, Hsp70, Hsp90, and Hsf1 (Cell Signaling Technology, Inc., USA), Hsp60 (Santa Cruz Biotechnology, Inc., USA),

Hsp27, Hsp32 (Enzo Life Science, Inc., USA), GAPDH (GeneTex, Inc., USA), and secondary antibodies conjugated with horseradish peroxidase (Merck, Darmstadt, Germany) were used in this study. All the detailed information of primary antibodies is listed in Table 2. Polyvinylidene fluoride (PVDF) membrane (Millipore Immobilon, Bedford, USA), Pierce™ DAB Substrate kit for Immunohistochemistry and Pierce™ TMB Substrate kit, First-strand complementary DNA (cDNA) synthesis kit (Thermo Fisher Scientific, TX, USA), and Total tissue RNA isolation kit (Favorgen Biotech Corp. Taiwan) were also used in this study. All the other chemicals used in the experiments were of molecular grade unless mentioned otherwise.

Animals

Adult (3 months and 2 weeks old) male Wistar rats, weighing (140–150 g) were chosen for the experiment. The animals were purchased from Karpagam University, Coimbatore, India. The animals were housed in the vivarium at standard temperature (22 ± 3 °C) and humidity (55 ± 5%) in a normal 12-h light/dark cycle and supplemented with standard commercially available pelleted rat chow (Krish Scientific, Pvt. Ltd., Bangalore, India), and the tap water were supplied ad libitum. All the animal experiments performed in this study were in consonance with the ethical guidelines certified by the committee for the purpose of control and supervision of experiments on animals (CPCSEA) and institutional animal ethical committee (Approval No.: 722/02/CPCSEA).

Experimental design

The animals were randomly divided into three groups of three rats in each group. Group 1: normal control rats were administered orally equivalent volume of tap water (0.2 ppm of F[−]). Group 2: rats were administered with F[−] 45 mg/kg body weight (b.w.), orally in a volume of 0.1 mL/100 g b.w. for 24 h. Group 3: rats were administered with F[−] 90 mg/kg b.w., orally in a volume of 0.1 mL/100 g b.w. for 24 h. The LD₅₀ value of acute F[−] toxicity was reported as 126.3 mg F[−]/kg in male rats (Whitford 1990). Based on our previous studies (Panneerselvam et al. 2015, 2017), we selected both doses of 45 and 90 mg F[−]/kg to assess the changes in the expression of myocardial Hsps in rat. After 24 h, the rats were sacrificed by cervical decapitation and the heart tissues were dissected out, rinsed in physiological saline, and stored in −80 °C for further experiments.

Reverse transcriptase polymerase chain reaction (RT-PCR)

The total RNA from the rat myocardium was isolated and then first-strand cDNA was synthesized using commercially available kit (First-strand cDNA synthesis kit, Thermo Fisher

Scientific, TX, USA) as per manufacturer's protocol. Polymerase chain reaction (PCR) analysis was performed in order to detect gene expressions of Hsps using the Eppendorf thermal cycler. Oligonucleotide primers used for amplification were shown in Table 1. The PCR amplification profile for each gene involved an initial denaturation step at 95 °C for 3 min and the annealing step was kept at a primer specific temperature for 45 s. The annealing temperature of 58 °C was selected for the primers Hsp27, Hsp40, Hsp70, and GAPDH and 63 °C for the primers Hsf1, Hsp32, Hsp60, and Hsp90. The PCR products were extended for 60 s at 72 °C, with a final elongation step at 72 °C for 7 min. After amplification, the RT-PCR products were electrophoresed on 2% agarose gels, and stained with ethidium bromide. Glyceraldehyde 3-phosphate dehydrogenase (GAPDH) was used as the internal control to ensure equal sample loading. Relative messenger RNA (mRNA) expression was quantified using *ImageJ* analysis software (NIH).

Western blot analysis

Western blot analyses were performed in the myocardial tissue of control and F⁻-intoxicated rats. The myocardial tissues were homogenized with tissue homogenizer in ice-cold RIPA lysis buffer (50 mM Tris-HCl, pH 7.4, 150 mM NaCl, 1% NP40, 0.25% sodium orthovanadate 1 mM PMSF, and 1x protease inhibitor cocktail, Sigma-Aldrich) and the total protein was measured using Bradford reagent with BSA as standard. Forty micrograms of protein sample was equally loaded in each lane separated by 10% SDS-PAGE gel electrophoresis and transferred to a PVDF membrane. The membrane was blocked using 5% fat-free BSA and incubated overnight with respective primary antibodies (Table 2). The proteins were detected using horseradish peroxidase-conjugated anti-rabbit secondary antibodies and visualized by the TMB staining kit. The membrane was not reprobed using stripping buffer. GAPDH was used as the internal control to ensure equal sample loading. Densitometry analyses of immunoblots were performed using *ImageJ* analysis software (NIH).

Immunohistochemical analysis

Immunohistochemistry was performed based on a standard optimized protocol of McNicol and Richmond (1998) with few modifications as standardized earlier in our previous study (Panneerselvam et al. 2015). The paraffin-embedded unstained section of myocardial tissues were deparaffinized in the xylene and followed by subsequent rehydration with descending series (100%, 90%, 70%, 50%, 30%) of ethanol. The sections were placed with laboratory stainless pressure cooker in order to unmask the antigen by heat-induced epitope retrieval method. Endogenous peroxidase activity was quenched by

Table 1 Primers used for the RT-PCR

Gene name ^a	GenBank accession number	Forward primer	Reverse primer	Product size (bp)	References
<i>HSF1</i>	XM_006241891	5'-CCAGCAGCAAAAAGTTGTCA-3'	5'-TGGTGACAGCATCAGAGGAG-3'	239	Reddy et al. 2014
<i>Hsp27 (HSPB1)</i>	NM_031970.4	5'-GATGAACATGGCTACATCTCTC-3'	5'-CTGATTGTGTGACTGCTTTG-3'	145	Reddy et al. 2014
<i>Hsp32 (HMOX1)</i>	NM_012580.2	5'-CGTGCAGAGAATCTGAGTTC-3'	5'-AGACGCTTTACGTAGTGCTG-3'	275	Mahmoud-Awny et al. 2015
<i>Hsp40 (DNAJA4)</i>	NM_001025411	5'-TGAAGGCATTGGTGGGAAGA-3'	5'-AGCGGTCTCTGGGGTTGATA-3'	167	Jeckel et al. 2014
<i>Hsp60 (HSPD1)</i>	NM_022229.2	5'-AAGGTTGGAAGAAAGGGTGTCA-3'	5'-CCTTCAACACAGCTACTCCATC-3'	643	Sarang et al. 2013
<i>Hsp70 (HSPA1B)</i>	L16764.1	5'-TGGTGTCTGACCAAGATGAAG-3'	5'-AGGTCGAAGATGAGCACGTT-3'	235	Cattelan et al. 2015
<i>Hsp90 (HSPC1)</i>	NM_175761.2	5'-TTATCACAGGTGAGACCAAGGA-3'	5'-CTCTCCATGTTGCTGTCCA-3'	358	Cattelan et al. 2015
<i>GAPDH</i>	NG_016258.2	5'-GCATGGCCTTCGCTGTTCC-3'	5'-GGGTGGTCCAGGGTTTCTTACTC-3'	336	Max et al. 2014

^a The acronyms used are related to database information and human homologs (<http://rgd.mcw.edu/nomen/homen.shtml>)

Table 2 Primary antibodies used in the study

Antibodies	Catalog no.	Manufacturer details	Species	Dilution (μ L)	Purpose
Hsf1	CST4356	Cell signaling Technology, Inc., USA	Rabbit Polyclonal	1:1000, 1:50	WB, IHC
Hsp27	ADI-SPA-803-F	Enzo Life Science, Inc., USA	Rabbit Polyclonal	1:1000, 1:200	WB, IHC
Hsp32	BML-HC3001	Enzo Life Science, Inc., USA	Rabbit Polyclonal	1:1000, 1:200	WB, IHC
Hsp40	CST4871	Cell signaling Technology, Inc., USA	Rabbit Polyclonal	1:1000, 1:100	WB, IHC
Hsp60	SC-1052	Santa Cruz Biotechnology, Inc., USA	Goat Polyclonal	1:2000, 1:500	WB, IHC
Hsp70	CST4872P	Cell signaling Technology, Inc., USA	Rabbit Polyclonal	1:1000, 1:50	WB, IHC
Hsp90	CST4877	Cell signaling Technology, Inc., USA	Rabbit Polyclonal	1:1000, 1:100	WB, IHC
GAPDH	GTX100118	GeneTex, Inc., USA	Rabbit Polyclonal	1:1000	WB

treating the slides with 1% H_2O_2 solution. The slides were blocked with 5% BSA for 1 h at room temperature to avoid the non-specific reactivity. The sections were incubated with respective primary antibodies in Table 2. The proteins were detected using horseradish peroxidase-conjugated anti-rabbit or anti-goat secondary antibodies. Immunoreactions were labeled using a DAB staining kit. Sections were counterstained with hematoxylin, dehydrated with ascending series of ethanol, and mounted with distrene plasticizer xylene. Immunoreactivity was quantitated in a blinded fashion for depicting the total number of positively stained cells.

Statistical analysis

All the values were expressed as mean $\pm$ SD of three rats in each group. The data were analyzed with one-way analysis of variance (ANOVA) followed by Tukey's multiple comparison test using SPSS (20.0) software package. $p < 0.05$ was considered to indicate statistical significance.

Results and discussion

The expression levels of cardiac Hsf1 and Hsps (Hsp27, Hsp32, Hsp40, Hsp60, Hsp70, and Hsp90) were determined using RT-PCR, western blotting, and immunohistochemical staining studies in the control and F^- -intoxicated rats. The relative mRNA (Fig. 1) and protein (Fig. 2) expression levels of Hsf1 and Hsps (Hsp27, Hsp32, Hsp60, and Hsp70) were found to be elevated in both the 45- and 90-mg F^- -treated rats when compared to control (Supplementary Figs. 1 and 2). The expression levels of Hsps (Hsp40 and Hsp90) were significantly declined in a dose-dependent fashion in these animals. The similar pattern of differential expression of all the above mentioned Hsps was also observed in the immunohistochemical studies (Fig. 3).

Hsps are among the major defensive stress proteins, which play an important role in the pathophysiology of various xenobiotic-induced cardiotoxicity models (Shan et al. 2003; Yu et al. 2012). Acute F^- poisoning is known to cause sudden cardiac failure in both humans (Vohra et al. 2008; Lech 2011) and experimental animals (McIvor et al. 1987). Our previous

Fig. 1 Effect of acute F^- intoxication on myocardial mRNA expression of Hsf1 and Hsps in rats. mRNA expression and its densitometric analysis of myocardial Hsf1 and Hsps (Hsp27, Hsp32, Hsp40, Hsp60, Hsp70, and Hsp90) in control and F^- -treated rats. GAPDH was used as loading

control. The values represent mean $\pm$ SD ($n = 3$). * $p < 0.05$ significantly different from control, + $p < 0.05$ significantly different from 45 mg F^- -treated rats (one-way ANOVA followed by Tukey's multiple comparison test)

Fig. 2 Effect of acute F⁻ intoxication on myocardial protein expression of Hsf1 and Hsps in rats. Western blotting and its densitometric analysis of myocardial Hsf1 and Hsps (Hsp27, Hsp32, Hsp40, Hsp60, Hsp70, and Hsp90) in control and F⁻-treated rats. GAPDH was used as loading

control. The values represent mean $\pm$ SD ($n = 3$). * $p < 0.05$ significantly different from control, ** $p < 0.05$ significantly different from 45 mg F⁻ treated rats (one-way ANOVA followed by Tukey's multiple comparison test)

study showed that acute F⁻ exposure impairs cardiac dysfunction through increased mitochondrial oxidative damage, apoptosis, and necrosis in rat myocardium (Panneerselvam et al. 2015). The present study was aimed to determine changes in the expression of cardiac Hsf1 and Hsps (Hsp27, Hsp32, Hsp40, Hsp60, Hsp70, Hsp90) and their crucial role in acute F⁻-induced cardiac dysfunction.

Hsf1 is the transcription factor, which regulates the expression of several Hsps (Christians et al. 2002). Upon the unfavorable state, Hsf1 migrate to the nucleus and binds in the promoter region of heat shock element, thereby initiating transcription and synthesis of Hsps. In general, activation of Hsf1 confers pro-survival mechanism in the cardiomyocytes exposed to various physiological stressors, including ROS (Nishizawa et al. 1999), oxidative stress (Kalmar and Greensmith 2009), and hypoxia (Snoeckx et al. 2001). Interestingly, in our study, a significant increase in the expression of Hsf1 and its downstream Hsp27, Hsp32, Hsp60, and Hsp70 with concomitant decrease in Hsp40 and Hsp90 were observed in the myocardium of F⁻-treated rats when compared to those in the control. Thus, the increased cardiac Hsf1 expression might be due to the induction of ROS-mediated oxidative stress in the myocardium of acute F⁻-intoxicated rats (Panneerselvam et al. 2015). Therefore, the elevated cardiac Hsf1 level endows defensive role in cardiac physiology of acute F⁻ intoxication. The activation of Hsf1 is accompanied by increased levels of Hsp27, Hsp32, Hsp60, and Hsp70, but decreased expression of Hsp40 and Hsp90. This decreased expression of Hsp40 and Hsp90 has to be investigated further.

It has been reported that increased expression of myocardial Hsp27 serves as a prognostic marker for myocardial ischemia (Ghayour-Mobarhan et al. 2012). Thus, increased cardiac Hsp27 expression in the present study was correlated with our previous report, which exhibits increased myocardial cTnI expression associated with myocardial necrosis and

impaired contractile function in acute F⁻ intoxication (Panneerselvam et al. 2015). Furthermore, Hsp27 has been demonstrated to play an important role in regulating intracellular redox homeostasis and anti-apoptosis (Arrigo et al. 2005). We speculate that increased expression of Hsp27 in the lower concentration of F⁻ exposure might be due to its reductive stress-associated cardiomyopathy (Zhang et al. 2010). Besides, decreased expression of Hsp27 in the higher concentration of F⁻ might be due to enhanced apoptotic stimulus in the rat heart. The results were in accordance with our previous findings, which revealed the increased myocardial expression of apoptotic markers such as terminal deoxynucleotidyl transferase (TdT) dUTP nick-end labeling (TUNEL) staining and pro-caspase 3 expression in a dose-dependent manner of acute F⁻-treated rats (Panneerselvam et al. 2015). Similarly, we reported the decreased levels of Hsp27 expression in the liver of F⁻-intoxicated rats, which could be due to oxidative stress-mediated apoptosis (Kanagaraj et al. 2015). It is well known that F⁻ is a potent inhibitor of myocardial antioxidant enzymes and increases the intracellular oxidants through the Nox4 expression in rat heart (Panneerselvam et al. 2015; Oyagbemi et al. 2016). The increased Hsp32 expression in the present study was directly associated with decreased glutathione content in diethyl maleate-treated brain (Ewing and Maines 1993) and F⁻-exposed heart (Panneerselvam et al. 2015) in rats. Increased levels of circulating Hsp60 were found in patients with acute myocardial infarction (AMI) suggesting a predictive marker for post-AMI adverse events (Novo et al. 2011). In our study, the increased levels of Hsp60 expression in acute F⁻ intoxication could be correlated with increased oxidative stress-mediated apoptosis in the myocardium (Panneerselvam et al. 2015). Our findings in the present were in accordance with our previous report, which exhibits increased expression of hepatic Hsp60 in rats exposed to chronic F⁻ intoxication (Kanagaraj et al. 2015).

Fig. 3 a Effect of acute F⁻ intoxication on myocardial immunohistochemical expression of Hsf1 and low molecular weight Hsps in rats. Myocardial sections were stained with antibodies specific for Hsf1, Hsp27, Hsp32, and Hsp40 in control and F⁻-intoxicated rats. Number of immunopositively stained cells in ten fields/section ($\times 200$) was quantified by three independent observers in blinded fashion. The *bar diagram* shows the quantification of positively stained cells of control and F⁻-treated rats. The values represent mean $\pm$ SD ($n = 3$). * $p < 0.05$ significantly different from control, + $p < 0.05$ significantly different from 45 mg F⁻-treated rats (one-way ANOVA followed by Tukey's multiple comparison test). **b** Effect of acute F⁻ intoxication on

myocardial immunohistochemical expression of high molecular weight Hsps in rats. Myocardial sections were stained with antibodies specific for Hsp60, Hsp70, and Hsp90 ($\times 200$) in control and F⁻ intoxicated rats. Number of immunopositively stained cells in ten fields/section ($\times 200$) was quantified by three independent observers in blinded fashion. The *bar diagram* shows the quantification of positively stained cells of control and F⁻ treated rats. The values represent mean $\pm$ SD ($n = 3$). * $p < 0.05$ significantly different from control, + $p < 0.05$ significantly different from 45 mg F⁻-treated rats (one-way ANOVA followed by Tukey's multiple comparison test)

Earlier studies showed that elevated extracellular levels of Hsp70 serve as a prognostic marker for inflammatory cytokine-mediated immune response and the progression of heart failure in patients with acute AMI (Jenei et al. 2013; Satoh et al. 2006). Our results are in accordance with the report of Chattopadhyay et al. (2011). Their findings showed the increased oxidative stress directly correlated with elevated Hsp70 expression in the mice kidney exposed to chronic F⁻ intoxication. Hayashi et al. (2006) reported that Hsp40 plays a crucial role in regulating the mitochondrial biogenesis and respiratory chain disorders in the progression of dilated cardiomyopathies. The inactivation of human DnaJ homolog Hdj2 was due to the oxidation of cysteine residue in the zinc finger domain leading to the loss of its chaperonic activity eventually increasing protein aggregation under oxidative stress (Choi et al. 2006). However, to our knowledge, this is the first-ever study to reveal the expression of Hsp40 in F⁻-intoxicated rats. Thus, the decreased expression of Hsp40 in the present study is in line with our previous findings that exhibit increased oxidative stress, apoptosis, and mitochondrial dysfunction in both in vivo (Panneerselvam et al. 2015) and in vitro (Ameeramja et al. 2016; Ameeramja and Perumal 2016). Further, Bouaziz et al. (2007) showed that F⁻ exposure inhibits the expression of Hsp90 in the liver and kidney of mother and offsprings. Further, it has been reported that iron oxide nanoparticle exposure decreases the Hsp90 expression in mice male accessory glands—seminal vesicle and prostate gland associated with oxidative stress (Sundarraj et al. 2017). Thus, the decreased expression of Hsp90 might be due to the increased oxidative stress and apoptotic and necrotic stimuli in both doses of the acute F⁻-intoxicated rats. In addition, the decreased expression of Hsp90 could be through the integration of other stress signaling pathways and post-translational modifications (Prodromou 2017). Similar decreased Hsp90 was also observed on exposure to steroid hormones with increased Hsf1 in the adult rat cardiomyocytes, which represents a unique pathway in the regulation of Hsps (Knowlton and Sun 2001). Besides, Hsp90 reduction resulted in Hsf1 activation revealing its negative regulation (Snoeckx et al. 2001). It has also been reported that pharmacological Hsp90 inhibitors activate Hsf1 stress response pathway and improve the glucose regulation in diabetic mice (Lee et al. 2013). However, the decreased expression of Hsp90 and upregulation of Hsf1 in the present study is subjected to further investigation.

Conclusion

To our knowledge, we are the first to reveal the differential expression of Hsps and their indispensable role in the pathophysiology of acute F⁻-induced cardiac dysfunction in rats. In our earlier studies, we observed that increased oxidative stress, mitochondrial dysfunction, apoptosis, impaired cytoskeleton

intermediate filaments, and energy deprivation-mediated AMPK signaling molecules might be one of the possible reasons for the differential expression of Hsps. Thus, acute F⁻ exposure alters myocardial Hsps expression at both transcript and protein levels. Further studies are needed to confirm with using Hsps-inducible or conditional knockout models.

Acknowledgements The authors are grateful to research fellows Ms. Kiruthika Sundarraj and Ms. Vinothini Subramaniam, Bharathiar University, Coimbatore, for their timely assistance throughout the work.

Compliance with ethical standards

Conflict of interest The authors declare that they have no conflict of interest.

Funding information This work was supported by the UGC-SAP DRS II:F-3-30/2013 and DST-FIST:SR/FST/LSI-618/2014, New Delhi, India, for the partial financial assistance. LP extends his thanks to the Indian Council of Medical Research, New Delhi, for financial assistance in the form of a senior research fellowship (No. 45/25/2013/BMS/TRM). AR acknowledges the UGC-BSR fellowship (UGC-BSR No. F7-25/2007) from UGC-BSR, New Delhi, India.

References

- Ameeramja J, Panneerselvam L, Govindarajan V, Jeyachandran S, Baskaralingam V, Perumal E (2016) Tamarind seed coat ameliorates fluoride induced cytotoxicity, oxidative stress, mitochondrial dysfunction and apoptosis in A549 cells. *J Hazard Mat* 301:554–565
- Ameeramja J, Perumal E (2016) Protocatechuic acid methyl ester ameliorates fluoride toxicity in A549 cells. *Food Chem Toxicol*. doi:10.1016/j.fct.2016.12.024
- Arrigo AP, Viot S, Chaufour S, Firdaus W, Kretz-Remy C, Diaz-Latoud C (2005) Hsp27 consolidates intracellular redox homeostasis by upholding glutathione in its reduced form and by decreasing iron intracellular levels. *Antioxid Redox Signal* 7:414–422
- ATSDR (2003) Agency for Toxic Substances and Disease Registry TP-91/17 ATSDR, toxicological profile for fluorides, hydrogen fluoride, and fluorine. Department of Health and Human Services, Atlanta
- Bouaziz H, Croute F, Boudawara T, Soleilhavoup JP, Zeghala N (2007) Oxidative stress induced by fluoride in adult mice and their suckling pups. *Exp. Toxicol Path* 58:339–349
- Brown IR, Rush SJ (1984) Induction of a “stress” protein in intact mammalian organs after the intravenous administration of sodium arsenite. *Biochem Biophys Res Commun* 120:150–155
- Cattelan A, Ceolotto G, Bova S, Albiero M, Kuppusamy M, De Martin S, Semplicini A, Fadini GP, de Kreutzenberg SV, Avogaro A (2015) NAD⁽⁺⁾-dependent SIRT1 deactivation has a key role on ischemia-reperfusion-induced apoptosis. *Vasc Pharmacol* 70:35–44
- Chattopadhyay A, Podder S, Agarwal S, Bhattacharya S (2011) Fluoride-induced histopathology and synthesis of stress protein in liver and kidney of mice. *Arch Toxicol* 85:327–335
- Choi HI, Lee SP, Kim KS, Hwang CY, Lee YR, Chae SK, Kim YS, Chae HZ, Kwon KS (2006) Redox-regulated co-chaperone activity of the human DnaJ homolog Hdj2. *Free Radic Biol Med* 40:651–659
- Christians ES, Yan LJ, Benjamin IJ (2002) Heat shock factor 1 and heat shock proteins: critical partners in protection against acute cell injury. *Crit Care* 30:S43–S50
- EHC (2002) Environment health criteria 227, fluorides. World Health Organization, WHO, Geneva

- Ewing JF, Maines MD (1993) Glutathione depletion induces heme oxygenase-1 (HSP32) mRNA and protein in rat brain. *J Neurochem* 60:1512–1519
- Gessner BD, Beller M, Middaugh JP, Whitford GM (1994) Acute fluoride poisoning from a public water system. *N Engl J Med* 330:95–99
- Ghayour-Mobarhan M, Saber H, Ferns GA (2012) The potential role of heat shock protein 27 in cardiovascular disease. *Clin Chim Acta* 413:15–24
- Hart FU, Bracher A, Hayer-Hart M (2011) Molecular chaperones in protein folding and proteostasis. *Nature* 475:324–332
- Hayashi M, Imanaka-Yoshida K, Yoshida T, Wood M, Fearn C, Tataka RJ, Lee JD (2006) A crucial role of mitochondrial Hsp40 in preventing dilated cardiomyopathy. *Nat Med* 12:128–132
- Jeckel KM, Bouma GJ, Hess AM, Petrilli EB, Frye MA (2014) Dietary fatty acids alter left ventricular myocardial gene expression in Wistar rats. *Nutr Res* 34:694–706
- Jenei ZM, Gombos T, Forhecz Z, Pozsonyi Z, Karádi I, Jánoskúti L, Prohászka Z (2013) Elevated extracellular HSP70 (HSPA1A) level as an independent prognostic marker of mortality in patients with heart failure. *Cell Stress Chaperones* 18:809–813
- Kalmar B, Greensmith L (2009) Induction of heat shock proteins for protection against oxidative stress. *Adv Drug Deliv Rev* 61:310–318
- Kanagaraj VV, Panneerselvam L, Govindarajan V, Ameeramja J (2015) Caffeic acid, a phyto polyphenol mitigates fluoride induced hepatotoxicity in rats: a possible mechanism. *Biofactors* 41:90–100
- Knowlton AA, Sun L (2001) Heat-shock factor-1, steroid hormones, and regulation of heat-shock protein expression in the heart. *Am J Physiol Heart Circ Physiol* 28:455–464
- Kumar A (2009) Fatality due to sodium fluoride ingestion—suicide or accident? *Can Soc Forensic Sci J* 42:69–74
- Lech T (2011) Fatal cases of acute suicidal sodium and accidental zinc fluorosilicate poisoning. Review of acute intoxications due to fluoride compounds. *Forensic Sci Int* 206:20–24
- Lee JH, Gao J, Kosinski PA, Elliman SJ, Hughes TE, Gromada J, Kemp DM (2013) Heat shock protein 90 (HSP90) inhibitors activate the heat shock factor 1 (HSF1) stress response pathway and improve glucose regulation in diabetic mice. *Biochem Biophys Res Commun* 430:1109–1113
- Lin L, Kim SC, Wang Y, Gupta S, Davis B, Simon SI, Torre-Amione G, Knowlton AA (2007) HSP60 in heart failure: abnormal distribution and role in cardiac myocyte apoptosis. *Am J Physiol Heart Circ Physiol* 293:H2238–H2247
- Mahmoud-Awny M, Attia AS, Abd-Ellah MF, El-Abhar HS (2015) Mangiferin mitigates gastric ulcer in ischemia/reperfused rats: involvement of PPAR-1, NF- κ B and Nrf2/HO-1 signaling pathways. *PLoS One* 10:e0132497
- Max D, Brandsch C, Schumann S, Kuhne H, Frommhagen M, Schutkowski A, Hirche F, Staeger MS, Stangl GI (2014) Maternal vitamin D deficiency causes smaller muscle fibers and altered transcript levels of genes involved in protein degradation, myogenesis, and cytoskeleton organization in the newborn rat. *Mol Nutr Food Res* 58:343–352
- McIvor ME, Cummings CE, Mower MM, Wenk RE, Lustgarten JA, Baltazar RF, Salomon J (1987) Sudden cardiac death from acute fluoride intoxication: the role of potassium. *Ann Emerg Med* 16:777–781
- McNicol AM, Richmond JA (1998) Optimizing immunohistochemistry: antigen retrieval and signal amplification. *Histopathology* 32:97–103
- Nishizawa J, Nakai A, Matsuda K, Komeda M, Ban T, Nagata K (1999) Reactive oxygen species play an important role in the activation of heat shock factor 1 in ischemic-reperfused heart. *Circulation* 99:934–941
- Novo G, Cappello F, Rizzo M, Fazio G, Zambuto S, Tortorici E et al (2011) Hsp60 and heme oxygenase-1 (Hsp32) in acute myocardial infarction. *Transl Res* 157:285–292
- Oyagbemi AA, Omobowale TO, Asegun ER, Adejumbi AO, Ajibade TO, Ige TM, Ogunpolu BS, Adedapo AA, Yakubu MA (2016) Sodium fluoride induces hypertension and cardiac complications through generation of reactive oxygen species and activation of nuclear factor kappa beta. *Environ Toxicol*. doi:10.1002/tox.22306
- Pacauskiene I, Kopustinskiene D, Paipaliene PP, Sadzeviciene R (2009) Effect of fluoride on the respiration rate of rat cardiac mitochondria. *Fluoride* 42:198–202
- Panneerselvam L, Govindarajan V, Ameeramja J, Nair HR, Perumal E (2015) Single oral acute fluoride exposure causes changes in cardiac expression of oxidant and antioxidant enzymes, apoptotic and necrotic markers in male rats. *Biochimie* 119:27–35
- Panneerselvam L, Raghunath A, Perumal E (2017) Acute fluoride poisoning alters myocardial cytoskeleton and AMPK signaling proteins in rats. *Int J Cardiol* 229:96–101
- Penman AD, Brackin BT, Embrey R (1997) Outbreak of acute fluoride poisoning caused by a fluoride overfeed, Mississippi, 1993. *Public Health Rep* 112:403–409
- Prodromou C (2017) Regulatory mechanisms of Hsp90. *Biochem Mol Biol J*. doi:10.21767/2471-8084.100030
- Reddy VS, Kumar CHU, Raghu G, Reddy GB (2014) Expression and induction of small heat shock proteins in rat heart under chronic hyperglycemic conditions. *Arch Biochem Biophys* 558:1–9
- Sarangi U, Singh MK, Abhijanya KVV, Reddy LPA, Prasad BS, Pitke VV, Paithankar K, Sreedhar AS (2013) Hsp60 chaperonin acts as barrier to pharmacologically induced oxidative stress mediated apoptosis in tumor cells with differential stress response. *Drug Target Insights* 7:35–51
- Satoh M, Shimoda Y, Akatsu T, Ishikawa Y, Minami Y, Nakamura M (2006) Elevated circulating levels of heat shock protein 70 are related to systemic inflammatory reaction through monocyte Toll signal in patients with heart failure after acute myocardial infarction. *Eur J Heart Fail* 8:810–815
- Shan YX, Liu TJ, Su HF, Samsamshariat A, Mestri R, Wang PH (2003) Hsp10 and Hsp60 modulate Bcl-2 family and mitochondria apoptosis signaling induced by doxorubicin in cardiac muscle cells. *Mol Cell Cardiol* 35:1135–1143
- Snoeckx LH, Cornelussen RN, Van Nieuwenhoven FA, Reneman RS, Van Der Vusse GJ (2001) Heat shock proteins and cardiovascular pathophysiology. *Physiol Rev* 81:1461–1497
- Su M, Chu J, Howland MN, Nelson LS, Hoffman RS (2003) Amiodarone attenuates fluoride-induced hyperkalemia in vitro. *Acad Emerg Med* 10:105–109
- Sundarraj K, Raghunath A, Panneerselvam L, Perumal E (2017) Iron oxide nanoparticles modulate heat shock proteins and organ specific markers expression in mice male accessory organs. *Toxicol Appl Pharmacol* 317:12–24
- Takase I, Kono K, Tamura A, Nishio H, Dote T, Suzuki K (2004) Fatality due to acute fluoride poisoning in the work place. *Leg Med (Tokyo)* 6:197–200
- Vohra R, Velez LI, Rivera W, Benitez FL, Delaney KA (2008) Recurrent life threatening ventricular dysrhythmias associated with acute hydrofluoric acid ingestion: observations in one case and implications for mechanism of toxicity. *Clin Toxicol* 46(46):79–84
- Whitford GM (1990) The physiological and toxicological characteristics of fluoride. *J Dent Res* 69:539–549
- Xu T, Zhang B, Yang F, Cai C, Wang G, Han Q, Zou L (2015) HSF1 and NF- κ B p65 participate in the process of exercise preconditioning attenuating pressure overload-induced pathological cardiac hypertrophy. *Biochem Biophys Res Commun* 460:622–627
- Yan X, Yang X, Hao X, Ren Q, Gao J, Wang Y (2015) Sodium fluoride induces apoptosis in H9c2 cardiomyocytes by altering mitochondrial membrane potential and intracellular ROS level. *Biol Trace Elem Res* 166:210–215
- Yu Y, Liu M, Zhang L, Cao Q, Zhang P, Jiang H, Zou Y, Ge J (2012) Heat shock transcription factor 1 inhibits H₂O₂-induced cardiomyocyte death through suppression of high-mobility group box 1. *Mol Cell Biochem* 364:263–269
- Zhang X, Min X, Li C, Benjamin JJ, Qian B, Zhang X, Ding Z, Gao X, Yao Y, Ma Y, Cheng Y, Liu L (2010) Involvement of reductive stress in the cardiomyopathy in transgenic mice with cardiac-specific overexpression of heat shock protein 27. *Hypertension* 55:1412–1417