Published in final edited form as:

J Dev Orig Health Dis. 2017 August ; 8(4): 474–482. doi:10.1017/S2040174417000150.

Neonatal fatty acid profiles are correlated with infant growth measures at six months

PF O'Tierney-Ginn^{1,*}, D Davina², M Gillingham³, DJP Barker^{4,5,6}, C Morris⁷, and KL Thornburg^{5,6}

¹Center for Reproductive Health, Department of Reproductive Biology, Case Western Reserve University, Cleveland, OH

²Peachtree Healthcare, Sacramento, CA

³Department of Molecular & Medical Genetics, Oregon Health and Science University (OHSU), Portland, OR

⁵Center for Developmental Health, OHSU Knight Cardiovascular Institute

⁶Bob and Charlee Moore Institute for Nutrition and Wellness, OHSU

⁷Department of Med Informatics & Clinical Epidemiology, OHSU

Abstract

Rapid weight gain in infancy and low levels of n-3 long chain polyunsaturated fatty acids (LCPUFA) at birth are associated with increased adiposity later in life. The association between placental LCPUFA delivery and weight gain in infancy is poorly understood. We sought to determine the relationships between maternal phenotype, placental fatty acid transporter expression and offspring growth patterns over the first 6 months. Placental tissue and cord blood were collected at term delivery from women with uncomplicated pregnancies. Offspring body composition measurements were recorded 1 day and 6 months after birth. Body mass index (BMI) z-scores were determined using WHO 2006 reference data. Body phenotype patterns were compared among offspring who had an increase in BMI z-score and those who had a decrease. High skinfold thickness at birth and positive change in BMI z-scores during infancy were associated with low neonatal n-3 LCPUFA plasma levels ($r=-0.46$, $P=0.046$) and high saturated fatty acids levels ($r=0.49$, $P=0.034$). Growth of skinfolds over 6 months of age was associated with placental fatty acid transporter gene expression. Change in BMI z-score in the first 6 months of life correlated with arm muscle area growth, a measure of lean mass ($r=0.62$, $P=0.003$), but not with growth in skinfold thickness. Early infancy weight gain was associated with poor plasma LCPUFA status at birth, and fat deposition in infancy was related to changes in placental lipid

*Corresponding Author: Perrie O'Tierney-Ginn, PhD, 2500 MetroHealth Drive, R358, Cleveland, Ohio, USA, 44118, poginn@metrohealth.org.

⁴Deceased

Conflicts of Interest

None

Ethical Standards

The authors assert that all procedures contributing to this work comply with the ethical standards of the US Department of Health & Human Services Regulations for the Protection of Human Subjects, with the Helsinki Declaration of 1975, as revised in 2008, and have been approved by the Oregon Health & Science University Institutional Review Board.

handling. Thus, neonatal fatty acid profiles may influence the trajectory of infant growth and fat and lean mass deposition.

Keywords

placenta; fatty acids; adiposity; infant growth

Introduction

The prevalence of childhood obesity is 16.9% in the United States according to the 2011–2012 National Health and Nutrition Examination Survey data¹. Weight gain, particularly in early infancy² is strongly associated with the risk of obesity in childhood and later life as demonstrated in several studies and multiple systematic reviews^{2–5}. Infants born small are more likely to gain weight quickly early in infancy as compared to those with higher birth weights, and this is associated with higher adiposity later in life³. Furthermore, rapid weight gain in the first year of life is associated with cardiovascular and metabolic dysregulation in adulthood⁶, and may be observed as early as 3 years of age⁷. Thus, growth in early infancy (<1 year) is an important influence on the future health of a child.

Nutrient exposure *in utero* influences infant growth after birth. Donahue *et al.*⁸ reported that high neonatal levels of the n-3 long chain polyunsaturated fatty acids (LCPUFA) docosahexanoate (DHA) and eicosapentanoate (EPA) predict a lower risk for obesity and adiposity at 3 years of age, as measured by skinfold thickness. Interestingly, maternal n-3 levels were not associated with infant adiposity⁸. Furthermore, n-3 LCPUFA supplementation during pregnancy and lactation in a randomized controlled trial, did not affect adiposity at 1 year of age⁹, suggesting that maternal fatty acid intake is not the primary determinant of infant adipose accumulation. The fetus – unlike the adult – has limited capacity to synthesize LCPUFA in adequate amounts to meet its developmental needs¹⁰, and thus depends upon both maternal supply and placental transport for acquiring LCPUFA.

In order to further investigate the association between the fatty acid profile in fetal blood and early infant adiposity, we utilized data from a cohort of women recruited during pregnancy and followed to term whereupon the placenta was collected and neonatal anthropometry were measured. Mother-infant pairs were followed to 6 months post-partum. We assessed the relationships between the change in body mass index (BMI) z-score in the first 6 months of life, and maternal and umbilical venous plasma fatty acid profiles. We hypothesized that infants with a greater positive change in BMI z-score would be physiologically different from those that had a smaller, or negative change. We predicted that their fatty acid profiles at birth would be different as would their placental lipid transporter gene expression levels.

Subjects and Methods

Subject recruitment and procedures

All aspects of this study were approved by the Oregon Health & Science University Institutional Review Board. Fifty women 18 years of age in their third trimester (26

weeks of pregnancy) who were patients at Cascades East Family Practice Center or Klamath Medical Clinic in Klamath Falls Oregon were recruited based upon screening at prenatal visits. Klamath Falls is a rural city of ~20,000 people on the Oregon-California border. Exclusion criteria were: women carrying multiple fetuses; fetuses with chromosomal anomalies; non-English speaking women.

Potential study participants from Cascades East Family Practice Center or Klamath Medical Clinic were identified based upon screening at prenatal visits and were asked to sign a preconsent document by clinic staff allowing the research nurse to contact them. Women were then scheduled for an initial clinical visit, where the overall purpose of the study and expectations were explained and informed consent was obtained.

Data and tissue collection

At the initial clinic visit, study participants were interviewed about their birth history, physical measurements were taken and venous blood was collected. Study participants were given a \$20 gift certificate for participating in the study.

Placental tissue and umbilical venous blood was collected at delivery. Placental weight (untrimmed), major and minor axial dimensions and thickness were measured and recorded. Placental biopsies were collected from four different cotyledons, avoiding calcified and underperfused areas. Maternal decidua was removed. Samples were snap frozen in liquid nitrogen and stored at -80°C . Birth weight, gestational age at birth and fetal sex were recorded. Umbilical venous blood was collected into an EDTA-treated tube and placed on ice until centrifuged at 1500g for 10 minutes at 4°C to isolate the plasma. Plasma was stored at -20°C .

The morning after delivery, the study nurse obtained physical measurements of the infant including length (crown-rump and crown-heel), chest, abdominal, arm and head circumferences and skinfold thicknesses (triceps, subscapular, thigh). Each measurement was taken three times to ensure accuracy. Six months after delivery, subjects were asked to come back to the clinic with their baby for another visit. The study nurse obtained additional physical measurements of the infants including length (crown-rump and crown-heel), chest, abdominal, arm and head circumferences and skinfold thicknesses (triceps, biceps, suprailiac, subscapular, thigh). Each measurement was taken three times to ensure accuracy. Mothers were also asked at this time about breast-feeding or bottle feeding practices.

We calculated age- and sex-specific weight-for-age *z*-score (WFAz), length-for-age *z*-score (LFAz), weight-for-length *z*-score (WFLz) and body mass index *z*-score (BMIz) from the World Health Organization 2006 growth chart data as recommended by the Center for Disease Control and Prevention for children in the U.S. ages 0 to 2 years (http://www.cdc.gov/growthcharts/who_charts.htm). Sum of skinfolds (triceps + subscapular + thigh) was calculated for neonates and infants at 6 months as an estimate of overall adiposity; from these data, growth of skinfolds in the first 6 months after birth was calculated. Arm muscle area ($\text{AMA} = [\text{mid-upper arm circumference} - (\pi \times \text{triceps skinfold})]^2 / [4 \pi]$ with measures in cm) was calculated for neonates and infants at 6 months as an estimate of muscle mass¹¹.

Quantitative plasma fatty acid analysis

Fatty acids were extracted and quantified by the method of Lagerstedt, *et al.*¹² as previously published¹³. Deuterated free fatty acids were added to samples prior to extraction as internal standards. The fatty acid-pentafluorobenzyl (PFB) esters were analyzed by gas chromatography mass spectrophotometry on a Trace DSQ (Thermoelectron). Each fatty acid was matched to the deuterated internal standard closest in length and retention time. Peak area ratios of known amounts of standard fatty acids and the internal standards were used to generate calibration curves to quantify unknowns using Xcalibur software.

Placental molecular analyses

RNA was isolated using TriReagent (Sigma) following the manufacturer's protocol. RNA was reverse transcribed to cDNA using MultiScribe Reverse Transcriptase (Applied Biosystems, Carlsbad, CA), and PCR performed using the Stratagene Mx3005P Thermocycler (Agilent, Santa Clara, CA). PCR amplicons were detected by fluorescent detection of SYBR Green (Power SYBR Green Master Mix, Applied Biosystems). For each primer pair, standard curves, no template controls and unknowns were run in triplicate. Following cycling, the melt curve of the resulting amplicon was analyzed to ensure that a single product was detected. Quantification of mRNA was achieved using the respective standard curves with manufacturer's software (MxPro QPCR; Agilent). Values were expressed as a ratio of the gene of interest:GAPDH control in each sample. Gene-specific primers, shown in the Supplementary Table (Table S1), were designed for fatty acid transport proteins (FATP)-4, FATP-6; fatty acid translocase (CD36); fatty acid binding proteins (FABP)-3, FABPpm, lipoprotein lipase (LPL) and endothelial lipase (EL). Placental gene expression data was log transformed to obtain normal distribution for group comparisons.

Statistical Analysis

Spearman correlation (ρ) was used to examine the relationship between plasma lipids, placental gene expression and infant growth measures. Two sample t-statistics and corresponding descriptive methodology were used to examine differences in maternal, placental and neonatal measures and fatty acid concentrations between infants who were or were not followed-up at 6 months of age. A P value of less than 0.05 was considered statistically significant.

Results

Participant characteristics are shown in Table 1. Six month infant follow-up was performed on 55% of the initial cohort (n=27). Subjects included in the postnatal analyses were similar to those without postnatal assessments, except for maternal age at enrollment (Supplementary Table S2). None of the mothers in this cohort developed gestational diabetes, but six women were diagnosed with pre-eclampsia. All analyses were performed with and without these six women, and were not found to be affected, thus the women were retained in all analyses. At birth, neonates in this study were thinner compared to the WHO reference values with an average BMI-z of -0.17. At 6 months, (average BMI-z of 0.09), these infants were similar to WHO reference values. Table 1 shows the relationships

between participant characteristics and change in BMI-z score during infancy. Of the 12 women who were breast-feeding at 6 months, 2 reported exclusive breast-feeding; 24 out of 27 women reported introducing solids by 6 months. We did not detect significant associations between infant feeding practices and growth measures. We did not have sufficient numbers to thoroughly investigate sex differences.

There was no association between maternal characteristics (parity, smoking status, breast-feeding practices or body type (height, weight, weight gain during pregnancy)) and change in BMI-z from 0 to 6 months. Placental measurements were not related to BMI-z changes in early infancy. BMI for age z-score at birth showed a trend ($P=0.063$) for a negative relationship with change in BMI-z during infancy, but other neonatal characteristics were not associated (e.g. neonatal skinfolds: $r=0.2541$; $P=0.24$). Infant weight at 6 months, WFA-z, WFL-z and BMI-z were strongly positively related to change in BMI z-score from 0 to 6 months. A positive BMI-z change was associated with greater weight gain in the first 6 months of life and a greater change in WFA-z and WFL-z, but not with crown-heel growth.

Neonatal and 6 month infant anthropometry values and their relationship to changes in BMI-z between 0 and 6 months are shown in Table 2. Head, arm, chest and abdominal circumference at birth were not correlated with change in BMI-z in early infancy. At 6 months of age, arm circumference and arm muscle area, an indicator of muscle mass, were larger in infants that had a greater positive change in BMI-z in early infancy. Growth of arm circumference and muscle area was also strongly positively related to change in BMI-z from birth to 6 months (Figure 1). Skin fold thicknesses were not associated with BMI-z change, at any age.

Plasma fatty acid profiles at birth were associated with weight gain in infancy

Umbilical venous plasma fatty acid profiles, and their relationship with BMI-z change in early infancy, are shown in Table 3. Infants with a positive change in BMI-z had a fatty acid profile at birth with a high proportion of saturated and low proportion of polyunsaturated fatty acids relative to infants with a small, or negative change in BMI-z (Fig. 2E & 2F). Figure 2 shows that BMI-z score at 6 months were positively correlated with the proportion of saturated fatty acids and negatively correlated with polyunsaturated fatty acids at birth. There was no correlation between fatty acid profiles at birth and neonatal growth measurements (Fig. 2A & 2B).

Fatty acid profiles at birth were also associated with skinfold thickness at birth. The sum of skinfolds at birth (thigh + biceps + subscapular) was negatively correlated with umbilical venous total polyunsaturated fatty acid levels ($r= -0.48$, $P=0.039$), total n-6 PUFA ($r= -0.47$, $P=0.041$), total n-3 PUFA ($r= -0.46$, $P=0.046$) and the proportion of n-6 PUFA ($r= -0.58$, $P=0.009$) and total PUFA ($r= -0.52$, $P=0.021$) in plasma. Neonatal skinfold thickness was positively correlated with the proportion of saturated fatty acids at birth ($r=0.49$, $P=0.034$). We were unable to detect any associations between fatty acid profiles at birth and infant skinfolds or skinfold growth in the first 6 months of life. We did not detect any correlations between fatty acid profiles at birth and neonatal or infant arm muscle area or arm muscle area growth.

Maternal fatty acid profiles in mid-pregnancy were not associated with weight gain in early infancy or skinfold thickness at birth, 6 months or growth in early infancy.

Placental fatty acid transporter gene expression was associated with infant adiposity measures

Quantitative PCR methodology was used to assess the association between placental fatty acid transporter gene expression and infant adiposity measures. Fatty acid transporter or lipase gene expression was not associated with changes in BMI-z. However several fatty acid transporters were correlated with skinfold thickness at 6 months of age and skinfold growth between birth and 6 months. Fatty acid transport protein (FATP)-4 mRNA levels were positively correlated with skinfold thickness at 6 months of age and the change in skinfold thickness from birth to 6 months (Figure 3). The expression of FATP-6 and CD36 were negatively associated with skinfold thickness in early infancy. Placental plasma membrane fatty acid binding protein (FABPpm) expression was positively correlated with 6 month skinfolds ($r=0.51$, $P=0.04$), but not with skinfold growth ($r=0.46$, $P=0.06$). Lipoprotein lipase mRNA expression was negatively associated with skinfold growth in the first 6 months of life ($r=-0.50$, $P=0.04$). We were unable to detect any statistically significant associations between placental fatty acid transporter/lipase gene expression and neonatal skinfolds, neonatal or infant arm muscle area, weight-for-length, BMI-z or umbilical cord fatty acid levels.

Discussion

A higher rate of weight gain in early infancy is associated with obesity in later life^{3,4}. We studied infants with negative and positive changes in BMI z-scores (BMI-z) within the first 6 months of life to better understand the nutritional factors associated with growth patterns representing slower and more rapid weight gain (and lower and higher risks for obesity), respectively. The main finding of our study was that a high proportion of saturated fatty acids and low DHA levels in umbilical vein plasma at term were associated with high skinfold thickness at birth, high infant BMI-z at 6 months of age and a positive change in BMI z-score in early infancy. Placental fatty acid transporter expression was correlated with skinfold growth in early infancy, a marker of fat deposition. Thus, fatty acid profiles at birth may be predictive of early infant growth and risk for obesity later in life; this may be related to differences in lipid transport function in the placenta.

We found that infants with a greater positive change in BMI-z in the first 6 months of life had a high proportion of saturated and low proportion of polyunsaturated plasma fatty acids at birth compared to those with a smaller or negative change in BMI-z over this time period. Low levels of n-3 fatty acids (DHA+EPA) in cord blood were found to be associated with higher adiposity at 3 years of age⁸ and rapid growth in the first 6 months related to obesity at both 3 years² and 6–11 years¹⁴ of age. We found a trend ($P=0.068$) for a lower proportion of n-3 LCPUFA in infants with greater BMI-z changes in the first 6 months. Together these findings suggest that low LCPUFA levels at birth may indicate a propensity for obesity later in life, perhaps through its association with higher rates of weight gain in early infancy.

Rapid growth within 6 months of birth may be related to increased fat deposition^{14,15}, which may continue to offer future obesity risk^{2,14}. However, changes in BMI z-scores during early development are not an accurate measure of adiposity¹⁶. Several authors have shown that early, even rapid, weight gain is positively associated with lean mass growth^{17,18} challenging the notion that lean mass tracks directly from fetal life to childhood and beyond¹⁹. However, as neonates are likely born with their full complement of muscle fibers²⁰, we speculate that increases in lean mass are due to an increase in skeletal muscle fiber size, not number. Thus, early infancy may be a time when myocyte enlargement occurs at varying rates among individuals – whether or not these changes are permanent is unknown. We found that early rapid weight gain was associated with increased arm muscle area growth – a marker of increased lean mass – rather than skinfold thickness growth. Importantly, those infants with a negative change in BMI-z during the first 6 months of life lost arm muscle area, indicating a loss in lean mass relative to body size, but continued to gain fat mass (as assessed by skinfold thickness), which could have negative connotations for future metabolic health²¹. These findings highlight the importance of assessing body composition, in addition to standard weight and length measures, during early growth studies in order to better understand potential long-term consequences of differences in rate of growth.

Early rapid weight gain is often associated with low birth weight – known as “catch-up growth”²². In our study, infants with a positive BMI-z change tended to have a low BMI-z at birth ($P=0.06$), suggesting they were thin at birth. However, consistent with the findings of Holzhauser *et al.*¹⁵, we observed that these two growth patterns (negative vs positive changes in BMI-z) were not related to birth weight. Thus, neonatal body composition is likely a better predictor of early growth patterns than birth weight. Weight gain over the first 6 months was positively associated with change in BMI-z, but change in skinfold thickness was not. As skinfolds represent subcutaneous fat, it is possible that intraabdominal fat deposition or soft tissue growth was greater in infants that had a larger positive change in BMI-z. Though absolute crown-heel growth was not related to BMI-z changes, we noted that the change in length-for-age z-score tended to be lower in infants who gained weight quickly ($P=0.09$). Less length growth (accounting for age and sex) and higher arm muscle area growth demonstrates a different pattern of lean mass distribution in infants that gain weight quickly. Future studies should assess the metabolic efficiency of lean mass in individuals with different distributions of fat and muscle. Thus, a high proportion of saturated fatty acids combined with low PUFA levels in cord plasma are associated with an early rapid weight gain, a positive change in BMI-z in the first 6 months, high BMI-z at 6 months and an overall greater lean mass.

Umbilical venous fatty acid profiles are affected by maternal fatty acid supply, placental delivery, and fetal lipid metabolism. Consistent with previous findings⁸, we did not see an association between maternal fatty acid profiles and neonatal or infant growth measures in this study. We found that gene expression levels of placental fatty acid transporters such as FAT/CD36, FATP-4 and FATP-6 were correlated with the growth of skinfolds from birth to 6 months of age, suggesting that early fat deposition and infant adiposity are related to placental lipid handling. As we only measured mRNA expression of these transporters, we cannot say whether the ability of the placenta to deliver fatty acids from mother to the fetus was related to early infant growth. However, umbilical venous, but not maternal, plasma

fatty acid levels were correlated with infant BMI-z, supporting the notion that placental lipid handling pathways were altered. Maternal fatty acids were measured at mid-pregnancy rather than at birth, therefore they may not accurately represent the maternal fatty acid supply at term.

Changes in placental fatty acid transporter gene expression have previously been shown to be associated with alterations in placental fatty acid uptake and transport to the fetus^{13,23–25}. Interestingly, placental CD36 and FATP-6 levels were negatively related to infant adiposity while FATP-4 levels were positively correlated to skinfold growth. The specific roles of each type and isoform of fatty acid transporter in the placenta are not well-understood. Placental FATP-4 expression correlates with cord blood DHA levels²⁶, while CD36 is thought to be a relatively non-specific transporter²⁷. Due to the cross-sectional nature of the gene expression data, we cannot speculate on whether the placental gene expression is representative of transporter activity at birth or reflects an upregulation/downregulation of expression in response to fetal fatty acid requirements and maternal supply.

Post-partum factors such as length of exclusive breast-feeding has been found to influence infant adiposity and risk of childhood obesity in a recent meta-analysis²⁸ but may not be protective long-term²⁹. We did not observe any associations between length of breast-feeding and weight gain trajectories, though we did not measure volume of milk or solid intake, which may confound these relationships. Our study was limited by collecting non-fasting maternal blood samples, which introduces variability into the lipid data. Also, by nature, this pilot study included a small number of subjects and we were only able to follow-up with 55% for infant growth measures. This affected our capacity to control for confounding variables and so we used conservative Spearman correlation analysis rather than multiple linear regression. Despite this limitation, we found strong relationships, many of which were consistent with findings of larger cohort studies.

This study is the first to link placental fatty acid transporter expression along with neonatal fatty acid profiles at birth to infant growth measures and adiposity. Our findings suggest that early infancy weight gain is associated with poor plasma LCPUFA status at birth and that fat deposition in infancy is related to changes in placental lipid handling. We speculate that neonatal fatty acid profiles – established by placental transport and maternal nutrition – determine the trajectory for infant growth and fat and lean mass deposition.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

None

Financial Support

This research was supported by the NIH-NICHD (R00HD062841 to PFOG), Edwards Endowment (KLT), Northwest Health Foundation, Holtzman Foundation, Audrey Love Foundation and Eagles Trust Foundation.

Reference List

1. Ogden CL, Carroll MD, Kit BK, Flegal KM. Prevalence of childhood and adult obesity in the united states, 2011–2012. *Jama*. 2014; 311:806–814. [PubMed: 24570244]
2. Taveras EM, Rifas-Shiman SL, Belfort MB, Kleinman KP, Oken E, Gillman MW. Weight status in the first 6 months of life and obesity at 3 years of age. *Pediatrics*. 2009; 123:1177–1183. [PubMed: 19336378]
3. Baird J, Fisher D, Lucas P, Kleijnen J, Roberts H, Law C. Being big or growing fast: Systematic review of size and growth in infancy and later obesity. *Bmj*. 2005; 331:929. [PubMed: 16227306]
4. Ong KK, Loos RJ. Rapid infancy weight gain and subsequent obesity: Systematic reviews and hopeful suggestions. *Acta paediatrica (Oslo, Norway: 1992)*. 2006; 95:904–908.
5. Monteiro PO, Victora CG. Rapid growth in infancy and childhood and obesity in later life—a systematic review. *Obesity reviews: an official journal of the International Association for the Study of Obesity*. 2005; 6:143–154. [PubMed: 15836465]
6. Leunissen RW, Kerkhof GF, Stijnen T, Hokken-Koelega A. Timing and tempo of first-year rapid growth in relation to cardiovascular and metabolic risk profile in early adulthood. *Jama*. 2009; 301:2234–2242. [PubMed: 19491185]
7. Belfort MB, Rifas-Shiman SL, Rich-Edwards J, Kleinman KP, Gillman MW. Size at birth, infant growth, and blood pressure at three years of age. *The Journal of pediatrics*. 2007; 151:670–674. [PubMed: 18035150]
8. Donahue SM, Rifas-Shiman SL, Gold DR, Jouni ZE, Gillman MW, Oken E. Prenatal fatty acid status and child adiposity at age 3 y: Results from a us pregnancy cohort. *The American journal of clinical nutrition*. 2011; 93:780–788. [PubMed: 21310834]
9. Hauner H, Much D, Vollhardt C, Brunner S, Schmid D, Sedlmeier EM, Heimberg E, Schuster T, Zimmermann A, Schneider KT, Bader BL, Amann-Gassner U. Effect of reducing the n-6:N-3 long-chain pufa ratio during pregnancy and lactation on infant adipose tissue growth within the first year of life: An open-label randomized controlled trial. *The American journal of clinical nutrition*. 2012; 95:383–394. [PubMed: 22205307]
10. Salem N Jr, Wegher B, Mena P, Uauy R. Arachidonic and docosaheaxaenoic acids are biosynthesized from their 18-carbon precursors in human infants. *Proc Natl Acad Sci USA*. 1996; 93:49–54. [PubMed: 8552667]
11. O'Tierney PF, Lewis RM, McWeeney SK, Hanson MA, Inskip HM, Morgan TK, Barker DJ, Bagby G, Cooper C, Godfrey KM, Thornburg KL. Immune response gene profiles in the term placenta depend upon maternal muscle mass. *Reprod Sci*. 2012; 19:1041–1056. [PubMed: 22534332]
12. Lagerstedt SA, Hinrichs DR, Batt SM, Magera MJ, Rinaldo P, McConnell JP. Quantitative determination of plasma c8-c26 total fatty acids for the biochemical diagnosis of nutritional and metabolic disorders. *Mol Genet Metab*. 2001; 73:38–45. [PubMed: 11350181]
13. O'Tierney-Ginn P, Roberts V, Gillingham M, Walker J, Glazebrook PA, Thornburg KL, Grove K, Frias AE. Influence of high fat diet and resveratrol supplementation on placental fatty acid uptake in the japanese macaque. *Placenta*. 2015; 36:903–910. [PubMed: 26145226]
14. Koontz MB, Gunzler DD, Presley L, Catalano PM. Longitudinal changes in infant body composition: Association with childhood obesity. *Pediatric obesity*. 2014; 9:e141–144. [PubMed: 25267097]
15. Holzhauser S, Hokken Koelega AC, Ridder M, Hofman A, Moll HA, Steegers EA, Witteman JC, Jaddoe VW. Effect of birth weight and postnatal weight gain on body composition in early infancy: The generation r study. *Early human development*. 2009; 85:285–290. [PubMed: 19091495]
16. Andres A, Hull HR, Shankar K, Casey PH, Cleves MA, Badger TM. Longitudinal body composition of children born to mothers with normal weight, overweight, and obesity. *Obesity (Silver Spring, Md)*. 2015; 23:1252–1258.
17. Brunner S, Schmid D, Huttinger K, Much D, Bruderl M, Sedlmeier EM, Kratzsch J, Amann-Gassner U, Bader BL, Hauner H. Effect of reducing the n-6/n-3 fatty acid ratio on the maternal

and fetal leptin axis in relation to infant body composition. *Obesity* (Silver Spring, Md). 2014; 22:217–224.

18. Chandler-Laney PC, Gower BA, Fields DA. Gestational and early life influences on infant body composition at 1 year. *Obesity* (Silver Spring, Md). 2013; 21:144–148.
19. Baker J, Workman M, Bedrick E, Frey MA, Hurtado M, Pearson O. Brains versus brawn: An empirical test of barker's brain sparing model. *Am J Hum Biol*. 2010; 22:206–215. [PubMed: 19701887]
20. Rowe RW, Goldspink G. Muscle fibre growth in five different muscles in both sexes of mice. *Journal of anatomy*. 1969; 104:519–530. [PubMed: 5804561]
21. Joglekar CV, Fall CH, Deshpande VU, Joshi N, Bhalerao A, Solat V, Deokar TM, Chougule SD, Leary SD, Osmond C, Yajnik CS. Newborn size, infant and childhood growth, and body composition and cardiovascular disease risk factors at the age of 6 years: The pune maternal nutrition study. *International journal of obesity*. 2007; 31:1534–1544. (2005). [PubMed: 17653070]
22. Eriksson JG, Forsen T, Tuomilehto J, Winter PD, Osmond C, Barker DJ. Catch-up growth in childhood and death from coronary heart disease: Longitudinal study. *Bmj*. 1999; 318:427–431. [PubMed: 9974455]
23. Xu Y, Agrawal S, Cook TJ, Knipp GT. Maternal di-(2-ethylhexyl)-phthalate exposure influences essential fatty acid homeostasis in rat placenta. *Placenta*. 2008; 29:962–969. [PubMed: 18829104]
24. Dube E, Gravel A, Martin C, Desparois G, Moussa I, Ethier-Chiasson M, Forest JC, Giguere Y, Masse A, Lafond J. Modulation of fatty acid transport and metabolism by maternal obesity in the human full-term placenta. *Biology of reproduction*. 2012; 87:14, 11–11. [PubMed: 22553224]
25. Brass E, Hanson E, O'Tierney-Ginn PF. Placental oleic acid uptake is lower in male offspring of obese women. *Placenta*. 2013; 34:503–509. [PubMed: 23602336]
26. Larque E, Krauss-Etschmann S, Campoy C, Hartl D, Linde J, Klingler M, Demmelmair H, Cano A, Gil A, Bondy B, Koletzko B. Docosahexaenoic acid supply in pregnancy affects placental expression of fatty acid transport proteins. *Am J Clin Nutr*. 2006; 84:853–861. [PubMed: 17023713]
27. Duttaroy AK. Transport of fatty acids across the human placenta: A review. *ProgLipid Res*. 2009; 48:52–61.
28. Yan J, Liu L, Zhu Y, Huang G, Wang PP. The association between breastfeeding and childhood obesity: A meta-analysis. *BMC public health*. 2014; 14:1267. [PubMed: 25495402]
29. O'Tierney PF, Barker DJ, Osmond C, Kajantie E, Eriksson JG. Duration of breast-feeding and adiposity in adult life. *The Journal of nutrition*. 2009; 139:422s–425s. [PubMed: 19106318]

Figure 1.

Relationships between infant growth measures and adiposity over the first 6 months of life.

A greater change in BMI z-score from birth to 6 months of age is correlated with a greater arm (A) and arm muscle area growth (B). N=18. *Spearman correlation coefficient (ρ). No differences in relationships were noted between male and female fetuses.

Figure 2. Relationship between umbilical venous fatty acid profiles and infant adiposity. BMI z-score at birth is not associated with umbilical venous (UV) plasma fatty acid profiles (A&B). A high BMI z-score at 6 months of age, and greater change in BMI z-score from birth to 6 months of age, is associated with a high proportion of saturated fatty acids (C&E) and a low proportion of polyunsaturated fatty acids (D&F) in umbilical venous plasma at birth. $N=22$. *Spearman correlation coefficient (ρ). No differences in relationships were noted between male and female fetuses.

Figure 3. Relationships between placental fatty acid transporter mRNA levels and growth of skinfolds. Placental fatty acid translocase (FAT/CD36) and fatty acid transport protein (FATP)-6 were negatively correlated with skinfold growth in early infancy (A&B). Placental FATP-4 was positively correlated with skinfold growth (C). $N=17$. *Spearman correlation coefficient (ρ). No differences in relationships were noted between male and female fetuses.

Maternal and placental characteristics and growth measures of infants and relationship with changes in BMI z-score between 0–6 months.

Table 1

		N	Overall	Correlation with change in BMI z-score birth-6 months		
				N	Spearman R	P-value
Maternal						
	Age (y)	44	26 ± 5	23	0.2176	n.s.
	Primiparous Yes/No (%)	49	14/35 (29%)	24	—	n.s.
	Height (m)	45	1.67 ± 0.08	24	0.2162	n.s.
	1 st Trimester Weight (kg)	41	74.1 ± 18.4	22	0.2565	n.s.
	1 st Trimester BMI (kg/m ²)	41	27.0 ± 6.7	20	0.2635	n.s.
	Total Weight Gain (kg)	40	11.4 ± 5.4	18	−0.1569	n.s.
	Currently smoking/non-smoking (%)	45	15/30 (33%)	24	—	n.s.
	Any breast-feeding Yes/No (%)	27	26/1 (96%)	24	—	n.s.
	Breast-feeding at 6 months Yes/No (%)	27	12/15 (44%)	24	—	n.s.
Placenta						
	Untrimmed weight (g)	47	776 ± 165	23	−0.0711	n.s.
	Width (cm)	47	16.8 ± 1.9	23	−0.1046	n.s.
	Length (cm)	47	19.5 ± 2.2	23	−0.2827	n.s.
	Aspect Ratio	47	1.2 ± 0.1	23	−0.2234	n.s.
Neonate						
	Gestational Age at birth (wk)	48	39.4 ± 1.2	24	−0.0192	n.s.
	Males/Females (% Male)	48	21/27 (44%)	24	—	n.s.
	Birth weight (kg)	48	3.5 ± 0.4	24	−0.1857	n.s.
	Crown-heel length (cm)	45	51.4 ± 3.0	24	0.1729	n.s.
	Weight-for-age z-score	48	0.37 ± 0.87	24	−0.1661	n.s.
	Length-for-age z-score	45	1.04 ± 1.57	24	0.1573	n.s.
	Weight-for-length z-score	44	−0.70 ± 1.59	23	−0.3034	n.s.
	BMI-for-age z-score	45	−0.17 ± 1.20	24	−0.3853	0.063
6 Month Infant						
	Weight at 6 months (kg)	27	7.6 ± 0.9	24	0.4765	0.0186

	N	Overall	Correlation with change in BMI z-score birth-6 months		
			N	Spearman R	P-value
Maternal					
	26	66.5 ± 3.4	24	-0.2000	n.s.
	27	-0.17 ± 0.99	24	0.4228	0.0396
	26	-0.28 ± 1.55	24	-0.2397	n.s.
	26	0.23 ± 1.04	24	0.7432	<0.0001
	26	0.09 ± 0.96	24	0.7458	<0.0001
Growth (0-6 mo)					
	27	4.2 ± 0.8	24	0.4252	0.0383
	23	17.2 ± 4.9	23	-0.1533	n.s.
	27	-0.45 ± 1.03	24	0.4136	0.0446
	24	-1.18 ± 1.95	24	-0.3514	n.s.
	23	0.98 ± 1.48	23	0.7421	0.0001
	24	0.24 ± 1.29	24	-----	-----

/ Data are mean ± SD unless otherwise noted. Relationship between change of BMI-z and categorical variables assessed via Kruskal-Wallis rank t-test.

Neonatal and infant anthropometry and growth of infants and relationship with changes in BMI z-score between 0–6 months.[†]

Table 2

Neonate	N	Overall	Correlation with change in BMI z-score birth-6 months		
			N	Spearman R	P value
Head circumference	44	35 ± 3	24	0.0491	n.s.
Arm circumference	44	10 ± 3	24	-0.1612	n.s.
Chest circumference	44	34 ± 6	24	0.0724	n.s.
Abdominal circumference	44	35 ± 4	24	0.3418	n.s.
Sum of skinfolds (mm)	43	12.7 ± 4.3	23	0.1160	n.s.
Arm muscle area (cm ²)	43	6.7 ± 3.8	23	-0.2260	n.s.
6 Month Infant					
Head circumference	24	43 ± 2	23	-0.0562	n.s.
Arm circumference	22	14 ± 3	21	0.6595	0.0011
Chest circumference	22	44 ± 4	21	0.2905	n.s.
Abdominal circumference	22	46 ± 4	21	0.2413	n.s.
Sum of skinfolds (mm)	21	24.9 ± 5.4	20	0.2216	n.s.
Arm muscle area (cm ²)	20	10.9 ± 6.7	19	0.5994	0.0067
Growth (0–6 mo)					
Head circumference	23	8 ± 4	23	0.0909	n.s.
Arm circumference	21	3.3 ± 5	21	0.6137	0.0031
Chest circumference	21	10 ± 8	21	0.1504	n.s.
Abdominal circumference	21	11 ± 4	21	-0.0593	n.s.
Sum of skinfolds (mm)	19	11.6 ± 7.0	19	0.1916	n.s.
Arm muscle area (cm ²)	18	3.6 ± 9.3	18	0.6677	0.0025

[†]Measures are in cm unless otherwise noted. Data are mean ± SD.

Table 3

Umbilical venous fatty acid profiles of infants and relationship with changes in BMI z-score between 0–6 months.¹

	N	Overall	Correlation with change in BMI z-score birth-6 months		
			N	Spearman R	P value
Myristic Acid, 14:0	41	233 ± 372	22	0.0932	n.s.
Palmitic Acid, 16:0	41	1941 ± 1407	22	0.2526	n.s.
Stearic Acid, 18:0	41	461 ± 255	22	0.1323	n.s.
Total SFA	41	2635 ± 2010	22	0.2331	n.s.
Myristoleic Acid, 14:1	41	4 ± 6	22	0.0286	n.s.
Palmitoleic Acid, 16:1	41	221 ± 125	22	0.0211	n.s.
Oleic Acid, 18:1	41	680 ± 1031	22	-0.0947	n.s.
Total MUFA	41	905 ± 1152	22	-0.0782	n.s.
Linoleic Acid, 18:2	41	373 ± 877	22	-0.0135	n.s.
Arachidonic Acid, 20:4	41	710 ± 431	22	-0.3609	n.s.
Total n-6	41	1083 ± 1027	22	-0.1248	n.s.
α-Linolenic Acid, 18:3	41	12 ± 25	22	0.0271	n.s.
Eicosapentanoic Acid, 20:5	41	35 ± 6	22	-0.1046	n.s.
Docosahexanoic Acid, 22:6	41	111 ± 60	22	-0.2090	n.s.
Total n-3	41	158 ± 74	22	-0.1534	n.s.
Total PUFA	41	1241 ± 1086	22	-0.1323	n.s.
n-6/n-3	41	6.1 ± 2.9	22	-0.2527	n.s.
Total FA	41	4780 ± 4117	22	0.0120	n.s.
%SFA	41	58 ± 10	22	0.5083	0.0221
%MUFA	41	17 ± 3	22	-0.1023	n.s.
%n-6	41	22 ± 8	22	-0.4979	0.0255
%n-3	41	4 ± 1	22	-0.4159	0.068
%PUFA	41	25 ± 8	22	-0.5669	0.0091

¹Data is μM unless otherwise noted. Data are mean ± SD.