Protein–nucleic acids interactions: new ways of connecting structure, dynamics and function

Maria Spies¹ • Brian O. Smith²

Received: 16 July 2017 / Accepted: 19 July 2017 / Published online: 4 August 2017

© International Union for Pure and Applied Biophysics (IUPAB) and Springer-Verlag GmbH Germany 2017

Molecular machines that act on nucleic acids, DNA and RNA are at the heart of the field of cellular information processing. A coherent description of the interactions involved in their assembly, activities and regulation affords a quantitative understanding of how transcription factors and DNA repair proteins find their unique targets among millions of nonspecific sequences and undamaged DNA bases, how the intricate choreography of DNA replication, recombination and repair and gene expression is regulated, how viral particles self-assemble and how chromosomes are organized inside living cells. These important questions are not easy to answer for the following reasons: (1) transactions between proteins and nucleic acids commonly involve extended surfaces with multiple interaction epitopes, and the resulting macromolecular assemblies are non-homogenous and dynamic; (2) the structures of multicomponent protein–DNA and protein–RNA complexes are often refractory to analysis by traditional X-ray crystallography and nuclear magnetic resonance (NMR). New techniques, clever adaptations and combinations of the state-of-the-art approaches are therefore needed.

Our selection of speakers cover exciting new developments, both technological and conceptual, in determining the structures of protein–DNA and protein–RNA

complexes, as well as in connecting the structure, dynamics and activity of these complexes using methods of single-molecule biophysics. While an understanding of nucleoprotein architecture at atomic resolution and how it changes in response to regulatory events can be achieved using structural biology techniques, single-molecule techniques allow us to visualize and interrogate biologically important molecules individually, in real time. For both structural biology and single-molecule techniques, the ability to validate observations made in vitro under physiological conditions is both more important and more feasible (Sakakibara et al. 2009; Ito et al. 2012; Hamatsu et al. 2013). The ability to observe and directly manipulate individual protein and nucleic acid molecules provides further unprecedented insight into the kinetics of the molecular recognition steps, which are often obscured in bulk biochemical and biophysical studies (Cornish and Ha 2007; Boehm et al. 2016). During the last decade or so, technological advances in structural biology and single-molecule biophysics have brought new vibrancy and excitement to the field of nucleoprotein interactions, which we hope we capture in the selection of this session's topics.

The three invited talks and the three oral presentations selected for this session exemplify the breadth of excellence at the frontiers of structural and single-molecule research:

The ParB protein of *Bacillus subtilis* is the subject of Fernando Moreno-Herrero's talk. Fernando Moreno-Herrero (Centro Nacional de Biotecnología, Spain) has been leading the field in atomic force microscopy studies of protein–nucleic acid interactions (Fuentes-Perez et al. 2013). ParB condenses and bridges DNA molecules in the context of bacterial chromosome organization and dynamics, which the Moreno-Herrero laboratory investigates in

This article is part of a Special Issue on 'IUPAB Edinburgh Congress' edited by Damien Hall.

✉ Brian O. Smith
Brian.Smith@glasgow.ac.uk

¹ Carver College of Medicine, University of Iowa, 51 Newton Rd., 4-532 BSB, Iowa City, IA 52242, USA

² Institute of Molecular, Cell and Systems Biology, College of Medical, Veterinary & Life Sciences, University of Glasgow, Joseph Black Building, Glasgow G12 8QQ, UK

collaboration with the Dillingham (University of Bristol), Crump (University of Bristol), Sobbot (University of Antwerp) and Murray (University of Newcastle) groups. In his talk, Fernando Moreno-Herrero shows that the C-terminal domain of *B. subtilis* ParB is critical for dynamic DNA binding and bridging using magnetic tweezers (MT) and a combined MT–total internal reflection microscopy setup (Taylor et al. 2015).

The CRISPR–Cas system is currently one of the hottest topics in biology due to its rapid adoption as the genome editing tool of choice, but there is still plenty to discover about its mechanism, and this knowledge will inform the development of more faithful and less promiscuous tools (Spies 2014). Ralf Seidel (Leipzig University, Germany) describes work using MT (Daldrop et al. 2015) to interrogate the mechanism of target recognition by the CRISPR–Cas system. By applying force and torque to the ends of the DNA molecule, his group were able to probe the binding affinity of Cascade for DNA targets with differently positioned imperfections (Szczelkun et al. 2014; Rutkauskas et al. 2017) and obtain a detailed energy profile for Cas9-mediated R-loop formation.

Metabolic disorders that originate from mitochondrial defects have been the subject of much public interest in recent years, and Hanna Yuan (Academia Sinica, Taiwan) presents her group's latest work on endonuclease G (EndoG), which digests paternal mitochondrial DNA during embryogenesis. EndoG also plays a role in mature mitochondria integrating DNA replication and repair, apoptosis and response to oxidative stress. Using X-ray crystallography and studies in *Caenorhabditis elegans*, the Yuan group reveals how EndoG degrades DNA without sequence specificity and shows how EndoG acts as a sensor of reactive oxygen species, thereby presenting new avenues for the prevention and treatment of the diseases that result from oxidative stresses (Lin et al. 2016; Zhou et al. 2016).

Electron paramagnetic resonance (EPR) has recently undergone a renaissance in the field of biomolecular structural biology, and Christoph Gmeiner (ETH Zurich, Switzerland) describes how the long-range distance restraints produced by combining site-directed spin labelling with pulsed EPR can be used in conjunction with short-range NMR restraints to determine the three-dimensional structure of the PTBP1/EMCV protein–RNA complex (Duss et al. 2014, 2015).

NMR spectroscopy remains the only technique that can simultaneously provide detailed structural and dynamic information on biomolecules in solution, and Bruno Kieffer (IGBMC, Illkirch-Graffenstaden, France) presents work that reveals the complex allosteric mechanism by which post-translational phosphorylation regulates the retinoic acid receptor's recognition of target DNA allosterically to control gene expression (Martinez-Zapien et al. 2014; Belorusova et al. 2016).

The assembly of virus particles is of fundamental interest and potentially a process that can be targeted to disrupt the spread of disease causing viral pathogens. Margherita Marchetti (Vrije Universiteit Amsterdam, the Netherlands) presents an elegant single-molecule study where a combination of optical trapping and confocal microscopy (Hashemi Shabestari et al. 2017) was used to visualize the self-assembly of artificial virus-like particles.

Compliance with ethical standards

Conflict of interest Maria Spies and Brian O. Smith declare that they have no conflicts of interest to declare.

References

- Belorusova AY, Osz J, Petoukhov MV, Peluso-Itlis C, Kieffer B, Svergun DI, Rochel N (2016) Solution behavior of the intrinsically disordered N-terminal domain of retinoid X receptor alpha in the context of the full-length protein. *Biochemistry* 55(12):1741–1748
- Boehm EM, Subramanyam S, Ghoneim M, Washington MT, Spies M (2016) Quantifying the assembly of multicomponent molecular machines by single-molecule Total internal reflection fluorescence microscopy. *Methods Enzymol* 581:105–145
- Cornish PV, Ha T (2007) A survey of single-molecule techniques in chemical biology. *ACS Chem Biol* 2(1):53–61
- Daldrop P, Brutzer H, Huhle A, Kauert DJ, Seidel R (2015) Extending the range for force calibration in magnetic tweezers. *Biophys J* 108(10):2550–2561
- Duss O, Yulikov M, Allain FH, Jeschke G (2015) Combining NMR and EPR to determine structures of large RNAs and protein–RNA complexes in solution. *Methods Enzymol* 558:279–331
- Duss O, Yulikov M, Jeschke G, Allain FH (2014) EPR-aided approach for solution structure determination of large RNAs or protein–RNA complexes. *Nat Commun* 5:3669
- Fuentes-Perez ME, Dillingham MS, Moreno-Herrero F (2013) AFM volumetric methods for the characterization of proteins and nucleic acids. *Methods* 60(2):113–121
- Hamatsu J, O'Donovan D, Tanaka T, Shirai T, Hourai Y, Mikawa T, Ikeya T, Mishima M, Boucher W, Smith BO, Laue ED, Shirakawa M, Ito Y (2013) High-resolution heteronuclear multidimensional NMR of proteins in living insect cells using a baculovirus protein expression system. *J Am Chem Soc* 135(5):1688–1691
- Hashemi Shabestari M, Meijering AE, Roos WH, Wuite GJ, Peterman EJ (2017) Recent advances in biological single-molecule applications of optical tweezers and fluorescence microscopy. *Methods Enzymol* 582:85–119
- Ito Y, Mikawa T, Smith BO (2012) In-cell NMR of intrinsically disordered proteins in prokaryotic cells. *Methods Mol Biol* 895:19–31
- Lin JL, Wu CC, Yang WZ, Yuan HS (2016) Crystal structure of endonuclease G in complex with DNA reveals how it nonspecifically degrades DNA as a homodimer. *Nucleic Acids Res* 44(21):10480–10490
- Martinez-Zapien D, Delsuc MA, Trave G, Lutz R, Rochette-Egly C, Kieffer B (2014) Production and characterization of a retinoic acid receptor RARgamma construction encompassing the DNA binding domain and the disordered N-terminal proline rich domain. *Protein Expr Purif* 95:113–120
- Rutkauskas M, Krivoy A, Szczelkun MD, Rouillon C, Seidel R (2017) Single-molecule insight into target recognition by CRISPR–Cas complexes. *Methods Enzymol* 582:239–273

- Sakakibara D, Sasaki A, Ikeya T, Hamatsu J, Hanashima T, Mishima M, Yoshimasu M, Hayashi N, Mikawa T, Walchli M, Smith BO, Shirakawa M, Guntert P, Ito Y (2009) Protein structure determination in living cells by in-cell NMR spectroscopy. *Nature* 458(7234):102–105
- Spies M (2014) Fulfilling the dream of a perfect genome editing tool. *Proc Natl Acad Sci USA* 111(28):10029–10030
- Szczelkun MD, Tikhomirova MS, Sinkunas T, Gasiunas G, Karvelis T, Pschera P, Siksnys V, Seidel R (2014) Direct observation of R-loop formation by single RNA-guided Cas9 and Cascade effector complexes. *Proc Natl Acad Sci USA* 111(27):9798–9803
- Taylor JA, Pastrana CL, Butterer A, Pernstich C, Gwynn EJ, Sobott F, Moreno-Herrero F, Dillingham MS (2015) Specific and non-specific interactions of ParB with DNA: implications for chromosome segregation. *Nucleic Acids Res* 43(2):719–731
- Zhou Q, Li H, Li H, Nakagawa A, Lin JL, Lee ES, Harry BL, Skeen-Gaar RR, Suehiro Y, William D, Mitani S, Yuan HS, Kang BH, Xue D (2016) Mitochondrial endonuclease G mediates breakdown of paternal mitochondria upon fertilization. *Science* 353(6297):394–399