bmj.com news roundup

Full versions of these stories are available at: bmj.com/content/vol330/issue7502/#NEWS_ROUNDUP

All IVF embryos should be checked for genetic defects, conference is told

All embryos resulting from in vitro fertilisation (IVF) should be routinely checked for genetic abnormalities before implantation, a conference on human genetics heard this week. New evidence indicates that screening embryos vastly improves the chances of producing a healthy baby in women aged over 35 who undergo IVF.

Dr Yury Verlinsky, a leader in the field of preimplantation genetic diagnosis and director of the Reproductive Genetics Institute, Chicago, told the sixth international symposium on preimplantation genetic diagnosis in London that the findings of screenings should lead to only normal embryos being used in treatment.

In a study of 709 couples Dr Verlinsky's team found that the percentage of couples who were able to have a baby after preimplantation genetic diagnosis was 81%, compared with 11% of couples who didn't undergo genetic diagnosis. By screening embryos Dr Verlinsky ensured that only embryos with a good chance of success were implanted.

Dr James Watson, president of Cold Spring Harbor Laboratory, New York, who is best known for his part in the discovery of the structure of DNA, said that couples should be allowed to choose a child's sex. Rebecca Coombes *London*

Israel allows sex selection of embryos for non-medical reasons

Israeli parents who have at least four children of the same sex and want one of the other sex can now apply to a health ministry committee for approval of preimplantation genetic diagnosis at their own expense.

Professor Avi Yisraeli, director general of the health ministry, who issued the

New guide ranks success of IVF treatment centres

Couples seeking fertility treatment in the United Kingdom will be able to compare the success rates of different centres by using statistics published in a new directory from the Human Fertilisation and Embryology Authority (HFEA). The figures will also be available in an online database.

"This is the most comprehensive and detailed information published about UK infertility services, and we believe it's unique globally," said Suzi Leather, the HFEA's chairwoman.

Many infertile women are not receiving help early enough, the report says. The chances of infertile women conceiving successfully after in vitro fertilisation (IVF) declines steeply after the age of 35 (figure). The average success rate for IVF treatment using fresh eggs (rather than frozen embryos) is 27.6% for women under 35; this figure falls to 18.3% for women aged 38 or 39 and to 10% for women between 40 and 42.

Colleen Shannon *London*

The HFEA Guide to Infertility and Directory of Clinics and the online database are at www.hfea.gov.uk

directive on the recommendation of experts on bioethics, said the new seven member body would approve sex selection of embryos for social reasons only in very unusual cases.

Except for one case officially approved by the ministry, all procedures done at Israeli hospitals for preimplantation genetic diagnosis have, until now, involved a family history indicating a high risk of serious genetic disorders. Such disorders include Tay-Sachs disease and familial dysautonomia (both of which occur mostly in Jews), as well as thalassaemia, myotonic dystrophy, neurofibromatosis, fragile X

syndrome, haemophilia, and Marfan's syndrome.

Now the committee, which comprises experts in law, medical genetics, and obstetrics, a social worker, and a clergyman, will decide whether a couple or single woman can choose, for social reasons, the desired sex among the resultant embryos through in vitro fertilisation and preimplantation genetic diagnosis.

Israel's policies on in vitro fertilisation are among the world's most liberal, with fertility units in every general hospital and with the health funds covering the cost of producing two babies for any infertile woman under 45 who wants them.

Judy Siegel-Itzkovich *Jerusalem*

UK government to set up small claims scheme for medical mishaps

An NHS scheme to deal outside the legal process with "low monetary value" claims for clinical negligence will be brought forward in this parliamentary session, the government announced in the Queen's speech last week.

The redress scheme will be based largely on recommendations in the 2003 *Making Amends* report from the chief medical officer, Professor Liam Donaldson (*BMJ* 2003;327:7), although it may include independent scrutiny of claims and provision for patients to have specialist representation when applying for redress, neither of which was included in the original proposals.

The scheme, which would be overseen by the NHS Litigation Authority, could provide compensation without litigation for claims up to a value of £30 000 (\$55 000; €44 000), although the government could opt for a lower figure. Legal aid would probably not be available for low value claims, and claimants would instead be expected to go through the scheme.

Clare Dyer *legal correspondent, BMJ*

Statins may reduce the risk of colorectal cancer

Taking statins for at least five years reduced the relative risk of colorectal cancer by 47%, a large study has shown.

The retrospective, case controlled study was conducted in northern Israel by epidemiologists at the Technion-Israel Institute of Technology's Medical School and Clalit Health Services National Cancer Control Centre in Haifa and published in the latest issue of the *New England Journal of Medicine* (2005;352:2184-92).

The team, led by Professor Gad Rennert, head of the Clalit Health Services National Cancer Registry and Professor Stephen Gruber of the University of Michigan, compared 1953