patients with colorectal cancer and 2015 control patients without colorectal cancer. Use of statins for at least five years was reported by 120 patients (6.1%) in the cancer group and 234 patients (11.6%) in the control group (odds ratio 0.50 (95% confidence interval 0.40 to 0.63).
Judy Siegel-Itzkovich *Jerusalem*

UK agency reports slight increase in radiation exposure

The average annual exposure to radiation among the British public has risen slightly, according to the Health Protection Agency. Most of the increase is due to the greater number of people undergoing radiography and the increased use of computed tomography (CT) scanners, which carry higher levels of radiation than conventional radiography, it says.

Although the national average levels of radiation exposure increased only very slightly from 2002 to 2003—from 2.6 mSv to just under 2.7 mSv—radiation from medical interventions had increased by 10%, from 0.37 mSv to 0.41 mSv.

"This is due to increases in the frequency of x rays carried out and to CT scans," said Stewart Hughes, principal scientist of the waste management group in the agency's radiation protection division.

Lynn Eaton *London*

Ionising Radiation Exposure of the UK Population: 2005 Review is available at www.hpa.org.uk/radiation

Drug chiefs ponder how to improve industry's reputation

Drug industry executives and advisers told a conference in Sydney last week that the poor public standing of the industry was hindering its ability to sell its products at higher prices to government agencies and was interfering with its attempts to address public criticism.

Introducing a panel discus-

sion on reputation management, Kristin Austin, director of the public relations company Ruder Finn, said that a review of coverage of the industry in the Sydney print media showed 30 positive stories in the preceding year but 40 critical stories.

Bill Curtis, the founder of the Sydney based advertising and medical communications company Curtis Jones & Brown, said that winning hearts and minds would increase the chances of being able to "support our pricing needs."

Bob Burton *Canberra*

Revalidation must serve doctors and the public

For it to succeed, the system of regulating doctors in the United Kingdom must do more than simply provide a means of identifying "bad" doctors, says the BMA. It should also promote high standards, provide a platform for reflecting on and developing practice, and provide a positive confirmation of doctors' registration, the BMA says, in a response to a review of revalidation being conducted by the chief medical officer, Professor Liam Donaldson.

The BMA supports the process of revalidation, proposed by the General Medical Council, whereby doctors in the UK are periodically checked for their fitness to practise. And it believes that medical registration should continue to be professionally led.

However, to gain public confidence, the process needs to be transparent and sufficiently robust to identify poorly performing doctors before their five yearly revalidation check is due, it says.

Revalidation, which was due to be launched in April this year, was suspended in January because of concerns that the scheme would not provide an adequate evaluation of a doctor's fitness to practise (*BMJ* 2005; 330:9, 1 Jan).

Zosia Kmietowicz *London*

The BMA's response to the chief medical officer's review is at www.bma.org.uk/ap.nsf/Content

Cancer is diagnosed more quickly in patients who bypass GPs

Roger Dobson *Abergavenny*

Delay in diagnosis of cancer in the United Kingdom varies widely, according to the type of cancer, new research shows. Patients with suspected prostate cancer face delays almost three times as long as those with suspected breast cancer, claims new research.

The study, which was published online (ahead of print publication) on 3 May in the *British Journal of Cancer* (www.bjcancer.com, doi: 10.1038/sj.bjc.6602587), looked at times to diagnosis for six types of cancer. It showed that the average time to diagnosis of prostate cancer is 148.5 days, while that for breast cancer is 55.2 days.

Patients who did not see their GP first waited much less time for a diagnosis, regardless of the type of cancer. Some patients bypass seeing their GP because an abnormality is discovered through screening, while others are already in hospital or are admitted through the emergency department.

"Reducing diagnostic delays with the intention of increasing the proportion of early stage cancers may improve cancer survival in the UK, which is poorer than most other European countries. Interventions aimed at reducing patient and primary care delays need to be developed and their effect on

diagnostic stage and psychological distress evaluated," write the authors, from the University of Leeds and Wales College of Medicine.

The research compared different elements of delay in diagnosis (delay by the patient and delays in primary care, referral, or secondary care) for six types of cancer (breast, colorectal, lung, ovarian, prostate, and non-Hodgkin's lymphoma) and compared delays in patients who saw their GP before diagnosis with those who did not.

The authors used data from completed questionnaires from 65 192 patients who took part in the national survey of NHS patients.

The mean overall times to diagnosis were 55.2 days for breast cancer, 88.5 for lung cancer, 90.3 for ovarian cancer, 102.8 for non-Hodgkin's lymphoma, 125.7 for colorectal cancer, and 148.5 for prostate cancer.

A total of 52 079 patients (80%) said they had visited their GP before diagnosis, 12 074 (19%) reported that they had not, and 1039 (2%) did not answer this question. These percentages varied according to the type of cancer, with 27% of patients with breast cancer but only 10% of those with colorectal cancer not seeing their GP first.