In brief

Pregnancies in England and Wales rose in 2003: The conception rate in England and Wales rose 2% between 2002 and 2003, from 72.2 to 73.7 pregnancies per 1000 women aged 15 to 44, provisional data from the Office for National Statistics have shown. Almost four fifths of conceptions resulted in a live birth, and the rest were aborted. The pregnancy rate for girls younger than 18 fell by 1.2% from 42.8 conceptions per 1000 girls aged 15-17 to 42.3 in 2003. See www.statistics.gov.uk.

Consultant and GP numbers rise: The numbers of consultants and GPs working in the NHS rose by 213 and 275 between September and December 2004, Department of Health figures for England show. There are now 7542 more consultants and 3331 more GPs working in the NHS than there were in 1999.

Refugee doctors unable to obtain posts in the NHS: More than eight out of 10 refugee medical doctors (85%) living in the UK are not working as doctors in the NHS, a survey by the Royal College of Physicians has found. More than half of refugee doctors (54%) are not employed in any capacity. See rplondon.ac.uk.

World Health Assembly adopts resolution for disaster response: The World Health Assembly has adopted a resolution on health action in crises and disasters, calling on WHO to provide early warning of disease outbreaks, improve access to clean water and sanitation, and increase the availability of health care. See www.who.int/gb.

European Commission launches antismoking campaign: The European Commission has launched a €72m (£50m; \$91m) campaign about the dangers of smoking under the slogan "help for a life without tobacco." The campaign is aimed at young people aged 15-30 across Europe. See www.help-eu.com.

Armed conflict is a leading cause of hunger, says FAO

Peter Moszynski *London*

Armed conflict is now the leading cause of hunger around the world, concludes a report from the United Nations Food and Agriculture Organization that was presented to the Committee on World Food Security in Rome last week.

A severely malnourished man in 1998, during Sudan's last outbreak of famine induced by war

The goal of halving hunger worldwide by the year 2015, set by the World Food Summit in 1996, will be missed by a wide margin if current trends persist, warns the report. But the goal of cutting the proportion of hungry and poor people may be achieved in most regions outside sub-Saharan Africa.

The committee, which met to review progress in reducing world hunger and achieving the UN's millennium development goals, was told that armed conflict was the major obstacle to progress, with the effects of HIV and AIDS and climate change not far behind.

"Conflict destroys lives, opportunities and environments and may be one of the most significant obstacles to sustainable development as it can destroy in hours and days what has taken years and decades to develop," says the report.

Speaking at a funding conference for Darfur last week, UN secretary general Kofi Annan described the situation as "unacceptable." He said that civilians were still at risk and subject to attacks and that violence was increasingly targeted at aid workers, hampering their work.

He also warned of a \$350m (£190m; €280m) shortfall in aid for the rest of Sudan, where more than three million people still rely on food aid.

Meanwhile, Paul Foreman, head of Médecins sans Frontières in Sudan was arrested on 30 May for publishing a report on rape in Darfur. The agency is "outraged by the charges" and insists on the accuracy of the report, which detailed 500 known cases. □

Assessment of the World Food Security Situation is at www.fao.org/docrep/meeting/009/j4968e/j4968e00.htm and *Linking Conflict and Development: A Challenge for the MDG Process* is at [ftp://ftp.fao.org/docrep/fao/meeting/009/j5292e.pdf](http://ftp.fao.org/docrep/fao/meeting/009/j5292e.pdf)

NICE guidance to prevent strokes and heart attacks lacks evidence

Susan Mayor *London*

An expert adviser for a report that was used to develop the guidance from the National Institute for Health and Clinical Excellence (NICE) claims that the guidance is not supported by adequate evidence from clinical trials.

The NICE guidance recommends that a combination of modified release dipyridamole and aspirin should be used in patients who have had an ischaemic stroke or a transient ischaemic attack for two years from the most recent event. After that time, or if dipyridamole is not tolerated, preventive therapy should revert to standard care (including long term treatment with low dose aspirin).

Cathie Sudlow, senior lecturer and honorary consultant neurologist at the University of Edinburgh, was one of the

experts who developed the report used by the NICE committee to develop its guidance. She said, "NICE appears to have based its recommendation... on one trial with around 3000 participants that I consider too small from which to draw conclusions for treating all patients in [the United Kingdom]. It also compared the combination with an inadequate dose of aspirin to make a fair comparison."

The trial showed that the combination of modified release dipyridamole and aspirin prevented further stroke, but this was in contrast to other studies, although these had used different preparations of dipyridamole, Dr Sudlow said. She added, "But NICE made a blanket recommendation on the basis of this one trial."

Dr Sudlow was concerned that the possibly premature recommendation of modified release dipyridamole plus aspirin for stroke prevention, rather than just aspirin on its own, could prove difficult for patients to comply with on a long term basis, particularly if they were also on several other drugs. "The decision was made in isolation from the realities of clinical practice," she argued.

"This is the first time experts are suggesting that NICE should not be going for the more expensive treatment option. It is usually the other way around," Dr Sudlow pointed out. "I think a much larger, randomised, prospective trial is required to compare modified release dipyridamole and aspirin with an adequate dose of aspirin alone to clarify the risk-benefit balance for each regimen." □

The guidance, *Vascular Disease: Clopidogrel and Dipyridamole: Guidance*, is available at www.nice.org.uk/TA090guidance.