Published in final edited form as:

Heart Fail Rev. 2017 March ; 22(2): 209–218. doi:10.1007/s10741-016-9572-5.

MicroRNAs, heart failure, and aging: potential interactions with skeletal muscle

Kevin A. Murach^{1,3} and John J. McCarthy^{1,2}

¹Center for Muscle Biology, University of Kentucky, Lexington, KY 40536, USA

²Department of Physiology, College of Medicine, University of Kentucky, Lexington, KY 40536, USA

³Department of Rehabilitation Sciences, College of Health Sciences, University of Kentucky, Lexington, KY 40536, USA

Abstract

MicroRNAs (miRNAs) are small noncoding RNAs that regulate gene expression by targeting mRNAs for degradation or translational repression. MiRNAs can be expressed tissue specifically and are altered in response to various physiological conditions. It has recently been shown that miRNAs are released into the circulation, potentially for the purpose of communicating with distant tissues. This manuscript discusses miRNA alterations in cardiac muscle and the circulation during heart failure, a prevalent and costly public health issue. A potential mechanism for how skeletal muscle maladaptations during heart failure could be mediated by myocardium-derived miRNAs released to the circulation is presented. An overview of miRNA alterations in skeletal muscle during the ubiquitous process of aging and perspectives on miRNA interactions during heart failure are also provided.

Keywords

MicroRNA; Heart failure; Exosomes; Skeletal muscle; Aging

Brief history

The original description of the first microRNA(miRNA) lin-4 was in studies seeking to better understand the genetic regulation of developmental timing in the nematode (*Caenorhabditis elegans*, or *C. elegans*) [67, 118]. These pioneering studies provided the first evidence that lin-14 expression was regulated by a posttranscriptional mechanism involving the interaction between the lin-14 3'-UTR and the small noncoding RNA lin-4. Almost a decade later, Pasquinelli et al. [85] reported the identification of let-7, a second miRNA from *C. elegans* that down-regulated lin-41 expression, and unlike lin-4, was expressed in a broad range of bilaterian animals including vertebrate, ascidian, hemichordate, mollusk, annelid, and arthropod. The revelation that let-7 was phylogenetically conserved in bilaterian animals

[✉] John J. McCarthy, jjmcca2@uky.edu.

Conflict of interest Drs. Kevin A. Murach and John J. McCarthy have no conflicts of interest of financial ties to disclose.

was a major milestone in the history of the miRNA field because it suggested the posttranscriptional regulation of gene expression by miRNAs was more widespread than just *C. elegans*. Shortly thereafter, three independent reports described the identification of 30–50 new miRNAs in the human, fly, and worm, providing additional support for the idea that miRNAs may have an important role in the regulation of gene expression in animals [62, 65, 66]. The prescient nature of these early findings is revealed in the latest release of miRBase (version 21, June 2014; www.mirbase.org) which catalogs 35,828 mature miRNA sequences from 223 species with 2588 and 1915 human and mouse mature miRNAs, respectively.

Biogenesis

The vast majority of miRNAs are produced from RNA polymerase II transcription, resulting in a primary miRNA (pri-miRNA) transcript that has the characteristic 5' m⁷G cap structure and 3' poly(A) tail [12, 68]. Recent genomic mapping confirmed an earlier study showing that roughly half of annotated miRNAs are intragenic (exon, intron, 3'-UTR or 5'-UTR), located within protein-coding or noncoding RNA genes [44, 88]. In general, miRNA expression parallels the host gene, though new experimental evidence indicates that up ~35 % of intronic miRNAs are expressed as an independent transcription unit under regulation of its own promoter [82]. Once transcribed, the pri-miRNA forms a stem-loop structure that is recognized by the microprocessor complex which contains two core components, the RNase III endonuclease Drosha, and the double-stranded RNA-binding protein DGCR8 (DiGeorge Syndrome critical region gene 8) [47, 64]. DGCR8 binds to the stem-loop structure and then guides Drosha into position, cleaving ~11 base pairs (bp) from the base of the stem-loop to produce a 60–70 bp hairpin RNA molecule designated the precursor miRNA (pre-miRNA) [51].

Following pri-miRNA processing, the 60–70 bp precursor pre-miRNA is transported from the nucleus by Exportin 5, a nuclear export receptor, to the cytoplasm [10, 120]. Once in the cytoplasm, a second RNase III endonuclease, Dicer, cleaves the pre-miRNA to produce a ~22 nucleotide double-stranded RNA molecule in which one strand, known as the guide strand, is transferred to the RISC (RNA-induced silencing complex) containing Argonaute 2 (Ago2) and the RNA-binding protein Tarbp2 [TAR (HIV) RNA-binding protein 2]; the other strand is typically targeted for degradation [71]. The mature miRNA directs RISC to 3'-UTR of target mRNA through complementary binding of the miRNA seed sequence which results in inhibition of translation and/or degradation of the target transcript [84].

Tissue-specific expression

As quickly as miRNAs were shown to be conserved in different species, it was recognized that the expression of some miRNAs was restricted to certain tissues. One of the first examples of a tissue-specific miRNA was miR-1, which was found to be expressed exclusively in the human heart [62, 66]. The finding that some miRNAs were expressed in a tissue-specific fashion was confirmed in a study by Lagos-Quintana et al. [62], showing that miR-1, miR-122a, and miR-124a expression was restricted to striated muscle, liver, and brain, respectively. In an effort to identify new miRNAs, Sempere et al. [94] identified 30 miRNAs that were enriched or specifically expressed within a particular tissue [94]. These

authors provided the first description of striated muscle-specific miR-1, miR-133a, and miR-206, which were later designated as myomiRs [75, 94].

The myomiR family has expanded since its original description to include miR-208a, miR-208b, miR-499, and, most recently, miR-486 [98, 107, 109]. Northern blot analyses showed that these new members of the myomiR family are strictly striated muscle specific (miR-208a, miR-208b and miR-499), being derived from the intron of different muscle-specific myosin heavy chain genes, or are highly enriched in muscle (miR-486) [98, 107]. Most myomiR family members are expressed in both the heart and skeletal muscle except for miR-208a, which is cardiac specific, and miR-206, which is skeletal muscle specific and enriched in slow-twitch muscles such as the soleus [77].

Circulating miRNAs

In the late 1960's, it became clear that substances from non-endocrine cells were released into circulation and affected the behavior of other cells in the body [30]. The discovery of these circulating proteins, termed "cytokines" in 1974 [17], opened up a new area of research related to how cells in different organ systems communicate with one another and affect whole-body homeostasis. It is now known that specific organs, such as skeletal muscle [34] and cardiac tissue [25], generate unique cytokines that are released into circulation under various conditions and have profound metabolic, mass regulation, and immunological effects on distant tissues. In 2007, Valadi et al. [104] discovered that miRNAs were released from mast cells and affected mast cells of differing origin. This discovery introduced a new non-hormone mechanism by which distant cells could influence one another at the molecular level. Moreover, similar to tissue-specific cytokines, tissue-specific miRNAs could potentially be released into circulation and exert distant effects [3].

The method by which miRNAs enter the circulation and are utilized by cells in other organ systems has not been thoroughly explored. It is known that ribonucleases are found in both plasma and serum [36], but extracellular miRNAs are stable and can be transmitted into and utilized by recipient cells [59, 81, 125]. Microvesicles, apoptotic bodies, non-vesicle-associated proteins (e.g., HDL), or RNA-binding proteins could mediate the transport process [3, 5, 24, 103, 111]. Valadi et al. [104] originally showed that miRNAs are packaged into exosomes in order to mediate intercellular communication. Arroyo et al. [5] then reported that the majority of circulating miRNAs are un-encapsulated and are chaperoned by protein complexes such as Argonaute 2, which maintains miRNA stability. In contrast to these findings [5, 103], Gallo et al. [41] showed that most miRNAs in human serum are within exosomes. Moreover, interference of the exosome biosynthesis process or disruption of exosomal membranes reduces extracellular miRNA content [59]. Most recently, it has been shown that serum miRNAs can be found in both vesicular and non-vesicular fractions in response to tissue damage in rats [95]. Although the manner in which miRNAs are transported in the circulation may be condition dependent, they appear to be abundantly expressed and likely exert effects in different tissue types.

miRNAs, heart failure, and skeletal muscle

Heart failure is a general term for the heart's inability to pump sufficiently and maintain blood flow to meet the body's needs. The condition is characterized by a constellation of cardiac function criteria (further stratified by preserved ejection fraction or reduced ejection fraction) [57, 72] and could be caused by a variety of underlying mechanisms [53]. In general, though, significant cardiac ultrastructural, myofibrillar, biochemical, and molecular abnormalities underlie the pathophysiology of all forms of heart failure [26, 69]. Detrimental changes in skeletal muscle morphology and metabolism may also parallel the progression of heart failure [16, 70, 100, 113]. The heart is always metabolically active and is among the most vital organs. It is possible that heart tissue alterations associated with cardiac disease progression could affect skeletal muscle via myocardially derived miRNAs released into circulation.

Due to the varying etiology of heart failure, it would be difficult to ascribe a global miRNA signature that characterizes all human cardiac dysfunction. However, a number of misregulated miRNAs are recurrent in the heart failure literature [79] and confirmed to be altered in various forms of diseased and damaged human cardiac tissue. Namely, miR-1 [11, 14, 63, 74, 83, 102, 119], miR-21 [56, 63, 74, 102, 108, 112, 119], miR-24 [55, 74, 108], miR-29b [63, 110], miR-133a and b [11, 14, 21, 29, 74, 91, 99, 108], miR-199 [55, 63, 108], miR-208 [11, 90], miR-214 [55, 83, 108], and miR-499 [55, 73, 74] are consistently affected in human heart failure of varying origin and degree. This panel of miRNAs may serve as a preliminary template for miRNA-mediated heart-skeletal muscle communication during heart failure.

Many of the aforementioned miRNAs present in diseased myocardium are also enriched in the circulation following acute cardiovascular trauma. For instance, circulating miR-1, miR-21, miR-29a, miR-133a, miR-208a and b, and miR-499 are elevated after acute myocardial infarction (MI) [1, 18, 19, 31, 61, 115, 117, 127]. MiRNA appearance in the circulation may simply be a consequence of large-scale cardiac membrane disturbance and subsequent release. However, miRNAs that are up regulated in heart tissue also manifest in the circulation during less traumatic chronic cardiovascular conditions. MiR-1 and miR-133a are elevated in the circulation with unstable angina, cardiomyopathy [61] and coronary atherosclerosis [9, 35]; miR-208a tends to be elevated with stable coronary artery disease [35]; miR-21 and miR-29 are up regulated with ventricular fibrosis [89, 112] and hypertrophic cardiomyopathy [89]; miR-499 and miR-208b are increased with viral myocarditis [18]; and miR-21 is up regulated with various degrees of diagnosed heart failure [13, 46, 101] as well as coronary artery disease [50]. Most recently, a panel of circulating miRNAs sensitive enough to distinguish between chronic heart failure with preserved versus reduced ejection fraction was identified, and circulating miR-221 and miR-328 levels increased the discriminatory power of circulating B-type natriuretic peptide for assessing heart failure [116]. Thus, circulating miRNAs can serve as biomarkers for both acute myocardial trauma as well as chronic heart conditions.

Direct release of miRNAs from the human heart is evident under various circumstances [23, 45]. Furthermore, myocardial and circulating miR-21 levels correspond in aortic stenosis

patients [112], miR-208a (a cardiac-specific miRNA) is enriched in serum-derived exosomes [8], and the relative abundance of circulating cardiac-specific miRNAs reflects the ratio found in heart tissue [2]. Cardiomyocytes in culture are shown to release extracellular vesicles [43, 49, 114] that contain miRNAs [42]. The content and quantity of exosomes are also responsive to external stimuli [42, 43]. More work is needed to describe the miRNA profile of human cardiomyocyte-derived extracellular vesicles, but consistent with what is found in myocardial tissue and the circulation with heart failure, cardiomyocytes cultured from human progenitor cells robustly express miR-1 and miR-499 [97]. In vitro experiments also show that cardiac fibroblasts secrete miR-enriched exosomes that affect cardiomyocyte gene expression and hypertrophy [6]. Cardiomyocyte-derived exosomes also facilitate communication with endothelial cells [42]. Various lines of evidence strongly indicate that cardiac tissue can synthesize and release miRNA-containing vesicles that change in response to stimuli and could affect gene expression in distant tissues. It is conceivable that sustained miRNA release from the heart during heart failure could facilitate communication with skeletal muscle.

Similar to what occurs between cardiac fibroblasts, endothelial cells, and cardiomyocytes, myoblasts and myotubes are shown to communicate with each other through exosome-derived miRNAs in vitro [37, 38]. It is therefore likely that mature skeletal muscle regulates its microenvironment via exosomes/miRNAs in vivo and that miRNAs from distant tissues (namely myocardium) could affect skeletal muscle molecular events. Of the miRNAs enriched in both cardiac muscle and the circulation during heart failure, myomiRs (namely miR-1, miR-133, miR-208, and miR-499 [78, 106, 107]) are shown to strongly affect developed skeletal muscle. Of these myomiRs, miR-1 and miR-133a are elevated in the circulation with various forms of cardiovascular disease [61] and remain elevated for 3 months post-MI [127]. It was also recently reported that circulating myomiRs (including miR-1) are elevated in the most advanced heart failure patients [2]. Suppressed miR-1 and miR-133a are implicated in mature skeletal muscle hypertrophy [77], likely through anti-repressive targeting of the IGF-1 growth signaling axis [32, 54]. Thus, increased circulating miR-1 and miR-133a could facilitate skeletal muscle atrophy in the wake of a traumatic cardiovascular event (Fig. 1). Interestingly, once heart failure is compensated with a left ventricular assist device, myomiR levels in the myocardium (specifically miR-1, miR-133a, and miR-133b) are decreased [91]. More work is needed to elucidate how different cardiac conditions affect specific myomiRs in the circulation.

Heart–skeletal muscle communication via miRNAs during heart failure is speculative at this point, but it is not without precedent. Other compounds such as myostatin released from cardiac tissue during heart failure can strongly regulate skeletal muscle mass [52]. Emerging evidence suggests that diseased cardiac tissue is a major source of circulating miRNAs [9] and that miRNAs (via extracellular vesicles or some other mechanism) participate in intercellular communication [3]. Further research on how miRNAs are released from cardiac tissue, under what conditions they are released, how skeletal muscle is potentially targeted by extracellular vesicles/miRNAs, and the specific effects of these miRNAs in skeletal muscle is warranted. The miRNA expression profile of skeletal muscle in heart failure patients should also be evaluated. Finally, elucidating the role of non-myomiR miRNAs, such as heart failure- [13] and skeletal muscle fibrosis-associated miR-21 [4, 60, 101], is

necessary to fully understand how circulating miRNAs can contribute to diseased conditions in various tissues.

miRNAs and the interaction between aging, heart failure, and skeletal muscle

Among the most pronounced and noticeable changes that occur with aging is an inevitable loss of skeletal muscle mass and subsequent function [15]. This condition, termed sarcopenia, is caused by a complex interplay of factors and is compounded by attenuated responsiveness to hypertrophic stimuli [86, 96]. Skeletal muscle is the largest organ in the body and synthesizes numerous miRNAs, including skeletal muscle-specific myomiRs that are responsive to external stimuli [75–78]. MicroRNA abundance, specificity, and plasticity suggest these molecules are important for skeletal muscle regulation.

In healthy older individuals, basal levels of intramuscular myomiRs and cellular proliferation-related miRNAs differ compared to younger individuals, which could conceivably contribute to age-associated atrophy [27, 28, 87]. Interestingly, basal miR-1 is lower in muscle of young physically active (~27 year) and lifelong-exercised older men (~70 year) versus inactive older men [122], indicating fitness level may influence baseline miRNA expression. However, following acute resistance exercise with amino acid ingestion in untrained individuals, miR-1 is reduced in young but not old skeletal muscle [27]. Similarly, age-associated alterations to miR-1 and miRNA-mediated IGF-1 signaling [87, 121] as well as attenuated global miRNA responsiveness to exercise in skeletal muscle [87] have been reported. No statistical difference in skeletal muscle or circulating myomiRs (miR-1, miR-133a, miR-206, miR-208b, and miR-499) after resistance training in older individuals (65–80 year) further suggests age-associated miRNA inflexibility at multiple anatomical levels [22, 124]. Blunted miRNA plasticity in response to exercise dovetails with attenuated skeletal muscle global gene expression seen with exercise training as age progresses [86]. The coexistence of reduced miRNA and mRNA expression with aging could be partly attributable to the fact that 35 % of mammalian miRNA are located within annotated genes [48] and are transcribed in parallel with their host transcripts [88].

In addition to miRNA misregulation with aging in skeletal muscle, basal cardiac miRNA expression appears to be altered with age-related heart dysfunction [92, 126]. It is challenging to tease apart the effects of aging and heart failure as aging is a risk factor for the development of cardiac dysfunction [20]. However, recent circulating miRNA findings provide some insight into the individual effects of these conditions. A distinct circulating miRNA signature characterizes the aging process [123], but circulating miRNAs profiles are unique between aged individuals with or without heart failure [93]. Interestingly, circulating miR-1 appears to be *reduced* as the severity of heart failure increases in elderly individuals (>68 year) [101]. Thus, an interaction between aging and aging with heart failure appears to manifest in the expression of circulating miRNAs.

The relationship between aging, heart failure, and skeletal muscle is largely unknown. However, age-related heart failure is characterized by alterations to microRNAs associated with fibrotic deposition and extracellular matrix accumulation in cardiac tissue [105].

Excessive fibrosis in any tissue is generally associated with reduced performance and negative health outcomes. In skeletal muscle, stem cells (satellite cells) regulate fibrotic deposition in some paracrine fashion [39], and a lack of satellite cells exacerbates fibrosis with aging [40]. Since cardiac tissue possesses stem cells [7, 80], it is possible that these cells regulate fibrosis similar to skeletal muscle via secreted miRNAs [58] and are negatively affected by heart failure and aging. These fibrosis-related miRNAs in circulation may conceivably influence skeletal muscle extracellular matrix remodeling as well. To this point, patients with diagnosed myocardial fibrosis resulting from hypertrophic cardiomyopathy have altered levels of circulating miRNAs implicated in fibrotic deposition [33]. Determining what tissues (e.g., skeletal muscle, cardiac) and cell types are releasing miRNAs into circulation and how these miRNAs affect other tissues is an important next step in defining the role of miRNAs in senescence and heart failure.

Conclusion

MiRNA biology is a burgeoning area of research since these small noncoding RNAs are known to elicit a powerful effect in a variety of cell types. Both heart failure and aging are characterized by distinct miRNA profiles in striated muscle as well as the circulation. myomiRs, or muscle-specific miRNAs, are noticeably affected with heart failure and aging and could mediate inter-organ communication through packaging and release into the circulation. Exercise appears to have an effect on miRNA expression during heart failure and aging, but more research on how miRNAs (both tissue- and non-tissue-specific) are altered during different physiological processes, their target mRNAs, and how they are packaged, released into circulation, and taken up and utilized by distant cells is warranted.

Acknowledgments

This publication was supported by the National Institute of Arthritis and Musculoskeletal and Skin Diseases of the National Institutes of Health under Award Number AR061939 to J.J.M.

References

1. Ai J, Zhang R, Li Y, Pu J, Lu Y, Jiao J, Li K, Yu B, Li Z, Wang R, Wang L, Li Q, Wang N, Shan H, Li Z, Yang B. Circulating microRNA-1 as a potential novel biomarker for acute myocardial infarction. *Biochem Biophys Res Commun*. 2010; 391(1):73–77. DOI: 10.1016/j.bbrc.2009.11.005 [PubMed: 19896465]
2. Akat KM, Moore-McGriff D, Morozov P, Brown M, Gogakos T, Correa Da Rosa J, Mihailovic A, Sauer M, Ji R, Ramarathnam A, Totary-Jain H, Williams Z, Tuschl T, Schulze PC. Comparative RNA-sequencing analysis of myocardial and circulating small RNAs in human heart failure and their utility as biomarkers. *Proc Natl Acad Sci USA*. 2014; 111(30):11151–11156. DOI: 10.1073/pnas.1401724111 [PubMed: 25012294]
3. Aoi W, Sakuma K. Does regulation of skeletal muscle function involve circulating microRNAs? *Front Physiol*. 2014; 5:39.doi: 10.3389/fphys.2014.00039 [PubMed: 24596559]
4. Ardite E, Perdiguero E, Vidal B, Gutarra S, Serrano AL, Munoz-Canoves P. PAI-1-regulated miR-21 defines a novel age-associated fibrogenic pathway in muscular dystrophy. *J Cell Biol*. 2012; 196(1): 163–175. DOI: 10.1083/jcb.201105013 [PubMed: 22213800]
5. Arroyo JD, Chevillet JR, Kroh EM, Ruf IK, Pritchard CC, Gibson DF, Mitchell PS, Bennett CF, Pogosova-Agadjanyan EL, Stirewalt DL, Tait JF, Tewari M. Argonaute2 complexes carry a population of circulating microRNAs independent of vesicles in human plasma. *Proc Natl Acad Sci USA*. 2011; 108(12):5003–5008. DOI: 10.1073/pnas.1019055108 [PubMed: 21383194]

6. Bang C, Batkai S, Dangwal S, Gupta SK, Foinquinos A, Holzmann A, Just A, Remke J, Zimmer K, Zeug A, Ponimaskin E, Schmiedl A, Yin X, Mayr M, Halder R, Fischer A, Engelhardt S, Wei Y, Schober A, Fiedler J, Thum T. Cardiac fibroblast-derived microRNA passenger strand-enriched exosomes mediate cardiomyocyte hypertrophy. *J Clin Invest*. 2014; 124(5):2136–2146. DOI: 10.1172/JCI70577 [PubMed: 24743145]
7. Beltrami AP, Barlucchi L, Torella D, Baker M, Limana F, Chimenti S, Kasahara H, Rota M, Musso E, Urbanek K, Leri A, Kajstura J, Nadal-Ginard B, Anversa P. Adult cardiac stem cells are multipotent and support myocardial regeneration. *Cell*. 2003; 114(6):763–776. [PubMed: 14505575]
8. Bi S, Wang C, Jin Y, Lv Z, Xing X, Lu Q. Correlation between serum exosome derived miR-208a and acute coronary syndrome. *Int J Clin Exp Med*. 2015; 8(3):4275–4280. [PubMed: 26064341]
9. Boeckel J-N, Reis SM, Leistner D, Thome CE, Zeiher AM, Fichtlscherer S, Keller T. From heart to toe: Heart's contribution on peripheral microRNA levels. *Int J Cardiol*. 2015; 172(3):616–617.
10. Bohnsack MT, Czaplinski K, Gorlich D. Exportin 5 is a RanGTP-dependent dsRNA-binding protein that mediates nuclear export of pre-miRNAs. *RNA*. 2004; 10(2):185–191. [PubMed: 14730017]
11. Bostjancic E, Zidar N, Stajer D, Glavac D. MicroRNAs miR-1, miR-133a, miR-133b and miR-208 are dysregulated in human myocardial infarction. *Cardiology*. 2010; 115(3):163–169. DOI: 10.1159/000268088 [PubMed: 20029200]
12. Cai X, Hagedorn CH, Cullen BR. Human microRNAs are processed from capped, polyadenylated transcripts that can also function as mRNAs. *RNA*. 2004; 10(12):1957–1966. DOI: 10.1261/rna.7135204 [PubMed: 15525708]
13. Cakmak HA, Coskunpinar E, Ikitimur B, Barman HA, Karadag B, Tiryakioglu NO, Kahraman K, Vural VA. The prognostic value of circulating microRNAs in heart failure: preliminary results from a genome-wide expression study. *J Cardiovasc Med (Hagerstown)*. 2015; 16(6):431–437. DOI: 10.2459/JCM.0000000000000233 [PubMed: 25643195]
14. Care A, Catalucci D, Felicetti F, Bonci D, Addario A, Gallo P, Bang ML, Segnalin P, Gu Y, Dalton ND, Elia L, Latronico MV, Hoydal M, Autore C, Russo MA, Dorn GW 2nd, Ellingsen O, Ruiz-Lozano P, Peterson KL, Croce CM, Peschle C, Condorelli G. MicroRNA-133 controls cardiac hypertrophy. *Nat Med*. 2007; 13(5):613–618. DOI: 10.1038/nm1582 [PubMed: 17468766]
15. Cederholm T, Morley JE. Sarcopenia: the new definitions. *Curr Opin Clin Nutr Metab Care*. 2015; 18(1):1–4. DOI: 10.1097/MCO.0000000000000119 [PubMed: 25207791]
16. Coats AJ, Clark AL, Piepoli M, Volterrani M, Poole-Wilson PA. Symptoms and quality of life in heart failure: the muscle hypothesis. *Br Heart J*. 1994; 72(2 Suppl):S36–S39. [PubMed: 7946756]
17. Cohen S, Bigazzi PE, Yoshida T. Commentary. Similarities of T cell function in cell-mediated immunity and antibody production. *Cell Immunol*. 1974; 12(1):150–159. [PubMed: 4156495]
18. Corsten MF, Dennert R, Jochems S, Kuznetsova T, Devaux Y, Hofstra L, Wagner DR, Staessen JA, Heymans S, Schroen B. Circulating MicroRNA-208b and MicroRNA-499 reflect myocardial damage in cardiovascular disease. *Circ Cardiovasc Genet*. 2010; 3(6):499–506. DOI: 10.1161/CIRCGENETICS.110.957415 [PubMed: 20921333]
19. D'Alessandra Y, Devanna P, Limana F, Straino S, Di Carlo A, Brambilla PG, Rubino M, Carena MC, Spazzafumo L, De Simone M, Micheli B, Biglioli P, Achilli F, Martelli F, Maggolini S, Marenzi G, Pompilio G, Capogrossi MC. Circulating microRNAs are new and sensitive biomarkers of myocardial infarction. *Eur Heart J*. 2010; 31(22):2765–2773. DOI: 10.1093/eurheartj/ehq167 [PubMed: 20534597]
20. Dai DF, Chen T, Johnson SC, Szeto H, Rabinovitch PS. Cardiac aging: from molecular mechanisms to significance in human health and disease. *Antioxid Redox Signal*. 2012; 16(12):1492–1526. DOI: 10.1089/ars.2011.4179 [PubMed: 22229339]
21. Danowski N, Manthey I, Jakob HG, Siffert W, Peters J, Frey UH. Decreased expression of miR-133a but not of miR-1 is associated with signs of heart failure in patients undergoing coronary bypass surgery. *Cardiology*. 2013; 125(2):125–130. DOI: 10.1159/000348563 [PubMed: 23711953]
22. Davidsen PK, Gallagher IJ, Hartman JW, Tarnopolsky MA, Dela F, Helge JW, Timmons JA, Phillips SM. High responders to resistance exercise training demonstrate differential regulation of

- skeletal muscle microRNA expression. *J Appl Physiol.* 2011; 110(2):309–317. DOI: 10.1152/japplphysiol.00901.2010 [PubMed: 21030674]
23. De Rosa S, Fichtlscherer S, Lehmann R, Assmus B, Dimmeler S, Zeiher AM. Transcoronary concentration gradients of circulating microRNAs. *Circulation.* 2011; 124(18):1936–1944. DOI: 10.1161/CIRCULATIONAHA.111.037572 [PubMed: 21969012]
 24. Diehl P, Fricke A, Sander L, Stamm J, Bassler N, Htun N, Ziemann M, Helbing T, El-Osta A, Jowett JB, Peter K. Microparticles: major transport vehicles for distinct microRNAs in circulation. *Cardiovasc Res.* 2012; 93(4):633–644. DOI: 10.1093/cvr/cvs007 [PubMed: 22258631]
 25. Doroudgar S, Glembotski CC. The cardiokine story unfolds: ischemic stress-induced protein secretion in the heart. *Trends Mol Med.* 2011; 17(4):207–214. DOI: 10.1016/j.molmed.2010.12.003 [PubMed: 21277256]
 26. Drexler H, Riede U, Munzel T, König H, Funke E, Just H. Alterations of skeletal muscle in chronic heart failure. *Circulation.* 1992; 85(5):1751–1759. [PubMed: 1315220]
 27. Drummond MJ, McCarthy JJ, Fry CS, Esser KA, Rasmussen BB. Aging differentially affects human skeletal muscle micro- RNA expression at rest and after an anabolic stimulus of resistance exercise and essential amino acids. *Am J Physiol Endocrinol Metab.* 2008; 295(6):E1333–E1340. DOI: 10.1152/ajpendo.90562.2008 [PubMed: 18827171]
 28. Drummond MJ, McCarthy JJ, Sinha M, Spratt HM, Volpi E, Esser KA, Rasmussen BB. Aging and microRNA expression in human skeletal muscle: a microarray and bioinformatics analysis. *Physiol Genomics.* 2011; 43(10):595–603. DOI: 10.1152/physiolgenomics.00148.2010 [PubMed: 20876843]
 29. Duisters RF, Tijssen AJ, Schroen B, Leenders JJ, Lentink V, van der Made I, Herias V, van Leeuwen RE, Schellings MW, Barenbrug P, Maessen JG, Heymans S, Pinto YM, Creemers EE. miR-133 and miR-30 regulate connective tissue growth factor: implications for a role of microRNAs in myocardial matrix remodeling. *Circ Res.* 2009; 104(2):170–178. 176p following 178. DOI: 10.1161/CIRCRESAHA.108.182535 [PubMed: 19096030]
 30. Dumonde DC, Wolstencroft RA, Panayi GS, Matthew M, Morley J, Howson WT. “Lymphokines”: non-antibody mediators of cellular immunity generated by lymphocyte activation. *Nature.* 1969; 224(5214):38–42. [PubMed: 5822903]
 31. Eitel I, Adams V, Dieterich P, Fuernau G, de Waha S, Desch S, Schuler G, Thiele H. Relation of circulating MicroRNA-133a concentrations with myocardial damage and clinical prognosis in ST-elevation myocardial infarction. *Am Heart J.* 2012; 164(5):706–714. DOI: 10.1016/j.ahj.2012.08.004 [PubMed: 23137501]
 32. Elia L, Contu R, Quintavalle M, Varrone F, Chimenti C, Russo MA, Cimino V, De Marinis L, Frustaci A, Catalucci D, Condorelli G. Reciprocal regulation of microRNA-1 and insulin-like growth factor-1 signal transduction cascade in cardiac and skeletal muscle in physiological and pathological conditions. *Circulation.* 2009; 120(23):2377–2385. DOI: 10.1161/CIRCULATIONAHA.109.879429 [PubMed: 19933931]
 33. Fang L, Ellims AH, Moore XL, White DA, Taylor AJ, Chin-Dusting J, Dart AM. Circulating microRNAs as biomarkers for diffuse myocardial fibrosis in patients with hypertrophic cardiomyopathy. *J Transl Med.* 2015; 13:314.doi: 10.1186/s12967-015-0672-0 [PubMed: 26404540]
 34. Febbraio MA, Pedersen BK. Contraction-induced myokine production and release: Is skeletal muscle an endocrine organ? *Exerc Sport Sci Rev.* 2005; 33(3):114–119. [PubMed: 16006818]
 35. Fichtlscherer S, De Rosa S, Fox H, Schwietz T, Fischer A, Liebetrau C, Weber M, Hamm CW, Roxel T, Muller-Ardogan M, Bonauer A, Zeiher AM, Dimmeler S. Circulating microRNAs in patients with coronary artery disease. *Circ Res.* 2010; 107(5):677–684. DOI: 10.1161/CIRCRESAHA.109.215566 [PubMed: 20595655]
 36. Fleischhacker M, Schmidt B. Circulating nucleic acids (CNAs) and cancer—a survey. *Biochim Biophys Acta.* 2007; 1775(1):181–232. DOI: 10.1016/j.bbcan.2006.10.001 [PubMed: 17137717]
 37. Forterre A, Jalabert A, Berger E, Baudet M, Chikh K, Errazuriz E, De Larichaudy J, Chanon S, Weiss-Gayet M, Hesse AM, Record M, Geloën A, Lefai E, Vidal H, Coute Y, Rome S. Proteomic analysis of C2C12 myoblast and myotube exosome-like vesicles: A new paradigm for myoblast-myotube cross talk? *PLoS ONE.* 2014; 9(1):e84153.doi: 10.1371/journal.pone.0084153 [PubMed: 24392111]

38. Forterre A, Jalabert A, Chikh K, Pesenti S, Euthine V, Granjon A, Errazuriz E, Lefai E, Vidal H, Rome S. Myotube-derived exosomal miRNAs downregulate Sirtuin1 in myoblasts during muscle cell differentiation. *Cell Cycle*. 2014; 13(1):78–89. DOI: 10.4161/cc.26808 [PubMed: 24196440]
39. Fry CS, Lee JD, Jackson JR, Kirby TJ, Stasko SA, Liu H, Dupont-Versteegden EE, McCarthy JJ, Peterson CA. Regulation of the muscle fiber microenvironment by activated satellite cells during hypertrophy. *FASEB J*. 2014; 28(4):1654–1665. DOI: 10.1096/fj.13-239426 [PubMed: 24376025]
40. Fry CS, Lee JD, Mula J, Kirby TJ, Jackson JR, Liu F, Yang L, Mendias CL, Dupont-Versteegden EE, McCarthy JJ, Peterson CA. Inducible depletion of satellite cells in adult, sedentary mice impairs muscle regenerative capacity without affecting sarcopenia. *Nat Med*. 2015; 21(1):76–80. DOI: 10.1038/nm.3710 [PubMed: 25501907]
41. Gallo A, Tandon M, Alevizos I, Illei GG. The majority of microRNAs detectable in serum and saliva is concentrated in exosomes. *PLoS ONE*. 2012; 7(3):e30679.doi: 10.1371/journal.pone.0030679 [PubMed: 22427800]
42. Garcia NA, Ontoria-Oviedo I, Gonzalez-King H, Diez-Juan A, Sepulveda P. Glucose starvation in cardiomyocytes enhances exosome secretion and promotes angiogenesis in endothelial cells. *PLoS ONE*. 2015; 10(9):e0138849.doi: 10.1371/journal.pone.0138849 [PubMed: 26393803]
43. Genneback N, Hellman U, Malm L, Larsson G, Ronquist G, Waldenstrom A, Morner S. Growth factor stimulation of cardiomyocytes induces changes in the transcriptional contents of secreted exosomes. *J Extracell Vesicles*. 2013; doi: 10.3402/jev.v2i0.20167
44. Godnic I, Zorc M, Jevsinek Skok D, Calin GA, Horvat S, Dovc P, Kovac M, Kunej T. Genome-wide and species-wide in silico screening for intragenic MicroRNAs in human, mouse and chicken. *PLoS ONE*. 2013; 8(6):e65165.doi: 10.1371/journal.pone.0065165 [PubMed: 23762306]
45. Goldraich LA, Martinelli NC, Matte U, Cohen C, Andrades M, Pimentel M, Biolo A, Clausell N, Rohde LE. Transcoronary gradient of plasma microRNA 423-5p in heart failure: evidence of altered myocardial expression. *Biomarkers*. 2014; 19(2):135–141. DOI: 10.3109/1354750X.2013.870605 [PubMed: 24506564]
46. Goren Y, Kushnir M, Zafir B, Tabak S, Lewis BS, Amir O. Serum levels of microRNAs in patients with heart failure. *Eur J Heart Fail*. 2012; 14(2):147–154. DOI: 10.1093/eurjhf/hfr155 [PubMed: 22120965]
47. Gregory RI, Yan KP, Amuthan G, Chendrimada T, Doratotaj B, Cooch N, Shiekhattar R. The microprocessor complex mediates the genesis of microRNAs. *Nature*. 2004; 432(7014):235–240. DOI: 10.1038/nature03120 [PubMed: 15531877]
48. Griffiths-Jones S, Grocock RJ, van Dongen S, Bateman A, Enright AJ. miRBase: microRNA sequences, targets and gene nomenclature. *Nucleic Acids Res*. 2006; 34:D140–D144. (Database issue). DOI: 10.1093/nar/gkj112 [PubMed: 16381832]
49. Gupta S, Knowlton AA. HSP60 trafficking in adult cardiac myocytes: role of the exosomal pathway. *Am J Physiol Heart Circ Physiol*. 2007; 292(6):H3052–H3056. DOI: 10.1152/ajpheart.01355.2006 [PubMed: 17307989]
50. Han H, Qu G, Han C, Wang Y, Sun T, Li F, Wang J, Luo S. MiR-34a, miR-21 and miR-23a as potential biomarkers for coronary artery disease: a pilot microarray study and confirmation in a 32 patient cohort. *Exp Mol Med*. 2015; 47:e138.doi: 10.1038/emmm.2014.81 [PubMed: 25656948]
51. Han J, Lee Y, Yeom KH, Nam JW, Heo I, Rhee JK, Sohn SY, Cho Y, Zhang BT, Kim VN. Molecular basis for the recognition of primary microRNAs by the Drosha-DGCR8 complex. *Cell*. 2006; 125(5):887–901. DOI: 10.1016/j.cell.2006.03.043 [PubMed: 16751099]
52. Heineke J, Auger-Messier M, Xu J, Sargent M, York A, Welle S, Molkentin JD. Genetic deletion of myostatin from the heart prevents skeletal muscle atrophy in heart failure. *Circulation*. 2010; 121(3):419–425. DOI: 10.1161/CIRCULATIONAHA.109.882068 [PubMed: 20065166]
53. Ho KK, Pinsky JL, Kannel WB, Levy D. The epidemiology of heart failure: the Framingham Study. *J Am Coll Cardiol*. 1993; 22(4):6A–13A. [PubMed: 8509564]
54. Huang MB, Xu H, Xie SJ, Zhou H, Qu LH. Insulin-like growth factor-1 receptor is regulated by microRNA-133 during skeletal myogenesis. *PLoS ONE*. 2011; 6(12):e29173.doi: 10.1371/journal.pone.0029173 [PubMed: 22195016]

55. Ikeda S, Kong SW, Lu J, Bisping E, Zhang H, Allen PD, Golub TR, Pieske B, Pu WT. Altered microRNA expression in human heart disease. *Physiol Genomics*. 2007; 31(3):367–373. DOI: 10.1152/physiolgenomics.00144.2007 [PubMed: 17712037]
56. Ikeda S, Pu WT. Expression and function of microRNAs in heart disease. *Curr Drug Targets*. 2010; 11(8):913–925. [PubMed: 20415651]
57. Kaminsky LA, Tuttle MS. Functional assessment of heart failure patients. *Heart Fail Clin*. 2015; 11(1):29–36. DOI: 10.1016/j.hfc.2014.08.002 [PubMed: 25432472]
58. Khanabdali R, Rosdahl AA, Dusting GJ, Lim SY. Harnessing the secretome of cardiac stem cells as therapy for ischemic heart disease. *Biochem Pharmacol*. 2016; doi: 10.1016/j.bcp.2016.02.012
59. Kosaka N, Iguchi H, Yoshioka Y, Takeshita F, Matsuki Y, Ochiya T. Secretory mechanisms and intercellular transfer of microRNAs in living cells. *J Biol Chem*. 2010; 285(23):17442–17452. DOI: 10.1074/jbc.M110.107821 [PubMed: 20353945]
60. Krichevsky AM, Gabriely G. miR-21: a small multifaceted RNA. *J Cell Mol Med*. 2009; 13(1):39–53. DOI: 10.1111/j.1582-4934.2008.00556.x [PubMed: 19175699]
61. Kuwabara Y, Ono K, Horie T, Nishi H, Nagao K, Kinoshita M, Watanabe S, Baba O, Kojima Y, Shizuta S, Imai M, Tamura T, Kita T, Kimura T. Increased microRNA-1 and micro- RNA-133a levels in serum of patients with cardiovascular disease indicate myocardial damage. *Circ Cardiovasc Genet*. 2011; 4(4):446–454. DOI: 10.1161/CIRCGENETICS.110.958975 [PubMed: 21642241]
62. Lagos-Quintana M, Rauhut R, Lendeckel W, Tuschl T. Identification of novel genes coding for small expressed RNAs. *Science*. 2001; 294(5543):853–858. DOI: 10.1126/science.1064921 [PubMed: 11679670]
63. Lai KB, Sanderson JE, Izzat MB, Yu CM. Micro-RNA and mRNA myocardial tissue expression in biopsy specimen from patients with heart failure. *Int J Cardiol*. 2015; 199:79–83. DOI: 10.1016/j.ijcard.2015.07.043 [PubMed: 26188824]
64. Landthaler M, Yalcin A, Tuschl T. The human DiGeorge syndrome critical region gene 8 and Its D. melanogaster homolog are required for miRNA biogenesis. *Curr Biol*. 2004; 14(23):2162–2167. DOI: 10.1016/j.cub.2004.11.001 [PubMed: 15589161]
65. Lau NC, Lim LP, Weinstein EG, Bartel DP. An abundant class of tiny RNAs with probable regulatory roles in *Caenorhabditis elegans*. *Science*. 2001; 294(5543):858–862. DOI: 10.1126/science.1065062 [PubMed: 11679671]
66. Lee RC, Ambros V. An extensive class of small RNAs in *Caenorhabditis elegans*. *Science*. 2001; 294(5543):862–864. DOI: 10.1126/science.1065329 [PubMed: 11679672]
67. Lee RC, Feinbaum RL, Ambros V. The *C. elegans* heterochronic gene lin-4 encodes small RNAs with antisense complementarity to lin-14. *Cell*. 1993; 75(5):843–854. [PubMed: 8252621]
68. Lee Y, Kim M, Han J, Yeom KH, Lee S, Baek SH, Kim VN. MicroRNA genes are transcribed by RNA polymerase II. *EMBO J*. 2004; 23(20):4051–4060. DOI: 10.1038/sj.emboj.7600385 [PubMed: 15372072]
69. Liew CC, Dzau VJ. Molecular genetics and genomics of heart failure. *Nat Rev Genet*. 2004; 5(11):811–825. DOI: 10.1038/nrg1470 [PubMed: 15520791]
70. Lipkin DP, Jones DA, Round JM, Poole-Wilson PA. Abnormalities of skeletal muscle in patients with chronic heart failure. *Int J Cardiol*. 1988; 18(2):187–195. [PubMed: 2830194]
71. Macrae IJ, Zhou K, Li F, Repic A, Brooks AN, Cande WZ, Adams PD, Doudna JA. Structural basis for double-stranded RNA processing by Dicer. *Science*. 2006; 311(5758):195–198. DOI: 10.1126/science.1121638 [PubMed: 16410517]
72. Marantz PR, Tobin JN, Wassertheil-Smoller S, Steingart RM, Wexler JP, Budner N, Lense L, Wachspress J. The relationship between left ventricular systolic function and congestive heart failure diagnosed by clinical criteria. *Circulation*. 1988; 77(3):607–612. [PubMed: 3342491]
73. Matkovich SJ, Hu Y, Eschenbacher WH, Dorn LE, Dorn GW 2nd. Direct and indirect involvement of microRNA-499 in clinical and experimental cardiomyopathy. *Circ Res*. 2012; 111(5):521–531. DOI: 10.1161/CIRCRESAHA.112.265736 [PubMed: 22752967]
74. Matkovich SJ, Van Booven DJ, Youker KA, Torre-Amione G, Diwan A, Eschenbacher WH, Dorn LE, Watson MA, Margulies KB, Dorn GW 2nd. Reciprocal regulation of myocardial microRNAs and messenger RNA in human cardiomyopathy and reversal of the microRNA signature by

- biomechanical support. *Circulation*. 2009; 119(9):1263–1271. DOI: 10.1161/CIRCULATIONAHA.108.813576 [PubMed: 19237659]
75. McCarthy JJ. MicroRNA-206: the skeletal muscle-specific myomiR. *Biochim Biophys Acta*. 2008; 1779(11):682–691. DOI: 10.1016/j.bbagr.2008.03.001 [PubMed: 18381085]
 76. McCarthy JJ. The MyomiR network in skeletal muscle plasticity. *Exerc Sport Sci Rev*. 2011; 39(3): 150–154. DOI: 10.1097/JES.0b013e31821c01e1 [PubMed: 21467943]
 77. McCarthy JJ, Esser KA. MicroRNA-1 and microRNA-133a expression are decreased during skeletal muscle hypertrophy. *J Appl Physiol*. 2007; 102(1):306–313. DOI: 10.1152/japplphysiol.00932.2006 [PubMed: 17008435]
 78. McCarthy JJ, Esser KA, Peterson CA, Dupont-Versteegden EE. Evidence of MyomiR network regulation of beta-myosin heavy chain gene expression during skeletal muscle atrophy. *Physiol Genomics*. 2009; 39(3):219–226. DOI: 10.1152/physiolgenomics.00042.2009 [PubMed: 19690046]
 79. Melman YF, Shah R, Das S. MicroRNAs in heart failure: Is the picture becoming less miRky? *Circ Heart Fail*. 2014; 7:203–214. [PubMed: 24449811]
 80. Messina E, De Angelis L, Frati G, Morrone S, Chimenti S, Fiordaliso F, Salio M, Battaglia M, Latronico MV, Coletta M, Vivarelli E, Frati L, Cossu G, Giacomello A. Isolation and expansion of adult cardiac stem cells from human and murine heart. *Circ Res*. 2004; 95(9):911–921. DOI: 10.1161/01.RES.0000147315.71699.51 [PubMed: 15472116]
 81. Montecalvo A, Larregina AT, Shufesky WJ, Stolz DB, Sullivan ML, Karlsson JM, Baty CJ, Gibson GA, Erdos G, Wang Z, Milosevic J, Tkacheva OA, Divito SJ, Jordan R, Lyons-Weiler J, Watkins SC, Morelli AE. Mechanism of transfer of functional microRNAs between mouse dendritic cells via exosomes. *Blood*. 2012; 119(3):756–766. DOI: 10.1182/blood-2011-02-338004 [PubMed: 22031862]
 82. Monteys AM, Spengler RM, Wan J, Tecedor L, Lennox KA, Xing Y, Davidson BL. Structure and activity of putative intronic miRNA promoters. *RNA*. 2010; 16(3):495–505. DOI: 10.1261/rna.1731910 [PubMed: 20075166]
 83. Naga Prasad SV, Duan ZH, Gupta MK, Surampudi VS, Volinia S, Calin GA, Liu CG, Kotwal A, Moravec CS, Starling RC, Perez DM, Sen S, Wu Q, Plow EF, Croce CM, Karnik S. Unique microRNA profile in end-stage heart failure indicates alterations in specific cardiovascular signaling networks. *J Biol Chem*. 2009; 284(40):27487–27499. DOI: 10.1074/jbc.M109.036541 [PubMed: 19641226]
 84. Pasquinelli AE. MicroRNAs and their targets: recognition, regulation and an emerging reciprocal relationship. *Nat Rev Genet*. 2012; 13(4):271–282. DOI: 10.1038/nrg3162 [PubMed: 22411466]
 85. Pasquinelli AE, Reinhart BJ, Slack F, Martindale MQ, Kuroda MI, Maller B, Hayward DC, Ball EE, Degnan B, Muller P, Spring J, Srinivasan A, Fishman M, Finnerty J, Corbo J, Levine M, Leahy P, Davidson E, Ruvkun G. Conservation of the sequence and temporal expression of let-7 heterochronic regulatory RNA. *Nature*. 2000; 408(6808):86–89. DOI: 10.1038/35040556 [PubMed: 11081512]
 86. Raue U, Trappe TA, Estrem ST, Qian HR, Helvering LM, Smith RC, Trappe S. Transcriptome signature of resistance exercise adaptations: mixed muscle and fiber type specific profiles in young and old adults. *J Appl Physiol*. 2012; 112(10):1625–1636. DOI: 10.1152/japplphysiol.00435.2011 [PubMed: 22302958]
 87. Rivas DA, Lessard SJ, Rice NP, Lustgarten MS, So K, Goodyear LJ, Parnell LD, Fielding RA. Diminished skeletal muscle microRNA expression with aging is associated with attenuated muscle plasticity and inhibition of IGF-1 signaling. *FASEB J*. 2014; 28(9):4133–4147. DOI: 10.1096/fj.14-254490 [PubMed: 24928197]
 88. Rodriguez A, Griffiths-Jones S, Ashurst JL, Bradley A. Identification of mammalian microRNA host genes and transcription units. *Genome Res*. 2004; 14(10A):1902–1910. DOI: 10.1101/gr.2722704 [PubMed: 15364901]
 89. Roncarati R, Viviani Anselmi C, Losi MA, Papa L, Cavarretta E, Da Costa Martins P, Contaldi C, Saccani Jotti G, Franzone A, Galastri L, Latronico MV, Imbriaco M, Esposito G, De Windt L, Betocchi S, Condorelli G. Circulating miR-29a, among other up-regulated microRNAs, is the only biomarker for both hypertrophy and fibrosis in patients with hypertrophic cardiomyopathy. *J Am Coll Cardiol*. 2014; 63(9):920–927. DOI: 10.1016/j.jacc.2013.09.041 [PubMed: 24161319]

90. Satoh M, Minami Y, Takahashi Y, Tabuchi T, Nakamura M. Expression of microRNA-208 is associated with adverse clinical outcomes in human dilated cardiomyopathy. *J Card Fail.* 2010; 16(5):404–410. DOI: 10.1016/j.cardfail.2010.01.002 [PubMed: 20447577]
91. Schipper ME, van Kuik J, de Jonge N, Dullens HF, de Weger RA. Changes in regulatory microRNA expression in myocardium of heart failure patients on left ventricular assist device support. *J Heart Lung Transpl.* 2008; 27(12):1282–1285. DOI: 10.1016/j.healun.2008.09.005
92. Seeger T, Boon RA. MicroRNAs in cardiovascular ageing. *J Physiol.* 2015; doi: 10.1113/JP270557
93. Seeger T, Haffez F, Fischer A, Koehl U, Leistner DM, Seeger FH, Boon RA, Zeiher AM, Dimmeler S. Immunosenescence-associated microRNAs in age and heart failure. *Eur J Heart Fail.* 2013; 15(4):385–393. DOI: 10.1093/eurjhf/hfs184 [PubMed: 23258801]
94. Sempere LF, Freemantle S, Pitha-Rowe I, Moss E, Dmitrovsky E, Ambros V. Expression profiling of mammalian microRNAs uncovers a subset of brain-expressed microRNAs with possible roles in murine and human neuronal differentiation. *Genome Biol.* 2004; 5(3):R13.doi: 10.1186/gb-2004-5-3-r13 [PubMed: 15003116]
95. Siracusa J, Koulmann N, Bourdon S, Goriot ME, Banzet S. Circulating miRNAs as biomarkers of acute muscle damage in rats. *Am J Pathol.* 2016; 186(5):1313–1327. DOI: 10.1016/j.ajpath.2016.01.007 [PubMed: 26952641]
96. Slivka D, Raue U, Hollon C, Minchev K, Trappe S. Single muscle fiber adaptations to resistance training in old (>80 yr) men: evidence for limited skeletal muscle plasticity. *Am J Physiol Regul Integr Comp Physiol.* 2008; 295(1):R273–R280. DOI: 10.1152/ajpregu.00093.2008 [PubMed: 18448613]
97. Sluijter JP, van Mil A, van Vliet P, Metz CH, Liu J, Doevendans PA, Goumans MJ. MicroRNA-1 and -499 regulate differentiation and proliferation in human-derived cardiomyocyte progenitor cells. *Arterioscler Thromb Vasc Biol.* 2010; 30(4):859–868. DOI: 10.1161/ATVBAHA.109.197434 [PubMed: 20081117]
98. Small EM, O'Rourke JR, Moresi V, Sutherland LB, McAnally J, Gerard RD, Richardson JA, Olson EN. Regulation of PI3-kinase/Akt signaling by muscle-enriched microRNA-486. *Proc Natl Acad Sci USA.* 2010; 107(9):4218–4223. DOI: 10.1073/pnas.1000300107 [PubMed: 20142475]
99. Sucharov C, Bristow MR, Port JD. miRNA expression in the failing human heart: functional correlates. *J Mol Cell Cardiol.* 2008; 45(2):185–192. DOI: 10.1016/j.yjmcc.2008.04.014 [PubMed: 18582896]
100. Sullivan MJ, Green HJ, Cobb FR. Skeletal muscle biochemistry and histology in ambulatory patients with long-term heart failure. *Circulation.* 1990; 81(2):518–527. [PubMed: 2297859]
101. Sygitowicz G, Tomaniak M, Blaszczyk O, Koltowski L, Filipiak KJ, Sitkiewicz D. Circulating microribonucleic acids miR-1, miR-21 and miR-208a in patients with symptomatic heart failure: preliminary results. *Arch Cardiovasc Dis.* 2015; 108(12):634–642. DOI: 10.1016/j.acvd.2015.07.003 [PubMed: 26498537]
102. Thum T, Galuppo P, Wolf C, Fiedler J, Kneitz S, van Laake LW, Doevendans PA, Mummery CL, Borlak J, Haverich A, Gross C, Engelhardt S, Ertl G, Bauersachs J. MicroRNAs in the human heart: a clue to fetal gene reprogramming in heart failure. *Circulation.* 2007; 116(3):258–267. DOI: 10.1161/CIRCULATIONAHA.107.687947 [PubMed: 17606841]
103. Turchinovich A, Weiz L, Langheinz A, Burwinkel B. Characterization of extracellular circulating microRNA. *Nucleic Acids Res.* 2011; 39(16):7223–7233. DOI: 10.1093/nar/gkr254 [PubMed: 21609964]
104. Valadi H, Ekstrom K, Bossios A, Sjostrand M, Lee JJ, Lotvall JO. Exosome-mediated transfer of mRNAs and micro-RNAs is a novel mechanism of genetic exchange between cells. *Nat Cell Biol.* 2007; 9(6):654–659. DOI: 10.1038/ncb1596 [PubMed: 17486113]
105. van Almen GC, Verhesen W, van Leeuwen RE, van de Vrie M, Eurlings C, Schellings MW, Swinnen M, Cleutjens JP, van Zandvoort MA, Heymans S, Schroen B. MicroRNA-18 and microRNA-19 regulate CTGF and TSP-1 expression in age-related heart failure. *Aging Cell.* 2011; 10(5):769–779. DOI: 10.1111/j.1474.9726.2011.00714.x [PubMed: 21501375]
106. van Rooij E, Liu N, Olson EN. MicroRNAs flex their muscles. *Trends Genet.* 2008; 24(4):159–166. DOI: 10.1016/j.tig.2008.01.007 [PubMed: 18325627]

107. van Rooij E, Quiat D, Johnson BA, Sutherland LB, Qi X, Richardson JA, Kelm RJ Jr, Olson EN. A family of microRNAs encoded by myosin genes governs myosin expression and muscle performance. *Dev Cell*. 2009; 17(5):662–673. DOI: 10.1016/j.devcel.2009.10.013 [PubMed: 19922871]
108. van Rooij E, Sutherland LB, Liu N, Williams AH, McAnally J, Gerard RD, Richardson JA, Olson EN. A signature pattern of stress-responsive microRNAs that can evoke cardiac hypertrophy and heart failure. *Proc Natl Acad Sci USA*. 2006; 103(48):18255–18260. DOI: 10.1073/pnas.0608791103 [PubMed: 17108080]
109. van Rooij E, Sutherland LB, Qi X, Richardson JA, Hill J, Olson EN. Control of stress-dependent cardiac growth and gene expression by a microRNA. *Science*. 2007; 316(5824):575–579. DOI: 10.1126/science.1139089 [PubMed: 17379774]
110. van Rooij E, Sutherland LB, Thatcher JE, DiMaio JM, Naseem RH, Marshall WS, Hill JA, Olson EN. Dysregulation of microRNAs after myocardial infarction reveals a role of miR-29 in cardiac fibrosis. *Proc Natl Acad Sci USA*. 2008; 105(35):13027–13032. DOI: 10.1073/pnas.0805038105 [PubMed: 18723672]
111. Vickers KC, Palmisano BT, Shoucri BM, Shamburek RD, Remaley AT. MicroRNAs are transported in plasma and delivered to recipient cells by high-density lipoproteins. *Nat Cell Biol*. 2011; 13(4):423–433. DOI: 10.1038/ncb2210 [PubMed: 21423178]
112. Villar AV, Garcia R, Merino D, Llano M, Cobo M, Montalvo C, Martin-Duran R, Hurle MA, Nistal JF. Myocardial and circulating levels of microRNA-21 reflect left ventricular fibrosis in aortic stenosis patients. *Int J Cardiol*. 2013; 167(6):2875–2881. DOI: 10.1016/j.ijcard.2012.07.021 [PubMed: 22882958]
113. von Haehling S. The wasting continuum in heart failure: from sarcopenia to cachexia. *Proc Nutr Soc*. 2015; doi: 10.1017/S0029665115002438
114. Waldenstrom A, Genneback N, Hellman U, Ronquist G. Cardiomyocyte microvesicles contain DNA/RNA and convey biological messages to target cells. *PLoS ONE*. 2012; 7(4):e34653. doi: 10.1371/journal.pone.0034653 [PubMed: 22506041]
115. Wang GK, Zhu JQ, Zhang JT, Li Q, Li Y, He J, Qin YW, Jing Q. Circulating microRNA: a novel potential biomarker for early diagnosis of acute myocardial infarction in humans. *Eur Heart J*. 2010; 31(6):659–666. DOI: 10.1093/eurheartj/ehq013 [PubMed: 20159880]
116. Watson CJ, Gupta SK, O'Connell E, Thum S, Glezeva N, Fendrich J, Gallagher J, Ledwidge M, Grote-Levi L, McDonald K, Thum T. MicroRNA signatures differentiate preserved from reduced ejection fraction heart failure. *Eur J Heart Fail*. 2015; 17(4):405–415. DOI: 10.1002/ehf.244 [PubMed: 25739750]
117. Widera C, Gupta SK, Lorenzen JM, Bang C, Bauersachs J, Bethmann K, Kempf T, Wollert KC, Thum T. Diagnostic and prognostic impact of six circulating microRNAs in acute coronary syndrome. *J Mol Cell Cardiol*. 2011; 51(5):872–875. DOI: 10.1016/j.yjmcc.2011.07.011 [PubMed: 21806992]
118. Wightman B, Ha I, Ruvkun G. Posttranscriptional regulation of the heterochronic gene *lin-14* by *lin-4* mediates temporal pattern formation in *C. elegans*. *Cell*. 1993; 75(5):855–862. [PubMed: 8252622]
119. Xu Y, Zhu W, Sun Y, Wang Z, Yuan W, Du Z. functional network analysis reveals versatile microRNAs in human heart. *Cell Physiol Biochem*. 2015; 36:1628–1643. [PubMed: 26160134]
120. Yi R, Qin Y, Macara IG, Cullen BR. Exportin-5 mediates the nuclear export of pre-microRNAs and short hairpin RNAs. *Genes Dev*. 2003; 17(24):3011–3016. DOI: 10.1101/gad.1158803 [PubMed: 14681208]
121. Zacharewicz E, Della Gatta P, Reynolds J, Garnham A, Crowley T, Russell AP, Lamon S. Identification of microRNAs linked to regulators of muscle protein synthesis and regeneration in young and old skeletal muscle. *PLoS ONE*. 2014; 9(12):e114009. doi: 10.1371/journal.pone.0114009 [PubMed: 25460913]
122. Zampieri S, Pietrangeli L, Loeffler S, Fruhmahn H, Vogelauer M, Burggraf S, Pond A, Grim-Stieger M, Cvecka J, Sedlak M, Tirpakova V, Mayr W, Sarabon N, Rossini K, Barberi L, De Rossi M, Romanello V, Boncompagni S, Musaro A, Sandri M, Protasi F, Carraro U, Kern H. Lifelong physical exercise delays age-associated skeletal muscle decline. *J Gerontol A Biol Sci Med Sci*. 2015; 70(2):163–173. DOI: 10.1093/gerona/glu006 [PubMed: 24550352]

123. Zhang H, Yang H, Zhang C, Jing Y, Wang C, Liu C, Zhang R, Wang J, Zhang J, Zen K, Zhang C, Li D. Investigation of microRNA expression in human serum during the aging process. *J Gerontol A Biol Sci Med Sci*. 2015; 70(1):102–109. DOI: 10.1093/gerona/glu145 [PubMed: 25165030]
124. Zhang T, Birbrair A, Wang ZM, Messi ML, Marsh AP, Leng I, Nicklas BJ, Delbono O. Improved knee extensor strength with resistance training associates with muscle specific miRNAs in older adults. *Exp Gerontol*. 2015; 62:7–13. DOI: 10.1016/j.exger.2014.12.014 [PubMed: 25560803]
125. Zhang Y, Liu D, Chen X, Li J, Li L, Bian Z, Sun F, Lu J, Yin Y, Cai X, Sun Q, Wang K, Ba Y, Wang Q, Wang D, Yang J, Liu P, Xu T, Yan Q, Zhang J, Zen K, Zhang CY. Secreted monocytic miR-150 enhances targeted endothelial cell migration. *Mol Cell*. 2010; 39(1):133–144. DOI: 10.1016/j.molcel.2010.06.010 [PubMed: 20603081]
126. Zhuo R, Fu S, Li S, Yao M, Lv D, Xu T, Bei Y. Des-regulated microRNAs in aging-related heart failure. *Front Genet*. 2014; 5:186.doi: 10.3389/fgene.2014.00186 [PubMed: 25009555]
127. Zile MR, Mehurg SM, Arroyo JE, Stroud RE, DeSantis SM, Spinale FG. Relationship between the temporal profile of plasma microRNA and left ventricular remodeling in patients after myocardial infarction. *Circ Cardiovasc Genet*. 2011; 4(6):614–619. DOI: 10.1161/CIRCGENETICS.111.959841 [PubMed: 21956146]

Fig. 1.
Theoretical mechanism by which myocardially derived miRNAs expressed during cardiac dysfunction and/or trauma may communicate with skeletal muscle to regulate mass and function