What is already known on this topic

Head louse infestation is widespread in children, and its prevalence has increased since the early 1990s

Treatment with insecticides may be affected by resistance, and combing has become more common as a treatment option

Evidence from randomised controlled trials for any form of treatment is limited

What this study adds

Dimeticone 4.0% lotion is efficacious at treating head louse infestation

Phenothrin 0.5% liquid is effective when properly applied

A high proportion of children with lice may be infested for several months despite parents' attempts to treat by various means

participants had either never used insecticides or had not used one for over three months. This may explain why we encountered little evidence for insecticide resistance through treatment failure with phenothrin, unlike recent studies in which participants were referred by general practices.³⁻⁹ The posology and formulation excipients of phenothrin liquid, however, probably contribute towards activity to overcome low levels of resistance.¹⁰ Overall, the efficacy for both products was comparable to that found for malathion lotions and permethrin.¹⁸

Our method of finding head lice by dry combing with a plastic detection comb is similar to that used by another study.¹¹ Our team is experienced in the technique. Consequently, we believe all treatment failures were identified and any potential bias due to under-reporting was eliminated.

This is the first randomised controlled trial of an insecticide-free treatment that does not require physical methods. Studies in vitro found dimeticone irreversibly immobilised lice within five minutes of application. The current treatment problems caused by resistance to neuroactive insecticides will not affect this product. The efficacy, lack of odour, and relative ease of use of 4% dimeticone lotion make it a viable alternative to conventional treatments. Most participants had used combing extensively, often with other products, but two thirds of participants had merely limited the number of lice. Half had had head lice continuously for over a year, a clear indication that current policies are not working.

Products used in this study worked well to kill lice, even when a cure was not achieved. Unlike phenothrin, however, dimeticone is not absorbed transdermally and could be used more than twice to effect a cure. Failure to cure may not be due to resistance but to application method. In some cases we found it difficult to ensure that the hair and scalp had been thoroughly covered. The amount of product used for each application of phenothrin liquid was, mostly, greater than the current 50 ml single

treatment pack. Consequently, under-dosing is probably widespread in the community. Better instructions for use and improved information at the primary care level could improve success.

Contributors: See bmj.com

Funding: Kerris Pharmaceuticals, Jersey, which had no role in the design, execution, or interpretation of the study.

Competing interests: IFB has been a consultant to various makers of pharmaceutical products, alternative therapies, and combs for treating louse infestations. PNL has analysed similar studies for other pharmaceutical companies.

Ethical approval: Ethical approval for this study was granted by Cambridge research ethics committee, and issues related to the locality were approved by Peterborough and Fenland and Huntingdon local research ethics committees. This clinical trial was monitored and audited by Covance.

- 1 Roberts RJ, Casey D, Morgan DA, Petrovic M. Comparison of wet combing with malathion for treatment of head lice in the UK: a pragmatic randomised controlled trial. *Lancet* 2000;356:540-4.
- 2 Bingham P, Kirk S, Hill N, Figueroa J. The methodology and operation of a pilot randomized control trial of the effectiveness of the bug busting method against a single application of insecticide product for head louse treatment. *Public Health* 2000;114:265-8.
- 3 Plastow L, Luthra M, Powell R, Wright J, Russell D, Marshall MN. Head lice infestation: bug busting vs. traditional treatment. *J Clin Nurs* 2001;10:775-83.
- 4 Burgess IF, Brown CM, Peock S, Kaufman J. Head lice resistant to pyrethroid insecticides in Britain. *BMJ* 1995;311:752.
- 5 Downs AMR, Stafford KA, Harvey I, Coles GC. Evidence for double resistance to permethrin and malathion in head lice. *Br J Dermatol* 1999;141:508-11.
- 6 Downs AMR, Stafford KA, Coles GC. Head lice prevalence in schoolchildren and insecticide resistance. *Parasitol Today* 1999;15:1-4.
- 7 Dodd CS. Interventions for treating head lice. Cochrane database of systematic reviews. In: *Cochrane library*, Issue 1, Oxford: Update Software; 2003.
- 8 Meinking TL, Clineschmidt CM, Chen C, Kolber MA, Tipping RW. An observer-blinded study of 1% permethrin crème rinse with and without adjunctive combing in patients with head lice. *J Pediatr* 2002;141:665-70.
- 9 Hill N, Moor G, Cameron MM, Butlin A, Preston M, Williamson MS. A single blind, randomised controlled trial comparing the bug buster kit with over the counter insecticide treatments for head lice in the UK. *BMJ* 2005;(in press).
- 10 Burgess IF. Dermatopharmacology of ectoparasitics and insect repellents. *Dermatopharmacology of topical preparations*. ed. In: B Gabard, P Elsner, C Surber, P Treffel, eds. Berlin: Springer, 1999:157-78.
- 11 Mumcuoglu KY, Friger M, Ioffe-Uspensky I, Ben-Ishai F, Miller J. Louse comb versus direct visual examination for the diagnosis of head louse infestations. *Pediatr Dermatol* 2001;18:9-12.

(Accepted 15 May 2005)

doi 10.1136/bmj.38497.506481.8F

Endpiece

An egotistical and dictatorial style

The author is aware that in some parts of his volume he has been betrayed into the adoption of an egotistic, in other parts of a dictatorial style. For this he craves the indulgence of the reader. Almost everything here recorded has passed under his immediate observation. Ardently and deeply interested in all that he has described, he so completely identified self with subject, that he found it impossible to separate them. Where he assumes a dictatorial language, it must be remembered that he is not insisting upon the efficacy of his individual practice, but upon that of Physicians who have long distinguished themselves by an eminent display of talent, and who now most deservedly enjoy the public confidence.

Beddingfield J. *A Compendium of Medical Practice, Illustrated by Interesting and Instructive Cases, and by Practical, Pathological, and Physiological Observations*. London: Highley, 1816: ix-x

Jeremy Hugh Baron, honorary professorial lecturer, Mount Sinai School of Medicine, New York