Published in final edited form as:

Drug Alcohol Depend. 2016 November 01; 168: 203. doi:10.1016/j.drugalcdep.2016.09.012.

Corrigendum to “Concordance between DSM-5 and DSM-IV nicotine, alcohol, and cannabis use disorder diagnoses among pediatric patients” [*Drug Alcohol Depend.* 140 (2014) 213–216]

Sharon M. Kelly^{a,*}, Jan Gryczynski^a, Shannon Gwin Mitchell^a, Arethusa Kirk^b, Kevin E. O’Grady^c, and Robert P. Schwartz^a

^aFriends Research Institute, Inc., 1040 Park Avenue, Suite 103, Baltimore, MD 21201, USA

^bTotal Health Care, 1501 Division Street, Baltimore, MD 21217, USA

^cDepartment of Psychology, University of Maryland, College Park, MD 20742, USA

We mistakenly stated that items from the Composite International Diagnostic Interview-2 Substance Abuse Module (CIDI-2 SAM) were used to assess substance use disorders. This should have read that items from the WHO World Mental Health-CIDI (WMH-CIDI; <http://www.hcp.med.harvard.edu/wmhcid/index.php>) were used to assess substance use disorders. Because the WMH-CIDI does not contain all the criteria for DSM-5 nicotine use disorder, the WMH-CIDI’s drug section items were adapted to assess nicotine use disorder. Readers of our article should note the mislabeling of the CIDI in the Abstract on page 213 and the Methods section on page 214 under Measures. This error does not alter the analysis or findings of the study.

*Corresponding author. skelly@friendsresearch.org (S.M. Kelly).