WHO takes up issue of child abuse

In efforts to stop violence against children, the World Health Organization is launching a campaign in which the health sector around the world will play a part. According to WHO, every year across Europe and Central Asia, at least 1300 children die after being beaten or otherwise mistreated by their parents, teachers, or carers. Other deaths are misreported as suicides or accidents.

Since 2002, WHO has been arguing that violence is a predictable and preventable health problem. But the overwhelming majority of health centres and many family doctors still see child abuse as mainly the concern of the police and social services.

"What we need is a change of mindset," said Roberto Bertollini, chief of WHO's environment and health programme for Europe. "The public health approach is a new way of looking at this problem."

Although police investigate criminals, Dr Bertollini said that health professionals are best placed to gather vital evidence of the harm that adults inflict on their small victims. Standardised procedures must be established internationally to register deaths and injuries.

Sophie Arie *Ljubljana*

Rules for human organs put out for consultation

New draft guidelines for the storage, use, and disposal of human organs and tissue, including donations for transplantation, were issued this week by the new Human Tissue Authority.

The authority, set up under the Human Tissue Act 2004, launched its first round of public consultation with five draft codes covering the key areas of consent; donation for transplantation; post mortem examination; anatomical examination; and removal, storage, and disposal. Consultation ends on 4 October.

The authority was set up in response to the revelation in the 1990s that doctors at several UK

hospitals had retained thousands of hearts, brains, and other body parts from dead children without the parents' knowledge.

The draft code on consent is intended to guide clinicians in communicating with patients and families to ensure that both sides appreciate exactly what is being consented to.

A consultation was also launched this week by the Department of Health on the draft regulations to implement the Human Tissue Act, including the scope of licensing of the storage of human tissue for research purposes and the circumstances in which live donations of tissues or organs, from both related and unrelated donors, will be allowed.

The draft codes of practice are available at www.hta.gov.uk. The draft regulations are available at www.dh.gov.uk.

Clare Dyer *legal correspondent, BMJ*

Smoking ban also cuts smoking at home

A total ban on smoking in enclosed public places would reduce smoking in the home and the effects of passive smoking, argued a study published last week by the Royal College of Physicians.

The study said that the 12 200 deaths each year in the United Kingdom from passive smoking were entirely preventable. It recommended that preventing passive smoking at home was a "public health priority" and that helping smokers to stop by having a ban in enclosed public places would lead to less smoking in the home.

Members of the Royal College of Physicians' Tobacco Advisory Group said that the previous health secretary, John Reid, was wrong to have claimed that a ban on smoking in public places would increase smoking in the home. They noted that the number of smoking households that became smoke free increased from 22% to 37% between 1996 and 2003. And they attributed this to increased restrictions on smoking in public places in that period.

Adrian O'Dowd *London*

Teenagers need sex education, not just abstinence advice

Janice Hopkins Tanne *New York*

In revised guidelines, the American Academy of Pediatrics has recommended that teenagers receive counselling to postpone sexual activity and also information about and access to contraception, including over the counter emergency contraception without a prescription (*Pediatrics* 2005;115:281-6). The academy's 60 000 members provide primary care to infants, children, and teenagers up to the age of 18 or 21.

Jonathan Klein, chairman of the committee that drew up the guidelines, told the *BMJ* that the new report was a routine update to consider newer options such as emergency contraception, not a reaction to the Bush administration's programmes promoting abstinence until marriage and monogamy thereafter. The report says that teenage pregnancy poses medical and psychosocial complications, and that children of very young mothers do not do as well as those of adult women.

Although sexual activity and teen pregnancy rates have declined, they remain a concern. More than 45% of girls and 48% of boys at high school (aged 13 or 14 to 17 or 18) have had sexual intercourse. The average age of first intercourse is 17 for girls and 16 for boys, but about a quarter of teenagers report having had intercourse by age 15.

Every year, 900 000 teenage girls become pregnant. Half of teenagers' pregnancies are with-

in six months of first intercourse. More than 40% of adolescent girls become pregnant at least once before the age of 20, although most pregnancies are in teenagers aged 18 or 19. Education about HIV and sexually transmitted disease has led more teenagers to use barrier contraceptives, but only 63% reported having used a condom the last time they had intercourse.

Births to teenagers aged 15 to 19 in the US are far higher than in other industrialised countries: 61 per 1000 in the US, compared with 41 in Canada, 33 in the United Kingdom, eight in the Netherlands, and four in Japan.

Dr Klein, of the University of Rochester School of Medicine, New York, said that the abstinence programme, which costs up to \$87.5m (£50.4m; €73.3m) a year and is shared by the federal government and the states, "was not based on medical or health information." He told the *BMJ*, "The academy believes young people and families ought to have access to comprehensive reproductive health information."

A report to the federal Department of Health and Human Services in June showed that abstinence programmes increased teenagers' awareness of abstinence but did not decrease teenagers' sexual activity, pregnancy, or infection with sexually transmitted diseases (<http://aspe.hhs.gov/hsp/05/abstinence/report.pdf>).

Margaret Hill, aged 19, from Mississippi, takes part in a rally on Capitol Hill to promote fidelity. Now doctors say that teaching abstinence is not enough