Published in final edited form as:

J Am Chem Soc. 2017 September 06; 139(35): 12259–12266. doi:10.1021/jacs.7b06319.

Detection of Fleeting Amine Radical Cations and Elucidation of Chain Processes in Visible Light-Mediated [3+2] Annulation by Online Mass Spectrometric Techniques

Yi Cai^{†,§}, Jiang Wang^{†,§}, Yuexiang Zhang^{†,§}, Zhi Li[†], David Hu[†], Nan Zheng^{*,‡}, and Hao Chen^{*,†}

[†]Department of Chemistry and Biochemistry, Center of Intelligent Chemical Instrumentation, Edison Biotechnology Institute, Ohio University, Athens, OH 45701

[‡]Department of Chemistry and Biochemistry, University of Arkansas, Fayetteville, AR 72701

Abstract

Visible-light mediated photoredox reactions have recently emerged as a powerful means for organic synthesis and thus have generated significant interest from the organic chemistry community. Although the reaction mechanisms have been probed by a number of techniques such as NMR, fluorescence quenching, and laser flash photolysis and various degree of success has been achieved, mechanistic ambiguity still exists (for instance, the involvement of the chain mechanism is still under debate) due to lack of structural information about the proposed and short-lived intermediates. Herein, we present the detection of transient amine radical cations involved in intermolecular [3+2] annulation reaction of *N*-cyclopropylaniline (CPA, **1**) and styrene **2** by electrospray ionization mass spectrometry (ESI-MS) in combination with online laser irradiation of the reaction mixture. In particular, the reactive CPA radical cation **1**⁺, the reduced photocatalyst Ru(I)(bpz)₃⁺, and the [3+2] annulation product radical cation **3**⁺ are all successfully detected and confirmed by high resolution MS. More importantly, the post-irradiation reaction with an additional substrate, the isotope-labeled CPA following photolysis of **1**, **2** and Ru catalyst provide strong evidences to support the chain mechanism in the [3+2] annulation reaction. Furthermore, the key step of the proposed chain reaction, the oxidation of CPA **1** to amine radical cation **1**⁺ by product radical cation **3**⁺ (generated using online electrochemical oxidation of **3**) is successfully established. Additionally, the coupling of ESI-MS with online laser irradiation has been successfully applied to probe the photostability of photocatalysts.

Graphical abstract

*Corresponding Author: nzhenh@uark.edu, chen2@ohio.edu.

§These authors contributed equally.

Author Contributions

The manuscript was written through contributions of all authors. All authors have given approval to the final version of the manuscript.

Supporting Information

Experimental details, apparatus photos and additional MS, MS/MS data and quantum yield experimental results The Supporting Information is available free of charge on the ACS Publications website.

Introduction

Photoredox catalysis has recently become a topic of intense study in the field of organic chemistry.¹ The heightened interest in this topic is likely attributed to photoredox catalysis' versatility in generating radicals using light under mild conditions, a specific mode of activation, which is different from how radicals are produced under non-photo conditions.² As such, it provides means to explore radical chemistry that is complementary to those under non-photo conditions. Moreover, photoredox catalysis' ability to merge with other types of catalysis to form dual catalysis further expands its utility far beyond the scope of traditional radical chemistry.³

In contrast to furious activity in method development, mechanistic studies of photoredox catalysis have been rather scarce.⁴ In those published and limited results, there exists controversy, such as whether photoredox reactions involve chain processes. This is an important mechanistic question in method development as different approaches are required to optimize the two processes separately. Light/dark experiments have been often used to differentiate them.⁴⁻⁶ However, there remains the ambiguity of interpreting light/dark experiments' results, in which the chain mechanism was established by other methods whereas light/dark experiments disapproved its involvement.^{6a, 6b} Quantum yield measurements are in principle a more general and sensitive method to differentiate the two competing processes. However, a wide range of quantum yields from 1.3 to 77 across several types of photoredox reactions have been reported,^{4g-i, 6b} which makes distinction of the two processes more difficult, particularly for those with quantum yields close to 1. Clearly, there is a strong need for development of new tools that allow for explicit elucidation of photoredox reaction mechanisms including identification of both the photoredox process and the chain mechanism.

Photoredox reactions pose a significant challenge to chemists who are interested in studying their mechanisms. They are typically fast and reversible, and the intermediates are short-lived. A variety of techniques such as NMR, EPR, fluorescence quenching, quantum yield measurement, and laser flash photolysis have been employed to study photoredox reactions.^{4,6} However, these spectroscopic methods usually lack structural information of the transient intermediates. Mass spectrometry (MS) has become a powerful technique for

detecting and characterizing short-lived reaction intermediates as well as studying reaction mechanisms, since the advent of soft ionization methods such as electrospray ionization (ESI)⁷ and desorption electrospray ionization (DESI)⁸. The inherent high sensitivity and rich mass information provided by MS measurements distinguish it from other commonly used methods. Moreover, in comparison to traditional spectroscopic approaches, it is a general technique and does not need chromophore-carrying substrates for investigating reaction mechanisms. However, the probed species such as substrates, catalysts, intermediates, or products must carry a charge, and sometimes the detection can suffer from ion suppression effect due to the matrix effect.

We recently developed an intermolecular [3+2] annulation of *N*-cyclopropylanilines with alkenes by photoredox catalysis (Scheme 1).⁹ This reaction displays some excellent features for a synthetic method such as broad functional group compatibility, 100% atom economy, and an overall redox-neutral process. The scope of this reaction was later expanded to include various pi bonds.¹⁰ These results have opened new avenues for the use of aniline-substituted cyclopropanes as synthetic building blocks.¹¹ Mechanistically, these methods are all believed to proceed through ring opening of the amine radical cations of *N*-cyclopropylanilines. This process has been previously studied by EPR^{12a} and electrochemical methods^{12b}. However, it has not been studied in the context of a complete catalytic cycle, which is more relevant to our efforts in expanding this reaction class to include other types of amine-substituted carbocycles. Therefore, we were eager to study the complete catalytic cycle of the [3+2] annulation reaction.

Herein, we designed a series of online ESI-MS apparatus (Figure 1) that allowed to directly detect the short-lived intermediates including the substrate radical cation **1**⁺ and the product radical cation **3**⁺ from the visible light-mediated [3+2] annulation catalyzed by Ru(II) (bpz)₃²⁺ (see structures of **1**⁺ and **3**⁺ in Scheme 2).⁹ Although MS in combination with online photolysis (e.g., using lamp or laser to irradiate a reaction mixture prior to MS detection) has been previously reported in literature,^{7q,13} this type of setup has been rarely used to study the mechanism of visible-light triggered photoredox reactions.^{7q} Moreover, none of the reported MS studies provided any conclusive evidence for the chain reaction mechanism involved in photoredox reactions. Our setup not only allows for the direct detection of elusive reaction intermediates via online photolysis using a laser pointer (Figure 1a), but also enables coupling with an online microreactor to study the reactivities of the reaction intermediates. For instance, another reagent such as CPA-d₅ was introduced downstream via a second channel (Figure 1b). In addition, for those intermediates that could not be generated photochemically (e.g., the radical cation of the [3+2] annulation product **3**⁺), electrolysis was employed as an alternative way to generate radical cation **3**⁺ whose reactivity could then be online monitored by MS (Figure 1c). Combination of electrochemistry (EC) with MS has been one of our research focuses which has applications for studying redox chemistry of both small organic molecules and large protein molecules.^{8d,14} In this study, using the setup, we obtained unequivocal evidence to support the involvement of the chain processes in the [3+2] annulation reaction. Since the setup is flexible and modular, we believe that it can be easily adopted to study other photoredox reactions.

Results and discussion

Apparatus

A high resolution Q Executive Plus hybrid quadrupole-Orbitrap mass spectrometer (Thermo Fisher Scientific, San Jose, CA) was used in this study. As shown in Figure 1a, to probe reaction intermediates, the commercial ESI ion source was removed to accommodate an in-house-made ESI source. A piece of fused Silica capillary (100 μm i.d., 198 μm o.d.) was used to deliver the reaction mixture solution for online photolysis and online electrospray ionization (Figure 1a). An irradiation window on the silica capillary (1 cm wide), located ca. 5 cm upstream from the ESI emitter tip, was made by burning away the organic capillary coating with flame. A laser pointer (403 nm, 50 mW, LaserPointerPro, HK) was employed as a light source to irradiate the reaction solution in the capillary to trigger the photoredox reaction as the solution flowed by the irradiation window. The laser pointer was held about 5 cm away from the irradiation window. A high voltage +5 kV was applied to ionize the reaction mixture, with the assistance of 170 psi N_2 nebulization gas. The reaction solutions were all rigorously degassed with argon before irradiation. The flow rate for infusing the reaction mixture solution through the silica capillary was 100 $\mu\text{L}/\text{min}$. The transportation time for the reaction mixture flowing from the irradiation window to the ESI emitter tip was about 236 ms (dead volume = 0.393 μL , flow rate = 100 $\mu\text{L}/\text{min}$).

To investigate the chain propagation process, the original MS setup described in Figure 1a was modified to perform a sequential reaction experiment (Figure 1b), in which a reaction mixture (e.g., a mixture of CPA **1**, styrene **2** and $\text{Ru(II)(bpz)}_3(\text{PF}_6)_2$ **4**) was introduced into the silica capillary via channel 1 (50 $\mu\text{L}/\text{min}$), photo-excited by the laser pointer, and then mixed with an isotope-labeled new substrate (introduced via channel 2, 50 $\mu\text{L}/\text{min}$). Upon mixing, the added substrate reacted with the photo-excited reaction mixture in the downstream capillary served as a microreactor, prior to MS detection. Because the isotope-labeled substrate was added after the photolysis of the reaction mixture, positive identification of the product derived from the added isotope-labeled substrate could be used as evidence to support the chain propagation process. MS' inherent high sensitivity enabled detection of the product in minute amount, which made this apparatus far more sensitive than dark/light experiments.

Alternatively, photolysis could be replaced by electrolysis to generate the amine radical cation of product **3**. As shown in Figure 1c, an electrochemical flow cell with a magic diamond electrode (details shown in the text below) was used in our experiment. Amine radical cation $\mathbf{3}^{+\bullet}$ was then mixed with CPA **1** introduced via channel 2 to examine its reactivity by online MS monitoring. In this case, no high voltage was applied to the ESI source in order to avoid the voltage conflict issue between the electrochemical cell and the ion source. Photos for all of the experimental set-ups mentioned above are included in the Supporting Information (Figure S1).

Detection of Amine Radical Cations

N-cyclopropylaniline (CPA) **1** and styrene **2** were chosen as the model substrates (Scheme 1) to investigate the intermolecular [3+2] annulation catalyzed by $\text{Ru(II)(bpz)}_3(\text{PF}_6)_2$ **4** using

the ESI-MS set-up with online photolysis shown in Figure 1a. We selected this pair of substrates because both were extensively studied in the initial development of the annulation reaction.^{9,10}

As previously proposed (Scheme 2),⁹ upon irradiation, $\text{Ru(II)(bpz)}_3^{2+}$ is promoted to the photoexcited triplet state $\text{Ru(II)*}(\text{bpz})_3^{2+}$, which oxidizes CPA **1** to amine radical cation $\mathbf{1}^{+\bullet}$ with the concomitant formation of Ru(I)(bpz)_3^+ . Amine radical cation $\mathbf{1}^{+\bullet}$ subsequently undergoes ring opening and then add intermolecularly to styrene **2** to produce the [3+2] annulation product radical cation $\mathbf{3}^{+\bullet}$. Reduction of $\mathbf{3}^{+\bullet}$ by $\text{Ru}(\text{bpz})_3^+$ completes the catalytic cycle. Alternatively, a chain mechanism in which amine radical cation $\mathbf{3}^{+\bullet}$ oxidizes CPA **1** to a new radical cation $\mathbf{1}^{+\bullet}$ while itself is reduced to the product **3**, allowing closure of the catalytic cycle.

To probe the above reaction mechanism, online irradiation and detection experiments using the aforementioned apparatus (Figure 1a) were first carried out. In this experiment, we started with the examination of the reaction between $\text{Ru(II)(bpz)}_3(\text{PF}_6)_2$ **4** and CPA **1** under the photolysis condition. A mixture of $[\text{Ru(II)(bpz)}_3](\text{PF}_6)_2$ **4** (0.0004 mmol, 100 μM) and CPA **1** (0.02 mmol, 5 mM) in 4 mL of CH_3NO_2 was infused at the flow rate of 100 $\mu\text{L}/\text{min}$ through the silica capillary for ESI-MS detection after argon degassing. Without irradiating the reaction solution flowing through the capillary by laser, an ESI-MS spectrum was acquired (Figure 2a). The ions of m/z 134 (measured m/z : 134.0966, theoretical m/z 134.0964, error +1.5 ppm) and m/z 288 (measured m/z : 288.0408, theoretical m/z 288.0405, error -1.0 ppm) were detected, corresponding to the protonated CPA $[\mathbf{1}+\text{H}]^+$ and $\text{Ru(II)(bpz)}_3^{2+}$, respectively. We noticed that there was a small peak for $\mathbf{1}^{+\bullet}$ (measured m/z : 133.0888, theoretical m/z 133.0886, error +1.5 ppm, Figure 2a inset) with intensity of 1.69E6 (manufacturer arbitrary unit). We attributed the formation of $\mathbf{1}^{+\bullet}$ to in-source oxidation of **1** during the ESI ionization process,¹⁵ as it was still observed in the absence of the Ru catalyst **4**. Furthermore, the reduced Ru(I) species was not observed.

Next, when the laser pointer was turned on to irradiate the mixture in the capillary through the irradiation window, the intensity of $\mathbf{1}^{+\bullet}$ peak increased by 17 fold (Figure 2b inset), which was ascribed to the photo-oxidation of **1**. Amine radical cation $\mathbf{1}^{+\bullet}$ could undergo ring opening to furnish the distonic radical cation. However, ESI-MS can't differentiate the two species because they have the same mass. Additionally, the reduced Ru catalyst peak of Ru(I)(bpz)_3^+ m/z 576 (measured m/z : 576.0815, theoretical m/z : 576.0815, error 0.0 ppm) was also detected. The inset in Figure 2b clearly showed the zoomed-in spectrum of Ru(I)(bpz)_3^+ at m/z 576 (in black), which matched well with its theoretical isotopic peak distribution (in red). This experiment result supported the photooxidation of CPA **1** by the photoexcited $\text{Ru(II)*}(\text{bpz})_3^{2+}$.^{2d} A control study in which CPA **1** from the reaction mixture was replaced by styrene **2** (Figure S2, Supporting Information) showed no formation of Ru(I)(bpz)_3^+ , excluding the possibility of the redox reaction between $\text{Ru(II)*}(\text{bpz})_3^{2+}$ and styrene **2**. Both results were consistent with the Stern Volmer studies in which CPA **1** was shown to be an effective quencher for $\text{Ru(II)*}(\text{bpz})_3^{2+}$ but styrene **2** was not (Figure S14 and Figure S16, Supporting Information).

Then, we monitored the [3+2] annulation of CPA **1** and styrene **2** in CH₃NO₂ with the Ru catalyst **4** using ESI-MS. A degassed mixture of [Ru(bpz)₃](PF₆)₂ **4** (0.0004 mmol, 100 μM), CPA **1** (0.02 mmol, 5 mM), and styrene **2** (0.4 mmol, 0.1 M) in 4 mL CH₃NO₂ was irradiated inside the capillary by the laser pointer, and the reaction was monitored online by ESI-MS. Under irradiation, the protonated [3+2] annulation product [3+H]⁺ (*m/z* 238) (measured *m/z*: 238.1593, theoretical *m/z* 238.1590, error +1.2 ppm) was clearly observed (Figure 2c). Upon collision-induced dissociation (CID), the ion *m/z* 238 gave rise to a fragment ion of *m/z* 145 by loss of C₆H₅NH₂ (Figure S3-a, Supporting Information), consistent with its assigned structure. Furthermore, the CID behavior of this ion was identical to that of the protonated product at *m/z* 238 generated from ESI of the authentic product compound **3** (Figure S3-b, Supporting Information), reinforcing the ion assignment. Although a small peak for the CPA radical cation **1**^{•+} was seen (Figure 2c inset, its intensity similar to that seen in Figure 2a), it was again probably due to the in-source ESI oxidation of the unreacted CPA **1**. There were a few unassigned peaks in Figure 2c, whose suggested chemical formulas were included in the Supporting Information (Table S1, Supporting Information). Surprisingly, the product radical cation **3**^{•+} was not observed in the acquired MS spectrum under this condition, which will be discussed further subsequently. Notably, Ru(I)(bpz)₃⁺ (*m/z* 576) could not be observed in Figure 2c, presumably because it was consumed in the annulation reaction. This data supported the photoredox process in which Ru(I)(bpz)₃⁺ reduced the product radical cation **3**^{•+} to complete the catalytic cycle (Scheme 2). The two processes, the photoredox process and the chain mechanism, likely competed against each other and intertwined throughout the reaction.

We encountered difficulties in detection of the product radical cation **3**^{•+} during our studies of the online photolysis of CPA **1** and styrene **2** in the presence of the Ru catalyst **4** (Figure 2c). We surmised that higher concentration of CPA **1** would result in a faster chain reaction between the product radical cation **3**^{•+} and CPA **1**, and thus would shorten the lifetime of **3**^{•+}. Alternatively, higher concentration of CPA **1** could lead to higher concentration of Ru(I) via reduction of the excited Ru(II) complex, which could also shorten the lifetime of **3**^{•+} via the photoredox process (Scheme 2).

To test this hypothesis and to observe the radical cation **3**^{•+}, we performed an ESI-MS monitoring experiment by lowering the concentration of CPA **1** from 5 mM (0.02 mmol in 4 mL CH₃NO₂) to 10 μM (0.04 μmol in 4 mL CH₃NO₂) for the online photoredox reaction while maintaining the concentrations of styrene **2** and Ru catalyst **4** as those experiments shown in Figure 2 (0.1 M styrene **2** and 100 μM Ru **4**). Indeed, when CPA **1** was diluted to 100 μM, the product radical cation **3**^{•+} at *m/z* 237.1512 (Figure 3c, measured *m/z*: 237.1512, theoretical *m/z* 237.1512, error 0.0 ppm) was observed. The signal of **3**^{•+} appeared only when the laser pointer was turned on, supporting that it was a photochemical product. Upon further dilution of CPA **1**, the peak intensity of the CPA radical cation **3**^{•+} decreased (Figure 3d). When the concentration of CPA **1** was lowered to 10 μM, the peak at *m/z* 237 was not observed (data not shown), probably because the concentration of the resulting **3**^{•+} was below the detection limit of our MS instrument. Notably, the observed **3**^{•+} was not an artifact of the insource ESI oxidation product. In a control experiment, ESI of the authentic product **3** did not give rise to observable signal of **3**^{•+}. Furthermore, direct irradiation of **3** in the

presence of Ru catalyst **4** did not produce $3^{+\cdot}$, either. All these results not only showed the existence of the proposed product radical cation $3^{+\cdot}$ in the [3+2] annulation reaction, but also verified our hypothesis that a low concentration of CPA **1** would favor the observation of $3^{+\cdot}$.

In order to further confirm the assignment of observed $3^{+\cdot}$ and $[3+H]^+$ ions, styrene **2** was replaced by styrene- α , β , β -d₃ in the annulation reaction under otherwise identical conditions. The protonated final product-d₃ **3**-d₃ at m/z 241 (measured m/z : 241.1778, theoretical m/z 241.1779, error -0.4 ppm) and the product-d₃ radical cation **3**-d₃ $^{+\cdot}$ at m/z 240.1699 (Figure S5b inset, measured m/z : 240.1699, theoretical m/z 240.1700, error -0.4 ppm) were both observed (Figure S5, Supporting Information).

Chain Mechanism investigation

a. Post-irradiation chain propagation—Encouraged by observing the dependence of detecting $3^{+\cdot}$ on CPA **1** concentration, we decided to investigate the chain propagation from a different angle, post-irradiation. The original MS setup described in Figure 1a was modified to perform a sequential reaction experiment (Figure 1b). A mixing Tee was used to introduce CPA-d₅ **1**-d₅ into the silica capillary microreactor to further react with the irradiated reaction mixture of CPA **1**, Ru(II) catalyst **4**, and styrene **2**. A degassed solution of CPA **1** (100 μ M), [Ru(bpz)₃](PF₆)₂ **4** (0.0004 mmol, 100 μ M), and styrene **2** (40 μ L, 0.4 mmol, 0.1 M) was infused via channel 1 for irradiation. The degassed solution of CPA-d₅ **1**-d₅ (100 μ M) was infused via channel 2. The injection flow rate was 50 μ L/min for both channels. This modified setup allowed that the channel 1 reaction solution was first irradiated by the blue laser pointer and then mixed with CPA-d₅ **1**-d₅ from channel 2 without light irradiation. If the chain process were involved, the radical cation intermediates $3^{+\cdot}$ would induce CPA-d₅ to participate in the [3+2] annulation reaction to afford the annulation product-d₅.

In a control study when there was no light irradiation (Figure 4a), m/z 134 and m/z 139 were observed, corresponding to the protonated CPA $[1+H]^+$ and CPA-d₅ $[1\text{-}d_5+H]^+$, respectively. When the laser pointer was turned on for irradiation, the protonated [3+2] annulation product $[3+H]^+$ peak m/z 238 was clearly observed. More importantly, a new peak corresponding to the protonated [3+2] annulation product-d₅ $[3\text{-}d_5+H]^+$ at m/z 243 was also detected (measured m/z : 243.1900, theoretical m/z : 243.1904, error 1.6 ppm, Figure 4b inset). Moreover, before adding CPA-d₅ through channel 2 (only solvent was introduced via channel 2), the product radical cation $3^{+\cdot}$ at m/z 237 was observed (Figure 4b inset). After the mixing, the product radical cation $3^{+\cdot}$ disappeared. The consumption of the product radical cation $3^{+\cdot}$ with concurrent formation of the CPA product-d₅ lent another strong evidence to support the chain propagation.

To rule out the possibility that the observed $[3\text{-}d_5+H]^+$ resulted from the oxidation of CPA-d₅ by the pre-excited photocatalyst, a second control experiment was carried out in which a degassed solution of [Ru(bpz)₃](PF₆)₂ **4** (0.0004 mmol, 100 μ M) and styrene **2** (0.4 mmol, 0.1 M) was infused via channel 1 for irradiation (CPA **1** was omitted), and the degassed solution of CPA-d₅ (100 μ M) was infused via channel 2. The injection flow rate for both channels was kept as 50 μ L/min. The protonated CPA-d₅ reaction product $[3\text{-}d_5+H]^+$ at m/z

243 was not observed (Figure S6, Supporting information). Presumably, because the lifetime of the photo-excited **4** is much shorter than the transportation time for the reaction mixture to flow from the irradiation window to the Tee mixer ($0.74 \mu\text{s}^{1a}$ vs. about 472 ms), the photochemically pre-excited Ru catalyst **4** could be deactivated before reaching CPA-d₅.

b. Oxidation of substrate 1 by 3⁺—The remaining task in fully establishing the chain propagation process was to validate the oxidation of CPA **1** by the radical cation of the annulation product **3⁺** (Figure 5a). We initially attempted the generation of **3⁺** by photolysis of the authentic reaction product **3** in the presence of Ru catalyst **4**. The targeted **3⁺** was not observed, even though Stern Volmer studies showed that product **3** was an effective quencher for the photoexcited **4** albeit somewhat less effective than CPA **1** (Figures S14 and S15, Supporting Information). We then attempted chemical oxidation of CPA **1** using a single-electron oxidant, tris(4-bromophenyl)ammoniumyl hexachloroantimonate. Although **3⁺** was successfully generated, a large amount of the oxidant remained. Because the unreactive oxidant could directly oxidize **1** and thus interfere with the targeted oxidation of **1** by **3⁺**, we had to abandon this approach, too.

Finally, we decided to focus on generation of **3⁺** using electrochemical oxidation of **3**. The setup described in Figure 1a was modified with addition of a thin-layer electrochemical flow cell to perform the electrochemical oxidation (Figure 1c). A magic diamond electrode ($12 \times 30 \text{ mm}$) was used with LiOTf as the supporting electrolyte. The flow electrochemical cell was connected to a Tee mixer through which CPA **1** was introduced. A solution of the annulation product **3** (1 mM) and LiOTf (1 mM) in CH₃CN was infused into the flow cell and CH₃CN was first infused via channel 2. A potentiostat was used to apply potential for electro-oxidation. The injection flow rate for both channels was 10 $\mu\text{L}/\text{min}$. When a +3.0 V voltage was applied to the flow cell to trigger the electrochemical oxidation of **3**, the targeted radical cation **3⁺** was observed (Figure 5b, black line). When the solvent CH₃CN, was replaced by CPA **1** in CH₃CN (100 μM) in channel 2, the signal of **3⁺** decreased (Figure 5b, red line) with concomitant formation of the CPA radical cation **1⁺** (Figure 5c). This set of data validated the oxidation of CPA **1** by the product radical cation **3⁺**, which was also supported by our cyclic voltammetric measurements of **1** and **3** (Figure S8, Supporting Information). Both have a very similar oxidation potential (ca. 0.8 V vs. SCE).

To rule out that the observed CPA radical cation **1⁺** was formed by reacting with other species in the oxidized solvent, a control experiment was performed, in which only LiOTf (1 mM) in CH₃CN (without **3**) was infused into the cell for oxidation under the same condition and then mixed with CPA **1** (100 μM). In this control experiment, no CPA radical cation **1⁺** was generated (Figure S7, Supporting Information), indicating that no other species from the electrochemical cell except **3⁺** could oxidize CPA **1**.

c. Quantum yield measurement—To corroborate our MS studies on elucidation of the chain mechanism, we measured the quantum yield of the [3+2] annulation of 3,5-dimethylcyclopropylaniline **5** and styrene **2**. Using potassium ferrioxalate as an actinometer and a 300 W Xenon lamp (50% of light intensity, $439 \pm 5 \text{ nm}$ bandpass filter high transmittance), we determined the quantum yield over the time interval of 15 min, 30 min,

and 60 min (Figure S9–S12, Supporting Information).^{4g,4i,6b} The quantum yield was calculated to be 1.64, 1.68, and 1.44, respectively, which maintained above one during the course of the experiment and supported the chain mechanism. The low quantum yield on its own could lead to ambiguity in deciphering the chain mechanism,^{4g} and therefore would require additional studies. However, our MS studies provided unequivocal evidence to support the chain mechanism.

Based on the low quantum yield we obtained for the [3+2] annulation, we anticipated that light/dark experiments might not reveal the chain mechanism. Not surprisingly, similar to Stephenson's,^{6a} Yoon's,^{6b} and others'^{4i,5} results, we observed that the annulation occurred only in the presence of light and stalled when the light source was off (Figure 6, details shown in Supporting Information). Again as with other reports, this result revealed the limitation of light/dark experiments to disapprove the chain mechanism. The failure of light/dark experiments in distinguishing photoredox mechanisms from chain mechanisms is presumably due to the premise that most of the reactions under photoredox catalysis contain a short radical chain. Unfortunately, light/dark experiments do not have response fast enough to detect the radical chain. Our apparatus (Figure 1) not only possesses the inherently fast response of online ESI-MS methods, but also enables coupling with an online photo microreactor or an electrochemical cell to monitor post-irradiation reaction products (e.g., the [3+2] annulation product of CPA-d₅ in the dark, Figure 4b; oxidation of CPA **1** by electrochemically generated **3**⁺, Figure 5). Both features make our apparatus a powerful tool to detect chain processes in photoredox reactions

Photostability of catalyst

Photostability is one of the key features for a good photocatalyst. However, because of the presence of various *insitu* generated radical species, deactivation of photocatalysts by adding one of these radical species to the catalyst's ligand is often encountered.¹⁶ Recently, the Stephenson group delineated a deactivation pathway for Ir(ppy)₃ via ligand functionalization using a combination of kinetic, synthetic, and spectroscopic tools.¹⁷ We were wondering whether we could use the online irradiation MS to study the potential deactivation pathway in the [3+2] annulation. This is particularly relevant to the catalyst used in the annulation reaction, [Ru(bpz)₃](PF₆)₂ **4**, which has shown to be less stable than other Ru complexes based on our observations. As shown in Figure 2b, in the absence of styrene **2**, the new peak at *m/z* 354 was observed, which was corresponding to the Ru complex monoalkylated by CPA. Upon CID (Figure S4, Supporting Information), this ion dissociated into fragment ions [CPA-H]⁺ (*m/z* 132), [354-CPA]²⁺ (*m/z* 288), [354-PhNHCH=CH₂]²⁺ (*m/z* 295), [354-PhNH₂]²⁺ (*m/z* 308), and [Ru(bpz)₂-H]⁺ (*m/z* 417), which indeed suggests that it is a covalently-bonded adduct ion of CPA and Ru catalyst. However, ligand shedding to the catalyst was not observed. In addition, to our surprise, in the presence of both CPA **1** and styrene **2**, the Ru complex **4** remained intact (Figure 2c). This interesting observation shed light on the premise that a less robust photocatalyst such as Ru(bpz)₃(PF₆)₂ **4** could still be used as long as there were viable reactions to be catalyzed.

Conclusions

The recent surge of activity in visible light photocatalysis has prompted organic chemists to study the mechanism of various reactions under photoredox catalysis. However, the mechanistic studies of these reactions prove challenging, as there is lack of analytic tools that are capable of detecting transient intermediates in low quantity while also providing rich structural information. The ESI-MS set-up with online photolysis, which requires little amount of materials (e.g., 400 nanomoles of both $[\text{Ru}(\text{bpz})_3](\text{PF}_6)_2$ **4** and CPA **1**) and possesses fast response and high detection sensitivity, has shown to be a powerful tool to study visible light photocatalysis mechanistically. The *in situ* ionization capability of ESI-MS with online irradiation allows us to detect key and short lived species such as amine radical cations and $\text{Ru}(\text{bpz})_3^+$, and examine deactivation of photocatalysts. Moreover, by combining electrochemistry with MS, we are able to prove the viability of the oxidation of CPA **1** by $3^{+\cdot}$, generated by electrochemical oxidation of **3**. This result along with the post-irradiation reaction study provides conclusive experimental support for the involvement of the chain mechanism in the [3+2] annulation reaction. Finally, this study provides another good example of using MS to detect elusive and transient reaction intermediates in low quantity.^{7,8} We believe that we have just scratched the surface of ESI-MS with online irradiation's and online electrolysis' potential in studying visible light photocatalysis, and more new applications are anticipated to come.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Funding Sources

This work was supported by NSF Career Award (CHE-1255539) and the National Institute of Health under grant number P30 GM103450 from the National Institute of General Medical Sciences to N.Z. as well as NSF IDBR (CHE-1455554), NSF (CHE-1709075) and NSF MRI (CHE-1428787) to H.C. Y.C. thanks the Ohio Third Frontier Fund for research assistantship.

We thank Dr. Theresa H. Nguyen for early synthetic support and Professor Bill Durham for advice on quantum yield measurements.

References

1. (a) Prier CK, Rankic DA, MacMillan DWC. Chem Rev. 2013; 113:5322–5363. [PubMed: 23509883] (b) Xuan J, Xiao WJ. Angew Chem, Int Ed. 2012; 51:6828–6838. (c) Fukuzumi S, Ohkubo K. Chem Sci. 2013; 4:561–574. (d) Yoon TP. ACS Catal. 2013; 3:895–902. [PubMed: 23691491] (e) Romero NA, Nicewicz DA. Chem Rev. 2016; 116:10075–10166. [PubMed: 27285582]
2. (a) Gentry EC, Knowles RR. Acc Chem Res. 2016; 49:1546–1556. [PubMed: 27472068] (b) Ghosh I, Marzo L, Das A, Shaikh R, Konig B. Acc Chem Res. 2016; 49:1566–1577. [PubMed: 27482835] (c) Staveness D, Bosque I, Stephenson CRJ. Acc Chem Res. 2016; 49:2295–2306. [PubMed: 27529484] (d) Goddard JP, Ollivier C, Fensterbank L. Acc Chem Res. 2016; 49:1924–1936. [PubMed: 27529633] (e) Nakajima K, Miyake Y, Nishibayashi Y. Acc Chem Res. 2016; 49:1946–1956. [PubMed: 27505299] (f) Chen JR, Hu XQ, Lu LQ, Xiao WJ. Chem Soc Rev. 2016; 45:2044–2056. [PubMed: 26839142]

3. (a) Ye Y, Sanford MS. *J Am Chem Soc.* 2012; 134:9034–9037. [PubMed: 22624669] (b) DiRocco DA, Rovis T. *J Am Chem Soc.* 2012; 134:8094–8097. [PubMed: 22548244] (c) Sahoo B, Hopkinson MN, Glorius F. *J Am Chem Soc.* 2013; 135:5505–5508. [PubMed: 23565980] (d) Zoller J, Fabry DC, Ronge MA, Rueping M. *Angew Chem, Int Ed.* 2014; 53:13264–13268. (e) Tellis JC, Primer DN, Molander GA. *Science.* 2014; 345:433–436. [PubMed: 24903560] (f) Zuo Z, Ahneman DT, Chu L, Terrett JA, Doyle AG, MacMillan DWC. *Science.* 2014; 345:437–440. [PubMed: 24903563] (g) Du J, Skubi KL, Schultz DM, Yoon TP. *Science.* 2014; 344:392–396. [PubMed: 24763585] (h) Xuan J, Zeng T-T, Feng Z-J, Deng Q-H, Chen J-R, Lu L-Q, Xiao W-J, Alper H. *Angew Chem, Int Ed.* 2015; 54:1625–1628. (i) Jeffrey JL, Terrett JA, MacMillan DWC. *Science.* 2015; 349:1532–1536. [PubMed: 26316601] (j) Terrett JA, Cuthbertson JD, Shurtleff VW, MacMillan DWC. *Nature.* 2015; 524:330–334. [PubMed: 26266976] (k) Uraguchi D, Kinoshita N, Kizu T, Ooi T. *J Am Chem Soc.* 2015; 137:13768–13771. [PubMed: 26456298] (l) Kainz QM, Matier CD, Bartoszewicz A, Zultanski SL, Peters JC, Fu GC. *Science.* 2016; 351:681–684. [PubMed: 26912852] (m) Corcoran EB, Pirmot MT, Lin S, Dreher DA, DiRocco DA, Davies IA, Buchwald SL, MacMillan DWC. *Science.* 2016; 353:279–283. [PubMed: 27338703]
4. For selected examples: Kayser RH, Young RH. *Photochem Photobiol.* 1976; 24:395–401. Delaive PJ, Foreman TK, Giannotti C, Whitten DG. *J Am Chem Soc.* 1980; 102:5627–5631. Andrews RS, Becker JJ, Gagne MR. *Org Lett.* 2011; 13:2406–2409. [PubMed: 21473567] Liu Q, Li YN, Zhang HH, Chen B, Tung CH, Wu LZ. *Chem–Eur J.* 2012; 18:620–627. [PubMed: 22162148] Pitre SP, McTiernan CD, Ismaili H, Scaiano JC. *J Am Chem Soc.* 2013; 135:13286–13289. [PubMed: 23952147] Romero NA, Nicewicz DA. *J Am Chem Soc.* 2014; 136:17024–17035. [PubMed: 25390821] Yayla HG, Peng F, Mangion IK, McLaughlin M, Campeau LC, Davies IW, DiRocco DA, Knowles RR. *Chem Sci.* 2016; 7:2066–2073. Meyer AU, Strakova K, Slanina T, König B. *Chem–Eur J.* 2016; 22:8694–8699. [PubMed: 27167454] Shen X, Harms K, Marsch M, Meggers E. *Chem–Eur J.* 2016; 22:9102–9105. [PubMed: 27145893] Bartling H, Eisenhofer A, König B, Gschwind RW. *J Am Chem Soc.* 2016; 138:11860–11871. [PubMed: 27541322] Pitre SP, McTiernan CD, Scaiano JC. *Acc Chem Res.* 2016; 49:1320–1330. [PubMed: 27023767] Neumann M, Fuldner S, König B, Zeitler K. *Angew Chem Int Ed.* 2011; 50:951–954. Nappi M, Bergonzini G, Melchiorre P. *Angew Chem Int Ed.* 2014; 53:4921–4925.
5. (a) Miyake Y, Nakajima K, Nishibayashi Y. *J Am Chem Soc.* 2012; 134:3338–3341. [PubMed: 22296639] (b) Jiang H, Cheng Y, Wang R, Zheng M, Zhang, Yu SY. *Angew Chem, Int Ed.* 2013; 52:13289–13292. (c) Zou YQ, Guo W, Liu FL, Lu LQ, Chen JR, Xiao WJ. *Green Chem.* 2014; 16:3787–3795. (d) Lu P, Hou T, Gu X, Li P. *Org Lett.* 2015; 17:1954–1957. [PubMed: 25837858]
6. (a) Wallentin CJ, Nguyen JD, Finkbeiner P, Stephenson CRJ. *J Am Chem Soc.* 2012; 134:8875–8884. [PubMed: 22486313] (b) Cismesia MA, Yoon TP. *Chem Sci.* 2015; 6:5426–5434. [PubMed: 26668708]
7. (a) Wilson SR, Perez J, Pasternak A. *J Am Chem Soc.* 1993; 115:1994–1997. (b) Arakawa R, Lu J, Yoshimura A, Nozaki K, Ohno T, Doe H, Matsuo T. *Inorg Chem.* 1995; 34:3874–8. (c) Sam JW, Tang X-J, Magliozzo RS, Peisach. *J Am Chem Soc.* 1995; 117:1012–1018. (d) Feichtinger D, Plattner DA. *Angew Chem Int Ed.* 1997; 36:1718–1719. (e) Feichtinger D, Plattner DA, Chen P. *J Am Chem Soc.* 1998; 120:7125–7126. (f) Ding W, Johnson KA, Amster IJ, Katal C. *Inorg Chem.* 2001; 40:6865–6866. [PubMed: 11754264] (g) Julian RR, May JA, Stoltz BM, Beauchamp JL. *J Am Chem Soc.* 2003; 125:4478–4486. [PubMed: 12683818] (h) Meyer S, Koch R, Metzger JO. *Angew Chem, Int Ed.* 2003; 42:4700–4703. (i) Wilson DJ, Konermann L. *Anal Chem.* 2003; 75:6408–6414. [PubMed: 16465695] (j) Sabino AA, Machado AHL, Correia CRD, Eberlin MN. *Angew Chem, Int Ed.* 2004; 43:2514–2518. (k) Chen P. *Angew Chem, Int Ed.* 2003; 42:2832–2847. (l) Fuermeier S, Griep-Raming J, Hayen A, Metzger JO. *Chem–Eur J.* 2005; 11:5545–5554. [PubMed: 16013032] (m) Bao H, Zhou J, Wang Z, Guo Y, You T, Ding K. *J Am Chem Soc.* 2008; 130:10116–10127. [PubMed: 18616252] (n) Meyer MM, Khairallah GN, Kass SR, O’Hair RAJ. *Angew Chem, Int Ed.* 2009; 48:2934–2936. (o) Yunker LPE, Stoddard RL, McIndoe JS. *J Mass Spectrom.* 2014; 49:1–8. [PubMed: 24446256] (p) Iacobucci C, Reale S, De Angelis F. *Angew Chem, Int Ed.* 2016; 55:2980–2993. (q) Chen S, Wan Q, Badu-Tawiah AK. *Angew Chem, Int Ed.* 2016; 55:9345–9349. (r) Widjaja F, Jin Z, Nash JJ, Kentamaa HI. *J Am Chem Soc.* 2012; 134:2085–2093. [PubMed: 22148169] (s) Furmeier S, Metzger JO. *J Am Chem Soc.* 2004; 126:14485–14492. [PubMed: 15521769]

8. (a) Takats Z, Wiseman JM, Gologan B, Cooks RG. *Science*. 2004; 306:471–473. [PubMed: 15486296] (b) Perry RH, Splendore M, Chien A, Davis NK, Zare RN. *Angew Chem, Int Ed*. 2011; 50:250–254. (c) Perry RH, Brownell KR, Chingin K, Cahill TJ III, Waymouth RM, Zare RN. *Proc Natl Acad Sci*. 2012; 109:2246–2250. [PubMed: 22315417] (d) Brown TA, Chen H, Zare RN. *J Am Chem Soc*. 2015; 137:7274–7277. [PubMed: 26030136] (e) Zheng Q, Liu Y, Chen Q, Hu M, Helmy R, Sherer EC, Welch CJ, Chen H. *J Am Chem Soc*. 2015; 137:14035–14038. [PubMed: 26498505]
9. Maity S, Zhu M, Shinabery RS, Zheng N. *Angew Chem, Int Ed*. 2012; 51:222–226.
10. (a) Nguyen TH, Maity S, Zheng N Beilstein. *J Org Chem*. 2014; 10:975–980. (b) Nguyen TH, Morris SA, Zheng N. *Adv Synth Catal*. 2014; 356:2831–2837. [PubMed: 26120292]
11. Rassadin VA, Six Y. *Tetrahedron*. 2016; 72:4701–4757.
12. (a) Qin XZ, Williams FJ. *Am Chem Soc*. 1987; 109:595–597. (b) Li X, Grimm ML, Igarashi K, Castagnoli N Jr, Tanko JM. *Chem Commun*. 2007:2648–2650.
13. (a) Keizer SP, Han W, Stillman MJ. *Inorg Chem*. 2002; 41:353–358. [PubMed: 11800625] (b) Ma X, Zhao X, Li J, Zhang W, Cheng JX, Ouyang Z, Xia Y. *Anal Chem*. 2016; 88:8931–8935. [PubMed: 27560604]
14. (a) Brown TA, Chen H, Zare RN. *Angew Chem Int Ed*. 2015; 54:11183–11185. (c) Brown TA, Hosseini-Nassab N, Chen H, Zare RN. *Chem Sci*. 2016; 7:329–332. [PubMed: 28791096] (d) Zheng Q, Chen H. *Annu Rev Anal Chem*. 2016; 9(1):411–448. (e) Zhang Y, Yuan Z, Dewald HD, Chen H. *Chem Commun*. 2011; 47:4171–4173. (f) Zheng Q, Zhang H, Tong L, Wu S, Chen H. *Anal Chem*. 2014; 86:8983–8991. [PubMed: 25141260] (g) Li J, Dewald HD, Chen H. *Anal Chem*. 2009; 81:9716–9722. [PubMed: 19873992]
15. Van Berkel GJ, McLuckey SA, Glish GL. *Anal Chem*. 1992; 64:1586–1593.
16. Schmidbauer S, Hohenleutner A, Beilstein Konig B. *J Org Chem*. 2013; 9:2088–2096.
17. Devery JJ III, Douglas JJ, Nguyen JD, Cole KP, Flowers RA II, Stephenson CRJ. *Chem Sci*. 2015; 6:537–541. [PubMed: 28706624]

Figure 1.

a) Apparatus showing direct monitoring of the photoredox reaction by online ESI-MS; b) apparatus allowing to investigate the post-irradiation reaction event: a reaction mixture is introduced into the silica capillary, photo-excited by the laser pointer, and then mixed with an isotope-labeled new substrate introduced via channel 2; c) apparatus for investigating reactivity of radical cation generated via online electrolysis: a thin-layer electrochemical flow cell is connected with ion source and a substrate is introduced via channel 2 for testifying the key electron transfer step involved in the chain mechanism; AE, RE and WE represent auxiliary electrode, reference electrode and working electrode, respectively.

Figure 2.

a) ESI-MS spectrum of Ru(II)(bpz)₃(PF₆)₂ and CPA in CH₃NO₂ acquired without light irradiation; b) ESI-MS spectrum of Ru(II)(bpz)₃(PF₆)₂ and CPA in CH₃NO₂ with light irradiation; c) ESI-MS spectrum of Ru(II)(bpz)₃(PF₆)₂, CPA and styrene in CH₃NO₂ with light irradiation.

Figure 3.

Zoomed-in ESI-MS spectra showing the influence of the CPA concentration on the observation of 3^{+} : a) 5 mM CPA, b) 0.5 mM CPA, c) 100 μ M CPA, and d) 25 μ M CPA. 3^{+} was observed when the CPA concentration was lowered down to 100 μ M. Absolute intensity for m/z 237.1512 in Figure 2c and 2d are 2.35E5 and 1.43E5 (manufacturer arbitrary unit), respectively. Note that no 1^{+} was observed in these experiments probably due to the presence of excess amount of styrene (except that a small peak of 1^{+} was seen for the case of 5 mM CPA due to in-source oxidation). Also, no Ru(I) species was observed.

Figure 4.

ESI-MS spectra of post-irradiation reaction: a mixture of degassed CPA 1 (100 μM), $[\text{Ru}(\text{bpz})_3](\text{PF}_6)_2$ 4 (100 μM), and styrene 2 (0.1 M) in CH_3NO_2 was first irradiated, then mixed with a degassed solution of CPA- d_5 (100 μM) in CH_3NO_2 introduced via channel 2 (see apparatus in Figure 1b), for testifying chain reaction mechanism. a) the light was turned off. b) the light turned on. The inset shows zoomed-in ESI-MS spectrum of irradiation without adding CPA- d_5 .

Figure 5.

ESI-MS for monitoring of the oxidation of CPA 1 by $3^{+\cdot}$: a degassed solution of product 3 (1 mM) and LiOTf (1 mM) in ACN was infused into the electrochemical flow cell to generate $3^{+\cdot}$, which reacted with CPA 1 (introduced via channel 2). a) Chemical equation showing the reaction between the $3^{+\cdot}$ and **1**. b) ESI-MS spectra (background subtracted) showing the formation of the radical cation $3^{+\cdot}$ when the oxidation potential was applied to the cell: black line (only solvent ACN was introduced via channel 2) and red line (CPA 1 was introduced via channel 2). c) ESI-MS spectrum (background subtracted) showing the formation of CPA radical cation $1^{+\cdot}$ when the cell was turned on and CPA 1 was introduced into channel 2.

Figure 6.
Light/dark experiments for the annulation

Scheme 1.

The intermolecular [3+2] annulation of *N*-cyclopropylaniline **1** and styrene **2** catalyzed by $\text{Ru}(\text{II})(\text{bpz})_3(\text{PF}_6)_2$.

Scheme 2.

Proposed mechanism for the [3+2] annulation reaction of *N*-cyclopropylaniline **1** and styrene **2** catalyzed by $\text{Ru}(\text{II})(\text{bpz})_3(\text{PF}_6)_2$.