Published in final edited form as:

Matters (Zur). 2017 ; 2017: .

Increased whole cerebellar serotonin in aged C57BL/6 mice

Nicholas W DeKorver, Dustin Lichty, Marieke van der Hart, Arash Rassoulpour, and Stephen J Bonasera

Internal Medicine, University of Nebraska Medical Center; Microdialysis, Brains Online

Abstract

Mobility and locomotor impairments have high prevalence, morbidity, and significant mortality in older adult populations. Cerebellar functional changes have been implicated in the pathogenesis of these age-related mobility and gait deficits unrelated to stroke, Parkinson's disease, or degenerative joint disease. We thus examined total cerebellar glutamate, glutamine, GABA, glycine, dopamine, norepinephrine, tryptophan, serotonin, alanine, threonine, and asparagine content from male 2–3-month (young, $n = 6$) and 21–24-month-old (aged, $n = 6$) C57BL/6 mice. Neurotransmitter and amino acid concentrations were determined by high-performance liquid chromatography followed with mass spectroscopy. We found a significant increase in cerebellar serotonin in aged versus young mice, but otherwise no significant phenotypic differences in measured neurotransmitter concentrations. Applying current thought about cerebellar aging and cerebellar serotonergic systems, we consider how this age-related increase in cerebellar serotonin may contribute to gait ataxia.

Keywords

Serotonin; Aging; Mouse; Aging; Cerebellum

Objective

To further explore potential synaptic deficits underlying age-related gait impairments, we assessed whole-cerebellum neurotransmitter and excitatory amino acid content in cohorts of young and aged C57BL/6 mice using HPLCmass spectroscopy (HPLC-MS). We report a

This observation is distributed under the terms of the Creative Commons Attribution 4.0 International License.

Correspondence to: Stephen J Bonasera.

Disciplines

Neuroscience

Type of Observation

Standalone

Type of Link

Standard Data

Triple Blind Peer Review

The handling editor, the reviewers, and the authors are all blinded during the review process.

Methods and Supplementary Material

Please see <https://sciencematters.io/artides/201702000011>.

Ethics Statement

Not applicable.

novel finding of increased whole cerebellum serotonin in aged, compared to young, C₅₇BL/6 mice.

Introduction

It is essential to understand how aging affects mobility, and find interventions to prevent or delay mobility, impairments. Mobility impairments often accompany advanced age [1]. They are major causes of morbidity in the “oldest old” the most rapidly expanding demographic group in the United States [2] [3] [4]. Gait speed is a sensitive biomarker of overall functional status [5], as well as a powerful predictor of overall mortality [6] [7]. Cerebellar dysfunction may be a potential source of age-related mobility impairments. In humans, overall cerebellar volume decreases with age as shown by voxel-based morphometry [8] and is associated with age-related gait impairments including slow gait speed [9]. Aging is accompanied by a marked change in cerebellar gene expression patterns in both rodent models and humans [10] [11]. No age-related decrease in cerebellar granule cell number or density has been noted in rodent models [12] [13]; small decreases have been described in humans [14]. Aging is also associated with a strain-specific increase in excitatory synaptic puncta in the internal granule cell layer; however, these synapses show deficits in excitatory amino acid neurotransmission [10]. Loss of visualized synaptic boutons on Purkinje cell dendritic networks also accompanies age-related Purkinje cell involution and loss [15]. Aging also evokes deficits in Purkinje cell function, including marked slowing of firing rates [16] and blunted responses to adrenergic stimulation [17]. These changes have been correlated with impairment in mobility-related behaviors, including walking on a runway and dynamic balance [18].

Results & Discussion

Results

All samples were suitable for HPLC-MS determination of cerebellar neurotransmitter and amino acid levels: glutamate, GABA, glycine, glutamine, dopamine, norepinephrine, tryptophan, serotonin, alanine, threonine, and asparagine. Our results suggest a 1.4-fold statistically significant increase in whole-cerebellar serotonin in aged C₅₇BL/6 mice compared to young conspecifics (Figure a). We otherwise found no other differences in whole cerebellum neurotransmitter or amino acid concentrations between the young and old mouse cohorts.

Discussion

Serotonin is an ancient regulator of cellular function [19], and has a coordinating role in the performance of movement behaviors across a wide variety of organisms [20] [21] [22] [23]. Aging has a clear impact on cerebellar serotonergic function. The activity of tryptophan hydroxylase, the enzyme catalyzing the rate-limiting conversion of tryptophan to serotonin, is significantly decreased in the midbrain and pons of aged rats [24] [25]. Concordantly, age-associated decreases in whole cerebellar serotonin have been reported in rats (assayed by a fluorometric reaction between serotonin and o-phthalaldehyde [26]) and senescence-accelerated mice (assessed by ³H-tryptophan injection [27]). However, no age-associated

differences in cerebellar serotonin content were noted among cohorts of 6–24-month-old BALB mice (assayed by HPLC [28]), and our study is the first to report an age-associated increase in cerebellar serotonin content in a mouse model. Multiple studies have also demonstrated a marked loss of serotonin reuptake transporter function in the cerebellum of older adults [29] [30].

Serotonin has its most profound effect on the Lugaro inhibitory interneurons of the internal granule cell layer. These cells are numerous, have strong inhibitory projections onto Golgi cell inhibitory interneurons, and are typically silent. Lugaro cells respond to serotonin (likely through a 5-HT₂ subfamily receptor) with a marked increase in spiking activity leading to increased inhibitory postsynaptic potentials onto Golgi cell membranes [31] [32]. Lugaro cell activity also directly inhibits Purkinje neuron firing [33]. A single Lugaro cell makes inhibitory synaptic contact with about 150 nearby Golgi cells [31]; each Golgi cell makes inhibitory synaptic contact with about 3,000–5,000 granule cells [34]. Lugaro cell activity thus has the potential to reduce both the tonic and phasic components of Golgi cell inhibition onto large populations of granule cells. Serotonergic neurotransmission also contributes more subtle effects on cerebellar input signal processing. Serotonin attenuates parallel fiber input onto Purkinje cells (both by increasing activity of inhibitory basket and stellate interneurons and by presynaptic decreases of granule cell glutamate [35] [36]). Serotonin also modulates cerebellar deep nuclei function in a complex, incompletely understood manner [37].

Our results suggest an age-related 1.4-fold increase in whole cerebellar serotonin content. The methods we employed do not allow us to further determine if this difference reflects increased intravesicular serotonin within cerebellar serotonergic projections, and/or increased extracellular serotonin. Similarly, we cannot determine if this increase occurs over the cerebellum as a whole, or preferentially affects either the cerebellar input regions (internal granule cell, molecular, and Purkinje layers) or the cerebellar output nuclei; nor can we determine whether this finding localizes to specific folia.

The predicted outcome of increasing cerebellar serotonergic activity may thus be to increase granule cell input sensitivity to mossy fiber input, decrease Purkinje cell sensitivity to parallel fiber input, and increase inhibition of Purkinje cell firing. These effects would combine to expand mossy fiber input participating in center-surround inhibition [38] [39]. In other words, increased serotonergic activity may allow a larger set of afferent sensory inputs concurrent access to the Purkinje cell network while preserving overall Purkinje cell firing properties. Of note, in normal human volunteers, increased extracellular serotonin (through a single dose of a selective serotonin uptake inhibitor) evoked a significant increase in resting-state fMRI centrality measures across the cerebellum, consistent with the hypothesis that increased serotonergic activity enhances cerebellar functional connectivity [40].

It is thus interesting to note in this context that inhibition of serotonergic tone (by treatment with the partial 5-HT_{1A}R agonist buspirone) improved cerebellar tremor [41] [42]. Further, prominent cerebellar ataxia has been observed in aged (21–24 months old) compared to young (2–3 months old) C₅₇BL/6 mice obtained from the same colony as the mice in this study [10]. Interestingly, this appeared to be strain-specific with no ataxia present in BALB

mice of identical ages. Locomotor ataxia thus accompanies the increased cerebellar serotonin content observed in C₅₇BL/6 mice, but not in BALB mice with no age-related changes in cerebellar serotonin content [28]. Gait ataxia may potentially be a behavior evoked by serotonergic increases in cerebellar functional connectivity. Further studies to determine if age-associated increases in cerebellar serotonin content are reflected in extracellular serotonin concentrations, and to determine the specific serotonin receptor subtypes associated with Lugaro cell activation, will better define the mechanisms through which serotonergic activity may influence cerebellar tremor, as well as identify potential therapeutic targets.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This study was supported by NIH/NIA R01-AG031158 to SJB.

The authors thank Ms. Tammy Chaudoin for her expert technical support.

Full Open Access

Supported by the Velux Foundation, the University of Zurich, and the EPFL School of Life Sciences.

Citations

1. Luppá, Melanie, et al. Age-related predictors of institutionalization: results of the German study on ageing, cognition and dementia in primary care patients (AgeCoDe). *Social Psychiatry and Psychiatric Epidemiology*. Dec; 2010 47(2):263–270. <https://doi.org/10.1007/s00127-010-0333-9>. DOI: 10.1007/s00127-010-0333-9 [PubMed: 21181110]
2. Vincent, et al. The next four decades: The older population in the United States: 2010 to 2050. 2010; (1138)
3. Hobbs, et al. 65+ in the United States (US Bureau of the Census Current Population Reports, Special Studies, P23–190). Washington and DC: US Government Printing Office. 1996
4. Ferrer, Assumpta, et al. Risk of falls in 85-year-olds is associated with functional and cognitive status: The Octabaix study. *Archives of Gerontology and Geriatrics*. Mar; 2012 54(2):352–356. <https://doi.org/10.1016/j.archger.2011.06.004>. DOI: 10.1016/j.archger.2011.06.004 [PubMed: 21722976]
5. Wong WL, et al. Reinvestment and Falls in Community-Dwelling Older Adults. *Neurorehabilitation and Neural Repair*. Dec; 2007 22(4):410–414. <https://doi.org/10.1177/1545968307313510>. DOI: 10.1177/1545968307313510
6. Van Kan, et al. Gait speed at usual pace as a predictor of adverse outcomes in community-dwelling older people an International Academy on Nutrition and Aging (IANA) Task Force. *JNHA - The Journal of Nutrition and Health and Aging*. 2009; 13(10):881–889.
7. Studenski S, et al. Gait Speed and Survival in Older Adults. *JAMA*. Jan; 2011 305(1):50–58. <https://doi.org/10.1001/jama.2010.1923>. DOI: 10.1001/jama.2010.1923 [PubMed: 21205966]
8. Andersen, Birgitte Bo, Gundersen, Hans Jørgen G., Pakkenberg, Bente. Aging of the human cerebellum: A stereological study. *The Journal of Comparative Neurology*. Oct; 2003 466(3):356–365. <https://doi.org/10.1002/cne.10884>. DOI: 10.1002/cne.10884 [PubMed: 14556293]
9. Rosano, et al. A regions-of-interest volumetric analysis of mobility limitations in community-dwelling older adults. *The Journals of Gerontology Series A: Biological Sciences and Medical Sciences*. 2007; 62(9):1048–1055.

10. Bonasera Stephen J, et al. Age-related changes in cerebellar and hypothalamic function accompany non-microglial immune gene expression, altered synapse organization, and excitatory amino acid neurotransmission deficits. *Aging*. Sep.2016 8:2153–2181. <https://doi.org/10.18632/aging.101040>. DOI: 10.18632/aging.101040 [PubMed: 27689748]
11. Weindruch, Richard, Prolla, Tomas A., Lee, Cheol-Koo. Gene-expression profile of the ageing brain in mice. *Nature Genetics*. Jul; 2000 25(3):294–297. <https://doi.org/10.1038/77046>. DOI: 10.1038/77046 [PubMed: 10888876]
12. Sturrock RR. A comparison of quantitative histological changes in different regions of the ageing mouse cerebellum. *Journal fur Hirnforschung*. 1989; 31(4):481–486.
13. Dlugos, et al. Morphometric analyses of Purkinje and granule cells in aging F344 rats. *Neurobiology of aging*. 1994; 15(4):435–440. [PubMed: 7969720]
14. Renovell, et al. Loss of granule neurons in the aging human cerebellar cortex. *International Journal of Developmental Biology*. 2001; 40(S1):S193–S194.
15. Chen, et al. Dying-back of Purkinje cell dendrites with synapse loss in aging rats. *Journal of neurocytology*. 1999; 28(3):187–196. [PubMed: 10617901]
16. Rogers, et al. Senescent changes in a neurobiological model system: Cerebellar Purkinje cell electrophysiology and correlative anatomy. *Neurobiology of aging*. 1980; 1(1):3–11. [PubMed: 7266732]
17. Gould, et al. Age-Related Deficits in the Cerebellar BetaAdrenergic Signal Transduction Cascade in Fischer 344 Rats. *Journal of Pharmacology and Experimental Therapeutics*. 1997; 281(2):965–971. [PubMed: 9152407]
18. Bickford, et al. Antioxidant-rich diets improve cerebellar physiology and motor learning in aged rats. *Brain research*. 2000; 866(1):211–217. [PubMed: 10825496]
19. Turlejski K. Evolutionary ancient roles of serotonin: long-lasting regulation of activity and development. *Acta Neurobiol Exp (Wars)*. 1996; 56:619–636. [PubMed: 8768313]
20. Flavell Steven W, et al. Serotonin and the Neuropeptide PDF Initiate and Extend Opposing Behavioral States in *C. elegans*. *Cell*. Aug; 2013 154(5):1023–1035. <https://doi.org/10.1016/j.cell.2013.08.001>. DOI: 10.1016/j.cell.2013.08.001 [PubMed: 23972393]
21. Murakami, Hana, et al. Manipulation of serotonin signal suppresses early phase of behavioral aging in *Caenorhabditis elegans*. *Neurobiology of Aging*. Jul; 2008 29(7):1093–1100. <https://doi.org/10.1016/j.neurobiolaging.2007.01.013>. DOI: 10.1016/j.neurobiolaging.2007.01.013 [PubMed: 17336425]
22. Harris-Warrick RM, Cohen AH. Serotonin modulates the central pattern generator for locomotion in the isolated lamprey spinal cord. *J Exp Biol*. 1985; 116:27–46. [PubMed: 4056654]
23. Yuan, Quan, Joiner, William J., Sehgal, Amita. A Sleep-Promoting Role for the *Drosophila* Serotonin Receptor 1A. *Current Biology*. Jun; 2006 16(11):1051–1062. <https://doi.org/10.1016/j.cub.2006.04.032>. DOI: 10.1016/j.cub.2006.04.032 [PubMed: 16753559]
24. Hussain, et al. Effect of aging on tryptophan hydroxylase in rat brain: implications on serotonin level. *Drug metabolism and disposition*. 2000; 28(9):1038–1042. [PubMed: 10950846]
25. Hussain, et al. Effect of reactive oxygen species on the metabolism of tryptophan in rat brain: influence of age. *Molecular and cellular biochemistry*. 2004; 258(1–2):145–153. [PubMed: 15030179]
26. Arivazhagan, et al. Neurochemical changes related to ageing in the rat brain and the effect of DL- α -lipoic acid. *Experimental gerontology*. 2002; 37(12):1489–1494. [PubMed: 12559418]
27. Kabuto, et al. Neurochemical changes related to ageing in the senescence-accelerated mouse brain and the effect of chronic administration of nimodipine. *Mechanisms of ageing and development*. 1995; 80(1):1–9. [PubMed: 7564556]
28. Magnone, et al. Neurochemical parameters of the main neurotransmission systems in aging mice. *Archives of gerontology and geriatrics*. 2000; 30(3):269–279. [PubMed: 10867170]
29. Van Dyck, et al. Age-related decline in central serotonin transporter availability with [123 I] β -CIT SPECT. *Neurobiology of aging*. 2000; 21(4):497–501. [PubMed: 10924762]
30. Kish Stephen J, et al. Regional distribution of serotonin transporter protein in postmortem human brain. *Nuclear Medicine and Biology*. Feb; 2005 32(2):123–128. <https://doi.org/10.1016/j.nucmedbio.2004.10.001>. DOI: 10.1016/j.nucmedbio.2004.10.001 [PubMed: 15721757]

31. Dieudonné, et al. Serotonin-driven long-range inhibitory connections in the cerebellar cortex. *The Journal of Neuroscience*. 2000; 20(5):1837–1848. [PubMed: 10684885]
32. Dumoulin, et al. IPSC kinetics at identified GABAergic and mixed GABAergic and glycinergic synapses onto cerebellar Golgi cells. *The Journal of Neuroscience*. 2001; 21(16):6045–6057. [PubMed: 11487628]
33. Dean, et al. Serotonin drives a novel GABAergic synaptic current recorded in rat cerebellar Purkinje cells: a Lugaro cell to Purkinje cell synapse. *The Journal of neuroscience*. 2003; 23(11): 4457–4469. [PubMed: 12805286]
34. Wen-Ke, Li, et al. Using a million cell simulation of the cerebellum: Network scaling and task generality. *Neural Networks*. Nov.2013 47:95–102. <https://doi.org/10.1016/j.neunet.2012.11.005>. DOI: 10.1016/j.neunet.2012.11.005 [PubMed: 23200194]
35. Mitoma, et al. Monoaminergic long-term facilitation of GABA-mediated inhibitory transmission at cerebellar synapses. *Neuroscience*. 1999; 88(3):871–883. [PubMed: 10363824]
36. Maura, et al. Serotonin inhibits the depolarization-evoked release of endogenous glutamate from rat cerebellar nerve endings. *Neuroscience letters*. 1986; 67(2):218–222. [PubMed: 2873535]
37. Saitow, et al. Serotonin and synaptic transmission in the cerebellum. *Handbook of the Cerebellum and Cerebellar Disorders*. 2013:915–926.
38. Dizon MJ, Khodakhah K. The Role of Interneurons in Shaping Purkinje Cell Responses in the Cerebellar Cortex. *Journal of Neuroscience*. Jul; 2011 31(29):10463–10473. <https://doi.org/10.1523/jneurosci.1350-11.2011>. DOI: 10.1523/jneurosci.1350-11.2011 [PubMed: 21775592]
39. Solinas, Sergio, Nieus, Thierry, D'Angelo, Egidio. A realistic large-scale model of the cerebellum granular layer predicts circuit spatio-temporal filtering properties. *Frontiers in Cellular Neuroscience*. 2010; 4 <https://doi.org/10.3389/fncel.2010.00012>. doi: 10.3389/fncel.2010.00012
40. Schaefer, Alexander, et al. Serotonergic Modulation of Intrinsic Functional Connectivity. *Current Biology*. Oct; 2014 24(19):2314–2318. <https://doi.org/10.1016/j.cub.2014.08.024>. DOI: 10.1016/j.cub.2014.08.024 [PubMed: 25242032]
41. Lou, et al. Use of buspirone for treatment of cerebellar ataxia: an open-label study. *Archives of neurology*. 1995; 52(10):982–988. [PubMed: 7575226]
42. Trouillas, et al. Buspirone, a 5-hydroxytryptamine_{1A} agonist, is active in cerebellar ataxia: Results of a double-blind drug placebo study in patients with cerebellar cortical atrophy. *Archives of neurology*. 1997; 54(6):749–752. [PubMed: 9193210]

Figure. Age-evoked increase in whole cerebellar serotonin in C57BL/6 mice

Boxplots provided for whole cerebellum neurotransmitter concentrations; data points superimposed upon boxplots. Glutamate, GABA, glutamine, tryptophan, and norepinephrine values correspond to the y-axis valued 0–10; dopamine values correspond to the y-axis labeled 0–100; glycine and serotonin values correspond to the y-axis labeled 0–1,000. Neurotransmitter concentration units provided on the x-axis. Blue dots depict measures from the young (2–3 months) cohort; green dots depict measures from the old (21–24 months) cohort.

For all comparisons, age $df = 1$, error $df = 10$, total $df = 11$. Respective F and p values are as follows: Glu $F_{1,10} = 1.13$, $p = 0.3131$; GABA $F_{1,10} = 1.64$, $p = 0.229$; Gln $F_{1,10} = 0.38$, $p = 0.5495$; Gly $F_{1,10} = 6.85$, $p = 0.0257$; Ala $F_{1,10} = 0.44$, $p = 0.52$; Thr $F_{1,10} = 3.8$, $p = 0.0799$; Trp $F_{1,10} = 3.13$, $p = 0.1073$; Asn $F_{1,10} = 2.9$, $p = 0.1192$; DA $F_{1,10} = 0.08$, $p = 0.7863$; NE $F_{1,10} = 0.03$, $p = 0.8665$; 5-HT (*) $F_{1,10} = 13.53$, $p = 0.0043$. Critical $p = 0.0063$ to achieve $\alpha = 0.05$ for Bonferroni correction over eight comparisons.

The MATLAB program used to generate the figures as well as to run ANOVA statistics can be found in the supplementary data.