Published in final edited form as:

J Clin Psychiatry. 2017 May ; 78(5): e529–e536. doi:10.4088/JCP.15m10113.

Symptoms of apathy independently predict incident frailty and disability in community-dwelling older adults

Emmeline Ayers, MPH¹, Miriam Shapiro, PhD¹, Roe Holtzer, PhD^{1,4}, Nir Barzilai, MD^{2,3}, Sofiya Milman, MD², and Joe Verghese, MBBS, MS^{1,2}

¹Department of Neurology, Albert Einstein College of Medicine, Bronx, NY 10461, USA

²Department of Medicine, Albert Einstein College of Medicine, Bronx, NY 10461, USA

³Department of Genetics, Albert Einstein College of Medicine, Bronx, NY 10461, USA

⁴Ferkauf Graduate School of Psychology, Yeshiva University, Bronx, NY, USA

Abstract

Objective—Although depressive symptoms are widely recognized as a predictor of functional decline among older adults, little is known about the predictive utility of apathy in this population. We prospectively examined apathy symptoms as predictors of incident slow gait, frailty and disability among non-demented, community-dwelling older adults.

Methods—We examined two independent prospective cohort studies - the LonGenity study (N = 625, 53% women, mean age 75.2 years) and the Central Control of Mobility in Aging (CCMA) study (N = 312, 57% women, mean age 76.4 years). Individuals were recruited from 2008 to 2014. Apathy was assessed using 3 items from the Geriatric Depression Scale. Slow gait was defined as 1 standard deviation or more below age and sex-adjusted mean values, frailty was defined using the Cardiovascular Health Study criteria, and disability was assessed with a well-validated disability scale.

Results—The prevalence of apathy was 20% in LonGenity cohort and 26% in the CCMA cohort. The presence of apathy at baseline, independent of depressive symptoms (besides apathy), increased risk of developing incident slow gait (hazard ratio [HR] = 2.10; CI = 1.36–3.24; $p = .001$), frailty (HR = 2.86; CI = 1.96–4.16; $p < .001$), and disability (HR = 3.43; CI = 1.73–6.79; $p < .001$) in the pooled sample. These associations remained significant when accounting for demographics, medical illnesses, and cognitive function.

Conclusions—Apathy is associated with increased risk of developing slow gait, frailty, and disability, independent of other established risk factors, in non-demented older adults. Apathy should be screened as a potentially preventable cause of functional decline in clinical psychiatric settings.

Address correspondence to: Joe Verghese, MBBS, Department of Neurology, Albert Einstein College of Medicine, 1225 Morris Park Avenue, Van Etten 308, Bronx, New York 10461, joe.verghese@einstein.yu.edu.

Conflict of interest: None declared

Keywords

Apathy; Depression; Frailty; Older adults

Apathy is one of the most common neuropsychiatric symptoms in older adults,^{1–4} and is increasingly prevalent with advancing age.^{1,3,4} Defined as concurrent reduction in cognitive, emotional, and motoric goal-directed behavior,⁵ apathy is hypothesized to result from damage to the frontal-subcortical pathways.^{5–7} Although nosologically similar to depression, apathy is now widely considered a distinct neuropsychiatric condition that can occur in the presence or absence of depression.^{8,9}

Apathy has been identified as a prominent behavioral symptom in Alzheimer's and Parkinson's diseases, and is associated with greater morbidity and mortality in these conditions.^{10,11} Additionally, it is associated with multiple adverse health outcomes among older adults with varying degrees of neurological impairment in both cross-sectional and longitudinal studies.^{12–14} For instance, apathy predicts gait dysfunction, disability, and cognitive decline in individuals with dementia.^{11–15} Furthermore, apathy predicts conversion to dementia among those with mild cognitive impairment (MCI),^{16–18} and increases risk of mortality in nursing home residents.¹⁹ However, despite apathy's widely researched function as a marker of decline among these vulnerable disease populations, little is known about the predictive validity of apathy in healthy older adults.

Identifying risk factors for functional decline in the community-dwelling, elderly population is particularly important, as these individuals are still easily able to participate in targeted psychiatric interventions. Behavioral markers, such as apathy and depression, can be screened quickly and inexpensively at routine clinical visits. While depression has been extensively studied as a predictor of functional decline in healthy older adults,^{20–25} few have looked at apathy,³ and, to our knowledge, no one has assessed apathy's longitudinal associations with frailty and motor status in this population.

To address these knowledge gaps, we examined the longitudinal relationship between apathy and subsequent functional decline in two prospective cohorts of community-dwelling, cognitively-intact, and non-disabled older adults. Researchers recommend utilizing complementary measurements of functioning to provide a more holistic understanding of health status,²⁶ therefore, we examined new onset of slow gait, frailty, and disability as our primary outcomes. Slow gait, frailty, and disability predict multiple adverse health outcomes in older adults, including falls, dementia, morbidity, and death, making them robust indicators of functioning.^{27–31} Expanding upon previous research,^{11–17} we hypothesized that baseline apathy, independent of depressive symptoms, will predict incident slow gait, frailty, and disability in community-dwelling older adults. If confirmed, our findings will lend support to apathy's clinical utility in psychiatric settings.

Methods

Participants

Participants were recruited from two independent ongoing cohort studies at the Albert Einstein College of Medicine - the LonGenity study³² and the Central Control of Mobility in Aging (CCMA) study.³³

LonGenity—The goal of the LonGenity study is to identify genotypes associated with longevity and successful aging in Ashkenazi Jewish seniors.³² Participants seen between March 2008 and September 2014 were included in this analysis. As previously described,³² Ashkenazi Jewish adults age 65 and older were systematically recruited using public records such as voter registration lists. A smaller sample was identified through contacts at synagogues, community organizations, and advertisements in Jewish newspapers. Potential participants were contacted by mail and then telephone to rule out the presence of baseline dementia. Participants received comprehensive neuropsychological, psychological, and quantitative gait assessments at baseline and annual follow-up visits. Exclusion criteria were: inability to speak English, dementia, significant loss of vision and/or hearing, and having a sibling already enrolled in the study. Of the 847 individuals recruited at baseline, 213 had not received follow-up assessments at the time of data analysis and nine were missing data. These participants were excluded from the study. Excluded individuals did not significantly differ with respect to age, gender, education, or illness index from eligible participants. Following exclusions, 625 individuals with median follow up of three years (range 1–6) were included in the current study (mean age 75.21 ± 6.4 years, 53.3% female).

Central Control of Mobility in Aging—The primary goal of the CCMA study is to determine cognitive predictors of mobility decline and disability in aging.³³ Participants seen between June 2011 and September 2014 were included in this analysis. The study procedure has been previously described.³³ Briefly, individuals age 65 and older residing in lower Westchester County were identified from population lists. Potential participants were contacted by mail and then telephone to rule out the presence of baseline dementia. Exclusion criteria were: inability to speak English, inability to ambulate independently, dementia, significant loss of vision and/or hearing, and major psychiatric disorders. Eligible participants received comprehensive neuropsychological, psychological, and quantitative gait assessments at baseline and annual follow-up visits. Of the 496 individuals recruited at baseline, 182 had not received follow-up assessments at the time of data analysis and two were missing data. These participants were excluded from the study. Excluded individuals did not significantly differ with respect to age, gender, education, or illness index from eligible participants. Following exclusions, 312 individuals with median follow up of 2 years (range 1–3) were included in the current study (mean age 76.4 ± 6.87 years, 56.7% female).

For both cohorts, the institutional review board of the Albert Einstein College of Medicine approved the experimental procedures and all participants provided written informed consent. Participants received complete clinical and neuropsychological evaluations annually in both studies and diagnoses of dementia and MCI were assigned according to the criteria of the *Diagnostic and Statistical Manual of Mental Disorders*, Fourth Edition,³⁴ using all

available clinical, neuropsychological and laboratory information at consensus case conferences attended by at least one study neurologist and one neuropsychologist, as previously described.³⁵

Outcome measures

We selected three common motor and functional outcomes (slow gait, frailty, and disability) associated with poor physical and cognitive aging.^{27–31,36} Each of these outcomes was assessed at baseline and annual follow-up visits in both cohorts. All of these assessments were conducted blinded to participants' apathy status at baseline and follow-up visits.

Slow Gait—Gait speed (cm/s) was measured while participants walked at their normal pace on an 8.5-m long computerized walkway (180×35.5×0.25 in.) with embedded pressure sensors (GAITRite, CIR systems, Havertown, PA). Excellent reliability and validity for GAITRite assessments have been reported.^{30,37} Slow gait was defined as one or more standard deviations below previously outlined age and sex-appropriate mean values.³⁸ Similar procedure was previously used to define slow gait in these two cohorts as well as other cohorts.³⁹

Frailty—Frailty diagnosis was operationalized using the Cardiovascular Health Study criteria,³⁶ defined as meeting three or more of the following: unintentional weight loss of more than 10 pounds in the past year, self-reported decrease in energy in the past year, muscle weakness (weak grip strength), slow gait speed (cut-scores described above), and self-reported decrease in physical activity in the past year.³⁶ Similar to previous reports,^{40–42} subjective grip strength (“do you feel as though your grip is weak?”) was used since objective measures were not available for all participants on follow-up. Among individuals in our sample with both objective and subjective measurements ($n = 451$), these two ratings showed substantial inter-rater agreement ($k = .681$).

Disability—Basic activities of daily living (ADLs) disabilities were assessed using standard questions with a scale developed by Gill and colleagues for use in community-based cohorts.⁴³ The scale has been used to assess functional status in our aging studies as previously described.³⁸ Excellent test retest reliability and inter-rater reliability has been reported for this scale in community samples.⁴³ The scale was administered by trained research assistants, and characterizes self-reported performance on basic but not instrumental ADLs. Disability was coded dichotomously and defined as needing assistance with or inability to perform any one of the following seven basic ADLs: bathing, dressing, walking inside the home, getting up from a chair, toileting, feeding, and grooming.

Apathy and depressive symptom measures

Symptoms of apathy and depressive affect were evaluated at baseline and yearly follow-up visits using the well-validated Geriatric Depression Scale (GDS).^{44,45} The 15-item GDS was used in the LonGenity study, and the corresponding 15 items were extracted from the 30-item GDS used in the CCMA study for symmetry of analysis. Similar procedure to harmonize depression assessments was previously applied in other cohorts.⁴⁶ Factor analyses have consistently identified an apathy dimension of the GDS, consisting of the

following three items: (1) Have you dropped many of your activities and interests ('yes' or 'no')? (2) Do you prefer to stay home, rather than go out and do new things ('yes' or 'no')? (3) Do you feel full of energy ('yes' or 'no')?^{4,47,48} Consistent with previous apathy research,^{2-4,18,49} we distinguished apathy (GDS-3; range 0–3) and depressive symptoms (GDS-12, remaining items, range 0–12) into two non-overlapping subscales, and utilized a cut-off score of two or more endorsed apathy items to indicate the presence of apathy. Previous studies have demonstrated good correlation between this measurement method and clinical diagnosis of apathy,^{2,4,18,50} with sensitivity and specificity of 69% and 85%, respectively.²

Other covariates

As previously reported,^{39, 40} a summary illness index was computed based on the presence or absence of physician diagnosed vascular diseases (diabetes, heart failure, hypertension, angina, myocardial infarction, and stroke) as well as other medical illnesses, including Parkinson's disease, chronic obstructive lung disease, and arthritis. An extensive cognitive battery was administered in both cohorts. For this investigation, we report executive functioning and processing speed, measured by verbal fluency (FAS) and digit symbol coding, respectively,^{51,52} as these cognitive processes are most commonly linked with apathy.^{6,7,10}

Data analysis

Baseline characteristics were summarized with descriptive statistics. To determine the longitudinal association of baseline apathy with slow gait, frailty, and disability, we used Cox proportional hazards models with each of the three outcome measures as dependent variables. Individuals who met criteria for one of the outcome measures at baseline were excluded from subsequent analyses for that outcome, but not from other analyses if they were eligible. The Cox models were examined separately in LonGenity (cohort 1) and CCMA (cohort 2), as well as in the pooled sample adjusted for cohort status.

We used four models in the analysis. Because one of our main aims was to show apathy's predictive value independent of comorbid depressive symptoms, we included the 12-item GDS (minus apathy score) as a continuous covariate in all models. In model 2, we additionally adjusted for age, sex, and education because apathy symptoms are influenced by these demographic characteristics.^{1,3,4} Apathy has also been associated with poor health status and cognitive decline;^{3,53-55} therefore, models 3 and 4 adjusted for summary illness index and cognitive functioning (FAS and Digit symbol coding), respectively. Associations are reported as hazard ratios (HR) with 95% confidence intervals (CI). Regression diagnostics for all models were examined and adequately met. All analyses in this study were carried out using SPSS version 20.

Results

Baseline characteristics

Table 1 displays baseline demographic and clinical data for subjects in both cohorts, as well as prevalence and incidence rates for study outcomes. Twenty percent of individuals in

cohort 1 endorsed baseline apathy versus 26% in cohort 2. In the pooled sample, 379 (40%) participants endorsed no symptoms of apathy, 351 (38%) endorsed one, 164 (18%) endorsed two, and 43 (5%) endorsed all three apathy symptoms. As expected, individuals with apathy were significantly older and had poorer health status, lower education, and higher depressive symptoms (Table 2).

Longitudinal association of baseline apathy symptoms with outcome measures

Table 3 presents the association of baseline apathy with incident slow gait, frailty, and disability for each cohort independently as well as the pooled sample. Cumulative hazard functions for the pooled sample, stratified by the presence of baseline apathy, are displayed in figure 1.

Baseline apathy increased risk of slow gait and frailty in cohort 1 (gait HR = 1.86; frailty HR = 3.39), cohort 2 (gait HR = 2.92; frailty HR = 1.97), and the pooled sample (gait HR = 2.10; frailty HR = 2.86). These associations did not change substantively after additional adjustments for demographics, health status, and cognitive functioning. Presence of baseline apathy increased disability risk in cohort 1 (HR = 4.36) and in the pooled sample (HR = 3.43); however, this relationship was not found in cohort 2. Additional adjustments for demographics, health status, and cognitive functioning did not affect the results.

Sensitivity analyses

We ran a series of sensitivity analyses on the pooled sample. Because apathy and depression are often comorbid conditions,⁵ we wanted to show that apathy accounted for the above relationships even in those without subclinical or minor depression. We reran model 1 for all three outcomes including only participants who endorsed one or fewer depressive symptoms (GDS-12 = 1) at baseline ($n = 654$, 105 (16%) of whom had apathy). This method is consistent, if not more stringent, than previous reports,^{4,49,50} and did not affect the associations between apathy and incident slow gait (HR = 2.09; CI = 1.25–3.48; $p = .005$), frailty (HR = 3.21; CI = 2.03–5.10; $p < .001$), and disability (HR = 5.16; CI = 2.36–11.31; $p < .001$).

To explore the possibility of a reverse-causation between apathy and study outcomes, we reran model 1 excluding all participants who met criteria for each outcome within two years of their baseline assessment. The relationships between apathy and incident slow gait (HR = 1.90; CI = .98–3.69; $p = .057$), frailty (HR = 3.74; CI = 2.22–6.31; $p < .001$), and disability (HR = 3.76; CI = 1.52–9.26; $p = .004$) remained strong.

There were two incident dementia cases in cohort 1 and five in cohort 2. Excluding these cases did not affect the relationship between apathy and the study outcomes (slow gait: HR = 2.11; CI = 1.37–3.25; $p = .001$; frailty: HR = 2.84; CI = 1.93–4.13; $p < .001$; disability: HR = 3.24; CI = 1.57–6.73; $p < .001$).

Lastly, to explore any ‘dose-dependent’ effect of apathy we reran model 1 with apathy as a categorical 4-level predictor (no symptoms, 1 symptom, 2 symptoms, and 3 symptoms). Figure 2 shows that cumulative risk of incident slow gait, frailty, and disability increases with increasing apathy score (full results in eAppendix 1).

Discussion

In this prospective cohort-based study, the presence of apathy in non-demented, non-disabled, community-dwelling older adults increased risk of functional and motoric decline. Individuals with baseline apathy had more than a two-fold risk of developing slow gait and frailty, and more than a three-fold risk of becoming disabled indicating the overall risk of functional decline associated with apathy in older adults. Furthermore, individuals' risk for these outcomes increased with apathy severity. These associations were independent of depressive symptoms and remained robust after adjustment of potential confounders, such as demographics, health status, and cognitive functioning. Excluding incident cases of dementia in our sensitivity analysis did not materially affect the relationship between apathy and the study outcomes, suggesting that despite the well-established links of apathy and cognitive impairment,^{7,8,13,56,57} the association of apathy with functional decline may act through other non-dementia pathways in non-demented older adults.

To our knowledge, this is the first study to show that apathy symptoms predict incident frailty and motoric decline among non-demented, community-dwelling older adults. With the exception of one study³ showing that apathy increases disability risk, most researchers have focused on depressive symptoms in this population. Depressive symptoms have been widely associated with frailty and disability in cross-sectional,^{20,22–24,58,59} and to a lesser extent, longitudinal,^{60,61} studies of healthy older adults. Although, studies have identified a lack of correlation between apathy and depression as well as disease specific differences in neuropathology.⁹ Many previous studies that have reported the association between depressive symptoms and functional decline do not differentiate between symptoms of apathy and depression, and it is unclear how much apathy symptoms contribute to the observed findings in these studies. Additionally, many instruments that measure depressive symptoms include several apathy items, inflating depression rates and conflating the two psychiatric constructs.⁹ It is quite possible that these previously reported associations were confounded by symptoms of apathy, and that it is in fact apathy that predicts these negative health outcomes. This discovery would have significant public health implications, as clinicians routinely screen for depression, but not apathy.

Indirect evidence points to several possible explanations for the relationship between apathy and subsequent motoric and functional decline. First, apathetic individuals may be more susceptible to frailty and disability due to lifestyle factors, such as reduced physical and leisure activity, poor social support, and medication non-compliance. However, there is also evidence for shared pathophysiological mechanisms, such as vascular pathways. Slow gait, frailty, and disability^{56,62,63} have all been previously associated with vascular disease. Healthy older adults with apathy have an almost three-fold risk of developing cardiovascular disease,⁵⁰ and show greater evidence of cerebrovascular disease on MRI.⁴⁹ As expected, higher numbers of vascular comorbidities were found among individuals with baseline apathy in our sample, but given that our summary illness index may not be sensitive to subtle vascular changes. However, the associations between apathy and slow gait, frailty, and disability remained significant when controlling for these comorbid conditions, indicating that these relationships are at least partly independent of vascular pathways.

Increased inflammatory activity is common in aging,^{57,64} and is hypothesized to be an important pathway for age-related physical decline.^{65–68} Among the inflammatory factors, elevated levels of interleukin 6 (IL-6) has been associated with frailty, disability, and gait disturbance among community-dwelling older adults.^{67–69} Elevated IL-6 is also implicated as an underlying mechanism for late-life depression;^{70–72} however, its role in late life apathy has yet to be examined. One study looking at depression among post-stroke individuals found that higher levels of IL-6 were associated with severity of apathetic-depressive symptoms,⁷³ indicating some evidence of an apathy-inflammatory link. Future studies should further explore this connection by examining apathy and inflammatory markers in the context of age-related motoric and functional decline. Furthermore, inflammation and vascular disease commonly co-occur;⁷⁴ therefore, apathy's relationship with functional and motoric decline may be through these pathways independently, additively, or synergistically. Future research on apathy should examine multiple biomarkers of inflammation and vascular pathology to elucidate the exact nature of these pathways and help inform potential interventions.

Yet another possible explanation for the observed relationship is the reverse-causation hypothesis. Perhaps apathy is an early behavioral manifestation of frailty and disability, such that individuals begin to display apathy symptoms during prodromal states of these conditions. However, our sensitivity analysis excluding pre-frail and pre-disabled individuals, suggests that apathy is not merely a byproduct, but rather, a predictor of these conditions.

There are several limitations of the current study that should be noted. Due to the clinical overlap between the three study outcomes, interpretation of apathy as a predictor of the three independent outcomes should be taken with caution. The disability instrument captures performance on physical or basic activities of daily living, and was not developed to identify performance on instrumental activities of daily living. We assessed apathy retrospectively in two cohorts using three items from the GDS that have consistently shown to reflect apathy;^{2–4,18} and with high agreement with more detailed apathy scales.² However, given the moderate sensitivity and specificity of the GDS, a more detailed apathy assessment might provide additional insights into specific apathy symptoms and our outcomes.⁷⁵ Furthermore, the fact that apathy was found to have a 'dose-dependent' effect on study outcomes, indicates that apathy symptoms themselves, and not necessarily a clinical diagnosis of apathy, places people at risk for future decline. Vascular conditions were clinically diagnosed and more direct measures of vascular function might have shown stronger relationships with apathy. Vascular and other biological markers should be further explored in future studies to provide mechanistic insights. Dementia was excluded through consensus case conference procedures. Future studies that examine the mediating effect of cognitive outcomes including subtypes of MCI on apathy would help delineate the relationship between apathy and functional decline.

To our knowledge, this is the first study to demonstrate apathy as an independent risk factor for frailty and motoric decline in non-disabled, non-demented, community-dwelling elderly. If our findings are corroborated by other studies, apathy screening should become a routine part of psychiatric visits with older adults. Furthermore, the underlying mechanisms of these

relationships should be explored via neuroimaging and biomarker techniques to inform the development of intervention strategies to prevent functional decline in high-risk apathy patients.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

The LonGenity study was funded by the National Institutes of Health (NIH) (R00AG037574, 1P01AG034906, R01AG046949, R01AG044829, 1R01AG042188, P30AG038072, and R37AG18381), Clinical and Translational Science Award KL2TR000088, Einstein Glenn Center, Paul Glenn Foundation, and the American Federation for Aging Research. The Central Control of Mobility in Aging study (from which patients were also recruited for the current study) was funded by NIH (R01AG036921 and R01AG044007).

References

1. Brodaty H, Altendorf A, Withall A, et al. Do people become more apathetic as they grow older? A longitudinal study in healthy individuals. *Int Psychogeriatr*. 2010; 22(3):426–436. [PubMed: 20003630]
2. van der Mast RC, Vinkers DJ, Stek ML, et al. Vascular disease and apathy in old age. The Leiden 85-plus study. *Int J Geriatr Psychiatry*. 2008; 23(3):266–271. [PubMed: 17621380]
3. Clarke DE, Ko JY, Lyketsos C, et al. Apathy and cognitive and functional decline in community-dwelling older adults: results from the Baltimore ECA longitudinal study. *Int Psychogeriatr*. 2010; 22(5):819–829. [PubMed: 20478091]
4. Ligthart SA, Richard E, Fransen NL, et al. Association of vascular factors with apathy in community-dwelling elderly individuals. *Arch Gen Psychiatry*. 2012; 69(6):636–642. [PubMed: 22664551]
5. Marin RS. Apathy: a neuropsychiatric syndrome. *J Neuropsychiatry Clin Neurosci*. 1991; 3:243–254. [PubMed: 1821241]
6. Cummings JL. Frontal-subcortical circuits and human behavior. *Arch Neurol*. 1993; 50:873–880. [PubMed: 8352676]
7. Tekin S, Cummings JL. Frontal-subcortical neuronal circuits and clinical neuropsychiatry: an update. *J Psychosom Res*. 2002; 53(2):647–654. [PubMed: 12169339]
8. Marin RS, Firinciogullari S, Biedrzycki RC. Group differences in the relationship between apathy and depression. *J Nerv Ment Dis*. 1994; 182(4):235–239. [PubMed: 10678322]
9. Levy ML, Cummings JL, Fairbanks LA, et al. Apathy is not depression. *J Neuropsychiatry Clin Neurosci*. 1998; 10(3):314–9. [PubMed: 9706539]
10. Kuzis G, Sabe L, Tiberti C, et al. Neuropsychological correlates of apathy and depression in patients with dementia. *Neurology*. 1999; 52:1403–1407. [PubMed: 10227625]
11. Starkstein SE, Merello M, Brockman S, et al. Apathy predicts more severe parkinsonism in Alzheimer's disease. *Am J Geriatr Psychiatry*. 2009; 17(4):291–298. [PubMed: 19322935]
12. Boyle PA, Malloy PF, Salloway S, et al. Executive dysfunction and apathy predict functional impairment in Alzheimer disease. *Am J Geriatr Psychiatry*. 2003; 11(2):214–221. [PubMed: 12611751]
13. Wadsworth LP, Lorus N, Donovan NJ, et al. Neuropsychiatric symptoms and global functional impairment along the Alzheimer's continuum. *Dement Geriatr Cogn Disord*. 2012; 34(2):96–111. [PubMed: 22922821]
14. Zawacki TM, Grace J, Paul R, et al. Behavioral problems as predictors of functional abilities of vascular dementia patients. *J Neuropsychiatry Clin Neurosci*. 2002; 14(3):296–302. [PubMed: 12154154]

15. Olazarán J, Hernández-Tamames JA, Molina E, et al. Clinical and anatomical correlates of gait dysfunction in Alzheimer's disease. *J Alzheimers Dis.* 2013; 33(2):495–505. [PubMed: 23011219]
16. Robert PH, Berr C, Volteau M, et al. Apathy in patients with mild cognitive impairment and the risk of developing dementia of Alzheimer's disease: a one-year follow-up study. *Clin Neurol Neurosurg.* 2006; 108(8):733–736. [PubMed: 16567037]
17. Palmer K, Di Iulio F, Varsi AE, et al. Neuropsychiatric predictors of progression from amnesic-mild cognitive impairment to Alzheimer's disease: the role of depression and apathy. *J Alzheimers Dis.* 2010; 20(1):175–183. [PubMed: 20164594]
18. Richard E, Schmand B, Eikelenboom P, et al. Symptoms of apathy are associated with progression from mild cognitive impairment to Alzheimer's disease in non-depressed subjects. *Dement Geriatr Cogn Disord.* 2012; 33(2–3):204–209. [PubMed: 22722671]
19. Hölttä EH, Laakkonen ML, Laurila JV, et al. Apathy: prevalence, associated factors, and prognostic value among frail, older inpatients. *J Am Med Dir Assoc.* 2012; 13(6):541–545. [PubMed: 22572553]
20. Beekman AT, Penninx BW, Deeg DJ, et al. Depression and physical health in later life: results from the Longitudinal Aging Study Amsterdam (LASA). *J Affect Disord.* 1997; 46(3):219–231. [PubMed: 9547118]
21. Lyness JM, Kim J, Tang W, et al. The clinical significance of subsyndromal depression in older primary care patients. *Am J Geriatr Psychiatry.* 2007; 15(3):214–223. [PubMed: 17213374]
22. Chopra MP, Zubritsky C, Knott K, et al. Importance of subsyndromal symptoms of depression in elderly patients. *Am J Geriatr Psychiatry.* 2005; 13(7):597–606. [PubMed: 16009736]
23. Forsell Y, Winblad B. Major depression in a population of demented and nondemented older people: prevalence and correlates. *J Am Geriatr Soc.* 1998; 46(1):27–30. [PubMed: 9434662]
24. Rog LA, Park LQ, Harvey DJ, et al. The independent contributions of cognitive impairment and neuropsychiatric symptoms to everyday function in older adults. *Clin Neuropsychol.* 2014; 28(2): 215–236. [PubMed: 24502686]
25. Nair V, Ayers E, Noone M, et al. Depressive symptoms and mild cognitive impairment: results from the Kerala-Einstein study. *J Am Geriatr Soc.* 2014; 62(1):197–199. [PubMed: 25180388]
26. Lutomski JE, Baars MA, Boter H, et al. Frailty, disability and multi-morbidity: the relationship with quality of life and healthcare costs in elderly people. *Ned Tijdschr Geneesk.* 2014; 158:A7297. [PubMed: 25204442]
27. Montero-Odasso M, Schapira M, Soriano ER, et al. Gait velocity as a single predictor of adverse events in healthy seniors aged 75 years and older. *J Gerontol A Biol Sci Med Sci.* 2005; 60(10): 1304–1309. [PubMed: 16282564]
28. Buchman AS, Boyle PA, Wilson RS, et al. Frailty is associated with incident Alzheimer's disease and cognitive decline in the elderly. *Psychosom Med.* 2007; 69(5):483–489. [PubMed: 17556640]
29. Solfrizzi V, Scafato E, Frisardi V, et al. Frailty syndrome and the risk of vascular dementia: the Italian Longitudinal Study on Aging. *Alzheimers Dement.* 2013; 9(2):113–122. [PubMed: 23245560]
30. Verghese J, Lipton RB, Hall CB, et al. Abnormality of gait as a predictor of non-Alzheimer's dementia. *N Engl J Med.* 2002; 347(22):1761–1768. [PubMed: 12456852]
31. Verghese J, Wang C, Lipton RB, et al. Quantitative gait dysfunction and risk of cognitive decline and dementia. *J Neurol Neurosurg Psychiatry.* 2007; 78(9):929–935. [PubMed: 17237140]
32. Ayers E, Barzilai N, Crandall JP, et al. Association of exceptional parental longevity and physical function in aging. *Age (Dordr).* 2014; 36(4):9677. [PubMed: 24997018]
33. Holtzer R, Wang C, Verghese J. Performance variance on walking while talking tasks: theory, findings, and clinical implications. *Age (Dordr).* 2014; 36(1):373–381. [PubMed: 23943111]
34. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders.* 4. Washington, DC: American Psychiatric Association; 1994.
35. Holtzer R, Verghese J, Wang C, et al. Within-person across-neuropsychological test variability and incident dementia. *JAMA.* 2008; 300(7):823–830. [PubMed: 18714062]
36. Fried TR, Bradley EH, Williams CS, et al. Functional disability and health care expenditures for older persons. *Arch Intern Med.* 2001; 161(21):2602–2607. [PubMed: 11718592]

37. Bilney B, Morris M, Webster K. Concurrent related validity of the GAITRite walkway system for quantification of the spatial and temporal parameters of gait. *Gait Posture*. 2003; 17(1):68–74. [PubMed: 12535728]
38. Verghese J, Wang C, Lipton RB, et al. Motoric cognitive risk syndrome and the risk of dementia. *J Gerontol A Biol Sci Med Sci*. 2013; 68(4):412–418. [PubMed: 22987797]
39. Verghese J, Ayers E, Barzilai N, et al. Motoric cognitive risk syndrome: Multicenter incidence study. *Neurology*. 2014; 83(24):2278–2284. [PubMed: 25361778]
40. Johansen KL, Dalrymple LS, Delgado C, et al. Comparison of self-report-based and physical performance-based frailty definitions among patients receiving maintenance hemodialysis. *Am J Kidney Dis*. 2014; 64(4):600–607. [PubMed: 24793033]
41. Woods NF, LaCroix AZ, Gray SL, et al. Frailty: emergence and consequences in women aged 65 and older in the Women's Health Initiative Observational Study. *J Am Geriatr Soc*. 2005; 53(8): 1321–1330. [PubMed: 16078957]
42. Barreto, PeS, Greig, C., Ferrandez, AM. Detecting and categorizing frailty status in older adults using a self-report screening instrument. *Arch Gerontol Geriatr*. 2012; 54(3):e249–254. [PubMed: 21889807]
43. Gill TM, Allore HG, Holford TR, et al. Hospitalization, restricted activity, and the development of disability among older persons. *JAMA*. 2004; 292(17):2115–2124. [PubMed: 15523072]
44. Yesavage JA, Brink TL, Rose TL, et al. Development and validation of a geriatric depression screening scale: a preliminary report. *J Psychiatr Res*. 1982; 17(1):37–49. [PubMed: 7183759]
45. Sheikh JI, Yesavage JA. Geriatric Depression Scale (GDS): Recent evidence and development of a shorter version. *Clinical Gerontologist*. 1986; 1(4):331–338.
46. Crawford GB, Robinson JA. The geriatric depression scale in palliative care. *Palliat Support Care*. 2008; 6(3):213–223. [PubMed: 18662414]
47. Adams KB. Depressive symptoms, depletion, or developmental change? Withdrawal, apathy, and lack of vigor in the Geriatric Depression Scale. *Gerontologist*. 2001; 41(6):768–777. [PubMed: 11723345]
48. Malakouti SK, Fatollahi P, Mirabzadeh A, et al. Reliability, validity and factor structure of the GDS-15 in Iranian elderly. *Int J Geriatr Psychiatry*. 2006; 21(6):588–593. [PubMed: 16783767]
49. Grool AM, Geerlings MI, Sigurdsson S, et al. Structural MRI correlates of apathy symptoms in older persons without dementia: AGES-Reykjavik Study. *Neurology*. 2014; 82(18):1628–1635. [PubMed: 24739783]
50. Eurelings LS, Ligthart SA, van Dalen JW, et al. Apathy is an independent risk factor for incident cardiovascular disease in the older individual: a population-based cohort study. *Int J Geriatr Psychiatry*. 2014; 29(5):454–463. [PubMed: 24105658]
51. Benton, AL., Hamsher, K. Multilingual Aphasia Examination. Iowa City: AJA Associates; 1989.
52. Wechsler, D. Wechsler Adult Intelligence Scale-III. San Antonio, TX: The Psychological Corporation; 1997.
53. Onyike CU, Sheppard JM, Tschanz JT, et al. Epidemiology of apathy in older adults: the Cache County Study. *Am J Geriatr Psychiatry*. 2007; 15(5):365–375. [PubMed: 17463187]
54. Feil D, Razani J, Boone K, Lesser I. Apathy and cognitive performance in older adults with depression. *Int J Geriatr Psychiatry*. 2003; 18(6):479–485. [PubMed: 12789667]
55. Shapiro ME, Mahoney JR, Zingman BS, et al. Apathy correlates with cognitive performance, functional disability, and HIV RNA plasma levels in HIV-positive individuals. *J Clin Exp Neuropsychol*. 2013; 35(9):934–945. [PubMed: 24090038]
56. Bastos-Barbosa RG, Ferrioli E, Coelho EB, et al. Association of frailty syndrome in the elderly with higher blood pressure and other cardiovascular risk factors. *Am J Hypertens*. 2012; 25(11): 1156–1161. [PubMed: 22810844]
57. Starr ME, Evers BM, Saito H. Age-associated increase in cytokine production during systemic inflammation: adipose tissue as a major source of IL-6. *J Gerontol A Biol Sci Med Sci*. 2009; 64(7):723–730. [PubMed: 19377014]
58. Brown PJ, Roose SP, Fieo R, et al. Frailty and depression in older adults: a high-risk clinical population. *Am J Geriatr Psychiatry*. 2014; 22(11):1083–1095. [PubMed: 23973252]

59. Buigues C, Padilla-Sánchez C, Fernández Garrido J, et al. The relationship between depression and frailty syndrome: a systematic review. *Aging Ment Health*. 2014;1–11. [PubMed: 24341516]
60. Paulson D, Lichtenberg PA. Vascular depression: an early warning sign of frailty. *Aging Ment Health*. 2013; 17(1):85–93. [PubMed: 22724516]
61. Lakey SL, LaCroix AZ, Gray SL, et al. Antidepressant use, depressive symptoms, and incident frailty in women aged 65 and older from the Women's Health Initiative Observational Study. *J Am Geriatr Soc*. 2012; 60(5):854–861. [PubMed: 22568404]
62. Moretti R, Cavressi M, Tomietto P. Gait and Apathy as Relevant Symptoms of Subcortical Vascular Dementia. *Am J Alzheimers Dis Other Dement*. 2014
63. Afilalo J, Karunanathan S, Eisenberg MJ, et al. Role of frailty in patients with cardiovascular disease. *Am J Cardiol*. 2009; 103(11):1616–1621. [PubMed: 19463525]
64. Bruunsgaard H, Pedersen M, Pedersen BK. Aging and proinflammatory cytokines. *Curr Opin Hematol*. 2001; 8(3):131–136. [PubMed: 11303144]
65. Penninx BW, Kritchevsky SB, Newman AB, et al. Inflammatory markers and incident mobility limitation in the elderly. *J Am Geriatr Soc*. 2004; 52(7):1105–1113. [PubMed: 15209648]
66. Penninx BW, Nicklas BJ, Newman AB, et al. Metabolic syndrome and physical decline in older persons: results from the Health, Aging And Body Composition Study. *J Gerontol A Biol Sci Med Sci*. 2009; 64(1):96–102. [PubMed: 19164274]
67. Li H, Manwani B, Leng SX. Frailty, inflammation, and immunity. *Aging Dis*. 2011; 2(6):466–473. [PubMed: 22396895]
68. Verghese J, Holtzer R, Oh-Park M, et al. Inflammatory markers and gait speed decline in older adults. *J Gerontol A Biol Sci Med Sci*. 2011; 66(10):1083–1089. [PubMed: 21719612]
69. Leng SX, Tian X, Matteini A, et al. IL-6-independent association of elevated serum neopterin levels with prevalent frailty in community-dwelling older adults. *Age Ageing*. 2011; 40(4):475–481. [PubMed: 21586579]
70. Bremner MA, Beekman AT, Deeg DJ, et al. Inflammatory markers in late-life depression: results from a population-based study. *J Affect Disord*. 2008; 106(3):249–255. [PubMed: 17716746]
71. Howren MB, Lamkin DM, Suls J. Associations of depression with C-reactive protein, IL1, and IL-6: a meta-analysis. *Psychosom Med*. 2009; 71(2):171–186. [PubMed: 19188531]
72. Dowlati Y, Herrmann N, Swardfager W, et al. A meta-analysis of cytokines in major depression. *Biol Psychiatry*. 2010; 67(5):446–457. [PubMed: 20015486]
73. Spalletta G, Cravello L, Imperiale F, et al. Neuropsychiatric symptoms and interleukin-6 serum levels in acute stroke. *J Neuropsychiatry Clin Neurosci*. 2013; 25(4):255–263. [PubMed: 24247852]
74. Hansson GK. Inflammation, atherosclerosis, and coronary artery disease. *N Engl J Med*. 2005; 352(16):1685–1695. [PubMed: 15843671]
75. Wancata J, Alexandrowicz R, Marquart B, et al. The criterion validity of the Geriatric Depression Scale: a systematic review. *Acta Psychiatr Scand*. 2006; 114(6):398–410. [PubMed: 17087788]

Clinical Points

- Although apathy is one of the most common neuropsychiatric symptoms in older adults, little is known of its effect on frailty and motoric decline.
- Presence of apathy was associated with increased risk of developing slow gait, frailty and disability in older adults.
- Apathy screenings may prevent functional decline and should become a routine part of psychiatric visits with older adults.

Figure 1.
Hazard curves for pooled sample, stratified by apathy status.

Figure 2.
Hazard curves for pooled sample, stratified by number of apathy symptoms

Table 1Baseline characteristics of study population and prevalence/incidence rates of study outcomes^a

<u>Baseline demographics</u>	Cohort 1^b (n = 625)	Cohort 2^c (n = 312)	<i>p</i> value
Age, y	75.21 (6.40)	76.4 (6.87)	0.01
Female, % (n)	53.3 (333)	56.7 (177)	0.26
Education, y	17.5 (2.66)	14.65 (3.08)	<0.01
Summary illness index (range, 0–9)	1.29 (1.04)	1.18 (.95)	0.15
No. of depressive symptoms (range, 0–12)	1 (1.33)	1 (1.12)	0.99
2 Apathy symptoms, % (n)	20.2 (126)	26 (81)	0.04
<u>Study outcomes</u>			
Follow-up, y	3.18 (1.43)	1.74 (0.66)	<0.01
Slow Gait, % (n)			
Prevalence	10.9 (68)	15.1 (47)	0.06
Incidence	14.5 (81)	8.3 (22)	0.03
Frailty, % (n)			
Prevalence	12.2 (76)	17.9 (56)	0.01
Incidence	17.9 (98)	14.8 (38)	0.31
Disability, % (n)			
Prevalence	3.8 (24)	4.5 (14)	0.62
Incidence	5.3 (32)	2.3 (7)	0.04

^aValues are reported as mean (SD) unless otherwise noted^bLonGenity cohort^cCentral Control of Mobility in Aging cohort

Table 2Baseline characteristics of study population according to apathy status^a

	Apathy (n = 207)	No Apathy (n = 730)	p value
Age, y	77.38 (7.00)	75.08 (6.39)	<0.01
Female, % (n)	55.07 (114)	53.70 (392)	0.73
Education, y	15.86 (2.96)	16.75 (3.14)	<0.01
Summary illness index (range, 0–9)	1.72 (1.09)	1.12 (0.95)	<0.01
No. of depressive symptoms (range, 0–12)	3.22 (3.33)	1.24 (1.87)	<0.01

^aValues are reported as mean (SD) unless otherwise noted

Table 3

Longitudinal association between baseline apathy symptoms and incident slow gait, frailty, and disability^a

	N	Events	Model 1 ^b	Model 2 ^c	Model 3 ^d	Model 4 ^e
Slow gait						
Cohort 1	557	81	1.86 (1.11–3.12) .018	1.83 (1.08–3.10) .025	1.59 (.94–2.7) .086	1.87 (1.10–3.18) .021
Cohort 2	265	22	2.92 (1.27–6.84) .012	2.73 (1.16–6.40) .021	3.02 (1.27–7.16) .012	2.97 (1.27–6.91) .012
Pooled sample	822	103	2.10 (1.36–3.24) .001	1.98 (1.27–3.07) .002	1.85 (1.19–2.89) .007	2.11 (1.36–3.92) .001
Frailty						
Cohort 1	549	98	3.39 (2.15–5.36) <.001	3.13 (1.98–4.93) <.001	2.87 (1.79–5.59) <.001	3.33 (2.09–5.31) <.001
Cohort 2	256	38	1.97 (1.19–3.82) .044	2.14 (1.08–4.24) .029	2.29 (1.16–4.52) .017	2.02 (1.04–3.93) .038
Pooled sample	805	136	2.86 (1.96–4.16) <.001	2.79 (1.92–4.06) <.001	2.55 (1.73–3.75) <.001	2.82 (1.93–4.12) <.001
Disability						
Cohort 1	601	32	4.36 (2.10–9.06) <.001	3.80 (1.80–8.02) <.001	3.78 (1.78–8.04) .001	3.93 (1.82–8.42) <.001
Cohort 2	298	7	.64 (.06–6.70) .681	.70 (.06–8.3) .777	.64 (.06–7.25) .722	1.00 (.09–11.46) .997
Pooled sample	899	39	3.43 (1.73–6.79) <.001	2.94 (1.48–5.85) .002	3.07 (1.52–6.22) .002	3.39 (1.66–6.93) .001

^a All analyses with pooled sample adjusted for cohort status Cohort 1 is LonGenity and Cohort 2 is Central Control of Mobility in Aging cohort. Results presented as: HR (95% CI) p.

^b Adjusted for depressive symptoms minus apathy (0–12).

^c Adjusted for baseline demographics (age, sex, and education level) and depressive symptoms minus apathy (0–12).

^d Adjusted for health status: summary illness index (0–9) and depressive symptoms minus apathy (0–12).

^e Adjusted for cognitive functioning (FAS and digit symbol coding) and depressive symptoms minus apathy (0–12).