Published in final edited form as:

Nat Med. 2017 January ; 23(1): 91–99. doi:10.1038/nm.4251.

Dickkopf-1 promotes hematopoietic regeneration via direct and niche-mediated mechanisms

Heather A Himburg^{1,7}, Phuong L Doan^{2,7}, Mamle Quarmyne^{1,3}, Xiao Yan^{1,3}, Joshua Sasine¹, Liman Zhao¹, Grace V Hancock⁴, Jenny Kan¹, Katherine A. Pohl¹, Evelyn Tran¹, Nelson J Chao², Jeffrey R Harris², and John P Chute^{1,5,6}

¹Division of Hematology and Oncology, Department of Medicine, University of California, Los Angeles (UCLA), Los Angeles, California, USA

²Division of Hematologic Malignancies and Cellular Therapy, Duke University, Durham, North Carolina, USA

³Department of Pharmacology and Cancer Biology, Duke University, Durham, North Carolina, USA

⁴Department of Cell and Developmental Biology, University of California, Los Angeles, Los Angeles, California, USA

⁵Broad Center for Regenerative Medicine and Stem Cell Research, University of California, Los Angeles, Los Angeles, California, USA

⁶Jonsson Comprehensive Cancer Center, University of California, Los Angeles, Los Angeles, California, USA

Abstract

The role of osteolineage cells in regulating hematopoietic stem cell (HSC) regeneration following myelosuppression is not well understood. Here we show that deletion of the pro-apoptotic genes *Bak* and *Bax* in osterix (*Osx*, also known as *Sp7* transcription factor 7)-expressing cells in mice promotes HSC regeneration and hematopoietic radioprotection following total body irradiation. These mice showed increased bone marrow (BM) levels of the protein dickkopf-1 (*Dkk1*), which was produced in *Osx*-expressing BM cells. Treatment of irradiated HSCs with *Dkk1 in vitro* increased the recovery of both long-term repopulating HSCs and progenitor cells, and systemic administration of *Dkk1* to irradiated mice increased hematopoietic recovery and improved survival. Conversely, inducible deletion of one allele of *Dkk1* in *Osx*-expressing cells in adult mice inhibited the recovery of BM stem and progenitor cells and of complete blood counts following irradiation. *Dkk1* promoted hematopoietic regeneration via both direct effects on HSCs, in which treatment with *Dkk1* decreased the levels of mitochondrial reactive oxygen species and suppressed

Reprints and permissions information is available online at <http://www.nature.com/reprints/index.html>.

Correspondence should be addressed to J.P.C. (jchute@mednet.ucla.edu).

⁷These authors contributed equally to this work.

Note: Any Supplementary Information and Source Data files are available in the online version of the paper.

AUTHOR CONTRIBUTIONS

COMPETING FINANCIAL INTERESTS

The authors declare no competing financial interests.

senescence, and indirect effects on BM endothelial cells, in which treatment with Dkk1 induced epidermal growth factor (EGF) secretion. Accordingly, blockade of the EGF receptor partially abrogated Dkk1-mediated hematopoietic recovery. These data identify Dkk1 as a regulator of hematopoietic regeneration and demonstrate paracrine cross-talk between BM osteolineage cells and endothelial cells in regulating hematopoietic reconstitution following injury.

Perivascular stromal cells and vascular endothelial cells (ECs) regulate HSC maintenance in the BM of mice¹⁻³. Deletion of nestin-expressing mesenchymal stromal cells (MSCs) has also been shown to decrease HSC content in the BM, which is associated with HSC mobilization⁴. Leptin receptor (Lepr)- and paired related homeobox 1 (Prx1)-expressing perivascular cells and nestin-expressing stromal cells have been postulated to represent overlapping perivascular populations which regulate HSC maintenance *in vivo*^{5,6}. Recently, maintenance of quiescent, long-term repopulating HSCs was suggested to be dependent on chondroitin sulfate proteoglycan 4 (Cspg4+ or NG2+) pericytes localized to the arteriolar vascular niche, which are distinct from Lepr-expressing peri-sinusoidal cells⁷. These studies have confirmed the importance of perivascular cells and vascular ECs in regulating HSC maintenance.

Expansion of the BM osteoblast population has been shown to amplify the HSC pool *in vivo*^{8,9}, and ganciclovir-inducible depletion of BM osteoblasts decreases phenotypic hematopoietic stem and progenitor cell content¹⁰. However, deletion of the genes encoding stem cell factor (SCF), the chemokine CXCL12 or N-cadherin in BM osteoblasts does not have effects on HSC content during homeostasis^{1-3,11}. Sp7 (Osx) is a transcription factor expressed by mesenchymal progenitor cells, which have been shown via lineage tracing to be osteolineage-restricted in the adult mouse^{6,12,13}. In the fetal BM, Osx-expressing cells contribute to nascent bone tissues and transient stromal cells, whereas perinatally, Osx-expressing cells contribute to long-lived MSCs and osteolineage cells^{6,13}. Notably, deletion of *Dicer1*, which encodes an RNaseIII endonuclease, in Osx-expressing cells promotes myelodysplasia in mice, suggesting that aberrant function of Osx-expressing cells may contribute to the development of hematologic malignancy¹⁴. Deletion of *CXCL12* from Osx-expressing cells was also shown to deplete B lymphoid progenitors in homeostasis, but no effect on HSC function was observed³.

Although genetic studies have provided insight into the function of BM niche cells in regulating hematopoiesis during homeostasis, important questions remain regarding the contributions of niche cells during stress or injury, as well as the effects of injury on niche-mediated regulation of HSCs. We and others have recently demonstrated the essential role of BM ECs in regulating HSC regeneration following myelotoxicity¹⁵⁻¹⁷, and we identified two BM EC-derived paracrine factors, pleiotrophin (PTN) and EGF, as regulators of HSC regeneration *in vivo*^{18,19}. However, the functions of BM mesenchymal and osteolineage cells in regulating HSC regeneration remain less well understood. Here we directly tested the function of Osx-expressing BM cells in regulating hematopoietic regeneration following myelosuppression and discovered that Osx-expressing BM cells promote hematopoietic regeneration via secretion of Dkk1.

RESULTS

Deletion of *Bak* and *Bax* in *Osx*-expressing cells radioprotects the hematopoietic system

To test whether radioprotection of these cells would alter the hematopoietic response to irradiation, we used *Cre-loxP* technology to delete *loxP*-flanked (floxed; FL) genes encoding the pro-apoptotic factors *Bak* and *Bax* in *Osx*-expressing cells (from *Sp7-Cre;Bak1^{-/-};Bax^{FL/-}* mice, hereafter referred to as *Osx-Cre;Bak1^{-/-};Bax^{FL/-}* mice)^{15,20}. To determine the proportion of *Osx*-labeled cells that expressed *Osx* in 8-week-old mice, we used *Sp7-Cherry* (hereafter referred to as *Osx-Cherry*) reporter mice, because these mice have a stronger reporter signal than mice with a GFP reporter driven by *Osx-Cre*. In 8-week-old *Osx-Cherry* reporter mice, approximately 60% of Cherry-labeled BM cells expressed *Osx*, as measured by flow cytometry, whereas in 5-d-old mice, 83% of Cherry-labeled BM cells expressed *Osx* (Fig. 1a). These results indicate that a subset of *Osx*-labeled BM cells lose *Osx* protein expression between birth and adulthood. Adult *Osx-Cre;Bak1^{-/-};Bax^{FL/-}* mice showed no baseline differences in the frequency of *Osx*⁺ BM cells, BM trabecular bone content, complete blood counts, HSC content or repopulating HSC function, as compared to those in *Osx-Cre;Bak1^{-/-};Bax^{FL/+}* control mice, which retain one wild-type allele of *Bax* (Supplementary Fig. 1a–g). Next we irradiated both strains of mice with 500 cGy total body irradiation (TBI) to assess the response of *Osx*-expressing BM cells and of hematopoietic stem and progenitor cells to injury. *Osx-Cre;Bak1^{-/-};Bax^{FL/-}* mice maintained *Osx*⁺ BM cells at day 3 after irradiation as compared to *Osx-Cre;Bak1^{-/-};Bax^{FL/+}* mice, which showed depletion of this population (Fig. 1b and Supplementary Fig. 1h). As compared to *Osx-Cre;Bak1^{-/-};Bax^{FL/+}* mice at day 7 after TBI, *Osx-Cre;Bak1^{-/-};Bax^{FL/-}* mice displayed increased BM cellularity, increased numbers of c-kit⁺Sca-1⁺lineage⁻ (KSL) stem-progenitors, SLAMF6⁺KSL HSCs, colony-forming cells (CFCs), and peripheral blood (PB) white blood cells (WBCs), neutrophils and lymphocytes (Fig. 1c–f). In competitive transplantation experiments, congenic B16.SJL mice transplanted with BM cells from *Osx-Cre;Bak1^{-/-};Bax^{FL/-}* mice showed a significant increase in multilineage hematopoietic cell reconstitution in both primary and secondary transplanted mice, as compared to mice that were transplanted with BM from *Osx-Cre;Bak1^{-/-};Bax^{FL/+}* mice (Fig. 1g,h). Taken together, these data suggest that the hematopoietic response to radiation injury is regulated by *Osx*-expressing BM cells and that deletion of the intrinsic pathway of apoptosis in these cells promotes radioprotection of the hematopoietic system.

Dkk1 is differentially expressed by *Osx*⁺ BM cells

Because *Osx-Cre;Bak1^{-/-};Bax^{FL/-}* mice showed enhanced hematopoietic recovery following TBI, we hypothesized that *Osx*-expressing BM cells might produce factors that promote hematopoietic regeneration. Analysis of BM supernatants from *Osx-Cre;Bak1^{-/-};Bax^{FL/-}* mice revealed enrichment for several candidate proteins as compared to *Osx-Cre;Bak1^{-/-};Bax^{FL/+}* mice (Supplementary Fig. 2a). Among these candidates, we focused on Dkk1, a Wnt inhibitor, because Dkk1 was previously shown to be expressed and secreted by osteolineage cells under the control of osterix²¹, although it has no established role in regulating hematopoietic regeneration. Furthermore, in preliminary screening experiments we found that Dkk1 promoted CFC recovery from BM KSL cells that were irradiated with 300 cGy *in vitro*, whereas two other candidates, tissue inhibitor of

metalloproteinase 2 (Timp2) and the chemokine Ccl20 (also known as MIP-3a), had no effect (data not shown). ELISA confirmed that *Osx-Cre;Bak1^{-/-};Bax^{FL/-}* mice contained significantly higher levels of Dkk1 protein in the BM than *Osx-Cre;Bak1^{-/-};Bax^{FL/+}* mice (Fig. 1i). Expression of *Dkk1* mRNA was also significantly increased in BM osteolineage cells from *Osx-Cre;Bak1^{-/-};Bax^{FL/-}* mice, as compared to that in *Osx-Cre;Bak1^{-/-};Bax^{FL/+}* mice; in both mouse strains, nearly all of the detectable Dkk1 protein was found in *Osx⁺* BM cells, as assessed by flow cytometry (Fig. 1j and Supplementary Fig. 2b). Moreover, we found that *Osx⁺* BM cells expressed significantly more Dkk1 than BM ECs in C57BL/6 mice, both at steady state and following TBI (Fig. 1k). These results indicate that *Osx⁺* BM cells are the primary source of Dkk1 in both *Osx-Cre;Bak1^{-/-};Bax^{FL/-}* and wild-type mice.

Dkk1 treatment accelerates hematopoietic regeneration *in vitro* and *in vivo*

We next sought to evaluate whether Dkk1 regulates hematopoietic regeneration following irradiation. First, we assessed the effect of Dkk1 on the function of non-irradiated HSCs *in vitro* and *in vivo*. Treatment of non-irradiated BM KSL cells with 500 ng/ml Dkk1 in medium supplemented with thrombopoietin (TPO), SCF and Flt-3 ligand had no effect on total cell expansion, KSL expansion or CFC content, as compared to cells treated with medium alone (Supplementary Fig. 3a,b). However, systemic treatment of C57BL/6 mice with 10 µg Dkk1 subcutaneously every other day for 4 weeks caused a significant increase in the numbers of BM KSL cells, BM c-kit⁺lin⁻ cells and BM CFCs, as well as in the PB neutrophil frequency, compared to saline-treated mice (Supplementary Fig. 3c–e). These data suggest that during homeostasis, Dkk1 promotes the *in vivo* expansion of the myeloid progenitor cell pool.

To evaluate the function of Dkk1 in regulating hematopoietic regeneration following irradiation, we irradiated BM KSL cells with 300 cGy and cultured them in cytokine-containing medium with or without 500 ng/ml Dkk1. Dkk1 treatment significantly increased the recovery of CFCs at day 7 of culture, as compared to treatment with cytokines alone (Fig. 2a). Furthermore, mice that had been competitively transplanted with the progeny of irradiated, Dkk1-treated KSL cells showed a significant increase in multilineage engraftment of donor CD45.2⁺ cells over the course of 16 weeks after transplantation (Fig. 2b). These results suggest that Dkk1 directly promotes the regeneration of HSCs capable of primary competitive repopulation following irradiation.

Because Dkk1 promotes the regeneration of irradiated hematopoietic stem and progenitor cells (HSPCs) *in vitro*, we next tested whether systemic administration of Dkk1 could accelerate hematopoietic regeneration *in vivo* following TBI. Adult C57BL/6 mice were irradiated with 800 cGy TBI and treated subcutaneously with 10 µg Dkk1 or saline every other day for 21 d, beginning at day 1 after TBI. Systemic administration of Dkk1 caused a greater than eightfold increase in Dkk1 levels in the BM of C57BL/6 mice within 1 h, as compared to untreated mice (Fig. 2c). Dkk1 treatment increased the recovery of WBCs, neutrophil counts, platelets and BM cell counts at day 21 as compared to saline-treated mice (Fig. 2d,e). Furthermore, the numbers of BM KSL cells and CFCs were significantly increased in Dkk1-treated mice as compared to those in saline-treated controls (Fig. 2f,g). Notably, following 800 cGy TBI, only 4 of 15 (27%) C57BL/6 mice that had been treated

with saline survived to day 30, as compared to 13 of 14 mice (93%) that had been treated with Dkk1, a nearly 350% increase in survival in response to Dkk1 treatment ($P = 0.0004$) (Fig. 2h).

We also evaluated whether Dkk1 treatment could affect long-term hematopoiesis in mice following radiation exposure. Adult C57BL/6 mice were irradiated with 600 cGy TBI, treated with 10 μ g Dkk1 every other day for 21 d and evaluated at 12 weeks after irradiation. As compared to saline-treated controls, Dkk1-treated mice showed no differences in complete blood counts and BM cell counts, percentages of SLAMF6⁺KSL or KSL cells, or frequencies of BM myeloid cells, B cells or T cells (Supplementary Fig. 4a–c). However, at 12 weeks following 600-cGy irradiation, saline-treated mice had substantially reduced numbers of BM CFCs as compared to irradiated, Dkk1-treated mice (Fig. 2i). These data suggest that TBI causes a long-term decline in myeloid progenitor cell function and that short-term treatment with Dkk1 abrogates radiation-induced damage to this progenitor cell population.

Inhibition of Dkk1 abrogates hematopoietic regeneration following TBI

Because systemic administration of Dkk1 augments hematopoietic regeneration and survival of mice following TBI, we next tested whether endogenous Dkk1 is necessary for normal hematopoietic regeneration following TBI. We irradiated adult C57BL/6 mice with a sublethal TBI dose of 500 cGy and then administered 100 μ g of a blocking Dkk1-specific antibody or an IgG control antibody every other day for 14 d. Anti-Dkk1 treatment significantly lowered Dkk1 levels in the BM at 24 h after treatment (Supplementary Fig. 5a). At day 10 following irradiation, mice that were treated with anti-Dkk1 had a significant decrease in BM cell counts and in the numbers of PB WBCs, neutrophils, lymphocytes, KSL cells and CFCs, as compared to those in irradiated, IgG-treated mice (Fig. 3a–c and Supplementary Fig. 5b). Furthermore, treatment with anti-Dkk1 caused a significant decrease in the 30-d survival of irradiated mice (6 of 12 alive, 50%) as compared to irradiated controls (12 of 12, 100%; $P = 0.006$) (Fig. 3d). Taken together, these results suggest that endogenous Dkk1 plays an important role in regulating hematopoietic regeneration and survival following myelosuppression.

Deletion of Dkk1 in Osx-labeled BM cells inhibits hematopoietic regeneration

We next tested whether deletion of a single allele of Dkk1 in Osx-labeled cells (*Osx-Cre;Dkk1^{FL/+}* mice) would affect the hematopoietic response to TBI as compared to *Dkk1^{+/+}* mice. 8-week-old *Osx-Cre;Dkk1^{FL/+}* mice showed no significant differences in hematologic profile or BM stem and progenitor cell content at baseline compared to that in *Dkk1^{+/+}* mice (Supplementary Fig. 6a–d). However, at day 10 following 500-cGy TBI, *Osx-Cre;Dkk1^{FL/+}* mice showed a significant decrease in the numbers of WBCs, neutrophils and lymphocytes, as compared to those in *Dkk1^{+/+}* mice (Fig. 3e). BM cell counts and the number of BM KSL cells, CFCs, BM myeloid cells, T cells and B cells were also decreased significantly in *Osx-Cre;Dkk1^{FL/+}* mice as compared to those in *Dkk1^{+/+}* mice (Fig. 3f–h and Supplementary Fig. 6e). These data suggest that endogenous Dkk1, produced by Osx-labeled BM cells, has a necessary role in regulating the hematopoietic response to irradiation.

Next we treated 500-cGy-irradiated *Osx-Cre;Dkk1^{FL/+}* mice with 10 µg of Dkk1 every other day for 10 d to determine whether Dkk1 treatment could rescue the hematopoietic phenotype of these mice. Although Dkk1 treatment did not alter the complete blood counts or BM cell counts in *Osx-Cre;Dkk1^{FL/+}* mice at day 10 after TBI, this treatment did significantly increase the numbers of BM KSL cells and BM CFCs, as compared to those in untreated mice (Fig. 3e–h). These results provide evidence that *Dkk1* deletion is responsible for the BM stem and progenitor cell abnormalities observed early after radiation injury in *Osx-Cre;Dkk1^{FL/+}* mice.

Our initial analysis of *Osx-Cre;Dkk1^{FL/+}* mice, in which the Cre protein was expressed from inception, did not allow for discrimination between the effects of *Dkk1* deletion in *Osx*-expressing osteoprogenitors and those in *Osx*-expressing long-lived MSCs in the BM, which can persist into adulthood⁶. In *Osx-Cre;Dkk1^{FL/+}* mice, expression of a tetracycline transactivator under the regulation of the osterix (*Sp7*) promoter controls the expression of a tetracycline response element (TRE)-controlled Cre fusion protein²², such that doxycycline (Dox) administration prevents Cre-mediated recombination in *Osx*-expressing cells (Dox-Off). To study the effects of *Dkk1* deletion starting at 8 weeks of age and older, we treated *Osx-Cre;Dkk1^{FL/+}* mice with Dox from inception to 8 weeks of age (hereafter referred to as 8-week-*Osx-Cre;Dkk1^{FL/+}* mice). Irradiation of 8-week-*Osx-Cre;Dkk1^{FL/+}* mice with 500 cGy TBI at 10 weeks of age caused a significant decrease in the numbers of PB WBCs, neutrophils and lymphocytes, as well as in BM cell counts and in the numbers of BM KSL cells and BM CFCs at day 10 following TBI, as compared to those in *Osx-Cre;Dkk1^{+/+}* mice (Fig. 3i–l). The enhanced hematopoietic toxicity following TBI that was observed in 8-week-*Osx-Cre;Dkk1^{FL/+}* mice was comparable to that observed in bearing *Dkk1* deletion from inception, *Osx-Cre;Dkk1^{FL/+}* mice, suggesting that BM osteoprogenitor cells are the primary source of Dkk1 in regulating hematopoietic regeneration.

Dkk1 suppresses reactive oxygen species generation and p38-mediated senescence in irradiated HSCs

Ionizing radiation can cause hematopoietic injury via generation of reactive oxygen species (ROS) and induction of HSC senescence²³. At 24 h following 300-cGy irradiation of cells *in vitro*, the levels of mitochondrial ROS increased substantially in BM CD34⁺-KSL cells, KSL cells and lineage-negative (lin[−]) cells (Fig. 4a and Supplementary Fig. 7a). Treatment with 500 ng/ml Dkk1 suppressed radiation-induced mitochondrial ROS generation in each cell population (Fig. 4a and Supplementary Fig. 7a). *In vivo*, systemic Dkk1 treatment of C57BL/6 mice irradiated with 500 cGy TBI resulted in a significant decrease in the levels of mitochondrial ROS in BM CD34⁺-KSL cells at 24 h after irradiation (Supplementary Fig. 7b). *In vitro* treatment with Wnt3a (a canonical Wnt agonist) or Wnt5a (a noncanonical Wnt ligand) did not significantly alter mitochondrial ROS levels in irradiated CD34⁺-KSL cells, KSL cells or lin[−] cells beyond the effects observed with radiation alone (Fig. 4a and Supplementary Fig. 7a). Consistent with its effects on decreasing ROS levels in irradiated HSCs, Dkk1 treatment of BM CD34⁺-KSL cells increased CFC recovery *in vitro* (Supplementary Fig. 7c). However, in the presence of a ROS scavenger, *N*-acetylcysteine (NAC), Dkk1 treatment did not increase CFC recovery from irradiated KSL cells, suggesting that Dkk1-mediated hematopoietic regeneration is dependent, at least in part, on suppression

of ROS (Supplementary Fig. 7c). Elevated ROS levels drive activation of p38 mitogen-activated protein kinase (MAPK), which mediates a cytokine release phenomenon associated with senescence^{23,24}. Irradiation significantly increased p38 phosphorylation in BM CD34⁺KSL cells at 24 h after exposure, and this increase was significantly attenuated by Dkk1 treatment (Fig. 4b). Of note, Dkk1 treatment had more modest effects in reducing p38 phosphorylation in irradiated BM KSL and lin⁺ cells (Supplementary Fig. 7e). Dkk1 treatment also significantly reduced senescence-associated β -galactosidase (SA- β -gal) levels in irradiated BM CD34⁺KSL cells, KSL cells and lin⁺ cells (Fig. 4c and Supplementary Fig. 7f). In contrast, treatment with Wnt3a or Wnt5a had no significant effects on either p38 phosphorylation or SA- β -gal levels in irradiated BM CD34⁺KSL cells (Fig. 4b,c). In keeping with its observed effects on SA- β -gal levels, Dkk1 suppressed radiation-induced expression of the p16-encoding isoform of cyclin-dependent kinase inhibitor 2a (*Cdkn2a*; hereafter referred to as *p16*) in BM CD34⁺KSL cells *in vitro* and attenuated *p16* expression in BM KSL cells in mice that were irradiated with 500 cGy (Fig. 4d). Dkk1 treatment also suppressed HSC apoptosis in irradiated BM KSL cells, as measured by caspase-3–caspase-7 activation (Supplementary Fig. 7g).

Next we extended our analysis of the effects of Dkk1 on radiation-induced senescence of BM HSC and progenitor cell populations by evaluating the long-term effects of systemic Dkk1 treatment on irradiated mice. At 12 weeks following 600 cGy TBI, C57BL/6 mice showed an increased frequency of SA- β -gal⁺ KSL cells, consistent with the idea that senescent BM stem and progenitor cells persist following radiation injury (Fig. 4e). This observation is in accord with a recent study that demonstrated that TBI induces persistent senescence of HSCs and premature aging of the hematopoietic system²⁵. Notably, mice that were treated with Dkk1 every other day through day 21 following TBI showed a significantly lower percentage of SA- β -gal⁺ KSL cells at 12 weeks, as compared to untreated, irradiated mice (Fig. 4e). Coupled with our observation that Dkk1 treatment led to an increase in BM CFCs at 12 weeks following irradiation (Fig. 2i), these data suggest that short-term treatment with Dkk1 promotes longer-term recovery of hematopoietic progenitor cells following radiation injury.

Dkk1 inhibits Wnt pathway activation in irradiated HSCs

Dkk1 is a member of the Dkk family of secreted proteins that can inhibit Wnt signaling via binding and sequestration of the Wnt co-receptors, LRP5 and LRP6²⁶. Using Axin2–LacZ reporter mice as previously described^{27,28}, we evaluated Wnt pathway activation in BM HSC and progenitor cell subsets in response to irradiation and Dkk1 treatment. Exposure of mice to 500 cGy TBI increased Wnt signaling in BM CD34⁺KSL cells, KSL cells and lin⁺ cells at 24 h after exposure (Fig. 4f). This activation of Wnt signaling was suppressed by Dkk1 treatment in these cell populations at 24 h after irradiation, as measured by the percentage of Axin2⁺ cells and the mean fluorescence intensity (MFI) of the Axin2–lacZ signal (Fig. 4f and Supplementary Fig. 8a). Of note, Wnt pathway activation in response to irradiation seemed to be stronger in the BM CD34⁺KSL cells and the KSL cells as compared to that in BM lin⁺ cells (Fig. 4f). As assessed by the percentage of Axin2⁺ cells, Dkk1-mediated suppression of Wnt signaling was comparable across the three cell populations, although MFI analysis suggested that Dkk1 had the strongest suppressive effect in BM KSL

cells (Supplementary Fig. 8a). *Axin2* mRNA levels increased concordantly in BM KSL cells in mice at 24 h after 500 cGy TBI, and this increase in *Axin2* mRNA was abrogated by Dkk1 treatment (Supplementary Fig. 8b).

To determine the functional relevance of Wnt inhibition to Dkk1-mediated hematopoietic regeneration, we used a viral Cre-based approach to delete *Ctnnb1*, the gene encoding β -catenin, in BM KSL cells from *Ctnnb1*^{FL/FL} mice (Supplementary Fig. 9a)²⁹. At 96 h following 300 cGy irradiation *in vitro*, Dkk1 treatment decreased ROS generation in *Ctnnb1*^{+/+} BM KSL cells but not in *Ctnnb1*^{-/-} KSL cells (Supplementary Fig. 9b). Similarly, Dkk1 treatment increased CFC recovery at 96 h after irradiation from *Ctnnb1*^{+/+} KSL cells but not from *Ctnnb1*^{-/-} KSL cells (Supplementary Fig. 9c). These data suggest that Wnt pathway inhibition contributes, at least in part, to the early effects of Dkk1 on ROS generation and hematopoietic progenitor recovery.

Dkk1 promotes hematopoietic regeneration via induction of EGF secretion by BM ECs

Next we tested the possibility that there might be cross-talk between Dkk1 and pleiotrophin (PTN) or EGF, two vascular-niche-derived proteins that we have previously shown to be HSC regenerative factors^{16,18,19}. We first tested whether treatment of primary BM osteolineage cells (*Osx*⁺*osteocalcin*⁺ cells cultured from femur bone explants) with PTN or EGF could induce Dkk1 secretion. However, this was not observed in 7-d culture; indeed, EGF treatment significantly reduced Dkk1 secretion (Supplementary Fig. 10). Next we asked whether Dkk1 could induce the secretion of these paracrine factors by primary BM ECs, as BM ECs are known to regulate HSC regeneration^{15,17,30,31}. Although Dkk1 treatment did not induce the secretion of PTN by primary BM ECs, it induced a greater than 5,000-fold increase in EGF levels after 7 d, as compared to that in untreated BM ECs (Fig. 4g). Systemic administration of Dkk1 to mice irradiated with 500 cGy caused a marked increase in EGF levels in the BM at days 10 and 14 following radiation (Fig. 4g). To test the functional relevance of Dkk1-mediated induction of EGF secretion to hematopoietic regeneration, we orally administered 10 μ g per g body weight of erlotinib, an EGF receptor antagonist, daily for 10 d to 500-cGy-irradiated, Dkk1-treated mice. Erlotinib treatment abrogated Dkk1-mediated recovery of PB WBCs, neutrophils, BM cell counts and BM KSL cells at day 10 and day 14, as compared to that in mice treated with Dkk1 alone (Fig. 4h). These results suggest that, in addition to its direct effects on hematopoietic cells, Dkk1 promotes hematopoietic regeneration via induction of EGF secretion by BM ECs.

DISCUSSION

The functions of BM osteolineage cells and MSCs in regulating HSC regeneration are incompletely understood^{32,33}. Here we demonstrate that deletion of a single allele of *Dkk1* in *Osx*-expressing BM cells abrogates hematopoietic regeneration following irradiation. Inducible deletion of *Dkk1* in *Osx*-expressing cells in 8-week-old mice caused a comparable defect in hematopoietic regeneration in response to TBI as observed in mice bearing constitutive deletion of *Dkk1* in *Osx*-expressing cells. These data suggest that BM osteoprogenitor cells, rather than long-lived MSCs, are the primary source of Dkk1 that regulates hematopoietic regeneration in adult mice.

In gain-of-function studies, systemic Dkk1 treatment accelerated hematopoietic regeneration and improved survival following lethal dose TBI. This marked improvement in survival may have been related to the augmented recovery of myeloid progenitor cells, which have been shown to mediate survival following lethal dose TBI³⁴. Dkk1 treatment also directly promoted the regeneration of HSCs with competitive repopulating capacity following irradiation *in vitro*. In a prior study of Dkk1 transgenic mice by Fleming *et al.*, overexpression of Dkk1 at steady state was associated with an increased frequency of HSCs capable of primary competitive repopulation, coupled with a deficiency in HSCs capable of serial transplantation³⁵. In the current study, we examined the effects of Dkk1 treatment or *Dkk1* deletion on the hematopoietic regenerative response to myelotoxic injury, whereas Fleming *et al.* described the hematopoietic effects of Dkk1 overexpression during homeostasis³⁵. It is possible that Dkk1-mediated signaling has distinct effects on HSCs in the setting of stress versus homeostasis, as has been demonstrated for Notch signaling³⁶. Furthermore, we found that Dkk1 treatment induces BM ECs to secrete EGF, a regenerative factor for HSCs¹⁹, an effect that likely also contributes to the differences in the results of our study as compared to the prior study by Fleming *et al.*³⁵.

Ionizing radiation causes the generation of ROS in HSCs, and ROS accumulation contributes to radiation-induced HSC apoptosis and senescence^{23,37,38}. ROS are also deleterious to short- and long-term HSC function and HSC self-renewal capacity^{39,40}. ROS promote HSC senescence via activation of p38 MAPK and p16^{23,24}. Conversely, FoxO proteins are essential for downregulating ROS levels in HSCs, and transfer of ROS via gap junctions containing the gap junction protein Gja1 (also known as connexin-43) to BM stromal cells can mitigate ROS-mediated HSC senescence^{39,40}. Notably, ROS generation occurs during BM or cord blood collection in ambient air, and this was recently shown to be highly damaging to HSC yield and function⁴¹. Furthermore, high-dose irradiation has been shown to markedly increase pO₂ concentrations in HSC niches via BM vascular disruption⁴². These findings underscore the therapeutic importance of suppressing ROS after myelosuppressive injury and may explain, in part, the potency of Dkk1 administration in promoting the survival of lethally irradiated mice.

Canonical Wnt signaling has been shown to regulate the fate of HSCs, myeloid progenitors and T cell subsets in a dosage-dependent manner^{26–28,43}. Using Axin2–lacZ reporter mice, we observed that TBI rapidly induced Wnt pathway activation in HSCs and progenitor cells and that systemic treatment with Dkk1 suppressed this response. Dkk1 also suppressed ROS generation and HSC senescence in irradiated mice, concordant with inhibitory effects on Wnt signaling. These findings are consistent with recent studies suggesting that Dkk1 treatment suppresses ROS generation in nonhematopoietic tissues, as well as a report demonstrating that Wnt pathway activation promotes mitochondrial ROS generation and senescence in myoblasts and fibroblasts^{44,45}. However, in the hematopoietic system, several studies have suggested that activation of canonical Wnt signaling promotes HSC self-renewal^{46–48}. A recent study by Lento *et al.* suggested that mice with constitutive deletion of *Ctnnb1* in hematopoietic Vav⁺ cells showed reduced hematopoietic progenitor cell recovery at day 14 following TBI⁴⁹. Several factors may explain the different observations between our study and that of Lento *et al.* First, acute inhibition of Wnt signaling with Dkk1 in the first 72 h after irradiation may have different effects on HSC regeneration as compared to

that in mice with constitutive deletion of *Ctnnb1*. Second, it is possible that Dkk1 mediates effects on hematopoietic regeneration via mechanisms that are independent from its inhibitory effects on canonical Wnt signaling, as observed in models of kidney epithelial repair and cardiogenesis^{50,51}. As we discovered, Dkk1 acts indirectly on BM ECs, yielding a marked increase in EGF secretion, which contributed to hematopoietic regeneration. In this regard, a recent study demonstrated that ECs express LRP6, a Dkk1 receptor, and that Dkk1 and Dkk2 may regulate angiogenesis⁵². Third, the dosage of canonical Wnt pathway activation has been demonstrated to substantially impact HSCs, progenitor cells and HSC differentiation^{28,53}, and high dose levels of Wnt pathway activation can result in decreased HSC repopulating capacity^{54,55}. Our data suggest that during the early period following radiation injury, strong induction of Wnt signaling in HSCs may be deleterious, whereas early inhibition of Wnt signaling is beneficial.

Canonical and noncanonical Wnt signaling have been shown to have discrete effects on hematopoiesis⁵⁶. Wnt5a expression in HSCs increases with aging, causing a shift to noncanonical Wnt signaling and a decline in HSC function in older mice⁵⁷. HSC senescence occurs over time in aged mice and is exacerbated by stress, including radiation exposure^{23,37}. Here we observed that acute radiation injury increased canonical Wnt pathway activation in HSCs and that treatment with Dkk1 suppressed this signaling, in association with repression of ROS generation and HSC senescence. However, treatment with Wnt5a did not exacerbate radiation-induced ROS generation or HSC senescence. We postulate that the effects of Wnt5a and noncanonical Wnt signaling during hematopoietic aging may be distinct from the effects of noncanonical or canonical Wnt signaling in the setting of acute injury.

Over the past decade, as the cellular composition of the HSC niche has been elucidated, important questions have emerged regarding the hierarchy of cellular niche components and the manner through which niche cells communicate during their orchestration of HSC maintenance and regeneration^{58,59}. Recent studies suggest that distinct BM ECs within arterial and sinusoidal blood vessels regulate the HSC maintenance and activation⁶⁰. Our results demonstrate that BM osteoprogenitors regulate hematopoietic regeneration via secretion of Dkk1, which promotes hematopoietic regeneration directly via inhibition of HSC senescence and indirectly via induction of EGF secretion by BM ECs. This latter mechanism provides a unique example of cross-talk between distinct niche cell types and highlights the importance of cooperative niche cell interactions in hematopoietic regeneration.

METHODS

Methods, including statements of data availability and any associated accession codes and references, are available in the online version of the paper.

ONLINE METHODS

Mice

All animal procedures were performed in accordance with animal use protocols approved by UCLA and Duke University. *Osx-Cre;Bak1^{-/-};Bax^{+/-}* mice were bred with *Bak1^{-/-};Bax^{FL/FL}* mice bearing a constitutive deletion of *Bak1* and *loxP*-flanked (floxed; FL) *Bax* alleles to generate *Osx-Cre;Bak1^{-/-};Bax^{FL/-}* mice and *Osx-Cre;Bak1^{-/-};Bax^{FL/+}* mice. *Bak1^{-/-};Bax^{+/-}* and *Bak1^{-/-};Bax^{FL/FL}* mice were generated as previously described^{15,20}. *Osx-Cre-Gfp* mice, *Ctnnb1^{FL/FL}* mice, *Osx-Cherry* mice⁶¹, and *Axin2-LacZ* mice⁶² were obtained from the Jackson Laboratory (Bar Harbor, ME). *Dkk1^{FL/+}* mice were generously provided by Dr. Seppo Vainio from the University of Oulu, Finland⁶³. *Dkk1^{FL/+}* mice were crossed with *Osx-Cre-Gfp* mice²² to generate tissue-specific deletion of *Dkk1*. Cre-mediated recombination in *Osx-Cre-Gfp* mice is regulated by treatment with doxycycline, as both tTA and tetO are expressed under regulation of the osterix (*Sp7*) promoter (Tet-off). *Dkk1* deletion was temporally controlled in some *Osx-Cre;Dkk1^{FL/+}* mice by doxycycline administration in drinking water from conception to 8 weeks of age (0.2 g doxycycline per liter water; Sigma Aldrich, St Louis, MO).

Histology and immunofluorescence staining

Hematoxylin and eosin (H&E) staining of femurs from mice was performed as previously described¹⁵. Femurs were fixed overnight in 4% paraformaldehyde, decalcified and paraffin-embedded. Immunofluorescence staining was performed on fixed, decalcified femurs embedded in OCT and sectioned with a cryostat as previously described¹⁵. Goat anti-osterix antibody (Santa Cruz Biotechnology, Santa Cruz, CA) was diluted 1:500 in blocking buffer and incubated overnight at 4 °C. Slides were rinsed and incubated with Alexa Fluor 488 (AF488)-conjugated donkey anti-goat antibody (Life Technologies, Grand Island, NY) at 1:500 dilution in blocking buffer. The slides were rinsed and mounted with DAPI anti-fade Gold (Life Technologies, Carlsbad, CA). Images were obtained using an Axiovert 200 microscope (Carl Zeiss, Thornwood, NY).

Flow cytometric analysis for Osx and Dkk1

Flow cytometric analysis for osterix protein was performed on whole BM following red blood cell (RBC) lysis. Cells were stained for V450-CD45 (BD Biosciences, Franklin Lakes, NJ) and then fixed and permeabilized using the BD Cytofix-Cytoperm kit (BD Biosciences). Cells were then stained for 1 h with rabbit anti-osterix (ab94744 1:500, Abcam, Cambridge UK) and goat anti-Dkk1 (ab188597, 1:500, Abcam), rinsed and stained with AF647-conjugated anti-goat antibody (Life Technologies) and AF488-conjugated anti-rabbit antibody (Life Technologies).

Isolation of Osx⁺ bone marrow cells

Osx-Cherry mice were used to isolate *Osx*-labeled BM cells. BM from femurs and tibias was isolated by crushing bones and then digesting them for 10 min at 37 °C in collagenase and dispase (Sigma-Aldrich). Cells were then rinsed and depleted of lineage-committed cells with the Miltenyi lineage-depletion kit (Miltenyi Biotec, San Diego, CA). The lineage-

negative fraction was stained for CD45 and sorted on a BD FACS ARIA. BM ECs were intravitally labeled by intravenous injection of 25 μ g of AF647-conjugated vascular endothelial (VE)-cadherin antibody (Biolegend 138006, San Diego, CA), Biolegend, San Diego, CA) into wild-type C57BL/6 mice. Mice were euthanized 15 min after injection. CD45⁺VE-cadherin⁺ cells were identified as BM ECs and CD45⁺Osx⁺ cells were identified as Osx⁺ BM cells.

Cytokine array and enzyme-linked immunosorbent assays (ELISAs)

Whole BM from one femur was collected from adult *Osx-Cre;Bak^{-/-};Bax^{FL/-}* mice and *Osx-Cre;Bak^{-/-};Bax^{FL/+}* mice at 7 d following 500 cGy TBI. After centrifugation, BM supernatants were collected into Iscove's modified Dulbecco's medium (IMDM) and analyzed for cytokine concentrations using the Quantibody mouse cytokine array C2000, according to the manufacturer's guidelines (RayBiotech, Inc., Norcross, GA). Dkk1 protein levels in the BM were confirmed by using a mouse Dkk1 ELISA kit (R&D systems, Minneapolis, MN).

Isolation and culture of BM osteoblasts and endothelial cells

To measure *Dkk1* gene and Dkk1 protein expression in BM osteolineage cells derived from *Osx-Cre;Bak^{-/-};Bax^{FL/-}* mice and *Osx-Cre;Bak^{-/-};Bax^{FL/+}* mice, BM osteolineage cells were isolated as previously described³². Briefly, ground bone fragments of mouse femurs and tibias were digested for 3 h with 0.2 mg/ml collagenase, rinsed, and then plated in osteolineage cell supportive medium in 24-well plates³². After 3 d, bone chips were removed, and adherent colonies were observed. RNA was isolated from adherent cells for analysis of *Bax*, *Dkk1*, *Osx*, and other markers. BM endothelial cells were grown by explanting vessels flushed from the bone marrow onto 40- μ m filters. Cells were grown to confluence in EGM-2 medium without EGF (Lonza, Walkersville, MD) with daily medium changes. For ELISA of PTN, EGF, and Dkk1 in the culture medium of BM osteolineage cells and BM ECs, medium was collected at day +7 of confluent cell culture. Cell culture supernatants were collected and concentrated 20-fold for ELISA analysis.

Gene expression analysis

For all studies, RNA was isolated using the Qiagen RNeasy micro kit (Qiagen, Valencia, CA). RNA was reverse-transcribed into cDNA using the iScript cDNA synthesis kit (Bio-Rad, Hercules, CA). Real-time PCR analysis for *Dkk1*, *Osx*, *Bax*, *p16*, and *Axin2* was performed using Applied Biosystems Taqman Gene Expression assays, as previously described (Life Technologies)¹⁵. The 2^{-CT} method⁶⁴ was used to quantify the change in expression of the target genes relative to that in a standard control sample. At least three technical replicates of each PCR measurement were performed. Each replicate was first normalized to the housekeeping gene *Gapdh*. Fold change relative to the control sample was then determined.

Radiation-mitigation studies

10- to 12-week-old female C57BL/6 mice were irradiated with 800 cGy TBI, which is lethal at our institution for approximately 50% of C57BL/6 mice by day +30 (LD₅₀/30), using a

Shepherd cesium-137 irradiator. Mice were subsequently treated with subcutaneous injections of 10 µg recombinant Dkk1 (R&D systems, Minneapolis, MN) or saline, beginning at 24 h post TBI and administered every other day through day +21. PB complete blood counts (CBC) were measured using a Hemavet 950 instrument (Drew Scientific, Miami Lakes, FL). For assessment of long-term effects of Dkk1 treatment, mice were irradiated with 600 cGy TBI and treated with the same dosage and schedule of Dkk1. At 12 weeks after TBI mice were euthanized, and hematopoietic content was analyzed. For anti-Dkk1 radiation-survival studies, 10- to 12-week-old female C57BL/6 mice were irradiated with 750 cGy TBI. Mice then received a subcutaneous injection of 100 µg goat anti-mouse-Dkk1 (AF1765, R&D Systems) or 100 µg goat IgG control (AB-108-C, R&D Systems) in a volume of 100 µl starting at 24 h after irradiation. Injections were administered every other day through day +21. To evaluate whether Dkk1 promoted survival via induction of EGF signaling, the EGF receptor inhibitor erlotinib (Sigma-Aldrich, St. Louis, MO) was administered to irradiated mice. Mice were irradiated with 500 cGy and treated subcutaneously with 10 µg Dkk1 or saline every other day, with or without 10 µg per g body weight erlotinib daily by oral gavage. Erlotinib treatment began 1 d before irradiation and continued through day +10 or +14.

Isolation of BM hematopoietic stem cells (HSCs)

BM HSCs were collected from mice as previously described^{18,19}. Briefly, BM cells were first treated with RBC lysis buffer (Sigma-Aldrich, St. Louis, MO), and lineage-committed cells were removed using a lineage-depletion column (Miltenyi Biotec, Auburn, CA). Lin⁻ cells were stained with allophycocyanin (APC)- and Cy7-conjugated anti-Sca-1, phycoerythrin (PE)-conjugated anti-c-kit, APC-conjugated anti-CD-34, and V450 lineage cocktail (Becton Dickinson (BD), San Jose, CA) or with isotype controls. Sterile cell sorting was conducted on a BD FACS ARIA cytometer. Purified c-kit⁺Sca-1⁺lin⁻ (KSL) and CD34⁻c-kit⁺Sca-1⁺lin⁻ (CD34⁻KSL) cells were collected into IMDM (Life Technologies) + 10% FBS + 1% penicillin–streptomycin.

CFC assays, HSC cultures, and competitive-repopulation assays

CFC assays (colony-forming unit–granulocyte monocyte (CFU-GM), burst-forming unit–erythroid (BFU-E), and colony-forming unit–mix (CFU-GEMM)) were performed as we have previously described^{15,18}. For *in vitro* cultures, BM CD34⁻KSL, KSL, and lineage-negative cells were cultured in TSF medium (IMDM, 10% FBS, 1% penicillin–streptomycin, 20 ng/ml thrombopoietin (TPO), 125 ng/ml SCF, 50 ng/ml Flt-3 ligand). SCF, Flt-3 ligand, and TPO were purchased from R&D Systems. For competitive-repopulation assays, BM cells were isolated from donor 10- to 12-week-old female CD45.2⁺ mice. Recipient 10-week-old female CD45.1⁺ B6.SJL mice were irradiated with 950 cGy TBI using a cesium-137 irradiator, and donor BM cells were then administered via tail vein injection along with a competing dose of 2×10^5 non-irradiated host BM cells. Multilineage donor hematopoietic cell engraftment was measured in the PB by flow cytometry, as previously described^{15,18,19}.

***In vitro* ROS and senescence assays**

BM CD34⁺KSL, KSL, and lineage-negative cells were isolated by FACS and irradiated with 300 cGy. Cells were cultured in the following conditions: TSF medium, TSF + 500 ng/ml mouse Dkk1 (R&D Systems), TSF + 100 ng/ml mouse Wnt3a (R&D Systems), or TSF + 200 ng/ml mouse Wnt5a (R&D Systems). After 24 h, cells were analyzed by flow cytometry for mitochondrial ROS, active caspase-3–caspase-7, p38 phosphorylation, SA- β -gal, and intracellular glutathione (GSH). MitoSOX Red detection reagent (Life Technologies) was incubated at 5 μ M in Hank's buffered saline solution (HBSS,) for 30 min at 37 °C. Cells were then rinsed and analyzed on a BD FACS CANTO-2 instrument to detect mitochondrial ROS. A Vybrant FAM Caspase-3 and Caspase-7 Assay Kit (Life Technologies) was used to detect active caspase-3 and caspase-7. Analysis of phospho-p38 was performed on cells that were fixed in 2% paraformaldehyde for 10 min and permeabilized in ice-cold methanol for 30 min. Cells were then stained with primary antibodies against phospho-p38 (50-191-935, Cell Signaling Technologies) followed by staining with appropriate secondary antibodies. SA- β -gal activity in live cells was quantified using the Enzo Life Sciences SA- β -gal kit (Enzo, Farmingdale NY). Intracellular GSH was detected using the Thiol Green intracellular GSH kit (Abcam) per the manufacturer's instructions.

Axin2–LacZ reporter analysis

Analysis of Wnt pathway activation was determined using the Axin2–LacZ reporter mouse (Jackson Laboratory), as previously described²⁷. Mice were irradiated with 500 cGy and treated with or without 10 μ g Dkk1 subcutaneously. At 24 h, the mice were euthanized and analyzed for LacZ signal by flow cytometry in BM CD34⁺KSL, KSL, and lineage-negative cells using the LacZ intracellular detection kit (Abcam). After staining for cell surface markers, cells were incubated for 1 min at 37 °C with the LacZ fluorescent substrate reagent and then placed immediately on ice before FACS analysis.

Analysis of β -catenin-deficient hematopoietic stem and progenitor cells

Loss-of-function β -catenin studies were performed by infecting KSL cells that were isolated from *Ctnnb1*^{FL/FL} mice²⁹ with viral Cre–GFP (FCT072 LV-CMV-Cre-GFP, Kerafast, Inc., Boston, MA) or sham GFP vectors (FCT003 LV-CMV-GFP, Kerafast). BM KSL cells were isolated by FACS from *Ctnnb1*^{FL/FL} mice and then plated overnight in X-Vivo 15 medium (Lonza) containing 125 ng/ml SCF, 100 ng/ml Flt-3 ligand, and 100 ng/ml TPO. After 24 h, the cells were transduced by the addition of 5 ng/ml polybrene (Fisher Scientific) and lentiviral particles for a multiplicity of infection (MOI) of 25. At 72 h, the cultures were sorted for GFP⁺ cells and *Ctnnb1* knockdown was confirmed by RT–PCR. The sorted cells were then irradiated with 300 cGy and placed in culture with TSF or TSF + 500 ng/ml Dkk1. For ROS studies, cultures were collected at 1 h and analyzed as previously described. For CFCs, cultures were collected 4 d after irradiation and plated in CFC assays.

ROS scavenger assay

For ROS inhibition with *N*-acetylcysteine (NAC, Life Technologies), NAC was added to cultures at a concentration of 10 μ M for 12 h before irradiation. KSL cells were irradiated

with 300 cGy and then cultured for 24 h with or without 500 ng/ml Dkk1 before plating in CFC assays.

Statistical analysis

GraphPad Prism 6.0 was used for all statistical analyses. All data were checked for normal distribution and similar variance between groups. Data were derived from multiple independent experiments from distinct mice or cell culture plates. Sample sizes for *in vitro* studies were chosen based on observed effect sizes and standard errors from prior studies. For all animal studies, a power test was used to determine the sample size needed to observe a twofold difference in means between groups with 0.8 power using a two-tailed Student's *t*-test. All animal studies were performed using sex- and age-matched animals, with wild-type littermates as controls. Animal studies were performed without blinding of the investigator, and no animals were excluded from the analyses. All comparisons performed were done using a two-tailed Student's *t*-test, unless otherwise indicated in the figure legends. Values are reported as mean $\pm$ s.e.m., unless stated otherwise.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank S. Vainio (University of Oulu) for the *Dkk1^{FL/+}* mice. This work was supported, in part, by NHLBI grant HL-086998-05 (J.P.C.), NIAID grant AI-067798-10 (J.P.C.), the California Institute for Regenerative Medicine Leadership Award LA1-08014 (J.P.C.) and the NIAID Centers for Medical Countermeasures Against Radiation (CMCR) Pilot Award 2U19AI067773-11 (H.A.H.).

References

1. Ding L, Saunders TL, Enikolopov G, Morrison SJ. Endothelial and perivascular cells maintain hematopoietic stem cells. *Nature*. 2012; 481:457–462. [PubMed: 22281595]
2. Ding L, Morrison SJ. Hematopoietic stem cells and early lymphoid progenitors occupy distinct bone marrow niches. *Nature*. 2013; 495:231–235. [PubMed: 23434755]
3. Greenbaum A, et al. CXCL12 in early mesenchymal progenitors is required for hematopoietic stem cell maintenance. *Nature*. 2013; 495:227–230. [PubMed: 23434756]
4. Méndez-Ferrer S, et al. Mesenchymal and hematopoietic stem cells form a unique bone marrow niche. *Nature*. 2010; 466:829–834. [PubMed: 20703299]
5. Pinho S, et al. PDGFR- α and CD51 mark human nestin⁺ sphere-forming mesenchymal stem cells capable of hematopoietic progenitor cell expansion. *J Exp Med*. 2013; 210:1351–1367. [PubMed: 23776077]
6. Mizoguchi T, et al. Osterix marks distinct waves of primitive and definitive stromal progenitors during bone marrow development. *Dev Cell*. 2014; 29:340–349. [PubMed: 24823377]
7. Kunisaki Y, et al. Arteriolar niches maintain hematopoietic stem cell quiescence. *Nature*. 2013; 502:637–643. [PubMed: 24107994]
8. Calvi LM, et al. Osteoblastic cells regulate the hematopoietic stem cell niche. *Nature*. 2003; 425:841–846. [PubMed: 14574413]
9. Zhang J, et al. Identification of the hematopoietic stem cell niche and control of the niche size. *Nature*. 2003; 425:836–841. [PubMed: 14574412]
10. Visnjic D, et al. Hematopoiesis is severely altered in mice with an induced osteoblast deficiency. *Blood*. 2004; 103:3258–3264. [PubMed: 14726388]

11. Kiel MJ, Acar M, Radice GL, Morrison SJ. Hematopoietic stem cells do not depend on N-cadherin to regulate their maintenance. *Cell Stem Cell*. 2009; 4:170–179. [PubMed: 19119091]
12. Park D, et al. Endogenous bone marrow MSCs are dynamic, fate-restricted participants in bone maintenance and regeneration. *Cell Stem Cell*. 2012; 10:259–272. [PubMed: 22385654]
13. Maes C, et al. Osteoblast precursors, but not mature osteoblasts, move into developing and fractured bones along with invading blood vessels. *Dev Cell*. 2010; 19:329–344. [PubMed: 20708594]
14. Raaijmakers MH, et al. Bone progenitor dysfunction induces myelodysplasia and secondary leukemia. *Nature*. 2010; 464:852–857. [PubMed: 20305640]
15. Doan PL, et al. Tie2⁺ bone marrow endothelial cells regulate hematopoietic stem cell regeneration following radiation injury. *Stem Cells*. 2013; 31:327–337. [PubMed: 23132593]
16. Himburg HA, et al. Pleiotrophin regulates the retention and self-renewal of hematopoietic stem cells in the bone marrow vascular niche. *Cell Rep*. 2012; 2:964–975. [PubMed: 23084748]
17. Hooper AT, et al. Engraftment and reconstitution of hematopoiesis is dependent on VEGFR2-mediated regeneration of sinusoidal endothelial cells. *Cell Stem Cell*. 2009; 4:263–274. [PubMed: 19265665]
18. Himburg HA, et al. Pleiotrophin regulates the expansion and regeneration of hematopoietic stem cells. *Nat Med*. 2010; 16:475–482. [PubMed: 20305662]
19. Doan PL, et al. Epidermal growth factor regulates hematopoietic regeneration after radiation injury. *Nat Med*. 2013; 19:295–304. [PubMed: 23377280]
20. Kirsch DG, et al. p53 controls radiation-induced gastrointestinal syndrome in mice independent of apoptosis. *Science*. 2010; 327:593–596. [PubMed: 20019247]
21. Zhang C, et al. Inhibition of Wnt signaling by the osteoblast-specific transcription factor osterix. *Proc Natl Acad Sci USA*. 2008; 105:6936–6941. [PubMed: 18458345]
22. Rodda SJ, McMahon AP. Distinct roles for Hedgehog and canonical Wnt signaling in specification, differentiation and maintenance of osteoblast progenitors. *Development*. 2006; 133:3231–3244. [PubMed: 16854976]
23. Cho J, et al. Purinergic P2Y receptor modulates stress-induced hematopoietic stem and progenitor cell senescence. *J Clin Invest*. 2014; 124:3159–3171. [PubMed: 24937426]
24. Freund A, Patil CK, Campisi J. p38 MAPK is a novel DNA damage response-independent regulator of the senescence-associated secretory phenotype. *EMBO J*. 2011; 30:1536–1548. [PubMed: 21399611]
25. Chang J, et al. Clearance of senescent cells by ABT263 rejuvenates aged hematopoietic stem cells in mice. *Nat Med*. 2016; 22:78–83. [PubMed: 26657143]
26. Mao B, et al. Kremen proteins are dickkopf receptors that regulate Wnt– β -catenin signaling. *Nature*. 2002; 417:664–667. [PubMed: 12050670]
27. Boudousquie C, et al. Differences in the transduction of canonical Wnt signals demarcate effector and memory CD8 T cells with distinct recall proliferation capacity. *J Immunol*. 2014; 193:2784–2791. [PubMed: 25127860]
28. Luis TC, et al. Canonical Wnt signaling regulates hematopoiesis in a dosage-dependent fashion. *Cell Stem Cell*. 2011; 9:345–356. [PubMed: 21982234]
29. Brault V, et al. Inactivation of the β -catenin gene by *Wnt1-Cre*-mediated deletion results in dramatic brain malformation and failure of craniofacial development. *Development*. 2001; 128:1253–1264. [PubMed: 11262227]
30. Poulos MG, et al. Endothelial Jagged-1 is necessary for homeostatic and regenerative hematopoiesis. *Cell Rep*. 2013; 4:1022–1034. [PubMed: 24012753]
31. Salter AB, et al. Endothelial progenitor cell infusion induces hematopoietic stem cell reconstitution *in vivo*. *Blood*. 2009; 113:2104–2107. [PubMed: 19141867]
32. Olson TS, et al. Megakaryocytes promote murine osteoblastic HSC niche expansion and stem cell engraftment after radioablative conditioning. *Blood*. 2013; 121:5238–5249. [PubMed: 23667055]
33. Dominici M, et al. Restoration and reversible expansion of the osteoblastic hematopoietic stem cell niche after marrow radioablation. *Blood*. 2009; 114:2333–2343. [PubMed: 19433859]

34. Na Nakorn T, Traver D, Weissman IL, Akashi K. Myeloerythroid-restricted progenitors are sufficient to confer radioprotection and provide the majority of day 8 CFU-S. *J Clin Invest.* 2002; 109:1579–1585. [PubMed: 12070305]
35. Fleming HE, et al. Wnt signaling in the niche enforces hematopoietic stem cell quiescence and is necessary to preserve self-renewal *in vivo*. *Cell Stem Cell.* 2008; 2:274–283. [PubMed: 18371452]
36. Varnum-Finney B, et al. Notch2 governs the rate of generation of mouse long- and short-term repopulating stem cells. *J Clin Invest.* 2011; 121:1207–1216. [PubMed: 21285514]
37. Shao L, et al. Total body irradiation causes long-term mouse BM injury via induction of HSC premature senescence in an Ink4a- and Arf-independent manner. *Blood.* 2014; 123:3105–3115. [PubMed: 24622326]
38. Ito K, et al. Regulation of oxidative stress by ATM is required for self-renewal of hematopoietic stem cells. *Nature.* 2004; 431:997–1002. [PubMed: 15496926]
39. Tothova Z, et al. FoxOs are critical mediators of hematopoietic stem cell resistance to physiologic oxidative stress. *Cell.* 2007; 128:325–339. [PubMed: 17254970]
40. Taniguchi Ishikawa E, et al. Connexin-43 prevents hematopoietic stem cell senescence through transfer of reactive oxygen species to bone marrow stromal cells. *Proc Natl Acad Sci USA.* 2012; 109:9071–9076. [PubMed: 22611193]
41. Mantel CR, et al. Enhancing hematopoietic stem cell transplantation efficacy by mitigating oxygen shock. *Cell.* 2015; 161:1553–1565. [PubMed: 26073944]
42. Spencer JA, et al. Direct measurement of local oxygen concentration in the bone marrow of live animals. *Nature.* 2014; 508:269–273. [PubMed: 24590072]
43. Luis TC, Ichii M, Brugman MH, Kincade P, Staal FJ. Wnt signaling strength regulates normal hematopoiesis, and its de-regulation is involved in leukemia development. *Leukemia.* 2012; 26:414–421. [PubMed: 22173215]
44. Kim IG, et al. Disturbance of DKK1 level is partly involved in survival of lung cancer cells via regulation of ROMO1 and γ -radiation sensitivity. *Biochem Biophys Res Commun.* 2014; 443:49–55. [PubMed: 24269823]
45. Yoon JC, et al. Wnt signaling regulates mitochondrial physiology and insulin sensitivity. *Genes Dev.* 2010; 24:1507–1518. [PubMed: 20634317]
46. Huang J, et al. Pivotal role for glycogen synthase kinase-3 in hematopoietic stem cell homeostasis in mice. *J Clin Invest.* 2009; 119:3519–3529. [PubMed: 19959876]
47. Li W, et al. Apc regulates the function of hematopoietic stem cells largely through β -catenin-dependent mechanisms. *Blood.* 2013; 121:4063–4072. [PubMed: 23547052]
48. Huang J, Nguyen-McCarty M, Hexner EO, Danet-Desnoyers G, Klein PS. Maintenance of hematopoietic stem cells through regulation of Wnt and mTOR pathways. *Nat Med.* 2012; 18:1778–1785. [PubMed: 23142822]
49. Lento W, et al. Loss of β -catenin triggers oxidative stress and impairs hematopoietic regeneration. *Genes Dev.* 2014; 28:995–1004. [PubMed: 24788518]
50. Lin SL, et al. Macrophage Wnt7b is critical for kidney repair and regeneration. *Proc Natl Acad Sci USA.* 2010; 107:4194–4199. [PubMed: 20160075]
51. Foley AC, Mercola M. Heart induction by Wnt antagonists depends on the homeodomain transcription factor Hex. *Genes Dev.* 2005; 19:387–396. [PubMed: 15687261]
52. Min JK, et al. The WNT antagonist dickkopf-2 promotes angiogenesis in rodent and human endothelial cells. *J Clin Invest.* 2011; 121:1882–1893. [PubMed: 21540552]
53. Famili F, et al. High levels of canonical Wnt signaling lead to loss of stemness and increased differentiation in hematopoietic stem cells. *Stem Cell Rep.* 2016; 6:652–659.
54. Kirstetter P, Anderson K, Porse BT, Jacobsen SE, Nerlov C. Activation of the canonical Wnt pathway leads to loss of hematopoietic stem cell repopulation and multilineage differentiation block. *Nat Immunol.* 2006; 7:1048–1056. [PubMed: 16951689]
55. Scheller M, et al. Hematopoietic stem cell and multilineage defects generated by constitutive β -catenin activation. *Nat Immunol.* 2006; 7:1037–1047. [PubMed: 16951686]
56. Malhotra S, et al. Contrasting responses of lymphoid progenitors to canonical and noncanonical Wnt signals. *J Immunol.* 2008; 181:3955–3964. [PubMed: 18768850]

57. Florian MC, et al. A canonical to noncanonical Wnt signaling switch in hematopoietic stem cell aging. *Nature*. 2013; 503:392–396. [PubMed: 24141946]
58. Mendelson A, Frenette PS. Hematopoietic stem cell niche maintenance during homeostasis and regeneration. *Nat Med*. 2014; 20:833–846. [PubMed: 25100529]
59. Morrison SJ, Scadden DT. The bone marrow niche for hematopoietic stem cells. *Nature*. 2014; 505:327–334. [PubMed: 24429631]
60. Itkin T, et al. Distinct bone marrow blood vessels differentially regulate hematopoiesis. *Nature*. 2016; 532:323–328. [PubMed: 27074509]
61. Strecker S, Fu Y, Liu Y, Maye P. Generation and characterization of osterix–Cherry reporter mice. *Genesis*. 2013; 51:246–258. [PubMed: 23180553]
62. Lustig B, et al. Negative feedback loop of Wnt signaling through upregulation of conductin (Axin2) in colorectal and liver tumors. *Mol Cell Biol*. 2002; 22:1184–1193. [PubMed: 11809809]
63. Pietilä I, et al. Secreted Wnt antagonist dickkopf-1 controls kidney papilla development coordinated by Wnt7b signaling. *Dev Biol*. 2011; 353:50–60. [PubMed: 21354128]
64. Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time quantitative PCR and the 2[−]CT method. *Methods*. 2001; 25:402–408. [PubMed: 11846609]

EDITORIAL SUMMARY**AOP**

The Wnt pathway inhibitor Dkk1, which is produced by bone marrow osteolineage cells, promotes hematopoietic recovery after radiation injury by both direct effects on hematopoietic cells and indirect effects on bone marrow endothelial cells.

Figure 1.

Deletion of *Bak* and *Bax* in *Osx*⁺ BM cells radioprotects hematopoietic stem and progenitor cells. (a) Mean percentages of *Osx*⁺ cells, as measured by flow cytometry, within *Osx*-labeled BM cells (*Osx*-cherry⁺) and *Osx*-unlabeled cells (*Osx*-cherry⁻) from 5-d-old (*n* = 4 mice/group) and 8-week-old (*n* = 8 mice/group) *Osx*-Cherry reporter mice. **P* < 0.001, ***P* < 0.01. (b) Left, representative FACS plots showing the percentage of *Osx*⁺ cells in CD45⁻ BM cells in *Osx-Cre;Bak1^{-/-};Bax^{FL/+}* (BAX FL^{+/+}) and *Osx-Cre;Bak1^{-/-};Bax^{FL/-}* (BAX FL^{-/-}) mice at day +3 following 500 cGy TBI. Right, the mean percentage of *Osx*⁺CD45⁻ BM cells (*n* = 4 mice/group). **P* = 0.004. (c) Representative images of H&E-stained femurs from BAX FL^{+/+} (left) and BAX FL^{-/-} (middle) mice at day +7 following 500 cGy TBI (40×; scale bars, 100 μm) and scatter plot of BM cell counts for mice in each group (*n* = 11 mice/group) (right). Horizontal lines represent means. **P* = 0.009. (d) Mean numbers of BM KSL cells (**P* = 0.03) (left) and SLAMF6⁺KSL cells (**P* = 0.04) (right) in BAX FL^{+/+} and BAX FL^{-/-} mice at day +7 following 500 cGy TBI (*n* = 11 mice/group). (e) Mean numbers of BM CFCs at day +7 (*n* = 21 assays/group). CFU-GM, colony-forming unit–granulocyte monocyte; BFU-E, burst-forming unit–erythroid; CFU-GEMM, colony-forming unit–mix. **P* < 0.0001. (f) Mean PB WBC (**P* = 0.01) (left), neutrophil (**P* = 0.02) (middle) and lymphocyte (**P* = 0.02) (right) counts at day +7 (FL^{+/+}, *n* = 14 mice; FL^{-/-}, *n* = 13 mice). (g) Donor (CD45.2⁺)

cell engraftment over time in recipient CD45.1⁺ mice that were transplanted with 1×10^6 BM cells from BAX FL^{+/+} or BAX FL^{-/-} mice and 1×10^5 competing, non-irradiated host BM cells. **P* = 0.03, **P* = 0.03, **P* = 0.02 and **P* = 0.04 for 4, 8, 12 and 20 weeks, respectively. Myeloid (Mac1/Gr1; **P* = 0.04), B cell (B220; **P* = 0.04) and T cell (CD3; **P* = 0.03) engraftment levels at 20 weeks are shown (4, 8 and 12 weeks: FL^{+/+}, *n* = 14 mice; FL^{-/-}, *n* = 15 mice; 20 weeks: FL^{+/+}, *n* = 9 mice; FL^{-/-}, *n* = 13 mice). (h) Mean levels of donor CD45.2⁺ cell and lineage engraftment in secondary recipient CD45.1⁺ mice at 20 weeks following competitive transplantation with BM cells from primary mice (FL^{+/+}, *n* = 7 mice; FL^{-/-}, *n* = 9 mice). CD45.2⁺, **P* = 0.04; B220⁺, **P* = 0.03; CD3⁺, **P* = 0.004. (i) Mean Dkk1 levels in the BM of non-irradiated or irradiated (500 cGy) BAX FL^{+/+} and BAX FL^{-/-} mice. **P* < 0.001 and ***P* < 0.001 (NI FL^{+/+}, *n* = 5 mice; NI FL^{-/-}, *n* = 9 mice; IRR FL^{+/+}, *n* = 8 mice; IRR FL^{-/-}, *n* = 7 mice). (j) Mean expression of Dkk1 in BM osteolineage cells in BAX FL^{-/-} mice relative to that in BAX FL^{+/+} mice (*n* = 6 mice/group). **P* = 0.02. (k) *Dkk1* expression in VE-cadherin⁺ BM ECs and Osx⁺ BM cells in mice before (0 cGy; VE-cadherin⁺, *n* = 3 mice; Osx⁺, *n* = 4 mice) and 48 h after 500 cGy TBI (*n* = 6 mice/group). *Dkk1* expression levels in all groups were normalized to that in VE-cadherin⁺ BM ECs at 48 h. **P* = 0.03 and ***P* = 0.003. Throughout, *P* values were derived by using a two-tailed Student's *t*-test.

Figure 2.

Dkk1 promotes hematopoietic regeneration *in vitro* and *in vivo*. **(a)** Mean numbers of CFCs recovered from 7-d cultures of irradiated (300 cGy) BM KSL cells that were treated with TPO, SCF and Flt-3 ligand (TSF)-containing medium (medium), with or without 500 ng/ml Dkk1 ($n = 3$ assays/group). $*P = 0.003$. **(b)** Left, mean percentage of donor CD45.2⁺ cell engraftment levels in recipient CD45.1⁺ mice over time following competitive transplantation with the progeny of 1×10^4 irradiated (300 cGy) BM KSL cells that were cultured with or without 500 ng/ml Dkk1 for 7 d. $*P = 0.01$, $*P = 0.03$, $*P = 0.03$, $*P = 0.01$ for 4, 8, 12 and 16 weeks, respectively. Right, mean percentages of donor myeloid ($*P = 0.02$), B cell ($*P = 0.01$) and T cell ($*P = 0.01$) engraftment at 16 weeks in recipient mice ($n = 17$ mice/group). **(c)** Dkk1 concentrations in the BM of mice at baseline, following 500 cGy treatment, and at 1 h and 24 h following administration of 10 µg Dkk1 administration (BL6 non-irradiated (NI), $n = 5$ mice; BL6, $n = 5$ mice; Dkk1 (1 h), $n = 6$ mice; Dkk1 (24 h), $n = 5$ mice). $*P < 0.001$, $**P < 0.001$. **(d)** Scatter plots of PB WBCs (left), neutrophils (middle) and platelet (right) counts at day +21 from mice irradiated with 800 cGy and then treated with saline or Dkk1 (saline, $n = 5$ mice; Dkk1, $n = 12$ mice). $*P = 0.008$, $*P = 0.02$, $*P = 0.01$; by Mann–Whitney U test. **(e)** Left, representative images of H&E-stained femurs from saline-treated (left) and Dkk1-treated (middle) mice at day +21 following treatment

with 800 cGy (40 $\times$; scale bars, 100 μ m). Right, mean BM cell counts ($n = 5$ mice/group). $*P = 0.02$. **(f)** Left, representative FACS plots showing the percentage of BM KSL cells in saline-treated and Dkk1-treated mice at day +21 following treatment with 800 cGy TBI. Middle and right, the mean percentage of KSL cells ($*P = 0.02$) (middle) and KSL cell numbers ($*P = 0.008$) (right) ($n = 5$ mice/group). P values determined by Mann–Whitney U test. **(g)** Mean numbers of BM CFCs at day +21 following treatment with 800 cGy TBI ($n = 6$ assays/group). $*P = 0.04$. **(h)** Survival of saline-treated (4/15) or Dkk1-treated (13/14) mice following treatment with 800 cGy TBI (27% versus 93%; $P = 0.0004$ by log-rank test). **(i)** Mean numbers of CFCs in C57BL/6 mice at 12 weeks after 600 cGy TBI and treatment with 10 μ g Dkk1 or saline, subcutaneously, every other day through day +21 ($n = 5$ assays/group). $*P = 0.001$. Throughout, unless otherwise noted, P values were derived by using a two-tailed Student's t -test.

Figure 3.

Inhibition or deficiency of Dkk1 in *Osx*-labeled BM cells suppresses hematopoietic regeneration. (a) PB WBCs, neutrophils and lymphocytes in mice treated with anti-Dkk1 or control IgG at day +10 following 500 cGy TBI. $*P = 0.0004$, $*P < 0.0001$, $*P = 0.002$, respectively ($n = 10$ mice/group). (b) Left, representative images of H&E-stained femurs stained with IgG (top) or anti-Dkk1 (bottom) at day +10 (40 $\times$; scale bars, 50 μ m). Right, BM cell counts at day +10 ($n = 5$ mice/group). $*P < 0.001$. (c) Mean KSL cell numbers at day +10 ($n = 5$ mice/group). $*P = 0.007$. (d) Survival analysis of mice irradiated with 750 cGy TBI and treated with anti-Dkk1 or IgG ($n = 12$ mice/group). $P = 0.006$. (e) The hematopoietic profiles (PB WBCs ($*P < 0.001$), neutrophils ($*P = 0.001$) and lymphocytes ($*P = 0.007$)) of *Osx-Cre;Dkk1^{FL/+}* (*Dkk1^{FL/+}*) and *Osx-Cre;Dkk1^{+/+}* (*Dkk1^{+/+}*) mice at day 10 after irradiation with 500 cGy TBI and of irradiated *Dkk1^{FL/+}* mice that were treated every other day with Dkk1 through day +10 (*Dkk1^{FL/+}* + Dkk1) and evaluated at day +10 (*Dkk1^{+/+}*, $n = 9$ mice; *Dkk1^{FL/+}*, $n = 10$ mice; *Dkk1^{FL/+}* + Dkk1, $n = 10$ mice). (f) BM cell counts at day +10 (*Dkk1^{+/+}*, $n = 9$ mice; *Dkk1^{FL/+}*, $n = 10$ mice; *Dkk1^{FL/+}* + Dkk1, $n = 10$ mice). $*P < 0.001$. (g) BM KSL cell numbers at day +10 (*Dkk1^{+/+}*, $n = 7$ mice; *Dkk1^{FL/+}*, $n = 10$ mice; *Dkk1^{FL/+}* + Dkk1, $n = 10$ mice). $*P = 0.002$, $**P = 0.03$. (h) Mean numbers of BM CFCs at day +10 (*Dkk1^{+/+}*, $n = 6$ assays; *Dkk1^{FL/+}*, $n = 6$ assays; *Dkk1^{FL/+}* + Dkk1, $n = 3$ assays). $*P = 0.008$, $**P = 0.005$. (i) Mean concentrations of PB WBCs ($*P = 0.001$), neutrophils ($*P < 0.001$) and lymphocytes ($*P < 0.001$) in *Osx-Cre;Dkk1^{+/+}* mice (*Dkk1^{+/+}*) and 8-week-*Osx-Cre;Dkk1^{FL/+}* mice (8-week-*Dkk1^{FL/+}*) at day +10 following 500 cGy TBI (*Dkk1^{+/+}*, $n = 11$ mice; 8-week-*Dkk1^{FL/+}*, $n = 7$ mice). (j) Left, representative images of H&E-stained femurs from *Dkk1^{+/+}* (top) and 8-week-*Dkk1^{FL/+}* (bottom) mice at day +10 (40 $\times$; scale bars, 50 μ m). Right, mean BM cell counts per femur (*Dkk1^{+/+}*, $n = 11$ mice; 8-week-*Dkk1^{FL/+}*, $n = 7$ mice). $*P < 0.001$. (k) Mean BM KSL cells at day +10 (*Dkk1^{+/+}*, $n = 7$ mice; 8-week-*Dkk1^{FL/+}*, $n = 6$ mice). $*P = 0.01$. (l) Mean BM CFCs at day +10 (*Dkk1^{+/+}*, $n = 6$ assays; 8-week-*Dkk1^{FL/+}*, $n = 3$ assays). $*P = 0.005$. Throughout, P values were derived by using a two-tailed Student's t -test.

Figure 4.

Dkk1 suppresses HSC senescence following irradiation and induces EGF secretion by BM ECs. **(a)** The percentage of MitoSOX⁺ cells in non-irradiated CD34⁺ KSL cells (NI) and at 24 h after 300 cGy TBI without treatment (medium) or with treatment with 500 ng/ml Dkk1, 100 ng/ml Wnt3a or 200 ng/ml Wnt5a; treatment was initiated 15 min before irradiation ($n = 4$ cell cultures/group). Medium included TPO, SCF and Flt-3 ligand. * $P < 0.001$, ** $P < 0.001$. **(b)** The percentage of phospho-p38-positive cells in CD34⁺ KSL cells at 24 h after 300 cGy irradiation (NI, $n = 8$ cultures; medium, $n = 7$ cultures; Dkk1, $n = 8$ cultures; Wnt3a, $n = 8$ cultures; Wnt5a, $n = 7$ cultures). * $P < 0.001$, ** $P < 0.001$. **(c)** The percentage of SA-β-gal⁺ CD34⁺ KSL cells at 24 h after 300 cGy irradiation ($n = 4$ cultures/group). * $P < 0.001$, ** $P < 0.001$. **(d)** Left, *p16* gene expression in BM CD34⁺ KSL cells at 24 h after 300 cGy irradiation in the indicated treatment groups. * $P = 0.005$, ** $P = 0.02$ ($n = 4$ cultures/group). Right, *p16* expression in BM KSL cells at 24 and 48 h after 500 cGy TBI in mice treated without (saline) or with 10 μg Dkk1 ($n = 4$ cultures/group). 24 h, * $P < 0.001$, ** $P < 0.001$; 48 h, * $P = 0.02$, ** $P = 0.02$. **(e)** Mean percentages of SA-β-gal⁺ BM KSL cells in non-irradiated mice and in mice at 12 weeks following 600 cGy TBI and treated with

saline or 10 µg Dkk1 every other day through day +21 (NI, $n = 4$ mice; saline, $n = 7$ mice; Dkk1, $n = 7$ mice). $*P = 0.04$, $**P = 0.02$. (f) Left, representative histograms of Axin2–LacZ⁺ cells within BM CD34⁺ KSL cells, KSL cells and lin⁺ cells in mice at 24 h after 500 cGy TBI in the treatment groups shown. Right, the mean percentage of Axin2⁺ cells in each cell population at 24 h following 500 cGy TBI with and without Dkk1 treatment ($n = 4$ mice/group). CD34⁺ KSL cells, $*P = 0.009$, $**P = 0.01$; KSL cells, $*P < 0.001$, $**P = 0.004$; lin⁺ cells, $*P < 0.001$, $**P = 0.002$. (g) Left, mean levels of PTN in cultures of BM ECs ($n = 5$ cultures/group). Middle, mean levels of EGF in BM ECs in response to 500 ng/ml Dkk1 treatment ($*P < 0.001$; BMEC, $n = 5$ cultures; BMEC + Dkk1, $n = 10$ cultures). Right, mean levels of EGF in C57BL/6 mice over time following 500 cGy TBI and treatment with saline or 10 µg Dkk1 every other day through day +14 ($n = 5$ mice/group) (day +10, $*P = 0.03$; day +14, $*P = 0.02$). (h) Mean values for PB WBCs, neutrophils, BM cell counts and BM KSL cells at days +10 and +14 following 500 cGy TBI and treatment with saline or 10 µg Dkk1, with and without 10 µg/g erlotinib ($n = 5$ mice/group). WBCs: day +10, $*P < 0.001$; day +14, $*P = 0.002$; neutrophils: day +10, $*P < 0.001$; day +14, $*P = 0.01$; BM cell counts: day +14, $*P = 0.02$; BM KSL cells: day +14, $*P = 0.01$. Throughout, P values were derived by using a two-tailed Student's t -test.