CORRESPONDENCE

Cardiovascular Comorbidity in Inflammatory Rheumatological Conditions

by Prof. Dr. med. Jürgen Braun, Prof. Dr. med. Klaus Krüger, Prof. Dr. med. Bernhard Manger, Prof. Dr. med. Matthias Schneider, Prof. Dr. med. Christopf Specker, and Prof. Dr. med. Hans Joachim Trappe in issue 12/2017

Rheumatic Heart Disease Still Relevant

We strongly object to one point of the comprehensive review by Braun et al. (1): contrary to the statement made by the authors (on page 199 of the article), rheumatic heart disease after recurrent rheumatic fever is certainly the most common cardiac complication in rheumatic diseases worldwide. Reliable sources estimate a prevalence of 33 million cases, of which a significant proportion are children aged 5 to 14 years (2.4 million) (2–4). About 300 000 new cases, and 233 000 to 294 000 deaths, occur each year due to the disease.

The high levels of morbidity and mortality are certainly due to the fact that many affected persons in less developed and newly industrialized countries have very limited diagnostic and therapeutic options. Access to an often life-saving heart valve surgery is not available to most of these people. However, the indigenous populations in the industrialized countries of Australia (Aboriginal) and New Zealand (Maori) have the highest incidence rates worldwide (3.5 per 1000 inhabitants) (3). These remarks should raise awareness that first, even in wealthy countries, rheumatic heart disease is by no means irrelevant, and second, in Europe (including Germany), the incidence of this disease can be expected to increase due to the immigration of many people from Africa and Asia.

DOI: 10.3238/arztebl.2017.0559a

REFERENCES

1. Braun J, Krüger K, Manger B, Schneider M, Specker C, Trappe HJ: Cardiovascular comorbidity in inflammatory rheumatological conditions. *Dtsch Arztebl Int* 2017; 114: 197–203.
2. GBD 2013 Mortality and Causes of Death Collaborators: Global, regional, and national age-sex specific all-cause and cause-specific mortality for 240 causes of death, 1990–2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet* 2015; 385: 117–71.
3. Carapetis JR, Steer AC, Mulholland EK, Weber M: The global burden of group A streptococcal diseases. *Lancet Infect Dis* 2005; 5: 685–92.
4. Nulu S, Bukhman G, Kwan GF: Rheumatic heart disease: the unfinished global agenda. *Cardiology Clinics* 2017; 35: 165–80.

Dr. med. Carsten Krüger, MIH, FRCPCH
St. Franziskus Hospital
Klinik für Kinder und Jugendliche, Ahlen
thea.carsten.krueger@web.de

Conflict of interest statement

The author declares that no conflict of interest exists.

Cardioprotective Effect of NSAIDs

In their review, Braun et al. (1) point out the increased cardiovascular risk for patients with rheumatoid arthritis (RA) due to long-term use of nonsteroidal anti-inflammatory drugs (NSAIDs) and glucocorticoids, referring to a meta-analysis of Roubille et al. (2). However, numerous studies have shown a cardioprotective effect of NSAIDs in RA (3, 4).

Observational studies, such as those on which the meta-analysis is based (2), are of limited value due to the influence of indication and selection biases, among other things. In the context of a high inflammatory background, NSAIDs indeed reduce cardiovascular risk, but not as much as methotrexate (MTX) or tumor necrosis factor (TNF) inhibitors. Patients who do not take NSAIDs are usually those who do not need NSAIDs and have a low cardiovascular risk per se (3).

Numerous publications have shown that NSAID the benefit from NSAID use increases with increasing inflammatory load, as these reduce inflammation. Thus, in patients with systemic inflammation (such as rheumatoid arthritis or systemic lupus erythematosus), risk of an acute myocardial infarction might first occur when NSAR use is discontinued after a prolonged period of time. Finally, studies of patients with ankylosing spondylitis (AS) show a cardiovascular advantage of long-term NSAID use (3); in consequence, NSAIDs are also included in AS therapy according to current guidelines.

The undifferentiated view that NSAIDs are cardiotoxic at all times is popular, but not supported by differential study analysis. As a further consequence of the discussion about the cardiovascular safety of NSAIDs, opioids are being increasingly prescribed for long-term treatment of chronic nonmalignant pain without evidence but with considerable risks.

DOI: 10.3238/arztebl.2017.0559b

REFERENCES

1. Braun J, Krüger K, Manger B, Schneider M, Specker C, Trappe HJ: Cardiovascular comorbidity in inflammatory rheumatological conditions. *Dtsch Arztebl Int* 2017; 114: 197–203.
2. Roubille C, Richter V, Starnino T, et al.: The effects of tumor necrosis factor inhibitors, methotrexate, non-steroidal anti-inflammatory drugs and corticosteroids on cardiovascular events in rheumatoid arthritis, psoriasis and psoriatic arthritis: a systematic review and meta-analysis. *Ann Rheum Dis* 2015; 74: 480–9.
3. Zingler G, Hermann B, Fischer T, Herdegen T: Cardiovascular adverse events by non-steroidal anti-inflammatory drugs: when the benefits outweigh the risks. *Expert Rev Clin Pharmacol* 2016; 9: 1479–92.
4. Goodson NJ, Brookhart AM, Symmons DP, et al.: Non-steroidal anti-inflammatory drug use does not appear to be associated with increased cardiovascular mortality in patients with inflammatory polyarthritis: results from a primary care based inception cohort of patients. *Ann Rheum Dis* 2009; 68: 367–72.

Prof. Dr. med. Thomas Herdegen
Institut für Experimentelle und Klinische Pharmakologie
UKSH, Campus Kiel
t.herdegen@pharmakologie.uni-kiel.de

Dr. nat. habil. Gerhard Zingler
Rostock

Conflict of interest statement

The authors declare that no conflict of interest exists.

Cardiovascular Comorbidities

I read the insightful review of Braun et al. (1) with interest, until I stumbled over a short sentence: “In patients with RA, statins reduce mortality by 20% (e18).” In other words, one in five deaths could be prevented or at least delayed. This number fits into the rising hype around statins since the publication of the 1994 4S study (Scandinavian Simvastatin Survival Study [2]).

Reading the literature cited in the article (e18) (3) reveals that the “20%” (to be precise, 21%) refers to the relative risk reduction (hazard ratio [HR] = 0.79). However, a brief calculation results in an absolute risk reduction of mortality of 2.7% (patients in the control group, 17.4%; patients in the statin group, 14.7%).

This results in a number needed to treat (NNT) of 37 over 4.5 years, corresponding to a number treated needlessly (NTN) of 36.

During a time of constant complaint about increasing polypharmacy, RA patients should not have to be burdened with statins as well—as would be obligated by the value of 20% cited by Braun et al. In my clinical-pharmacological lectures and presentations, I have always urged my listeners to ask and to insist, when stakeholders of all stripes offer a success rate of more than 5%.

From my point of view, it is unfair to offer the unsuspecting reader a reduction in mortality “of 20%”, when in fact only 2.7% can be identified from the literature.

DOI: 10.3238/arztebl.2017.0560a

REFERENCES

1. Braun J, Krüger K, Manger B, Schneider M, Specker C, Trappe HJ: Cardiovascular comorbidity in inflammatory rheumatological conditions. *Dtsch Arztebl Int* 2017; 114: 197–203.
2. Scandinavian Simvastatin Survival Study Group: Randomised trial of cholesterol lowering in 4444 patients with coronary heart disease: the Scandinavian Simvastatin Survival Study (4S). *Lancet* 1994; 344: 1383–9.
3. Schoenfeld SR, Lu L, Rai SK, Seeger JD, Zhang Y, Choi HK: Statin use and mortality in rheumatoid arthritis: a general population—based cohort study. *Ann Rheum Dis* 2016; 75: 1315–20.

Prof. Dr. Frank P. Meyer
Wanzleben-Börde
U_F_Meyer@gmx.de

Conflict of interest statement

The author declares that no conflict of interest exists.

In Reply

We are pleased with the high level of interest in our review (1).

The comments from Krüger highlight that, despite all the evidence for a dramatic decline of rheumatic fever in the developed world, one should not forget that there is a higher prevalence in other regions and that an increase of such problems in Germany due to further refugee flows is possible.

We appreciate the contribution of Herdegen and Zingler who stress the possibility of a cardioprotective effect of nonsteroidal anti-inflammatory drugs (NSAIDs). We explicitly point out in our review that “two independently conducted studies with AS patients [...] showed that not the intake of high NSAID doses, but

of low doses, was associated with increased mortality.” Further, we state that “these results indicate that there may also be beneficial effects of NSAIDs that should be taken into account [...] in patients with chronic inflammatory disease.” Nevertheless, there is also currently an increasing body of evidence that would suggest caution when using NSAIDs (2). That NSAIDs are the first-choice treatment for patients with axial spondyloarthritis, and that the risk of cardiovascular events is indeed significantly lower in young patients not affected by other risk factors, was also stated.

Meyer criticizes our findings: “In patients with RA, statins reduce mortality by 20% (e18).” He prefers a calculation of the absolute risk reduction rather than the relative risk reduction, as the effects are less impressive. This has advantages and disadvantages; both are correct. Overall, we absolutely do not recommend treating all RA patients with statins.

It should be stressed at this point that the evidence for the cardioprotective effect of statins is increasingly stable for the entire population. This applies to both primary and secondary prevention (3). The (cardiovascular) mortality of RA patients is clearly increased—with persistent inflammatory activity as the driving force. Classical risk factors for atherosclerosis, such as hypercholesterolaemia, are associated with this increase (4). These occur twice as frequently in RA patients as in the normal population. Not taking into account known risk factors leads to an increased incidence of myocardial infarction. For management of RA, it is important that statins not only have a favorable effect on lipid metabolism but also are anti-inflammatory (5).

Patients with RA should be regularly evaluated for risk factors and symptoms of cardiovascular disease and be treated adequately and consequently.

DOI: 10.3238/arztebl.2017.0560b

REFERENCES

1. Braun J, Krüger K, Manger B, Schneider M, Specker C, Trappe HJ: Cardiovascular comorbidity in inflammatory rheumatological conditions. *Dtsch Arztebl Int* 2017; 114: 197–203.
2. Bally M, Dendukuri N, Rich B, et al.: Risk of acute myocardial infarction with NSAIDs in real world use: Bayesian meta-analysis of individual patient data. *BMJ* 2017; 357: j1909.
3. Mihaylova B, Emberson J, Blackwell L, et al.: Cholesterol Treatment Trialists' (CTT) Collaborators. The effects of lowering LDL cholesterol with statin therapy in people at low risk of vascular disease: meta-analysis of individual data from 27 randomised trials. *Lancet* 2012; 380: 581–90.
4. Boyer JF, Gourraud PA, Cantagrel A, Davignon JL, Constantin A: Traditional cardiovascular risk factors in rheumatoid arthritis: a meta-analysis. *Joint Bone Spine* 2011; 78: 179–83.
5. McCarey DW, McInnes IB, Madhok R, et al.: Trial of Atorvastatin in Rheumatoid Arthritis (TARA): double-blind, randomised placebo-controlled trial. *Lancet* 2004; 363: 2015–21.

On behalf of the authors

Prof. Dr. med. Jürgen Braun
Rheumazentrum Ruhrgebiet, Herne
j.braun@rheumazentrum-ruhrgebiet.de

Conflict of interest statement

Prof. Braun has received fees for a publication related to the topic.