CORRESPONDENCE

The First Aid and Hospital Treatment of Gunshot and Blast Injuries

by PD Dr. med. Axel Franke, Dr. med. Dan Bieler, Prof. Dr. med. Benedikt Friemert, Prof. Dr. med. Robert Schwab, PD Dr. med. Erwin Kollig, and Dr. med. Christoph Gsgen in issue 14/2017

Andrew Dickinson

The Institute of General Practice and Multi-Disciplinary Healthcare
Universittsklinikum Tbingen, Germany
andykdickinson@icloud.com

Matt Gunning

Netcare Umhlanga Hospital, Durban, South Africa

Structured Approach in Prehospital Care

Franke et al. succinctly and skillfully deal with a broad theme that is very relevant, but more clarity is required with respect to two areas of the prehospital care described.

Firstly, the authors' proposition that, "There is a lack of uniform treatment recommendations in the sense of standards or guidelines for the prehospital [...] treatment of gunshot or blast injuries [...] especially for major events," is misleading. Two British physicians developed a rigorous Major Incident Medical Management System (MIMMS) in 1995 and it is now an established and well-tested civilian course used internationally (1). Its guidelines are applicable and adaptable for all major incidents, which include blast or ballistic injury, or a combination. Treatment options during the first response (triage sieve) in a major incident are application of a tourniquet ('C') and putting the patient in the recovery position ('A').

Secondly, resuscitation guidelines need to be standardized, extremely clear, and easily communicated. The authors agree with that sentiment, but the message as to which algorithm should be used in a pre-hospital context could be clearer. C-ABC was developed by the British military following a stark lesson learnt on the battlefield; catastrophic hemorrhage ('C') kills before an airway ('A') can be secured, and thus the paradigm has shifted to C-ABC during prehospital trauma resuscitation (2, 3). If 'C' (active and massive bleeding) is subjectively ruled out then 'A' is by default the priority but it is important to clearly communicate that C-ABC should be the cognitive flow of the first responder in a trauma situation.

DOI: 10.3238/arztebl.2017.0561a

REFERENCES

1. Franke A, Bieler D, Friemert B, Schwab R, Kollig E, Gsgen C: The first aid and hospital treatment of gunshot and blast injuries. *Dtsch Arztebl Int* 2017; 114: 237–43.
2. Hodgetts TJ, Mackway-Jones K (eds): Major incident medical management and support. The practical approach. Plymouth: BMJ 1995.
3. Hodgetts TJ, Mahoney PF, Russell MQ, Byers M: ABC to CABC: redefining the military trauma paradigm. *Emerg Med J* 2006; 23: 745–6.
4. Lee C, Porter KM, Hodgetts TJ: Tourniquet use in the civilian prehospital setting. *Emerg Med J* 2007; 24: 584–7.

Hospital Preparedness

In their article, Franke et al. give an important overview of treatments and treatment procedures for major casualty incidents (MCI) (1). In addition to the fastest possible identification and treatment of severely injured patients at the site of the incident, a further recommendation of the authors is that hospitals are prepared for a major incident. With regard to the latter point, we would like to broaden the discussion by adding the following important aspects.

In the hospital, the early phase of transitioning from routine activities to MCI procedures is essential for obtaining the most structured response possible. For this purpose, protocols for standard operating procedures that contain clearly defined roles with their responsibilities, chain of command, and clear guidelines for logistical structures must be compiled and available at all times. This includes for example the establishment of a triage sieve, and the definition of the sub-areas of the individual MCI triage categories by the emergency room team.

In addition, each individual patient must be clearly identifiable at all times during care. A paper-based form of documentation and/or the use of a hospital information system (HIS) with so-called MCI dummies (e.g., pre-defined cases prepared in advance) could be used for this.

From an interdisciplinary emergency and clinical laboratory perspective, developing a protocol in advance is necessary to be able to successfully allocate the required laboratory supplies (blood values and blood groups; blood units and products) to the patient during a MCI. The same applies to imaging diagnostics and the transmission of information to interface areas.

In addition to the repetitive training measures in the individual sub-areas (for example, central emergency reception area, laboratory, operative unit, intensive care unit), each concept has to be evaluated for its functionality and regularly tested for all clinics. For this, joint exercises of the rescue services have been established to carry out mock drills using persons who simulate injuries.

DOI: 10.3238/arztebl.2017.0561b

REFERENCES

1. Franke A, Bieler D, Friemert B, Schwab R, Kollig E, Gsgen C: The first aid and hospital treatment of gunshot and blast injuries. *Dtsch Arztebl Int* 2017; 114: 237–43.

Dr. med. Ramona Dolscheid-Pommerich
Prof. Dr. med. Birgit Stoffel-Wagner
 Institut für Klinische Chemie und Klinische Pharmakologie

Dr. med. Ingo Gräff
 Interdisziplinäres Notfallzentrum
 Universitätsklinikum Bonn, Germany
 ramona.dolscheid-pommerich@ukbonn.de

4. Hauser CJ, Boffard K, Dutton A, et al.: Results of the CONTROL trial: efficacy and safety of recombinant activated Factor VII in the management of refractory traumatic hemorrhage. *J Trauma* 2010; 69: 489–500.

Prof. Dr. med. Dr. -Ing. Holger Kieseewetter
 Hämostaseologikum
 Berliner Zentrum für Blutgerinnungserkrankungen, Germany
 2.haemo-mitte@haemostaseologikum-steglitz.de

Measures for Bleeding Control

The article by Franke et al. (1) recommends wound packing or tamponade to stop bleeding; that this should be combined with the administration of a hemostyptic is only stated in parentheses. Unfortunately, the hemostyptics are not further defined.

The recommended treatment for dealing with massive bleeding and associated perioperative coagulation disorders is described in Grottke et al. (2). As transfusion triggers are often not immediately available, pragmatic rapid therapy should be used (3). For blood loss of approximately 5 liters, approximately 10 units of red blood cell concentrates of the blood group 0 (if the patient has an unknown blood group) should be transfused, together with 4 to 5 units of platelet concentrates of any blood group.

The required volume therapy should be essentially carried out with crystalloid solutions if fresh plasma is not sufficiently available in liquid form, and otherwise, in combination with fresh plasma (using 2 to 3 liters of either crystalloid solution or fresh plasma of blood group AB).

Once a patient has lost 5 liters of blood, the coagulation activity is only about 30% active. Therefore, 4000 IU of the coagulation factors II, VII, IX, and X (prothrombin complex concentrate, PCC) should be co-administered if only the crystalloid and colloidal solution is used, and about 2000 IU of PCC if fresh plasma is used. For massive bleeding, activated PCC should be used. However, 4 g of fibrinogen should be given intravenously in parallel so that sufficient fibrin is available for blood clotting.

Positive experiences have been reported for treating dilutional coagulopathy due to large blood loss with recombinant factor VIIa (for example, administering 7 mg for an 80 kg patient) (4). This factor can be used in addition to volume therapy and administration of non-activated PCC.

Once the bleeding has been stopped, it is imperative to administer heparin for anticoagulation, due to the high risk of thrombosis, until complete mobilization is achieved.

DOI: 10.3238/arztebl.2017.0562a

REFERENCES

1. Franke A, Bieler D, Friemert B, Schwab R, Kollig E, Günsen C: The first aid and hospital treatment of gunshot and blast injuries. *Dtsch Arztebl Int* 2017; 114: 237–43.
2. Grottke O, Frietsch T, Maas M, et al.: Handlungsempfehlung, Umgang mit Massivblutungen und assoziierten perioperativen Gerinnungsstörungen. *Anästh Intensivmed* 2013; 54: 147–57.
3. Henrich W, Surbek D, Kainer F, et al.: Diagnosis and treatment of peripartum bleeding. *Perinat Med* 2008; 36: 467–78.

Supplement to Forensic Analysis

Franke et al. present a relevant overview of primary care and hospital treatment of gunshot and blast injuries (1). However, the measures given for forensic analysis, inaccurately presented under “Goals of further treatment,” require further clarification.

Time constraints when treating patients with such injuries often make it unrealistic for a crime scene technician or forensic expert to be consulted for evidence collection. Therefore, every surgeon should have basic knowledge about the evidence collection for gunshot injuries.

In addition to the four evidence objects mentioned in the text (projectiles, fragments, clothing, and potentially relevant objects), the following are also important for allowing event reconstruction (2, 3):

- Skin sample taken on self-adhesive dressing material.
- Dressing material used during first aid.
- Excised material from the wound.
- Imaging examinations
- Photographs of injuries

In general, the physician giving first aid cannot determine which of the listed evidence objects will be relevant during event reconstruction.

When dealing with gunshot injuries, numerous pieces of evidence can be lost relatively quickly (for example, gunshot residue lost by skin disinfection); therefore, evidence should be collected as soon as possible. Importantly, following recommendations for safeguarding gunshot evidence in the clinical area can contribute decisively to determining criminal acts and to classifying unclear gunshot injuries (Accident? Suicide? Homicide?).

In the case of non-self-inflicted wounds, an appropriate forensics analysis is in the mutual interests not only of the patient, law enforcement, and the legal system but also of the attending physician, who would be expected to testify as an expert witness in court. If permitted by time constraints during the first aid response, it is advisable to consult a forensic expert.

DOI: 10.3238/arztebl.2017.0562b

REFERENCES

1. Franke A, Bieler D, Friemert B, Schwab R, Kollig E, Günsen C: The first aid and hospital treatment of gunshot and blast injuries. *Dtsch Arztebl Int* 2017; 114: 237–43.
2. Schyma C, Madea B: Schussspurensicherung – Praktischer Umgang mit Schuss- und Schmauchspuren. *Rechtsmedizin* 2010; 20: 123–33.
3. Zack F, Manhart J, Rummel J, Büttner A: Die operierte Schussverletzung. Schussspurensicherung im klinischen Bereich ohne primäre

Beteiligung der Polizei oder der Rechtsmedizin. Unfallchirurg 2015; 118: 468–71.

PD Dr. med. Fred Zack
Prof. Dr. med. Andreas Büttner
 Institut für Rechtsmedizin der Universitätsmedizin Rostock, Germany
 fred.zack@med.uni-rostock.de

In Reply:

We would like to thank the overall positive reactions to our work (1). Due to reasons of space, it was not possible to address this complex topic to its full extent.

Dickinson et al. criticize, among other things, the fact that we did not recommend a clearly structured approach to first aid for injured persons in a mass casualty incident, such as a terrorist attack with numerous gunshot and blast injuries. Here, we deliberately tried to highlight only the medical priorities.

Within our German federal system, preparation is the responsibility of the local authorities and federal states. Thus, the establishment of management, observation, and treatment structures adequate for terrorist situations, and the adaptation of existing emergency care algorithms, are currently being discussed. Local solutions adapted to the existing care structures are currently being developed. It is correct and important to emphasize that the C–ABC algorithm, for instance, must be prioritized in these considerations.

These preparations include, as correctly pointed out by Dolscheid-Pommerich et al., a clear allocation of roles and definitions of responsibilities and command hierarchy. They also include predestinated triage and documentation areas immediately outside the hospital, as well as the adaptation and testing of the Hospital Emergency Operations Plan (*Krankenhaus Alarm- und Einsatzplan*, KAEP). These intra-hospital organizational structures have to become hospital-specific and must be trained at each hospital with realistic resource usage. Such training exercises are essential for identifying ways to improve both the pre-hospital and hospital treatment, yet the cost burden means that hospitals, local authorities, and federal states face both technical and political challenges.

We deliberately did not include recommendations for individual hemostyptics, since we consider that the scientific discussion about the advantages and disadvantages of individual preparations has not yet been finalized. Nevertheless, the procurement and

stocking of these supplies are recommended. The explanation of Kieseewetter for differentiated volume therapy during life-threatening (perforating) bleeding represents a possible procedure and includes key elements of the current S3 guideline on treatment of polytrauma injuries. These are part of the individualized emergency room procedures and are established, for example, in regional and supra-regional trauma centers. For resource planning and storage in the event of mass casualties due to gunshot and blast injuries, these valuable recommendations must be taken into account.

The aim of our article (1) was to provide an overview and to sensitize all those involved in treatment in an emergency situation to the importance of rapidly stopping bleeding and of identifying injured persons who are bleeding from the trunk or into body cavities. In the case of the latter, mortality can only be reduced by timely transportation to allow bleeding to be stopped surgically and the additional administration of differentiated volume therapy.

Zack et al. rightly point out that obtaining forensic evidence is an important aspect for all parties concerned. This is easier to do with individual cases than with mass casualties. However, all practitioners should be aware of this commitment and take this into account when treating patients, even though ensuring the survival of as many people as possible must initially be given priority at the beginning of a mass attack or a local emergency situation.

Our work focused specifically on providing the basic information about gunshot and blast injuries that are required for a coordinated action from all parties in the different levels involved in the emergency response and supply chain. This required us to limit our work to the essentials. Thus, the compromises and limitations of this article should be evaluated within this framework.

DOI: 10.3238/arztebl.2017.0563

REFERENCES

1. Franke A, Bieler D, Friemert B, Schwab R, Kollig E, Günsen C: The first aid and hospital treatment of gunshot and blast injuries. *Dtsch Arztebl Int* 2017; 114: 237–43.

On behalf of the authors

PD Dr. med. Axel Franke
 Klinik für Unfallchirurgie, Orthopädie, Rekonstruktive- und Handchirurgie,
 Verbrennungsmedizin
 Bundeswehrzentral Krankenhaus Koblenz, Germany
 axel1franke@bundeswehr.org

Conflict of interest statement

The authors of the contributions declare that no conflict of interest exists.