Published in final edited form as:

Curr Treat Options Psychiatry. 2017 March ; 4(1): 1–12. doi:10.1007/s40501-017-0100-6.

Psychosocial Treatment Options for Major Depressive Disorder in Older Adults

Brenna N. Renn, PhD¹ and Patricia A. Areán, PhD¹

¹Department of Psychiatry & Behavioral Sciences, University of Washington, Seattle, WA

Opinion Statement

Late-life depression (LLD) is a public health concern with deleterious effects on overall health, cognition, quality of life, and mortality. Although LLD is relatively common, it is not a normal part of aging and is often under-recognized in older adults. However, psychotherapy is an effective treatment for LLD that aligns with many patients' preferences and can improve health and functioning. This review synthesized the current literature on evidence-based psychotherapies for the treatment of depression in older adults. Findings suggest that active, skills-based psychotherapies (cognitive behavioral therapy [CBT] and problem-solving therapy [PST]) may be more effective for LLD than non-directive, supportive counseling. PST may be particularly relevant for offsetting skill deficit associated with LLD, such as in instances of cognitive impairment (especially executive dysfunction) and disability. Emerging treatments also consider contextual factors to improve treatment delivery, such as personalized care, access, and poverty. Tele-mental health represents one such exciting new way of improving access and uptake of treatment by older adults. Although these strategies hold promise, further investigation via randomized controlled trials and comparative effectiveness are necessary to advance our treatment of LLD. Priority should be given to recruiting and training the geriatric mental health workforce to deliver evidence-based psychosocial interventions for LLD.

Keywords

Geriatric psychiatry; geropsychology; late-life depression; older adults; treatment; psychotherapy

Corresponding Author: Patricia A. Areán, PhD, 1959 NE Pacific St. Box 356560, Seattle, WA 98195, parean@uw.edu.
Brenna N. Renn, PhD, 1959 NE Pacific St. Box 356560, Seattle, WA 98195

Compliance with Ethics Guidelines

Conflict of Interest

Dr. Renn has nothing to disclose.

Dr. Areán's research is funded by NIMH and United Health Care. Dr. Areán is also a member of the American Psychological Association Advisory Board for Treatment Guidelines, and also works with the UW AIMS Center, designing educational programs to train care managers in evidence-based psychotherapy, including problem solving treatment, behavioral activation, cognitive processing therapy, and cognitive behavioral therapy for anxiety and insomnia.

Human and Animal Rights and Informed Consent

This article does not contain any studies with animal subjects performed by any of the authors. All reported studies with human subjects performed by the authors have been previously published and complied with all applicable ethical standards (including the Helsinki declaration and its amendments, institutional/national research committee standards, and international/national/institutional guidelines).

Introduction

Changing life circumstances and aging-related processes place older adults at unique risk of encountering factors that contribute to late-life depression (LLD), including chronic disease, disability and frailty, caregiving demands, experiences related to grief and loss, and cognitive impairment. In addition to the personal impact of depression, this condition also confers an economic burden, as it is associated with increased overall health service utilization and healthcare expenditure among older adults(1). Although depression is not a normal aspect of aging, LLD is relatively common. Prevalence data range depending on diagnostic severity (e.g., diagnosis of major depressive disorder [MDD] versus depressive symptomatology); US nationally representative estimates of the 12-month prevalence of MDD for adults aged 55 and older are between 4(2) – 5.63%(3), and that for any mood disorder at 4.9(2) – 6.77(3). These estimates may be conservative due in part to methodological and diagnostic issues(2, 4). While the DSM-5 criteria are the same for MDD in older and younger patients, the clinical presentation may differ, with older adults endorsing more somatic symptoms and less mood complaints(5). LLD may be more chronic than presentation in younger adults, and is often associated with or exacerbated by chronic physical health conditions, cognitive impairment, and dementia(6). As such, LLD often goes unrecognized and undertreated.

When treating depression in older adults, it is essential to identify co-occurring medical conditions that may be contributing to mood symptoms or complicate its treatment. Psychiatric treatment guidelines recommend psychotherapy and/or pharmacotherapy for initial treatment of individuals with MDD(7). However, treatment with antidepressant medication may be problematic, as nonadherence is common among older adults for myriad reasons including polypharmacy, cost, and beliefs about medication(8). A recent meta-analysis found that cognitive impairment (particularly executive dysfunction) is associated with poor antidepressant treatment response in LLD(9). Furthermore, mounting evidence suggests a greater patient preference for psychological treatment compared to medication(10).

A recent systematic review and meta-analysis(11) confirmed that psychotherapy is an effective treatment for late-life depression. Magnitudes of effect vary by type of control group, with large effects in comparison to waitlist and attention controls, and small-to-moderate effects compared to treatment-as-usual and non-specific supportive therapy. Given the deleterious effect of LLD on psychological well-being, physical health, and quality of life, it is imperative to broad access to care and deliver effective, evidence-based treatment for LLD among a growing, diverse aging population. This paper summarizes the evidence base for psychosocial treatments of LLD and reports the most recent findings. The review focuses on the psychotherapies with the strongest evidence base and discusses emerging trends and future directions in the literature.

Psychosocial Treatments

Cognitive behavior therapy

Cognitive behavior therapy (CBT) is one of the most rigorously tested psychosocial treatments for depression across the lifespan, with consistent evidence as efficacious for late-life depression. This treatment approach, as pioneered by Beck(12) and Ellis(13), posits that maladaptive cognitions precipitate and maintain depression and other forms of emotional distress. Specifically, depression results from a triad of negative views about oneself, one's future, and the world. Thus, this structured treatment actively engages older adults to identify and modify, or "restructure," their maladaptive cognitions while also incorporating behavioral components; namely behavioral activation, relaxation training, and skill rehearsal. Treatment may be relatively brief in duration, ranging from 12–16 sessions. CBT for older adults may be modified to include gerontologically-relevant elements such as physical and cognitive changes, cohort beliefs, losses and role transitions, and there are multiple resources to inform CBT practice with older adults(14, 15).

Meta-analyses typically find that CBT yields large effects on depression in older adult samples compared to non-active (e.g., waitlist) controls(16, 17), although weaker effects have been found when comparing CBT to other active treatment comparison groups, and among studies of depressed older adults with physical illness or cognitive impairment(16). Age-relevant settings such as the US Department of Veterans Affairs classify CBT as an evidence-based practice and emphasize clinical training and implantation of this treatment(18). Given the extensive literature base supporting the clinical dissemination of CBT, recent research has demonstrated interest in emerging trends of improving access to CBT among older adults and addressing concomitant cognitive deficits associated with depression. For example, Scogin et al.(19) augmented CBT with memory training in an attempt to ameliorate deficits in encoding strategies; although CBT was effective at reducing depressive symptoms in this study, the additional training did not produce cognitive effects. However, depression is more often associated with executive dysfunction than memory deficits; as such, Goodkind and colleagues(20) used neuropsychological assessment to investigate cognitive outcomes of psychotherapy. Among depressed older adults, worse executive function at baseline predicted better treatment response to CBT. Findings suggest that CBT, and likely other structured therapies that teach cognitive techniques for improving psychiatric symptoms, may be especially beneficial in LLD for those with executive dysfunction.

Low-intensity CBT interventions for older adults with depression have received increasing attention. One such study in Australian adults over the age of 60 demonstrated efficacy of self-guided internet-based CBT compared to clinician-guided treatment, noting large clinical improvements in symptoms of depression(21) and cost effectiveness(22). A low-intensity peer-facilitated workbook-based CBT intervention in older adults has demonstrated preliminary efficacy for reducing depressive symptoms in older adults(23). However, these recent works have tended to focus on depressive symptoms, rather than frank clinical presentation of MDD, in older adults. Much of this work is still in its infancy of demonstrating feasibility and preliminary efficacy; further work is needed to demonstrate

non-inferiority of novel treatments with traditional care and develop scalable implementation strategies.

Behavior therapy

The core principle of behavioral treatments (BT) for depression hinges on Lewinsohn's(24) model which posited that as depressed individuals lose contact with meaningful and rewarding activities, they eventually find fewer activities pleasant and increasingly disengage, thereby reducing positive reinforcement, effectively precipitating and maintaining a depressed state. BT is related to CBT and focused on assisting patients to see the connection between mood and daily activities via self-monitoring and activity scheduling. As depressed individuals learn how to confront avoidance and engage in more positive interactions with their environment, their depressive symptoms remit and quality of life improves. Contemporary BT centers on behavioral activation (BA(25)) and includes contingency management and social skills training. BA may be particularly relevant to older adults, as later life is associated with changes and losses, including transition of occupation and social roles; navigating changing family roles and caregiving; grief related to death, loss, and physical decline; and reduced functioning. Therefore, targeting engagement in rewarding and purposeful activities is pertinent to these very transitions older adults are navigating. There is recent meta-analytic support for BA for depression in adults(26); however, few studies have implemented BA among older adults with major depressive disorder. Recently, Egede et al.(27) conducted an open non-inferiority trial of telemedicine and in-person BA for older Veterans meeting diagnostic criteria for MDD. Both treatment modalities resulted in large, meaningful effects in depression symptom reduction and remission, and telemedicine was judged not inferior to in-person treatment.

Other so-called “third wave” behavior therapies, such as acceptance and commitment therapy (ACT(28)) have generated increased interest as treatments for depression across the lifespan, with applicability for LLD. ACT is a behavioral approach based on relational frame theory(29) that emphasizes acceptance of events that are beyond one's control—in the context of LLD, this may include chronic or terminal illness, pain and disability, and grief and loss. A preliminary trial of ACT among long-term care residents suggested acceptability and preliminary efficacy on depressive symptoms(30). However, participants had elevated depressive symptoms rather than diagnosed LLD. As with BA, future directions necessitate establishing the efficacy and effectiveness of ACT or other third wave therapies with largescale clinical trials.

Problem-solving therapy

Originally introduced in 1971 by D’Zurilla and Goldfried(31), problem-solving therapy (PST) aims to reduce psychopathology and maximize quality of life by developing constructive problem-solving skills for more effective coping. PST is based on the premise that depression is maintained by ineffective problem-solving and the resultant poor coping, low self-efficacy, and negative emotions. The PST therapist teaches effective ways of addressing problems while maximizing well-being by utilizing a 7-step approach, encompassing (1) selecting a specific problem and defining it in concrete terms; (2) selecting a goal that is feasible to reach between sessions, (3) identifying different ways to

reach that goal, (4) evaluating each solution based on the likelihoods the patient can actually implement it, (5) choosing the best solution, (6) creating a plan to implement the solution, and (7) circling back around to ascertain the effectiveness of the solution. While PST shares some similarities with CBT and BT, it focuses on ineffective problem solving rather than maladaptive cognitions or lack of activity. To date, PST is the only psychotherapy for late-life depression that has demonstrated efficacy relative to a supportive therapy control(32) as well as CBT among medically ill older adults(33).

Although it has demonstrated efficacy across the lifespan, PST is particularly effective and relevant for issues of late-life depression(34). Recent meta-analysis of PST interventions for LLD suggested that PST may be more effective for treatment of LLD than depression in younger populations(35). PST treatment can address skill deficit in the context of age-related issues including disability, executive dysfunction, and cognitive impairment. Due in part to the treatment's structured approach of teaching straightforward, discrete steps for solving real problems, PST has an established record of treating LLD among individuals with comorbid mild cognitive impairment, particularly among those with deficits in executive function, with very large effect sizes (e.g., number needed to treat [NNT] = 4). While there is preliminary evidence of improved executive functioning associated with reduction in depression, there is no evidence of superiority of PST on other cognitive outcomes(36).

With regard to access, one of the most effective way to increase older adult participation in psychotherapy is through collaborative care approaches and integration into primary care(6). Perhaps the most widely recognized and cited collaborative care mode is IMPACT(37), in which care managers worked with supervising mental health specialists to deliver psychiatric services, including PST, to depressed older adults in primary care. The IMPACT trial showcased the effectiveness of PST in the context of collaborative care(38). At the time of this writing, a trial is underway (GermanIMPACT(39)) to investigate and replicate IMPACT in German primary care context for adults 60 and older with depressive symptoms consistent with a unipolar mood disorder.

In addition to collaborative care, PST has been investigated in the context of home-based treatment. The Program to Encourage Active, Rewarding Lives for Seniors (PEARLS(40) trained social service workers to deliver PST to depressed adults over the age of 60. Promising new work by Kiosses and colleagues(41) presented a modified, home-delivered PST intervention for older adults with LLD, cognitive impairment, and disability. Named problem adaptation therapy (PATH), this approach supplements PST with age-relevant compensatory and environmental strategies, emphasis on emotion regulation, and caregiver involvement in treatment. Future work is needed to reduce depression and disability in this vulnerable population, whom have not been adequately investigated in existing psychosocial interventions.

New research in PST has also extended its efficacy for LLD to a specific symptom of depression: suicidal ideation. Given the interconnectedness of depression, hopelessness, and wishes to die, problem-solving skills may also reduce suicidal ideation among depressed older adults by generating alternatives to what may feel like hopeless, unsolvable problems.

Recent trials have supported the role of PST at increasing hopefulness and attenuating suicidal ideation. For example, Choi and colleagues(42) offer preliminary support for telehealth PST at reducing suicidal ideation among homebound older adults. PST was more effective than supportive therapy at reducing suicidal ideation in older adults with both executive dysfunction and MDD(43).

Interpersonal psychotherapy

Interpersonal psychotherapy (IPT) was developed by Klerman, Weissman, and colleagues(44) and draws on psychodynamic theory to focus on interpersonal causes of depression. Like CBT and PST, treatment is typically time-limited (e.g., 12–16 weeks). IPT is structured into three phases: (1) the initial phase, in which a psychoeducation about depression is reviewed, recent interpersonal issues are identified, and treatment goals are collaboratively outlined; (2) the intermediate phase, in which individual works to address one or two interpersonal problems, which may focus on role transitions, issues of grief and loss, interpersonal deficits, or interpersonal conflicts; and (3) the termination phase of treatment, in which treatment success is reviewed. The IPT protocol has been expanded upon for use with older adults with depression(45). Unlike other psychotherapies, IPT frames depression as a medical illness, which may appeal to older cohorts who perceive depression and its treatment as stigmatizing. Furthermore, IPT is well-positioned to address the interpersonal and grief-related issues, such as complicated bereavement, common in aging. Like the other structured, brief treatments reviewed herein, IPT has demonstrated efficacy in the treatment of depression. Meta-analytic findings of IPT for adults demonstrated moderate-to-large effects for acute depression compared to nonnative control groups, efficacy comparable to other psychological treatments or pharmacotherapy(46). However, only one trial of IPT for LLD(47) was included in this review. Despite the dearth of literature on IPT for depression in older adults, promising recent findings include a pilot examination of IPT's role in reducing suicidal ideation in depressed older adults(48). Thus, although IPT has a strong evidence base for depression in adults, little is known about its effectiveness as a stand-alone treatment for LLD. Further investigation of IPT in older adults would inform interventions for LLD that explicitly leverage social capacities in a developmental context.

Reminiscence therapy and life review

Reminiscence therapy is a term encompassing a variety of approaches that entail review of one's past experiences. While reminisce may include unstructured storytelling, life-review therapy is more structured and typically focuses on a therapeutic process of promoting a positive view of one's past among older adults with depression(49). Such treatment tends to be brief (e.g., 6–8 sessions) and in addition to LLD may be used more broadly for end-of-life, life satisfaction, and social integration. Meta-analytic findings suggest that reminiscence produces small to moderate effects on depressive symptoms and may be appropriate to LLD in the context of chronic physical illness(49). Evidence from a recent systematic review synthesized three trials of life review therapy among frail older adults; all trials demonstrated large clinical effects of treatment on depressive symptoms in international samples (Spain(50), Switzerland(51), and Taiwan(52)). However, the trials to date have not used confirmed diagnoses of depression (rather, symptoms of depression), and entailed

comparison of a nonactive control condition that did not receive an active psychological treatment; thus, it is not clear how this type of treatment compares to other psychological approaches for LLD, and results may be explained by regression to the mean.

Emerging Treatments & Future Directions

There is no doubt that psychosocial treatments are effective treatment approaches for older adults with depression, and the extant literature suggests that these interventions tend to be treatments of choice for most older adults(53–55). Unfortunately, older adults face serious challenges accessing psychosocial interventions(56), and when they do, there is little guidance as to which intervention should be the best fit(57). Additionally, many clinicians who offer psychosocial treatments are not trained in the evidence-based treatment described above. Indeed, a recent Institute of Medicine (IOM) report highlights the challenges faced in the delivery of high quality psychosocial interventions, including the poor preparation clinicians receive while in training and the lack of system level supports and incentives to ensure clinician continue to deliver these evidence based practices (58). Finally, in our experience working with low-income older adults, environment is a critical factor driving depression, yet most interventions are not designed to address the environmental contribution to depression(59).

Access to psychosocial interventions

The most common barriers to utilization of psychological services among older adults are affordability of care, difficulty traveling, and lack of transportation(53). Although the PEARLS program mentioned herein is designed to deliver in-home care, clinician travel to people's home is not a reimbursable activity. Recently, researchers have been studying the use of technology to facilitate access to this care. Emerging work has demonstrated the feasibility of internet-delivered CBT among Australian older adults with elevated depressive symptoms(60) and shown that tele-PST was well-accepted and efficacious for depression among low-income homebound adults(61). However, the latter study focused on a “young old” population (50 to 64 years old). As reviewed herein, recent work by Egede and colleagues(27) has supported the noninferiority of telemedicine using BA for LLD. Related to the growing interest in internet-based treatment delivery, there is a growing interest in novel use of mobile technology for depression treatment, such as mental health apps for smartphones. Although no studies to date have demonstrated the efficacy of apps on LLD, trials are underway to change the current models of face-to-face psychotherapy. Apps have the potential to play an important role in the future of psychological service delivery. Similarly, therapeutic videogames are of interest, but neither technology has been investigated in clinical trials. Given the large proportion of older adults who use the internet and smartphone technology, there is a general paradigm shift toward tele-mental health as acceptable and efficacious, with the power to improve access.

Personalized care

Selectively targeting treatment based on mechanisms of change may improve treatment response. The personalized treatment approach is beginning to bear fruit in regards to late life depression. As we discussed above under our review of PST, decades of research has

demonstrated that deficits in the cognitive control network are associated with poor response to selective serotonin reuptake inhibitors (SSRI) medications, but PST seems uniquely suited to this variant of late life depression. In a study interrogating the role that cognitive deficits play in CBT, investigators including Gallagher-Thompson discovered that pretreatment deficits in cognitive flexibility appears to be associated with better outcomes(20). We have recently been funded to study the impact of a stepped behavioral intervention, called Engage, on LLD. This intervention was developed using patient-centered and clinician centered design, as well as recent information regarding reward seeking and processing in LLD (62, 63). Preliminary data suggests that not only is this intervention easy for clinicians and patient to use, but maybe more effective than PST(62). The role of cognitive processing variables, particularly executive function, has garnered increased attention as moderators of treatment effects. While in-depth discussion of LLD and cognitive impairment (especially dementia) is beyond the scope of this review, LLD has been implicated with deficits in executive functioning, processing speed, and psychomotor speed. To date, all randomized controlled trials (RCTs) for LLD and cognitive deficits have involved a PST approach. More work is needed to ascertain treatment targets for the relationship between mood and cognitive impairment. The interested reader is directed to recent comprehensive reviews of LLD and cognitive impairment in older adults(64–67). Further investigation of cognitive, neurological, and other mechanisms of treatment response are warranted to develop new, personalized treatments, and to enhance clinical decision-making.

Clinician training

Despite numerous policy reports on the need to build the geriatric mental health workforce(68–72), we still have an deficit of clinicians prepared to delivery high quality psychosocial interventions in general(73), and specifically for depression in late life(74). The HRSA Geriatric Workforce Enhancement Program is designed to specifically training clinicians in working with older adults, and includes training of nurses, psychologists and social workers (<http://bhpr.hrsa.gov/grants/geriatricalliedhealth/gwep.html>). Newly awarded programs are focusing on training in the evidence-based practices detailed in this paper. What will be needed, beyond training, is the development of quality measures post training, to ensure these interventions continue to be deployed in the field. As is discussed at length in the IOM report on standards for psychosocial interventions, quality measures not only of clinician behavior, but system level support to provide evidence-based care is sorely needed. Systems supports include access to experts or peer mentoring to trouble-shoot when clinicians face a challenge implementing an evidence-based practice, and technology based decision-making tools to help clinicians adapt evidence-based treatments for individual patients(75).

Poverty

Low-income older adults tend to have poor response to antidepressants, even when combined with psychotherapy(76). We speculate that the reason for poor response to medications is that these treatments do not address the drivers of depression in the context of poverty. When people are faced with a daily struggle to find food, shelter, safety from harm and medical care, we hypothesize that a stress reaction takes place that creates a syndrome similar to a major depression in presentation(59). Our group has taken the lead on two trials

to ascertain the relative merit of clinical case management on depression in low-income older adults. Clinical case management is an advocacy-based intervention that addresses older adult distress by assisting with linkage to social and medical services, stable and safe housing, and other needs. We have found in two large scale RCTs that clinical case management is a highly successful intervention for late life depression in the context of poverty, and that the additional of CBT(77), or PST(78, 79) produced no added benefit.

Conclusions

In summary, many of the same psychotherapies with evidence for treating depression in the general adult population demonstrate efficacy for LLD. Direct comparisons of psychological interventions suggest that active, skills-based psychotherapies (CBT and PST) may be more effective for LLD than non-directive, supportive counseling(80). More RCTs comparing interventions with active controls are necessary before drawing firm conclusions about targeting treatment in LLD.

Regardless of modality, psychological treatments are most effective when they incorporate patient preferences and goals. Clinical decisions should be made in collaboration with the patient and utilize a multimodal, biopsychosocial approach to address concomitant physical, cognitive, and neurological disorders while also appreciating sociocultural factors. At present, older adults with LLD utilize psychotherapy at low rates, and the geriatric workforce is ill equipped to manage the “silver tsunami”(70) of aging adults. As such, increased attention is needed to develop and implement effective, scalable, and accessible interventions to reduce the morbidity of LLD and improve the quality of life of older adults.

Acknowledgments

Funding: This research was partially supported by a training grant from the National Institutes of Health (Grant 5T32MH073553).

References

* Of importance

** Of outstanding importance

1. Luppá M, Sikorski C, Motzek T, et al. Health service utilization and costs of depressive symptoms in late life — a systematic review. *Curr Pharm Des.* 2012; 18(36):5936–57. [PubMed: 22681171]
2. Byers AL, Covinsky KE, Friedman MB, et al. High occurrence of mood and anxiety disorders among older adults: the National Comorbidity Survey Replication. *Arch Gen Psychiat.* 2010; 67(5): 489–96. [PubMed: 20439830]
3. Reynolds K, Pietrzak RH, El-Gabalawy R, et al. Prevalence of psychiatric disorders in US older adults: findings from a nationally representative survey. *World Psychiatry.* 2015; 14(1):74–81. [PubMed: 25655161]
4. Snowdon J, Almeida OP, O'Connor DW. Dubious findings concerning the prevalence of late-life depression. *Aust NZ J Psychiat.* 2016; 50(3):284–5.
5. Hegeman JM, de Waal MW, Comijs HC, et al. Depression in later life: a more somatic presentation? *J Affect Disorders.* 2015; 170:196–202. [PubMed: 25254617]
- 6 *. Hall CA, Reynolds CF III. Late-life depression in the primary care setting: challenges, collaborative care, and prevention. *Maturitas.* 2014; 79(2):147–52. *These authors review LLD in*

the context of primary care, enhanced by the description of three pivotal studies of the management of depression in this setting. [PubMed: 24996484]

- 7 *. Qaseem A, Barry MJ, Kansagara D, et al. Nonpharmacologic versus pharmacologic treatment of adult patients with major depressive disorder: a clinical practice guideline from the American College of Physicians. *Ann Intern Med.* 2016; 164(5):350–9. *These updated practice guidelines from the American College of Physicians reviews the evidence for treatment of major depressive disorder among adults and is intended for clinicians working with this population.* [PubMed: 26857948]
8. Zivin K, Kales HC. Adherence to depression treatment in older adults: a narrative review. *Drug Aging.* 2008; 25(7):559–71.
9. Pimontel MA, Rindskopf D, Rutherford BR, et al. A meta-analysis of executive dysfunction and antidepressant treatment response in late-life depression. *Am J Geriatr Psychiatry.* 2016; 24(1):31–41.
10. McHugh RK, Whitton SW, Peckham AD, et al. Patient preference for psychological vs pharmacologic treatment of psychiatric disorders: a meta-analytic review. *J Clin Psychiatr.* 2013; 74(6):595–602.
11. Huang AX, Delucchi K, Dunn LB, et al. A systematic review and meta-analysis of psychotherapy for late-life depression. *Am J Geriatr Psychiatry.* 2015; 23(3):261–73.
12. Beck A. Nature and relation to behavior therapy. *Behav Ther.* 1970; 1:184–200.
13. Ellis, A. Reason and emotion in psychotherapy. New York, NY: Stuart; 1962.
14. Laidlaw K. Comprehensive conceptualization of cognitive behaviour therapy for late life depression. *Behav Cogn Psychoth.* 2004; 32(4):389–99.
15. Gallagher-Thompson, D., Thompson, LW. Treating late-life depression: a cognitive-behavioral therapy approach Therapist guide. New York, NY: Oxford University Press; 2010.
16. Pinquart M, Duberstein PR, Lyness JM. Effects of psychotherapy and other behavioral interventions on clinically depressed older adults: a meta-analysis. *Aging Ment Health.* 2007; 11(6):645–57. [PubMed: 18074252]
17. Gould RL, Coulson MC, Howard RJ. Cognitive behavioral therapy for depression in older people: a meta-analysis and meta-regression of randomized controlled trials. *J Am Geriatr Soc.* 2012; 60(10):1817–30. [PubMed: 23003115]
18. Veterans Health Administration. Local implementation of evidence-based psychotherapies for mental and behavioral health conditions. Washington DC: Department of Veterans Affairs; 2012.
19. Scogin F, Fairchild JK, Yon A, et al. Cognitive bibliotherapy and memory training for older adults with depressive symptoms. *Aging Ment Health.* 2014; 18(5):554–60. [PubMed: 24073847]
- 20 *. Goodkind MS, Gallagher-Thompson D, Thompson LW, et al. The impact of executive function on response to cognitive behavioral therapy in late-life depression. *Int J Geriatr Psychiatry.* 2016; 31(4):334–9. *This manuscript presents an investigation of neuropsychological function in LLD and the relationship to CBT treatment response. These results suggested that older adults with relative weaknesses in executive functioning may have improved treatment outcomes. The authors posited that the cognitive skills imparted by CBT that may be particularly helpful for this population.* [PubMed: 26230057]
21. Titov N, Fogliati VJ, Staples LG, et al. Treating anxiety and depression in older adults: randomised controlled trial comparing guided v. self-guided internet-delivered cognitive-behavioural therapy. *Brit J Psychiatr.* 2016; 2(1):50–8.
22. Titov N, Dear BF, Ali S, et al. Clinical and cost-effectiveness of therapist-guided internet-delivered cognitive behavior therapy for older adults with symptoms of depression: a randomized controlled trial. *Behav Ther.* 2015; 46(2):193–205. [PubMed: 25645168]
23. Seeley JR, Manitsas T, Gau JM. Feasibility study of a peer-facilitated low intensity cognitive-behavioral intervention for mild to moderate depression and anxiety in older adults. *Aging Ment Health.* 2016:1–7.
24. Lewinsohn, PM. A behavioral approach to depression. In: Friedman, RJ., Katz, MM., editors. *The psychology of depression: contemporary theory and research.* New York, NY: John Wiley; 1974. p. 157-78.

25. Lejuez CW, Hopko DR, Acierno R, et al. Ten year revision of the brief behavioral activation treatment for depression: revised treatment manual. *Behav Modif.* 2011; 35(2):111–61. [PubMed: 21324944]
26. Ekers D, Webster L, Van Straten A, et al. Behavioural activation for depression; an update of meta-analysis of effectiveness and sub group analysis. *PLoS One.* 2014; 9(6):e100100. [PubMed: 24936656]
- 27 **. Egede LE, Acierno R, Knapp RG, et al. Psychotherapy for depression in older veterans via telemedicine: a randomised, open-label, non-inferiority trial. *Lancet Psychiatry.* 2015; 2(8):693–701. *This is the first RCT assessing tele-mental health using a manualized, evidence-based psychotherapy for LLD. Investigators compared BA delivered in-person or via in-home teleconferencing and found reduction of symptom severity in both treatment groups, as well as high study retention and session attendance, establishing preliminary noninferiority of telemedicine for the psychological treatment of LLD.* [PubMed: 26249300]
28. Hayes SC, Luoma JB, Bond FW, et al. Acceptance and commitment therapy: model, processes and outcomes. *Behav Res Ther.* 2006; 44(1):1–25. [PubMed: 16300724]
29. Barnes-Holmes Y, Hayes SC, Barnes-Holmes D, et al. Relational frame theory: a post-Skinnerian account of human language and cognition. *Adv Child Dev Behav.* 2001; 28:101–38. [PubMed: 11605362]
30. Davison TE, Eppingstall B, Runci S, et al. A pilot trial of acceptance and commitment therapy for symptoms of depression and anxiety in older adults residing in long-term care facilities. *Aging Ment Health.* 2016:1–8.
31. D’Zurilla TJ, Goldfried MR. Problem solving and behavior modification. *J Abnorm Psychol.* 1971; 78(1):107–26. [PubMed: 4938262]
32. Arean PA, Raue P, Mackin RS, et al. Problem-solving therapy and supportive therapy in older adults with major depression and executive dysfunction. *Am J Psychiatry.* 2010; 167(11):1391–8. [PubMed: 20516155]
33. Sharpe L, Gittins CB, Correia HM, et al. Problem-solving versus cognitive restructuring of medically ill seniors with depression (PROMISE-D trial): study protocol and design. *Bmc Psychiatry.* 2012; 12:207. [PubMed: 23173830]
34. *Kiosses DN, Alexopoulos GS. Problem-solving therapy in the elderly. *Curr Treat Options Psychiatry.* 2014; 1(1):15–26. *This systematic review of PST in older adults highlights the efficacy of PST for reducing depressive symptoms in older adults with LLD. The authors highlight the utility of PST in issues of aging, including disability (e.g., macular degeneration, stroke) and cognitive impairment, and offer preliminary evidence for PST delivered via the Internet.* [PubMed: 24729951]
35. Kirkham JG, Choi N, Seitz DP. Meta-analysis of problem solving therapy for the treatment of major depressive disorder in older adults. *Int J Geriatr Psychiatry.* 2016; 31(5):526–35. [PubMed: 26437368]
36. Mackin RS, Nelson JC, Delucchi K, et al. Cognitive outcomes after psychotherapeutic interventions for major depression in older adults with executive dysfunction. *Am J Geriatr Psychiatry.* 2014; 22(12):1496–503. [PubMed: 24378255]
37. Unutzer J, Katon W, Callahan CM, et al. Collaborative care management of late-life depression in the primary care setting: a randomized controlled trial. *JAMA.* 2002; 288(22):2836–45. [PubMed: 12472325]
38. Arean P, Hegel M, Vannoy S, et al. Effectiveness of problem-solving therapy for older, primary care patients with depression: results from the IMPACT project. *Gerontologist.* 2008; 48(3):311–23. [PubMed: 18591356]
39. Wernher I, Bjerregaard F, Tinsel I, et al. Collaborative treatment of late-life depression in primary care (GermanIMPACT): study protocol of a cluster-randomized controlled trial. *Trials.* 2014; 15:351. [PubMed: 25195020]
40. Ciechanowski P, Wagner E, Schmalting K, et al. Community-integrated home-based depression treatment in older adults: a randomized controlled trial. *JAMA.* 2004; 291(13):1569–77. [PubMed: 15069044]

41. Kiosses DN, Ravdin LD, Gross JJ, et al. Problem adaptation therapy for older adults with major depression and cognitive impairment: a randomized clinical trial. *JAMA Psychiatry*. 2015; 72(1): 22–30. [PubMed: 25372657]
42. Choi NG, Marti CN, Conwell Y. Effect of problem-solving therapy on depressed low-income homebound older adults' death/suicidal ideation and hopelessness. *Suicide Life Threat Behav*. 2016; 46(3):323–36. [PubMed: 26456016]
43. Gustavson KA, Alexopoulos GS, Niu GC, et al. Problem-solving therapy reduces suicidal ideation in depressed older adults with executive dysfunction. *Am J Geriatr Psychiatry*. 2016; 24(1):11–7.
44. Klerman, GL., WM, Rounsaville, BJ., Chevron, E. Interpersonal psychotherapy of depression. New York, NY: Basic Books; 1984.
45. Hinrichsen, GAC., Kathleen, F. Interpersonal psychotherapy for depressed older adults. Washington DC: American Psychological Association; 2006.
46. Cuijpers P, Donker T, Weissman MM, et al. Interpersonal psychotherapy for mental health problems: a comprehensive meta-analysis. *Am J Psychiat*. 2016; 173(7):680–7. [PubMed: 27032627]
47. Reynolds CF III, Dew MA, Pollock BG, et al. Maintenance treatment of major depression in old age. *New Engl J Med*. 2006; 354(11):1130–8. [PubMed: 16540613]
48. Heisel MJ, Talbot NL, King DA, et al. Adapting interpersonal psychotherapy for older adults at risk for suicide. *Am J Geriatr Psychiatry*. 2015; 23(1):87–98.
49. Pinquart M, Forstmeier S. Effects of reminiscence interventions on psychosocial outcomes: a meta-analysis. *Aging Ment Health*. 2012; 16(5):541–58. [PubMed: 22304736]
50. Serrano JP, Latorre JM, Gatz M, et al. Life review therapy using autobiographical retrieval practice for older adults with depressive symptomatology. *Psychol Aging*. 2004; 19(2):272–7.
51. Preschl B, Maercker A, Wagner B, et al. Life-review therapy with computer supplements for depression in the elderly: A randomized controlled trial. *Aging Ment Health*. 2012; 16(8):964–74. [PubMed: 22788983]
52. Hsu YC, Wang JJ. Physical, affective, and behavioral effects of group reminiscence on depressed institutionalized elders in Taiwan. *Nurs Res*. 2009; 58(4):294–9. [PubMed: 19609181]
53. Gum AM, Iser L, Petkus A. Behavioral health service utilization and preferences of older adults receiving home-based aging services. *Am J Geriatr Psychiatry*. 2010; 18(6):491–501. [PubMed: 21217560]
54. Feldman DM, Gum A. Multigenerational perceptions of mental health services among deaf adults in Florida. *Am Ann Deaf*. 2007; 152(4):391–7. [PubMed: 18257508]
55. Gum AM, Arean PA, Hunkeler E, et al. Depression treatment preferences in older primary care patients. *Gerontologist*. 2006; 46(1):14–22. [PubMed: 16452280]
56. Arean PA, Raue PJ, Sirey JA, et al. Implementing evidence-based psychotherapies in settings serving older adults: challenges and solutions. *Psychiatr Serv*. 2012; 63(6):605–7. [PubMed: 22638006]
57. Arean PA. Personalizing behavioral interventions: the case of late-life depression. *Neuropsychiatry (London)*. 2012; 2(2):135–45. [PubMed: 23646065]
58. Institute of Medicine. Psychosocial interventions for mental and substance use disorders: a framework for establishing evidence-based standards. Washington DC: The National Academies Press; 2015.
59. Alexopoulos GS. Depression in the elderly. *Lancet*. 2005; 365(9475):1961–70. [PubMed: 15936426]
60. Dear BF, Zou J, Titov N, et al. Internet-delivered cognitive behavioural therapy for depression: a feasibility open trial for older adults. *Aust NZ J Psychiatry*. 2013; 47(2):169–76.
61. Choi NG, Hegel MT, Marti N, et al. Telehealth problem-solving therapy for depressed low-income homebound older adults. *Am J Geriatr Psychiatry*. 2014; 22(3):263–71. [PubMed: 23567376]
62. Alexopoulos GS, Raue PJ, Gunning F, et al. “Engage” therapy: behavioral activation and improvement of late-life major depression. *Am J Geriatr Psychiatry*. 2016; 24(4):320–6. [PubMed: 26905044]

63. Alexopoulos GS, Arean P. A model for streamlining psychotherapy in the RDoC era: the example of 'Engage'. *Mol Psychiatry*. 2014; 19(1):14–9. [PubMed: 24280983]
64. Byers AL, Yaffe K. Depression and risk of developing dementia. *Nat Rev Neurol*. 2011; 7(6):323–31. [PubMed: 21537355]
65. Koenig AM, Bhalla RK, Butters MA. Cognitive functioning and late-life depression. *J Int Neuropsych Soc*. 2014; 20(5):461–7.
66. Morimoto SS, Alexopoulos GS. Cognitive deficits in geriatric depression clinical correlates and implications for current and future treatment. *Psychiat Clin N Am*. 2013; 36(4):517. +
67. Wang S, Blazer DG. Depression and cognition in the elderly. *Annu Rev Clin Psycho*. 2015; 11:331–60.
68. Driessen J, Bellon JE, Stevans J, et al. Innovative approaches to interprofessional care at the University of Pittsburgh Medical Center. *J Interprof Care*. 2015; 29(5):520–1. [PubMed: 26171868]
69. Bartels SJ, Gill L, Naslund JA. The Affordable Care Act, accountable care organizations, and mental health care for older adults: implications and opportunities. *Harv Rev Psychiatry*. 2015; 23(5):304–19. [PubMed: 25811340]
70. Bartels SJ, Naslund JA. The underside of the silver tsunami--older adults and mental health care. *N Engl J Med*. 2013; 368(6):493–6. [PubMed: 23343039]
71. Bartels SJ, Lebowitz BD, Reynolds CF III, et al. Programs for developing the pipeline of early-career geriatric mental health researchers: outcomes and implications for other fields. *Acad Med*. 2010; 85(1):26–35. [PubMed: 20042817]
72. Eden, J, Le, M., Blazer, D., editors. Institute of Medicine. The mental health and substance use workforce for older adults: in whose hands?. Washington DC: The National Academies Press; 2012.
73. Weissman S. American psychiatry in the 21st century: the discipline, its practice, and its work force. *Bull Menninger Clin*. 1994; 58(4):502–18. [PubMed: 7812263]
74. Carpenter BD, Sakai E, Karel MJ, et al. Training for research and teaching in geropsychology: preparing the next generation of scholars and educators. *Gerontol Geriatr Educ*. 2016; 37(1):43–61. [PubMed: 26652756]
75. Aarons GA, Green AE, Willging CE, et al. Mixed-method study of a conceptual model of evidence-based intervention sustainment across multiple public-sector service settings. *Implement Sci*. 2014; 9:183. [PubMed: 25490886]
76. Cohen A, Houck PR, Szanto K, et al. Social inequalities in response to antidepressant treatment in older adults. *Arch Gen Psychiatr*. 2006; 63(1):50–6. [PubMed: 16389196]
77. Arean PA, Gum A, McCulloch CE, et al. Treatment of depression in low-income older adults. *Psychol Aging*. 2005; 20(4):601–9. [PubMed: 16420135]
- 78 **. Arean PA, Raue PJ, McCulloch C, et al. Effects of problem-solving therapy and clinical case management on disability in low-income older adults. *Am J Geriatr Psychiatry*. 2015; 23(12):1307–14. *Results from this trial offers evidence that clinical case management for low-income older adults with disability is a powerful intervention for LLD, and that the addition of PST offers no additional benefit. This manuscript presents evidence that clinical case management resulted in improved functioning among participants, with maintenance of intervention effects as long as 24 weeks after beginning treatment.* [PubMed: 26628206]
- 79 **. Alexopoulos GS, Raue PJ, McCulloch C, et al. Clinical case management versus case management with problem-solving therapy in low-income, disabled elders with major depression: a randomized clinical trial. *Am J Geriatr Psychiatry*. 2016; 24(1):50–9. *As above, this trial provides evidence of clinical case management as a crucial component of depression management for low-income older adults with disability and LLD. The results of this RCT showcases clinical case management as a stand-alone intervention that evidenced similar rates of symptom reduction and LLD remission as PST.* [PubMed: 25794636]
80. Cuijpers P, Karyotaki E, Pot AM, et al. Managing depression in older age: psychological interventions. *Maturitas*. 2014; 79(2):160–9. [PubMed: 24973043]