Received:
12 October 2016

Revised:
29 December 2016

Accepted:
24 January 2017

<https://doi.org/10.1259/bjr.20160802>

Cite this article as:

Haquin A, Sigovan M, Si-Mohamed S, Mabrut J-Y, Manichon A-F, Bakir M, et al. Phase-contrast MRI evaluation of haemodynamic changes induced by a coeliac axis stenosis in the gastroduodenal artery. *Br J Radiol* 2017; **90**: 20160802.

FULL PAPER

Phase-contrast MRI evaluation of haemodynamic changes induced by a coeliac axis stenosis in the gastroduodenal artery

¹AUDREY HAQUIN, MD, ²MONICA SIGOVAN, PhD, ^{1,2}SALIM SI-MOHAMED, MD, ³JEAN-YVES MABRUT, PhD, MD, ¹ANNE-FRÉDÉRIQUE MANICHON, MD, ¹MELISA BAKIR, MD, ¹AGNÈS RODE, MD and ^{1,2}LOÏC BOUSSEL, PhD, MD

¹Department of Radiology, Hôpital de la Croix-Rousse, University of Lyon, Lyon, France

²CREATIS, CNRS UMR 5220—INSERM U1206—University of Lyon, Lyon, France

³Department of Surgery, Hôpital de la Croix-Rousse, University of Lyon, Lyon, France

Address correspondence to: Dr Salim Si-Mohamed

E-mail: salimaymeric@gmail.com

Objective: To evaluate the correlation between the gastroduodenal artery (GDA) haemodynamic changes and the degree of coeliac axis (CA) stenosis using phase-contrast MRI.

Methods: The study was institutional review board approved, and written informed consent was obtained from patients included prospectively. A two-dimensional phase-contrast MRI was performed in 23 patients scheduled for a potential complex supramesocolic surgery, in a plane perpendicular to the GDA, during inspiration and expiration. The peak and mean velocities (V_p and V_m), mean flow rate (Q_m) and flow direction at inspiration and at expiration have been correlated with the degree of CA stenosis evaluated by CT.

Results: 13 of 23 patients presented CA stenosis due to the median arcuate ligament (34–80% of stenosis), 4 of them had associated atheromatous calcifications. V_p , V_m and Q_m of GDA presented a significant and linear relationship with the degree of CA stenosis, at inspiration as well as at expiration ($r > 0.74$, $p < 0.001$). Velocities and flow rates presented a different pattern depending on the degree of CA stenosis: $<34\%$ ($n = 10$),

flow direction remained physiological with low velocities and flow rates; $>60\%$ of stenosis ($n = 5$), a reverse flow direction with increased velocities and flow rates were observed; variable patterns between 34% and 60%.

Conclusion: Phase-contrast MRI permits the evaluation of haemodynamic changes in GDA induced by CA stenosis, including median arcuate ligament compression, and could be of great interest in therapeutic decision making in supramesocolic surgery, such as liver transplantation or duodenopancreatectomy, by detecting haemodynamically significant stenoses.

Advances in knowledge: Physiological phase-contrast MRI detects haemodynamically significant stenoses of the CA by evaluating haemodynamic parameters in the GDA. Physiological phase-contrast MRI demonstrates that flow in the GDA can be a direct marker of the real haemodynamic impact of a CA stenosis on the hepatic vascularization and could thus participate in the pre-surgical work-up of duodenopancreatectomy or hepatic graft, whereas existing pre-operative imaging are only morphological.

INTRODUCTION

During supramesocolic surgery, collateral pathways from the superior mesenteric artery to the hepatic artery, mainly the gastroduodenal artery (GDA), are ligated. In case of coeliac axis (CA) stenosis, a decrease of $>50\%$ of blood flow in the common hepatic artery^{1,2} can be observed. A main cause of CA stenosis is represented by median arcuate ligament (MAL) compression in $>25\%$ of patients.^{3–6} Also, during liver transplantation, the presence of CA compression by a MAL or an atheromatous stenosis may decrease hepatic arterial flow and lead to graft loss because bile duct vascularization depends exclusively on the arterial

flow of the hepatic artery. In order to maintain hepatic artery flow,^{7–9} a surgical vascular bypass can be mandatory during pancreaticoduodenectomy or liver transplantation which leads to an increase in procedure duration and complexity.

It is therefore important to assess the haemodynamics in the coeliac area during the pre-surgical work-up in these patients. Currently, this is performed mainly by anatomical measurement (X-ray angiography, CT or MRI¹⁰) of the degree of CA stenosis and the diameter of the GDA, the main collateral pathway that can compensate for the

decrease of flow due to the CA with retrograde flow from the superior mesenteric artery.¹¹ Nevertheless, these measurements only indirectly reflect the complex haemodynamics in the coeliac area: Doppler hepatic artery flow monitoring^{12,13} or clamping trial of the GDA^{14–16} are often necessary during the surgical procedure.¹⁷ Thus, a simple and robust non-invasive tool would be of great interest to directly assess the effect of CA stenosis on the haemodynamics before surgery. Pre-operative Doppler has been proposed to handle this task, but it suffers from technical difficulties such as visualization of the vessels in overweight patients, and operator-related variability.¹⁸ We thus propose to investigate the potential of phase-contrast MRI in this pre-surgical work-up. Indeed, phase-contrast MRI is a reliable technique that uses the change in the phase shifts of the flowing protons in the region of interest to create an image in a quantitative fashion, allowing evaluation of haemodynamic condition in the arteries.¹⁹ Furthermore, we propose to focus on the GDA. Indeed, understanding the global haemodynamics in the coeliac arteries is complex because of the anatomy of the coeliac vasculature and the variations of flow with breathing^{20,21} and with the degree of CA stenosis from one patient to another. Moreover, absolute values of normal blood flow in the hepatic artery for a given patient are difficult to establish since flow depends on many parameters.²² Hence, the direction and intensity of the flow in the GDA could be a marker of the real impact of CA stenosis on hepatic vascularization.

The purpose of this study was thus to evaluate the correlation between GDA haemodynamic changes (flow, velocity and direction) and the degree of CA stenosis, at inspiration and at expiration, using phase-contrast MRI.

METHODS AND MATERIALS

Patients

Between October 2012 and December 2013, 23 consecutive patients (17 males and 6 females; mean age: 61 ± 14 years; range: 28–84 years), scheduled for a potential complex hepatic surgery were prospectively included in this study. Hepatic surgery included liver transplantation and large hepatic resection, associated or not with cephalic pancreaticoduodenectomy. Institutional review board IRB (Hospices civils de Lyon, CPP) approval and informed consent were obtained for each patient. There were no conflicts of interest.

CT imaging

CT acquisitions were performed before the MRI examination on CT systems with at least a 4-cm Z-coverage, at 120 kV and after intravenous injection of iodine-based contrast agent at arterial phase at deep inspiration [Brilliance 64 (Philips, Cleveland, OH), Aquilion™ 64 (Toshiba; Tustin, CA), LightSpeed Pro 32, (GE, Healthcare, OH) General Electric Medical System]. Reconstruction parameters for axial slices were all of 1.5 mm.

MRI

MR examinations were performed on a Siemens Avanto system (Erlangen, Germany) at 1.5 T. The complete MRI protocol included a liver examination with T_2 weighted acquisition, T_1 weighted acquisition in and out of phase, diffusion-weighted imaging with three b -values (50, 400 and 800) and dynamic

contrast material-enhanced imaging ($0.2 \text{ cm}^3 \text{ kg}^{-1}$, MultiHance®; Bracco, Milan, Italy). Blood flow evaluation in the GDA was performed after the liver MRI protocol. First of all, the GDA was located using a two-dimensional balanced steady-state free precession (TrueFISP) sequence at deep inspiration and expiration in the axial plane with the following acquisition parameters: repetition time/echo time = 4/1.7 ms, flip angle = 52° , adapted field of view, slice thickness = 4 mm, in-plane resolution = $0.78 \times 0.78 \text{ mm}$, number of averages = 1, bandwidth = 448 Hz/pixel and acquisition duration of 12 s. Phase-contrast sequences for flow quantification consisted of a retrospective electrocardiogram-gated Cartesian two-dimensional cine fast low angle shot performed perpendicular to the GDA using the following acquisition parameters: echo time = 2.5 ms, temporal resolution = approximately 50–65 ms, depending on the heart rate, flip angle = 30° , adapted field of view, slice thickness = 5.5 mm, in-plane resolution = $1.25 \times 1.25 \text{ mm}$, number of averages = 1, bandwidth = 558 Hz/pixel. 20 cardiac phases were reconstructed. The acquisition was performed in a breath-hold interval with an average acquisition time of 24 s and was repeated twice: once at the end of a deep inspiration and the second at the end of deep expiration. An initial velocity encoding of 100 cm s^{-1} was used and subsequently modified as necessary when velocity aliasing was present, which resulted in some cases in multiple repetitions to obtain correct GDA depiction and velocity measurement. The total examination time was 45–60 min.

Image analysis

Analysis was performed by a radiologist with 4 years' experience in image analysis. The degree of CA stenosis was measured only on CT, in parasagittal views, along the main axis of the vessel, using a dedicated software (AW Server 2® software; General Electrics) as the smaller diameter of the vessel divided by the diameter of the healthy vessel downstream. Furthermore, the diameter of the GDA was also recorded after multiplanar reconstruction, perpendicularly to the centreline of the vessel. The cause of CA stenosis was recorded. It was considered to be related to a MAL when a focal narrowing of the CA with a hooked appearance was present.⁴ Flow analysis was performed on the phase-contrast sequences using dedicated software (Argus®; Siemens, Germany). The flow direction (anterograde or retrograde), peak velocity (cm s^{-1}), mean velocity (cm s^{-1}) and mean flow rate (ml min^{-1}) in the GDA were recorded at deep inspiration and expiration. Morphological evaluation of the liver was systematically performed, but the results were not included in the study.

Clinical data

Surgical procedure, hepatic artery bypass graft, surgical release of the MAL, ischaemic cholangitis, sepsis, bile leakage, hepatic arterial and portal venous thrombosis were recorded over a 1-month follow-up period.

Statistical analysis

Linear regression and Pearson correlation test were used to assess the relation between the degree of CA stenosis and mean velocity, maximum velocity, mean flow and diameter of the GDA, at inspiration and at expiration. The difference between the mean flow rate at inspiration and at expiration was also

studied using paired *t*-test. The differences were considered to be significant when the *p*-value was <0.05 . All statistical analysis was performed using Intercooled Stata v. 10.0 (StataCorp LP, College Station, TX). Data were expressed as mean $\pm$ standard deviation.

RESULTS

All the 23 patients completed the MRI examination (17 males and 6 females; mean age: 61 ± 14 years; range: 28–84 years).

Coeliac axis stenosis

13 patients presented MAL compression with a mean degree of CA stenosis of $54.9 \pm 16.3\%$ (range: 33.8–80.7%), including 4 patients with atheromatous calcifications presenting a mean degree of CA stenosis of $54.4 \pm 12.9\%$ (range: 47.2–73.8%). 7 of all the 23 patients presented atheromatous calcifications at the origin of the CA, including the 4 herein described and 3 with stenosis of the CA $<13\%$ ($8.7 \pm 4.7\%$, range 3.3–12.3%). No severe stenosis due to atheromatous calcification only has been found.

Relation between coeliac axis stenosis degree and gastroduodenal artery diameter

The mean diameter of the GDA was 4.2 ± 1.9 mm (range 2.1–6.8 mm) with a significant correlation between the diameter of the GDA and the degree of CA stenosis ($R = 0.7$; $p = 0.0002$) (Figure 1).

Relation between coeliac axis stenosis degree and flow direction

Out of these 13 patients, 8 patients presented with an inversion in the GDA at expiration and inspiration, all with degree of CA stenosis $>35\%$ ($62 \pm 16.7\%$, range: 34.9–80.7%) and 1 patient had a GDA flow inversion at expiration only, with a degree of stenosis of 34%. One should note that this patient presented with an anatomical variant, with a right hepatic artery from the superior mesenteric artery, and an iatrogenic dissection of the CA associated with MAL compression. All the other patients¹⁰ presented with a degree of CA narrowing $<18\%$ ($7.9 \pm 5.8\%$, range: 0.6–18.4%), no MAL compression and no GDA flow

inversion was observed at expiration and inspiration (Figure 2). All the patients presenting with CA stenosis $>60\%$ (5 patients) had a retrograde flow in the GDA at inspiration and at expiration, whereas all the patients with stenosis $<34\%$ (10 patients) had an antegrade flow in the GDA. Between 34% and 60%, the flow was retrograde in four patients and antegrade in four patients. Table 1 reports the values of peak velocity, mean velocity and flow in these three groups.

Relationship between coeliac axis stenosis degree and V_p , V_m and Q_m

A significant and linear relation was found between the degree of CA stenosis and GDA peak velocity, mean velocity and flow with coefficient of correlation of, respectively, 0.82, 0.74 and 0.77 at inspiration and 0.81, 0.76 and 0.80 at expiration ($p < 0.0001$ in any cases) (Figure 3).

Difference between Q_m at inspiration and at expiration.

No significant difference was found between mean flow rate at inspiration and at expiration ($p = 0.24$) (Figure 4).

Clinical data

Out of the 23 patients, 16 did not undergo surgery during the study. Five patients received liver transplantation (26.1%). Two patients with CA stenosis $>60\%$ and GDA reversal flow underwent hepatic artery bypass graft during liver transplantation with good outcome for one and sepsis for the second. One patient with CA stenosis $>60\%$ and GDA reversal flow did not profit from hepatic artery bypass graft due to the high surgical risk according to the surgical team and presented hepatic arterial thrombosis at 8 days after transplantation treated by a revision surgery. Two patients with CA stenosis $<34\%$ without GDA reversal flow presented good outcome without the need of hepatic bypass graft. Two patients underwent liver resection associated with cephalic pancreaticoduodenectomy. One patient with CA stenosis of 47.6% and GDA reversal flow underwent a surgical release of the CA and presented hepatic arterial haemorrhage 1 week after with bad outcome (death). The second with CA stenosis $<34\%$ without GDA reversal flow did not need hepatic bypass graft with good outcome.

No ischaemic cholangitis was recorded. Five patients received liver transplantation (26.1%).

DISCUSSION

The GDA plays an important role in hepatic vascularization. It acts as a bypass between the hepatic and the mesenteric vascularization, with a physiological flow direction going from the hepatic (merging from the CA) to the mesenteric artery. The flow can be nulled or the direction reversed in case of CA stenosis. Flow in the GDA can thus be a direct marker of the real haemodynamic impact of CA stenosis on the hepatic vascularization and could participate in the pre-surgical work-up of duodenopancreatectomy (in which a ligation of the GDA is often mandatory) or liver transplantation^{7–9} in order to avoid unnecessary revascularization of a non-haemodynamically significant CA stenosis.

Figure 1. Variation of the diameter (mm) of the gastroduodenal artery (GDA), with the severity of coeliac axis stenosis (%).

Figure 2. Amplitude and phase-coded sequences focused on gastroduodenal artery (GDA; red circle) for two patients at inspiration and at expiration: one with no significant coeliac axis (CA) stenosis and one with degree of CA stenosis of 78%. The patient with no significant CA stenosis showed a physiological direction of GDA flow (black in phase-coded sequence, same as the abdominal aorta—blue arrowhead). The patient with 78% of CA stenosis showed an inversion in GDA flow at inspiration as well as at expiration.

CA Stenosis Degree	Sagittal CT View	Inspiration		Expiration	
		Amplitude	Phase	Amplitude	Phase
2%
78%

Currently, this pre-operative assessment is mainly based on morphological assessment of the degree of CA stenosis and diameter of the GDA, preferentially performed by CT.^{11,23–26} This led Sugae *et al*,¹⁰ based on a retrospective study comparing the pre-operative morphological grading of CA stenosis with required revascularization procedures during pancreatic surgery, to recommend: <50% stenosis, no additional procedure;

between 50% and 80%, release of the MAL if present; and >80% stenosis, reconstruction of the coeliac arterial axis. This recommendation could be modified according to our results. Indeed, we observed a reverse flow in the GDA for some CA stenoses <50% (starting from 35%) and a very variable pattern of flow between 34% and 60% stenosis. Furthermore, the flow was systematically inverted >60% CA stenosis.

Table 1. Results: values of peak velocity, mean velocity and flow in the three groups of coeliac axis (CA) stenosis

CA stenosis	<34%		34–60%		>60%	
Number of patients (n)	10		8		5	
Parameters	Inspiration	Expiration	Inspiration	Expiration	Inspiration	Expiration
Number of patients with reversed GDA flow (n)	0	0	3	4	5	5
Mean of maximum velocity (cm s^{-1})	−33.94 ^a	−31.41 ^a	−6.95 ^a	6.97	44.52	55.88
Mean of mean velocity (cm s^{-1})	−7.96 ^a	−8.4 ^a	0.35	4.55	11.22	16.98
Mean of mean flow (ml min^{-1})	−90 ^a	−140 ^a	10	10	400	440

GDA, gastroduodenal artery.

^aNegative values are related to an antegrade flow in the GDA, from the hepatic (merging from the CA) to the mesenteric artery.

Figure 3. Variation of gastroduodenal artery maximum velocity (cm s^{-1}), mean velocity (cm s^{-1}) and mean flow (ml min^{-1}) with degree of coeliac axis stenosis at inspiration and at expiration.

The linear relationship between CA stenosis and flow in the GDA suggests a progressive mechanism of adaptation of the flow to CA stenosis but with important interindividual variations between 34% and 60% stenosis. This highlights the necessity of measuring GDA flow, as the haemodynamic and anastomosis complexity can lead to different patterns of flow in GDA for mid-grade stenoses.

Doppler has been proposed to handle this task by providing haemodynamic information on the CA itself.^{22,27} However, to our knowledge, no studies of the GDA haemodynamic parameters using Doppler are available. Moreover, performing a Doppler examination is often difficult, with the GDA not or poorly seen in 81% of cases,²⁸ depending on the operator's

experience and the patient's echogenicity, whereas the rate of success for cine phase-contrast acquisitions in our study was 100%.

Another criterion to assess haemodynamically significant CA stenosis could be the diameter of the GDA, since flow is related to the vessel diameter. However, despite a significant relationship between the diameter of the GDA and degree of CA stenosis, we found a very weak correlation between the two parameters.

Cine phase-contrast MR technique has already been used for the quantification of abdominal arterial flow, mainly in renal artery stenosis evaluation.^{29–31} Mano et al³² used it successfully on GDA, to assess the haemodynamic changes of coeliacomesenteric anastomosis in patients with pancreaticoduodenal artery aneurysms and CA occlusion, also finding a GDA retrograde flow associated with severe CA stenosis.

Figure 4. Mean flow difference (ml min^{-1}) at inspiration and at expiration.

Because breathing changes the relative position of the CA and diaphragm, it increases the degree of CA stenosis in case of the MAL, the most common cause of CA stenosis. Nevertheless, we found no significant difference in GDA flow between inspiration and expiration, except for one of our patients, who presented a physiological flow direction at inspiration and an inversion at expiration. Since CA stenosis increases at expiration,²⁰ with an increase of 10% reported by Lee et al²¹ in a Doppler study, we recommend performing the phase-contrast MRI study only in expiration in order to decrease examination time.

Limitations

First, the spatial resolution of the phase-contrast MRI sequence was relatively low, resulting in difficulties for closely contouring the GDA during the flow measurement. This could be overcome

by decreasing the voxel size, which would, however, lead to an increase in acquisition time, incompatible with breath-holding. Furthermore, velocities were only encoded in one direction which can lead to errors in flow measurement. Four-dimensional phase-contrast MRI could be used to overcome this problem but again at the price of an increased acquisition time that is not suitable for clinical practice. Finally, we did not correlate the measurement using phase-contrast MRI with a reference standard such as intraoperative Doppler ultrasound due to the fact that these measurements were not achievable during surgery in our centre.

In conclusion, we report a significant and strong correlation ($r > 0.74$) between the degree of CA stenosis and the haemodynamic parameters in the GDA (flow, peak velocity and mean

velocity). Hence, phase-contrast MRI allows the evaluation of haemodynamic changes in the GDA induced by CA stenosis, mostly related to MAL compression. Moreover, our results show that despite a linear correlation between the degree of CA stenosis and haemodynamics in the GDA, interindividual variations are important for mid-grade stenosis, opening in the future to a specific study of this population.

Phase-contrast MRI could become an objective tool for predicting the necessity of a revascularization intervention during surgery by detecting haemodynamically significant stenoses if the limitation of acquisitions time can be overcome. Furthermore, it may help predict the potential indication of an additional surgical revascularization procedure during liver transplantation or supramesocolic surgery.

REFERENCES

- Jurim O, Shaked A, Kiai K, Millis JM, Colquhoun SD, Busuttil RW. Celiac compression syndrome and liver transplantation. *Ann Surg* 1993; **218**: 10–12. doi: <https://doi.org/10.1097/0000658-199307000-00003>
- Douard R, Ettorre GM, Chevallier JM, Delmas V, Cugnenc PH, Belghiti J. Celiac trunk compression by arcuate ligament and living-related liver transplantation: a two-step strategy for flow-induced enlargement of donor hepatic artery. *Surg Radiol Anat* 2002; **24**: 327–31.
- Bron KM, Redman HC. Splanchnic artery stenosis and occlusion. Incidence; arteriographic and clinical manifestations. *Radiology* 1969; **92**: 323–8. doi: <https://doi.org/10.1148/92.2.323>
- Horton KM, Talamini MA, Fishman EK. Median arcuate ligament syndrome: evaluation with CT angiography. *Radiographics* 2005; **25**: 1177–82.
- Nara S, Sakamoto Y, Shimada K, Sano T, Kosuge T, Takahashi Y, et al. Arterial reconstruction during pancreatoduodenectomy in patients with celiac axis stenosis—utility of Doppler ultrasonography. *World J Surg* 2005; **29**: 885–9. doi: <https://doi.org/10.1007/s00268-005-7878-x>
- Loukas M, Pinyard J, Vaid S, Kinsella C, Tariq A, Tubbs RS. Clinical anatomy of celiac artery compression syndrome: a review. *Clin Anat* 2007; **20**: 612–17. doi: <https://doi.org/10.1002/ca.20473>
- Kurosaki I, Hatakeyama K, Nihei K, Oyamatsu M. Celiac axis stenosis in pancreatoduodenectomy. *J Hepatobiliary Pancreat Surg* 2004; **11**: 119–24. doi: <https://doi.org/10.1007/s00534-003-0871-6>
- Vilatobá M, Zamora-Valdés D, Guerrero-Hernández M, Romero-Talamás H, Leal-Villalpando RP, Mercado MA. Arcuate ligament compression as a cause of early-onset thrombosis of the hepatic artery after liver transplantation. *Ann Hepatol* 2011; **10**: 88–92.
- Jiang ZJ, Liang TB, Feng XN, Wang WL, Shen Y, Zhang M, et al. Arcuate ligament syndrome inducing hepatic artery thrombosis after liver transplantation. *Hepatobiliary Pancreat Dis Int* 2008; **7**: 433–6.
- Sugae T, Fujii T, Kodera Y, Kanzaki A, Yamamura K, Yamada S, et al. Classification of the celiac axis stenosis owing to median arcuate ligament compression, based on severity of the stenosis with subsequent proposals for management during pancreatoduodenectomy. *Surgery* 2012; **151**: 543–9. doi: <https://doi.org/10.1016/j.surg.2011.08.012>
- Song SY, Chung JW, Kwon JW, Joh JH, Shin SJ, Kim HB, et al. Collateral pathways in patients with celiac axis stenosis: angiographic-spiral CT correlation. *Radiographics* 2002; **22**: 881–93.
- Agnes S, Avolio AW, Magalini SC, Frongillo F, Castagneto M. Celiac axis compression syndrome in liver transplantation. *Transplant Proc* 2001; **33**: 1438–9.
- Kohler TR, Debas H, Cramers M, Strandness DE. Pancreatoduodenectomy and the celiac artery compression syndrome. *Ann Vasc Surg* 1990; **4**: 77–80. doi: <https://doi.org/10.1007/BF02042695>
- Bull DA, Hunter GC, Crabtree TG, Bernhard VM, Putnam CW. Hepatic ischemia, caused by celiac axis compression, complicating pancreatoduodenectomy. *Ann Surg* 1993; **217**: 244–7. doi: <https://doi.org/10.1097/0000658-199303000-00005>
- Berney T, Pretre R, Chassot G, Morel P. The role of revascularization in celiac occlusion and pancreatoduodenectomy. *Am J Surg* 1998; **176**: 352–6. doi: [https://doi.org/10.1016/S0002-9610\(98\)00195-0](https://doi.org/10.1016/S0002-9610(98)00195-0)
- Pfeiffenberger J, Adam U, Drognitz O, Kröger JC, Makowiec F, Schareck W, et al. Celiac axis stenosis in pancreatic head resection for chronic pancreatitis. *Langenbecks Arch Surg* 2002; **387**: 210–15.
- Farma JM, Hoffman JP. Nonneoplastic celiac axis occlusion in patients undergoing pancreatoduodenectomy. *Am J Surg* 2007; **193**: 341–4; discussion 344. doi: <https://doi.org/10.1016/j.amjsurg.2006.09.027>
- AbuRahma AF, Stone PA, Srivastava M, Dean LS, Keiffer T, Hass SM, et al. Mesenteric/ celiac duplex ultrasound interpretation criteria revisited. *J Vasc Surg* 2012; **55**: 428–36. doi: <https://doi.org/10.1016/j.jvs.2011.08.052>
- Lotz J, Meier C, Leppert A, Galanski M. Cardiovascular flow measurement with phase-contrast MR imaging: basic facts and implementation. *Radiographics* 2002; **22**: 651–71.
- van Petersen AS, Meerwaldt R, Kolkman JJ, Huisman AB, van der Palen J, van Bockel JH, et al. The influence of respiration on criteria for transabdominal duplex examination of the splanchnic arteries in patients with suspected chronic splanchnic ischemia. *J Vasc Surg* 2013; **57**: 1603–11. doi: <https://doi.org/10.1016/j.jvs.2012.11.120>
- Lee VS, Morgan JN, Tan AGS, Pandharipande PV, Krinsky GA, Barker JA, et al. Celiac artery compression by the median arcuate ligament: a pitfall of end-expiratory MR imaging. *Radiology* 2003; **228**: 437–42. doi: <https://doi.org/10.1148/radiol.2282020689>
- Gruber H, Loizides A, Peer S, Gruber I. Ultrasound of the median arcuate ligament syndrome: a new approach to diagnosis. *Med Ultrason* 2012; **14**: 5–9.

23. Ilica AT, Kocaoglu M, Bilici A, Ors F, Bukte Y, Senol A. Median arcuate ligament syndrome: multidetector computed tomography findings. *J Comput Assist Tomogr* 2007; **31**: 728–31.
24. Gümüş H, Gümüş M, Tekbaş G, Önder H, Ekici F, Çetinçakmak MG, et al. Clinical and multidetector computed tomography findings of patients with median arcuate ligament syndrome. *Clin Imaging* 2012; **36**: 522–5.
25. Schaefer PJ, Pfarr J, Trentmann J, Wulff AM, Langer C, Siggelkow M, et al. Comparison of noninvasive imaging modalities for stenosis grading in mesenteric arteries. *Rofo* 2013; **185**: 628–34. doi: <https://doi.org/10.1055/s-0033-1335212>
26. Kopecky KK, Stine SB, Dalsing MC, Gottlieb K. Median arcuate ligament syndrome with multivessel involvement: diagnosis with spiral CT angiography. *Abdom Imaging* 1997; **22**: 318–20. doi: <https://doi.org/10.1007/s002619900199>
27. Erden A, Yurdakul M, Cumhuri T. Marked increase in flow velocities during deep expiration: a duplex Doppler sign of celiac artery compression syndrome. *Cardiovasc Intervent Radiol* 1999; **22**: 331–2. doi: <https://doi.org/10.1007/s002709900399>
28. Spaulding KA, Kissner ME, Kim EK, Pretorius DH, Rose SC, Garroosi K. Three-dimensional gray scale ultrasonographic imaging of the celiac axis: preliminary report. *J Ultrasound Med* 1998; **17**: 239–48.
29. Schoenberg SO, Knopp MV, Bock M, Kallinowski F, Just A, Essig M, et al. Renal artery stenosis: grading of hemodynamic changes with cine phase-contrast MR blood flow measurements. *Radiology* 1997; **203**: 45–53. doi: <https://doi.org/10.1148/radiology.203.1.9122415>
30. Schoenberg SO, Knopp MV, Londy F, Krishnan S, Zuna I, Lang N. Morphologic and functional magnetic resonance imaging of renal artery stenosis: a multireader tricenter study. *J Am Soc Nephrol* 2002; **13**: 158–69.
31. Silverman JM, Friedman ML, van Allan RJ. Detection of main renal artery stenosis using phase-contrast cine MR angiography. *AJR Am J Roentgenol* 1996; **166**: 1131–7. doi: <https://doi.org/10.2214/ajr.166.5.8615257>
32. Mano Y, Takehara Y, Sakaguchi T, Alley MT, Isoda H, Shimizu T, et al. Hemodynamic assessment of celiaco-mesenteric anastomosis in patients with pancreatoduodenal artery aneurysm concomitant with celiac artery occlusion using flow-sensitive four-dimensional magnetic resonance imaging. *Eur J Vasc Endovasc Surg* 2013; **46**: 321–8. doi: <https://doi.org/10.1016/j.ejvs.2013.06.011>