Published in final edited form as:

Clin Sci (Lond). 2015 September ; 129(6): 489–504. doi:10.1042/CS20140852.

Inhibition of the lymphocyte metabolic switch by the oxidative burst of human neutrophils

Philip A. Kramer^{*,†,§}, Lynn Prichard[‡], Balu Chacko^{*,†,§}, Saranya Ravi^{*,†,§}, E. Turner Overton^{‡,§}, Sonya L. Heath^{‡,§}, and Victor Darley-USmar^{*,†,§}

^{*}Mitochondrial Medicine Laboratory, University of Alabama at Birmingham, Birmingham, AL 35294

[†]Department of Pathology, University of Alabama at Birmingham, Birmingham, AL 35294

[‡]Department of Medicine- Infectious Diseases, University of Alabama at Birmingham, Birmingham, AL 35294

[§]Center for Free Radical Biology, University of Alabama at Birmingham, Birmingham, AL 35294

Abstract

Activation of the phagocytic NADPH oxidase (NOX-2) in neutrophils is a critical process in the innate immune system and is associated with elevated local concentrations of superoxide, hydrogen peroxide and hypochlorous acid. Under pathological conditions, NOX-2 activity has been implicated in the development of autoimmunity, indicating a role in modulating lymphocyte effector function. Notably, T cell clonal expansion and subsequent cytokine production requires a metabolic switch from mitochondrial respiration to aerobic glycolysis. Previous studies demonstrate that hydrogen peroxide generated from activated neutrophils suppresses lymphocyte activation but the mechanism is unknown. We hypothesized that activated neutrophils would prevent the metabolic switch and suppress the effector functions of T cells through a hydrogen peroxide-dependent mechanism. To test this, we developed a model co-culture system using freshly isolated neutrophils and lymphocytes from healthy human donors. Extracellular flux analysis was used to assess mitochondrial and glycolytic activity and FACS analysis to assess immune function. The neutrophil oxidative burst significantly inhibited the induction of lymphocyte aerobic glycolysis, caused inhibition of oxidative phosphorylation, and suppressed lymphocyte activation through a hydrogen peroxide-dependent mechanism. The impact of hydrogen peroxide on bioenergetics in the lymphocytes was confirmed using authentic reagent and a redox cycling agent. In summary, we have shown that the lymphocyte metabolic switch from mitochondrial respiration to glycolysis is prevented by the oxidative burst of neutrophils. This

Corresponding author: Victor Darley-USmar, Ph.D, Department of Pathology, University of Alabama at Birmingham, Biomedical Research Building II, 901 19th Street South, Birmingham, Alabama 35294, Tel: 205-975-9686, Fax: 205-934-1775, Darley@uab.edu.

Authorship:

PAK, LP, VDU experimental design, analysis of data, and interpretation of results and writing and revising manuscript; PAK conducted Seahorse experiments; LP and PAK conducted flow cytometry experiments; ETO and SH interpreted results and contributed to the writing and revising of manuscript. BC and SR contributed to the design and performance of experiments and manuscript revision.

Conflict of Interest Disclosure: Victor Darley-USmar is a recipient of funding from Seahorse Bioscience but this is not related to this study.

direct inhibition of the metabolic switch is then a likely mechanism underlying the neutrophil-dependent suppression of T cell effector function.

Keywords

T cells; hydrogen peroxide; glycolysis; oxidative phosphorylation

Introduction

Leukocyte and platelet metabolism have been linked to disease states with bioenergetic biomarkers being used to assess progression of human disease [1–4]. In addition, recent data indicate that monocytes and lymphocytes modulate both glycolysis and oxidative phosphorylation during activation of both the innate and adaptive immune systems [5–7]. The metabolism of circulating neutrophils, monocytes, lymphocytes and platelets are distinctive, and these differences can be used to evaluate of the relationship between bioenergetics and function [2, 8]. Under basal conditions, neutrophils possess minimal, if any, mitochondrial function and meet their energy requirements through glycolysis [2, 8, 9]. In contrast, lymphocytes isolated from peripheral blood, primarily utilize oxidative phosphorylation for energy production [2]. However, lymphocyte metabolic pathways are highly adaptive. Notably, the transition of lymphocytes from a naïve to an activated state requires a metabolic switch from oxidative phosphorylation to aerobic glycolysis [10, 11]. Similarly, this change is also observed in many highly proliferative cells, and is thought to provide the necessary energy and molecular building blocks for DNA, protein, and lipid biosynthesis as well as cell signaling [11]. Disruption of the metabolic switch in T lymphocytes has been shown to inhibit cellular clonal expansion and cytokine production [12, 13]. Furthermore, in contrast to regulatory and memory T cells, glycolysis is required to support T cell effector cell function including cytokine production and proliferation [10, 11, 13, 14].

At sites of inflammation, CD4 (Th1 and Th2) and CD8 (cytotoxic) effector T cells are recruited to combat viral or bacterial infection. Neutrophils are key components of innate immunity, and use reactive oxygen species (ROS) to eliminate pathogens. This is a coordinated process which also involves extrusion of neutrophil extracellular traps (NETs) which aid in the clearance of pathogens by direct histone, chromatin, and antimicrobial protein interaction [15–17]. T cells recognize peptides presented by MHC class 1 and 2 molecules from the host cells as part of the T cell surveillance program. T cells actively recruit macrophages and neutrophils to the site of inflammation as well as facilitate B cell antibody production. In the absence of T regulatory cells or their associated cytokines (TGF- β , IL-4 and IL-10), T effector cells can promote chronic inflammation and even autoimmunity. While reactive oxygen species (ROS) derived from the neutrophil oxidative burst have long been considered a promoter of inflammation and disease progression, new evidence suggests that these ROS have an immunoregulatory role [18–21]. The production of superoxide from NADPH oxidase (NOX-2) generates high μM concentrations of superoxide and hydrogen peroxide (H_2O_2) which then raises the interesting possibility that NOX-2 dependent ROS formation is an essential regulatory modulator of T-cell function. On

the other hand this appears paradoxical since exposure of a broad range of eukaryotic cells to H₂O₂, including those in the vasculature and the cardiomyocyte, results in damage to key metabolic pathways including oxidative phosphorylation and glycolysis [22–26]. Importantly, this has not been examined in lymphocytes which are frequently in close proximity to neutrophils at sites of inflammation [19].

Neutrophil-lymphocyte interactions have been investigated in the context of cytokine production, antigen presentation, and oxidant generation and some studies suggest that levels of the oxidative burst which are too low fail to regulate the T-cell effector function and can thereby contribute to the development of autoimmune disease [19–21, 27]. In fact, many autoantibodies associated with Systemic Lupus Erythematosus (SLE) are targeted towards neutrophilic debris and associated double-stranded DNA [15]. In support of this concept, decreasing the neutrophil oxidative burst has effectively inhibited NETosis but has failed to improve SLE outcomes in animal models [28]. Further, some patients with Chronic Granulomatous Disease (CGD), who have a defective oxidative burst caused by mutations in the NCF1 gene which encodes the p47 protein in the NADPH oxidase enzyme, were shown to be more susceptible to autoimmune conditions and have an elevated inflammatory state [29, 30]. T cell driven pathologies such as Rheumatoid Arthritis and Encephalomyelitis were enhanced in mouse models with NCF1 mutations [18]. The same group suppressed arthritis severity by addition of compounds that restored the neutrophil oxidative burst [31]. Finally, the suppressive effects of some neutrophil subsets have been shown to contribute to the pathogenesis of HIV, particularly in the setting of advanced AIDS [27, 32]. These studies and others suggest a direct immunoregulatory role of neutrophil derived ROS on lymphocytes; however, a mechanism has yet to be determined [33–35]. Taken together these data led us to hypothesize that ROS from the neutrophil oxidative burst modulates lymphocyte effector function through its inhibitory effects on the metabolic switch from oxidative phosphorylation to glycolysis. This immunoregulatory effect of neutrophil derived ROS may then be essential to the prevention of chronic inflammation and autoimmunity, especially in the presence of neutrophil necrotic debris and NETs. Here we utilize extracellular flux technology in a novel co-culture system to investigate, for the first time, the metabolic adaptations and perturbations in lymphocytes cultured with neutrophils.

Methods

Isolation and plating of human lymphocytes and neutrophils

All protocols and procedures for the collection, isolation, analysis, and storage of blood or its components have been reviewed and approved by the Institutional Review Board at the University of Alabama at Birmingham. Neutrophils and lymphocytes were isolated from whole blood of healthy individuals as previously described [36]. Plating of the cells following cell counting and suspension was performed in Extracellular Flux assay medium (XF-DMEM). Lymphocytes were plated at 15×10^4 cells/well in the Seahorse 96 well microplate coated with Cell-Tak (CB-40242, Fisher). Neutrophils were added in co-culture at the cell densities specified. As there is some significant variation in the degree of the oxidative burst and mitochondrial function between donors, representative data from a single donor is reported for each experiment presented in this study. Key findings were verified on

a minimum of 3 donors with 3–6 technical replicates per experiment. Overall, cells were obtained from 10 healthy individual donors.

Neutrophil and lymphocyte immune function assessment

Neutrophils were plated at 7.5×10^4 cells per well and PMA injected in the Extracellular flux analyzer and the oxidative burst measured as described below. Three hours after PMA treatment, 100ul of XF-DMEM media was replaced with media containing 2μM SytoxGreen (S-7020, Life technologies). NETosis in neutrophils was monitored by SytoxGreen fluorescence after 10 min at 37°C using 480nm/530nm (excitation/emission) on a microplate reader (PerkinElmer).

Lymphocyte survivability, cytokine production, and clonal expansion were monitored by FACs analysis after an initial stimulation with 100ng/ml PMA for the neutrophil oxidative burst on non-CellTak coated Seahorse Extracellular flux microplates. Lymphocytes were washed from the plate by gentle pipetting, centrifuged, and resuspended in 500ul R10 media, with PMA and Ionomycin added at the above concentrations at 37°C and 5% humidified CO₂ for 4 days. Golgistop and GolgiPlug (10 μg/ml; BD Biosciences) were added during the last 12 hours. Cells were harvested and washed once with PBS before being labeled with fluorescent LIVE/DEAD fixable dead cell dye (Molecular Probes, Invitrogen). Fluorochrome conjugated monoclonal antibodies antiCD-3 Alexa 780 (Clone: UCHT1), CD8 V500 (Clone: RPA-T8), and CD4 Qdot 655 (Clone: S3.5) (BD Biosciences) were used for surface staining. Following fixation and permeabilization with Cytotfix and Cytoperm (BD Biosciences), cells were washed and stained with intracellular markers IFN-γ Alexa Fluor 700 (Clone: B27), and IL-2 PE (Clone: MQ1-17H12) (BD Biosciences). Following staining, cells were washed and fixed in 2% paraformaldehyde (Sigma-Aldrich) and analyzed on an LSRII flow cytometer within 24 h (BD Biosciences).

At least 100,000 CD3+ events were acquired from each lymphocyte sample. Data analysis was performed using FlowJo version 9.7.6 software (Tree Star). CD3+ lymphocytes were gated based on forward and side scatter properties after the exclusion of doublets. Gates were set relative to positive controls and negative controls.

Lymphocyte clonal expansion was assessed by labeling with 1.25 μm CFSE (Molecular Probes, Eugene OR) for 4 min at room temp. After washing in PBS, cells were resuspended in 1 ml of complete RPMI with 10% Ab serum. Phorbol-12-myristate-13-acetate (PMA), a Protein Kinase C activator, and the calcium ionophore, ionomycin, were added at indicated concentrations and the lymphocytes were incubated 4 days at 37° C and 5% CO₂. On the 5th day, the cells were centrifuged and resuspended in 1 ml RPMI with 10% Ab serum with GolgiPlug and GolgiStop and more PMA and Ionomycin. They were incubated an additional 6 hours at 37°C then kept overnight at 4°C. Surface and intracellular staining followed by flow cytometric analysis was performed as above.

Assessment of bioenergetic function and the Oxidative Burst

The induction of aerobic glycolysis in lymphocytes and the oxidative burst in neutrophils was accomplished by the addition of 100ng/ml phorbol-12-myristate-13-acetate (PMA) or as otherwise indicated. PMA was loaded into port A at a 10x concentration and was injected

after three basal oxygen consumption and extracellular acidification rate readings (OCR/ECAR). OCR and ECAR measurements were taken every 8 min until the oxidative burst response of neutrophils ceased, approximately 160 min, and then the bioenergetic profile of lymphocytes was obtained. The bioenergetic profile consists of basal OCR/ECAR measurements followed by the injection of 1 µg/ml Oligomycin, 0.6 µM FCCP, and 10 µM Antimycin A as previously described [36, 37]. Briefly, oligomycin inhibits mitochondrial ATP synthase and the resulting drop in OCR and rise in ECAR are attributed to ATP-linked OCR and the compensation of glycolysis for the loss of mitochondrial ATP production. The protonophore, FCCP, uncouples the mitochondrial proton gradient and oxygen consumption from ATP synthase and drives maximal OCR. Antimycin A inhibits complex III of the electron transport chain and suppresses all mitochondrial oxygen consumption and the remaining OCR is considered non-mitochondrial [38].

In the co-culture experiment cells were centrifuged on to the Cell Tak-coated plate at 200g, the plate rotated, and then centrifuged at 300g as previously described [36]. The volume of the well was brought up to 180 µl with XF-DMEM and the plate incubated for 10–30 min in a non-CO₂ incubator. The Seahorse cartridge was loaded at 10x the final concentration of 100 ng/ml PMA in port A and 1 µg/ml oligomycin, 0.6 µM FCCP, and 10 µM Antimycin A in ports B, C, and D, respectively. The effect of catalase (C9322, Sigma-Aldrich) on lymphocyte bioenergetics was assessed by incubation at 1000 U/ml for 30 min prior to inducing the oxidative burst of neutrophils.

Glycolytic function of lymphocytes in co-culture was determined by the glucose stress test [38]. Glucose is essential for the oxidative burst of neutrophils, and for that reason, the lymphocyte and neutrophils were treated with PMA outside of the Extracellular Flux Analyzer for a period of 160 min before the media was exchanged with XF-DMEM containing no glucose or pyruvate. The media exchange consisted of two washes of replacement medium to effectively lower glucose to <1% of its original concentration. The cells were allowed 20 min to equilibrate before being placed on the XF analyzer. Three basal measurements were obtained before the injection of 5 mM glucose, 1 µg/ml oligomycin, and 100 mM 2-deoxy-glucose (2DG). Glucose stimulated ECAR was termed glycolysis, while oligomycin sensitive ECAR represented the glycolytic reserve. 2DG, the competitive inhibitor of hexokinase, inhibited all glycolytic activity and the remaining ECAR was termed non-glycolytic.

Lymphocytes were exposed to bolus hydrogen peroxide and the intracellular redox cycling agent, 2,3-Dimethoxy-1,4-naphthoquinone (DMNQ) to determine the bioenergetic consequences of hydrogen peroxide in a neutrophil-free system [37]. Lymphocytes were isolated and plated as previously described and 50, 100, and 500 µM bolus hydrogen peroxide was added to the cells for 30 minutes at 37°C prior to the first assay measurement. A 1, 3, and 5 µM DMNQ dose response was obtained under the same conditions prior to PMA injection and bioenergetics assessment.

Statistics

All OCR/ECAR traces and analyses were analyzed for each representative donor with 3–6 replicate wells. The data are presented as mean $\pm$ S.E.M. A Student's t-test was used to determine statistical significance (P 0.05) using the standard Excel statistics package.

Results

Isolation and characterization of lymphocyte and neutrophils

To assess the ability of isolated neutrophils to undergo an oxidative burst and NETosis, CD15⁺ neutrophils were isolated from whole blood of healthy volunteers as previously described [36]. The conditions required to elicit PMA-dependent oxidative burst, measured as the amount of oxygen consumed over 100 min, and NET formation, measured as Sytoxgreen fluorescence, in neutrophils were first established (Figure 1A, B). The concentration of PMA required to elicit the maximal oxygen consumption was 50–100ng/ml and this corresponded well with the extent of NETosis (Figure 1B) as reported in the literature [39]. PMA was selected for these studies because of its consistency in stimulation of NOX-2 and because it is one of the few agents which can activate both neutrophils and lymphocytes. Negatively selected lymphocytes were activated to assess cytokine production and clonal expansion, key characteristics of activated lymphocytes. Lymphocytes cultured for 4 days with 50ng/ml PMA and 100ng/ml Ionomycin showed an increase in IFN- γ production as compared to the media control (Figure 1C). Lymphocyte proliferation was measured using the cell permeable stain, CFSE, and demonstrated nearly 100% proliferation of lymphocytes as seen by loss of CFSE (Figure 1D).

Lymphocyte and neutrophil oxidative and glycolytic metabolism

The lymphocyte 'metabolic switch' from oxidative phosphorylation to aerobic glycolysis was determined in cell culture using Extracellular Flux analysis [40]. In panel 2A the OCR and in Panel 2C ECAR in lymphocytes alone are shown with and without the addition of PMA (100ng/ml). Within 8 min of addition of PMA, both OCR and ECAR are stimulated. These effects are sustained for the subsequent 160 min after which a mitochondrial stress test was performed. Oligomycin, which inhibits the mitochondrial ATP synthase, resulted in the expected decrease in OCR which was stimulated on the addition of the uncoupler, FCCP. The addition of Antimycin A inhibits all mitochondrial respiration and shows a slightly elevated non-mitochondrial respiration in PMA-treated lymphocytes. The addition of oligomycin stimulates ECAR with or without PMA. Interestingly, the overall glycolytic capacity is greater after addition of PMA consistent with increased glycolytic flux. The modest stimulation of ECAR on addition of FCCP and its inhibition by Antimycin A occurs to the same extent with or without PMA and most likely represents proton production from the TCA cycle. These data indicate that the lymphocytes are metabolically active and demonstrate the anticipated switch to aerobic glycolysis on addition of PMA.

Similarly, neutrophil OCR and ECAR were measured over the same time course (Figure 2B, D). As we have reported previously, neutrophils have minimal mitochondrial function and utilize a small amount of glycolysis under the basal state and show no significant response to the addition of mitochondrial inhibitors [2, 8]. PMA induces a rapid increase in OCR which

is due to the activation of NOX2 and the production of superoxide which is maximal at approximately 30 min after injection of PMA and progressively decreases over the subsequent 100 min [8]. Incubating the neutrophils or lymphocytes in glucose free medium or with the competitive inhibitor of hexokinase, 2-deoxy-glucose, resulted in complete suppression of the oxidative burst and glucose oxidation assessed by the ECAR measurement (data not shown).

As reported in the literature, low dose PMA and ionomycin had a synergistic effect on inducing lymphocyte activation [5, 41]. However, ionomycin led to a transient elevation in ECAR and OCR in lymphocytes, and resulted in loss of maximal mitochondrial function (Supplementary Figure 1). Additionally, ionomycin completely suppressed the oxidative burst of neutrophils (data not shown). While ionomycin and PMA are often used together to activate lymphocytes, these data demonstrate that the calcium ionophore suppresses h metabolic processes in both cell types. Therefore, in this study, PMA alone was used to stimulate lymphocyte and neutrophil activation unless otherwise specified.

Lymphocyte metabolic function in the presence of activated neutrophils

To determine the effect of neutrophil activation on lymphocyte metabolism, a co-culture method was developed. Neutrophils were plated at 5–75K cells/well on the Extracellular flux microplate with 150K lymphocytes. Neutrophils (25k) co-cultured with lymphocytes without PMA had no effect on basal lymphocyte ECAR or OCR (data not shown). After basal OCR/ECAR was established, PMA was injected and the cells monitored for 160 min. With the addition of PMA, lymphocyte OCR and ECAR was stimulated as observed with either cell type alone (Figure 3A, B). At 160 min after PMA injection, at which point oxygen consumption and extracellular acidification due to the oxidative burst was complete, the mitochondrial stress test was performed as shown in Figure 3C. It is important to note that at the time of assessing lymphocyte mitochondrial function the neutrophils are not responsive to the mitochondrial inhibitors and the oxidative burst is essentially complete. The detailed analysis of the lymphocyte OCR and ECAR parameters are shown in Figure 4. The activated neutrophils decreased basal, ATP linked, maximal and reserve capacity with a significant effect detected for 10–25k neutrophils (i.e. 1–2.5 neutrophils for every 15 lymphocytes). The PMA and oligomycin sensitive ECAR for the lymphocytes was sensitive to activated neutrophils over a similar range of cell densities (Figure 4E, F).

Activated neutrophils release a broad range of mediators on activation including proteases and ROS. To determine whether hydrogen peroxide (H_2O_2) is mediating the effects on ECAR and OCR in the co-culture of neutrophils (25k) and lymphocytes, 1000U/ml of catalase, which removes H_2O_2 , was added 30 min prior to the start of the assay and was present for the duration of the oxidative burst. Figure 5A and 5B show the PMA-stimulated lymphocyte OCR and ECAR following the oxidative burst in the presence of neutrophils. Interestingly, the extent of the oxidative burst measured by OCR and ECAR was prolonged in the presence of catalase, suggesting the H_2O_2 is also damaging the oxidative burst components in the neutrophil. The decline in mitochondrial and glycolytic function following the oxidative burst as shown in Figure 4 was clearly evident and prevented by

catalase (Figure 5C). In contrast, superoxide dismutase (SOD-100U/ml) administration failed to prevent the metabolic dysfunction in the lymphocytes (data not shown).

The extracellular acidification arises from any processes in the cell generating protons that are released and change the pH of the medium. To determine the contribution of glycolysis to these values the glycolytic stress test was performed in the neutrophil and lymphocyte co-cultures with PMA and the media replaced for the glucose stress test at 160 min. In this assay, glucose is omitted from the medium and three basal measurements are obtained (Figure 6A). Next, 5mM glucose is returned to the media in the first injection and results in a rapid increase in ECAR in the control lymphocytes treated with PMA which is suppressed in a neutrophil dependent fashion. Addition of oligomycin further stimulates glycolysis as mitochondrial ATP production is inhibited and addition of the inhibitor of hexokinases 2-DG suppresses the ECAR to basal levels prior to the addition of glucose. Glycolysis was measured as the difference between ECAR after glucose injection and the non-glycolytic rate following 2-DG. Glycolytic reserve is the oligo-sensitive ECAR. Increasing neutrophil number significantly suppressed aerobic glycolysis and the glycolytic reserve in lymphocytes (Figure 6A, B). Notably, the neutrophil-dependent inhibition of glycolysis was prevented by catalase (Figure 6C) suggesting that this process is H_2O_2 mediated.

Lymphocyte metabolic function in a neutrophil free system with bolus hydrogen peroxide or DMNQ

To assess the metabolic regulation of H_2O_2 in the absence on neutrophils, lymphocytes were exposed to 50, 100, and 500 μ M bolus doses 30 min prior to the first OCR/ECAR measurement. The ability of PMA to increase basal OCR and ECAR was suppressed in a dose dependent manner with 100 μ M H_2O_2 causing suppression of ATP-linked OCR, reserve capacity, and oligomycin-sensitive ECAR to a similar degree as the same subject's neutrophils (25k) (Figure 7). Interestingly, the H_2O_2 produced by 25k neutrophils (40 μ M) was approximately twice as effective as the bolus reagent in inducing inhibition of lymphocyte bioenergetics. To address if this difference was due to proximity of H_2O_2 production 2,3-Dimethoxy-1,4-naphthoquinone (DMNQ), an intracellular generator of superoxide and hydrogen peroxide was exposed to lymphocytes 30 min prior to the assay. No significant increase in OCR was observed over the untreated control, indicating that, at the highest dose (5 μ M), low levels of these oxidants were being generated [37]. However, this still resulted in significant suppression of basal, PMA-sensitive, and oligomycin-sensitive changes in OCR and ECAR. This suggests that the modulation of lymphocyte metabolism by the neutrophil oxidative burst is particularly sensitive to local concentration of H_2O_2 .

Perturbation of lymphocyte metabolism coincides with immune dysfunction

FACs analysis was used to elucidate the functional consequences of the neutrophil oxidative burst on lymphocyte cytokine production and proliferation after 4 days. Lymphocytes and neutrophils were plated in co-culture as described above on non-Cell Tak coated microplates to maintain the same cell conditions and proximity as implemented during the bioenergetics assessments. After 160 min PMA treatment lymphocytes were washed from the plate and centrifuged to remove neutrophil cellular debris and treated with PMA and ionomycin as

described above. Following the 4 day incubation, 11% of CD3+ lymphocytes were positive for high IFN- γ production compared to control (Figure 8). Cytokine production was reduced to 8% when lymphocytes were co-cultured with 2.5×10^4 (25k) activated neutrophils (Figure 8D). Increasing neutrophil (50–75k) number lead to a greater and significant dose dependent suppression of IFN- γ production (data not shown). Catalase markedly inhibited the neutrophil-dependent suppression of lymphocyte IFN- γ production and had a moderate stimulatory effect on IFN- γ production in activated lymphocytes (Figure 8F).

Clonal expansion was stimulated with PMA/Ionomycin and measured by CFSE staining 4 days later. Figure 9 demonstrates a representative FACs profile showing significant lymphocyte proliferation in all groups compared to media alone; however a neutrophil-dependent decline in lymphocyte proliferation was observed (Figure 9C, D). This effect was prevented by addition of 1000U/ml catalase during the 160 min neutrophil oxidative burst at the start of the 4 day incubation (Figure 9F). Interestingly, the short duration of catalase exposure also stimulated lymphocyte proliferation, which we ascribe to scavenging of endogenously generated hydrogen peroxide (Figure 9F).

Discussion

Neutrophils have a life span measured in hours to days before they migrate through the vasculature in response to local inflammation, undergo programmed cell death and NETosis, or are cleared by the spleen or phagocytosis [42, 43]. As expected the neutrophil oxidative burst and NETosis occur rapidly after PMA stimulation under the conditions used for bioenergetic measurements (Figure 1A, B). These processes are associated with chronic inflammation and the generation of autoantibodies recognizing NET DNA or other neutrophil intracellular components [15, 44, 45]. The role of neutrophils in chronic inflammation and autoimmunity appears to be highly dependent on their interaction with effector T cells [19]. Monocytes are the typical means by which lymphocytes are activated, and are also capable of generating an oxidative burst, though to a lesser extent than neutrophils. Upon MHC-T cell receptor interaction and antigen recognition, naïve lymphocytes activate, proliferate, and secrete cytokines, or undergo cytotoxic degranulation. PMA can stimulate lymphocyte activation in the absence of antigenic stimuli as evidenced by the increased IFN- γ production and cell proliferation (Figure 1C, D). Neutrophils and lymphocytes have been shown to co-exist at sites of inflammation as well as lymphoid organs resulting in decreased lymphocyte activation and proliferation [19, 44]. This suggests that there are mechanisms through which T cell activation is inhibited in the presence of activated neutrophils, and possibly monocytes, which may be important in preventing the development of autoimmune conditions or controlling inflammation.

Aerobic glycolysis is a means by which highly proliferating cells can generate ATP and the molecular building blocks for cell growth and DNA synthesis. High-dose PMA stimulated aerobic glycolysis in freshly isolated lymphocytes within 8 min and was stable for over two hours as seen by the rapid rise in ECAR in Figure 2 and 3. This rapid induction is likely associated with the reported Glut1 trafficking to the cell surface through PI3K/AKT and/or MAPK-dependent signaling with possible activation of glycolytic enzymes (e.g. HK, IPFK) [11, 46–48]. The role of PMA in lymphocyte activation and the induction of aerobic

glycolysis, however, is not well understood. A significant increase in OCR is observed after PMA stimulation, which is consistent with the increased energetic demand associated with activation (Figure 2). PMA was used in place of other stimuli in part to demonstrate that many of these metabolic changes occur quickly and independently of nuclear transcription, and to disentangle the confounding roles of cell surface receptor expression and cytokine signaling which would ultimately depend on the inflammatory environment.

Lymphocyte mitochondrial and glycolytic function showed a >50% decrease in activity when 25K neutrophils were present, (a 1:6 ratio of neutrophils to lymphocytes) suggesting a relatively small number of activated neutrophils can impact the metabolism of lymphocytes and perhaps other cell types in close proximity (Figure 4). The oxidant, H_2O_2 , is generated rapidly from the dismutation of superoxide during the neutrophil oxidative burst. Hydrogen peroxide is a non-radical species that can freely diffuse through membranes, and inhibit both glycolytic and mitochondrial machinery [25, 26]. Catalase prevented the neutrophil-mediated bioenergetic dysfunction and allowed lymphocytes to undergo the metabolic switch even in the presence of activated neutrophils (Figure 5, 6). Bolus H_2O_2 effectively suppressed oxidative and glycolytic metabolism in a neutrophil-free system (Figure 7). Interestingly, assuming oxygen consumption resulted in a molar equivalent H_2O_2 production by the neutrophil oxidative burst, the total amount of H_2O_2 generated was less than half that provided by bolus H_2O_2 to achieve the same effect on lymphocyte bioenergetics. Similarly, the intracellular redox cycling agent, DMNQ, modulated lymphocyte metabolism to a similar degree as neutrophils in a dose dependent manner (Figure 7). The possibility that neutrophils could facilitate a much greater degree of metabolic suppression due to the conversion of hydrogen peroxide to hypochlorous acid (HOCL) by Myeloperoxidase was also considered, however the HOCL scavenger, taurine (5mM), did not prevent inhibition of lymphocyte bioenergetics (data not shown).

Lymphocyte activation and modulation of the glycolysis is essential for effector cell function. Recently GAPDH, a glycolytic enzyme, was shown to post-transcriptionally regulate IFN- γ expression by direct binding of mRNA when engagement of glycolysis was blocked [13]. The same group indicated that T-cell proliferation could precede with ATP obtained from oxidative phosphorylation or aerobic glycolysis, depending on the substrates available (e.g. galactose vs. glucose). The neutrophil dependent ROS formation resulted in modulation of both mitochondrial function and glycolysis and was associated with changes in T-effector cell function and proliferation (Figure 8, 9).

Clonal expansion and cytokine production reflect the ability of lymphocytes to respond to an activation stimulus and perform their immune-modulating effects locally and systemically. IFN- γ production is stimulated upon lymphocyte activation with PMA; however, 25k neutrophils were able to suppress IFN- γ production by >20% (Figure 8C, D). Lymphocyte proliferation is suppressed after only a short exposure to neutrophil-derived H_2O_2 (Figure 9C, D). Interestingly, catalase treatment in the absence of neutrophils increases IFN- γ production and proliferation (Figure 8E, F; 9E, F). This is consistent with studies showing that a ROS generating system, possibly a NADPH oxidase, is present and functional in activated lymphocytes which would allow for auto-suppression of effector immune functions [31, 49]. Indeed, the immediate increase in OCR observed with PMA in lymphocytes alone

appears to be due to an increasing non-mitochondrial oxygen consumption, consistent with activation of an NADPH oxidase. Furthermore, catalase treatment reversed the inhibitory effects of the neutrophil oxidative burst on lymphocyte IFN- γ production and caused an even greater degree of proliferation, further supporting this hypothesis. To determine if endotoxin contamination could also be the cause, a Limulus ameobocyte lysate (LAL) was performed and revealed a negligible amount of endotoxin (35pg) exposure to the cells (data not shown). Furthermore, endotoxin is not a superantigen and requires an antigen presenting cell for lymphocyte activation, which is absent in our co-culture system. Taken in the context of the metabolic switch, and their suppressed cytokine production, these data indicate that H₂O₂ is regulating the immune response by preventing these cells from undergoing a metabolic switch to support these essential effector functions.

In conclusion, impaired metabolism and reduced activation potential in lymphocytes is mediated by H₂O₂ generated from the neutrophil oxidative burst. The relative proportions of neutrophils and lymphocytes at the site of inflammation are subject to duration, severity, and type of infection [19]. Our study shows the coincident disruption of the metabolic switch and loss of activation, prompting future studies to investigate the specific metabolic targets. This mechanism may be in place to prevent the unrestrained activation of lymphocytes in the presence of neutrophil necrotic debris, and thus any impaired oxidative burst capacity could enhance lymphocyte activation and so contribute to chronic inflammation and autoimmunity.

Clinical Perspectives

This study was undertaken to discern the means by which neutrophils suppress lymphocyte immune function, an interaction which is becoming increasingly appreciated in current literature. Here we demonstrate that NADPH oxidase-derived hydrogen peroxide from activated neutrophils results in significant mitochondrial and glycolytic dysfunction in lymphocytes. Disruption of the lymphocytes metabolic switch from oxidative phosphorylation to a more glycolytic metabolism corresponds to decreased cytokine production and clonal expansion. This may be an important mechanism by which neutrophil-derived oxidative signaling regulates lymphocyte effector function, a process which may be essential to suppressing the inflammation associated with autoimmune disorders and chronic inflammation.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Funding:

The authors appreciate support from the UAB CFAR Flow Core (NIH grant P30 AI027767) and funding from NIH T32 T32HL07918 (PAK), and the O'Brien Center P30 DK079337.

Abbreviations

2DG

2-deoxy-glucose

AIDS	Acquired Immunodeficiency Syndrome
ATP	Adenosine triphosphate
CO₂	Carbon dioxide
CFSE	Carboxyfluorescein succinimidyl ester
CGD	Chronic Granulomatous Disease
DNA	Deoxyribonucleic acid
DMNQ	2,3-Dimethoxy-1,4-naphthoquinone
ECAR	Extracellular Acidification Rate
XF-DMEM	Extracellular Flux assay medium
FCCP	Trifluorocarbonyl cyanide Phenylhydrazine
Glut1	Glucose transporter 1
HK	Hexokinase
HIV	Human Immunodeficiency Virus
HLADR	Human leukocyte antigen-DR
H₂O₂	Hydrogen peroxide
HOCl	Hypochlorous acid
IPFK	Inducible Phosphofructokinase
IFN-γ	Interferon gamma
IL-2	Interleukin 2
MHC	Major Histocompatibility Complex
mRNA	Messenger Ribonucleic acid
MAPK	Mitogen-activated protein kinases
NOX-2	NADPH oxidase-2
NK	Natural Killer Cells
NCF1	Neutrophil Cytosolic Factor 1
NET or Netosis	Neutrophil extracellular trap
OCR	Oxygen Consumption Rate
PMA	Phorbol 12-myristate 13-acetate
PBS	Phosphate buffered saline

PI3K/AKT	Phosphoinositide 3-kinase/AKT
ROS	Reactive oxygen species
SLE	Systemic Lupus Erythematosus
SOD	Superoxide dismutase
Th1 and Th2	T-Helper cells 1 and 2

References

1. Ravi S, Mitchell T, Kramer PA, Chacko B, Darley-USmar VM. Mitochondria in monocytes and macrophages-implications for translational and basic research. *The international journal of biochemistry & cell biology*. 2014; 53C:202–207.
2. Kramer PA, Ravi S, Chacko B, Johnson MS, Darley-USmar VM. A review of the mitochondrial and glycolytic metabolism in human platelets and leukocytes: Implications for their use as bioenergetic biomarkers. *Redox Biology*. 2014; 2:206–210. [PubMed: 24494194]
3. Zharikov S, Shiva S. Platelet mitochondrial function: from regulation of thrombosis to biomarker of disease. *Biochem Soc Trans*. 2013; 41:118–123. [PubMed: 23356269]
4. Chacko BK, Kramer PA, Ravi S, Benavides GA, Mitchell T, Dranka BP, Ferrick D, Singal AK, Ballinger SW, Bailey SM, Hardy RW, Zhang J, Zhi D, Darley-USmar VM. The Bioenergetic Health Index: a new concept in mitochondrial translational research. *Clinical science*. 2014; 127:367–373. [PubMed: 24895057]
5. van der Windt GJ, O'Sullivan D, Everts B, Huang SC, Buck MD, Curtis JD, Chang CH, Smith AM, Ai T, Faubert B, Jones RG, Pearce EJ, Pearce EL. CD8 memory T cells have a bioenergetic advantage that underlies their rapid recall ability. *Proc Natl Acad Sci U S A*. 2013; 110:14336–14341. [PubMed: 23940348]
6. Pearce EL, Poffenberger MC, Chang CH, Jones RG. Fueling immunity: insights into metabolism and lymphocyte function. *Science*. 2013; 342:1242454. [PubMed: 24115444]
7. Pearce EL, Pearce EJ. Metabolic pathways in immune cell activation and quiescence. *Immunity*. 2013; 38:633–643. [PubMed: 23601682]
8. Chacko BK, Kramer PA, Ravi S, Johnson MS, Hardy RW, Ballinger SW, Darley-USmar VM. Methods for defining distinct bioenergetic profiles in platelets, lymphocytes, monocytes, and neutrophils, and the oxidative burst from human blood. *Lab Invest*. 2013; 93:690–700. [PubMed: 23528848]
9. Maiani NA, Geissler J, Srinivasula SM, Alnemri ES, Roos D, Kuipers TW. Functional characterization of mitochondria in neutrophils: a role restricted to apoptosis. *Cell Death Differ*. 2004; 11:143–153. [PubMed: 14576767]
10. Gerriets VA, Rathmell JC. Metabolic pathways in T cell fate and function. *Trends in immunology*. 2012; 33:168–173. [PubMed: 22342741]
11. MacIver NJ, Michalek RD, Rathmell JC. Metabolic regulation of T lymphocytes. *Annual review of immunology*. 2013; 31:259–283.
12. Cham CM, Driessens G, O'Keefe JP, Gajewski TF. Glucose deprivation inhibits multiple key gene expression events and effector functions in CD8+ T cells. *European journal of immunology*. 2008; 38:2438–2450. [PubMed: 18792400]
13. Chang CH, Curtis JD, Maggi LB Jr, Faubert B, Villarino AV, O'Sullivan D, Huang SC, van der Windt GJ, Blagih J, Qiu J, Weber JD, Pearce EJ, Jones RG, Pearce EL. Posttranscriptional control of T cell effector function by aerobic glycolysis. *Cell*. 2013; 153:1239–1251. [PubMed: 23746840]
14. Sukumar M, Liu J, Ji Y, Subramanian M, Crompton JG, Yu Z, Roychoudhuri R, Palmer DC, Muranski P, Karoly ED, Mohny RP, Klebanoff CA, Lal A, Finkel T, Restifo NP, Gattinoni L. Inhibiting glycolytic metabolism enhances CD8+ T cell memory and antitumor function. *The Journal of clinical investigation*. 2013; 123:4479–4488. [PubMed: 24091329]

15. Kaplan MJ. Role of neutrophils in systemic autoimmune diseases. *Arthritis research & therapy*. 2013; 15:219. [PubMed: 24286137]
16. Remijsen Q, Kuijpers TW, Wirawan E, Lippens S, Vandenabeele P, Vanden Berghe T. Dying for a cause: NETosis, mechanisms behind an antimicrobial cell death modality. *Cell Death Differ*. 2011; 18:581–588. [PubMed: 21293492]
17. Al Ghouleh I, Khoo NK, Knaus UG, Griendling KK, Touyz RM, Thannickal VJ, Barchowsky A, Nauseef WM, Kelley EE, Bauer PM, Darley-USmar V, Shiva S, Cifuentes-Pagano E, Freeman BA, Gladwin MT, Pagano PJ. Oxidases and peroxidases in cardiovascular and lung disease: new concepts in reactive oxygen species signaling. *Free Radic Biol Med*. 2011; 51:1271–1288. [PubMed: 21722728]
18. Hultqvist M, Olofsson P, Holmberg J, Backstrom BT, Tordsson J, Holmdahl R. Enhanced autoimmunity, arthritis, and encephalomyelitis in mice with a reduced oxidative burst due to a mutation in the Ncf1 gene. *Proc Natl Acad Sci U S A*. 2004; 101:12646–12651. [PubMed: 15310853]
19. Muller I, Munder M, Kropf P, Hansch GM. Polymorphonuclear neutrophils and T lymphocytes: strange bedfellows or brothers in arms? *Trends in immunology*. 2009; 30:522–530. [PubMed: 19775938]
20. Sareila O, Kelkka T, Pizzolla A, Hultqvist M, Holmdahl R. NOX2 complex-derived ROS as immune regulators. *Antioxidants & redox signaling*. 2011; 15:2197–2208. [PubMed: 20919938]
21. Hultqvist M, Olsson LM, Gelderman KA, Holmdahl R. The protective role of ROS in autoimmune disease. *Trends in immunology*. 2009; 30:201–208. [PubMed: 19356981]
22. Szabo C, Ischiropoulos H, Radi R. Peroxynitrite: biochemistry, pathophysiology and development of therapeutics. *Nat Rev Drug Discov*. 2007; 6:662–680. [PubMed: 17667957]
23. Radi R, Cassina A, Hodara R. Nitric oxide and peroxynitrite interactions with mitochondria. *Biol Chem*. 2002; 383:401–409. [PubMed: 12033431]
24. Brookes PS, Kraus DW, Shiva S, Doeller JE, Barone M-C, Patel RP, Lancaster JR Jr, Darley-USmar V. Control of Mitochondrial Respiration by NO., Effects of Low Oxygen and Respiratory State. *J Biol Chem*. 2003; 278:31603–31609. [PubMed: 12788936]
25. Colussi C, Albertini MC, Coppola S, Rovidati S, Galli F, Ghibelli L. H₂O₂-induced block of glycolysis as an active ADP-ribosylation reaction protecting cells from apoptosis. *FASEB journal: official publication of the Federation of American Societies for Experimental Biology*. 2000; 14:2266–2276. [PubMed: 11053248]
26. Hyslop PA, Hinshaw DB, Halsey WA Jr, Schraufstatter IU, Sauerheber RD, Spragg RG, Jackson JH, Cochrane CG. Mechanisms of oxidant-mediated cell injury. The glycolytic and mitochondrial pathways of ADP phosphorylation are major intracellular targets inactivated by hydrogen peroxide. *The Journal of biological chemistry*. 1988; 263:1665–1675. [PubMed: 3338986]
27. Mantovani A, Cassatella MA, Costantini C, Jaillon S. Neutrophils in the activation and regulation of innate and adaptive immunity. *Nature reviews. Immunology*. 2011; 11:519–531.
28. Campbell AM, Kashgarian M, Shlomchik MJ. NADPH oxidase inhibits the pathogenesis of systemic lupus erythematosus. *Science translational medicine*. 2012; 4:157ra141.
29. De Ravin SS, Naumann N, Cowen EW, Friend J, Hilligoss D, Marquesen M, Balow JE, Barron KS, Turner ML, Gallin JI, Malech HL. Chronic granulomatous disease as a risk factor for autoimmune disease. *The Journal of allergy and clinical immunology*. 2008; 122:1097–1103. [PubMed: 18823651]
30. Song E, Jaishankar GB, Saleh H, Jithpratuck W, Sahni R, Krishnaswamy G. Chronic granulomatous disease: a review of the infectious and inflammatory complications. *Clinical and molecular allergy: CMA*. 2011; 9:10. [PubMed: 21624140]
31. Hultqvist M, Olofsson P, Gelderman KA, Holmberg J, Holmdahl R. A new arthritis therapy with oxidative burst inducers. *PLoS medicine*. 2006; 3:e348. [PubMed: 16968121]
32. Bowers NL, Helton ES, Huijbregts RP, Goepfert PA, Heath SL, Hel Z. Immune suppression by neutrophils in HIV-1 infection: role of PD-L1/PD-1 pathway. *PLoS pathogens*. 2014; 10:e1003993. [PubMed: 24626392]

33. Schmielau J, Finn OJ. Activated granulocytes and granulocyte-derived hydrogen peroxide are the underlying mechanism of suppression of t-cell function in advanced cancer patients. *Cancer research*. 2001; 61:4756–4760. [PubMed: 11406548]
34. Vasconcelos ZF, Santos BM, Costa ES, Lima M, Tabak DG, Bouzas LF, Azevedo WM, Barcinski MA, Bonomo A. T-lymphocyte function from peripheral blood stem-cell donors is inhibited by activated granulocytes. *Cytotherapy*. 2003; 5:336–345. [PubMed: 12944240]
35. Pillay J, Kamp VM, van Hoffen E, Visser T, Tak T, Lammers JW, Ulfman LH, Leenen LP, Pickkers P, Koenderman L. A subset of neutrophils in human systemic inflammation inhibits T cell responses through Mac-1. *The Journal of clinical investigation*. 2012; 122:327–336. [PubMed: 22156198]
36. Kramer PA, Chacko BK, Ravi S, Johnson MS, Mitchell T, Darley-Usmar VM. Bioenergetics and the oxidative burst: protocols for the isolation and evaluation of human leukocytes and platelets. *Journal of visualized experiments: JoVE*. 2014
37. Dranka BP, Benavides GA, Diers AR, Giordano S, Zelickson BR, Reily C, Zou L, Chatham JC, Hill BG, Zhang J, Landar A, Darley-Usmar VM. Assessing bioenergetic function in response to oxidative stress by metabolic profiling. *Free Radic Biol Med*. 2011; 51:1621–1635. [PubMed: 21872656]
38. Hill BG, Benavides GA, Lancaster JR Jr, Ballinger S, Dell'Italia L, Jianhua Z, Darley-Usmar VM. Integration of cellular bioenergetics with mitochondrial quality control and autophagy. *Biol Chem*. 2012; 393:1485–1512. [PubMed: 23092819]
39. Remijsen Q, Vanden Berghe T, Wirawan E, Asselbergh B, Parthoens E, De Rycke R, Noppen S, Delforge M, Willems J, Vandenabeele P. Neutrophil extracellular trap cell death requires both autophagy and superoxide generation. *Cell research*. 2011; 21:290–304. [PubMed: 21060338]
40. van der Windt GJ, O'Sullivan D, Everts B, Huang SC, Buck MD, Curtis JD, Chang CH, Smith AM, Ai T, Faubert B, Jones RG, Pearce EJ, Pearce EL. CD8 memory T cells have a bioenergetic advantage that underlies their rapid recall ability. *Proc Natl Acad Sci U S A*. 2013; 110:14336–14341. [PubMed: 23940348]
41. Chatila T, Silverman L, Miller R, Geha R. Mechanisms of T cell activation by the calcium ionophore ionomycin. *Journal of immunology*. 1989; 143:1283–1289.
42. Bratton DL, Henson PM. Neutrophil clearance: when the party is over, clean-up begins. *Trends in immunology*. 2011; 32:350–357. [PubMed: 21782511]
43. Pillay J, den Braber I, Vrisekoop N, Kwast LM, de Boer RJ, Borghans JA, Tesselaar K, Koenderman L. In vivo labeling with $^2\text{H}_2\text{O}$ reveals a human neutrophil lifespan of 5.4 days. *Blood*. 2010; 116:625–627. [PubMed: 20410504]
44. Nauseef WM, Borregaard N. Neutrophils at work. *Nature immunology*. 2014; 15:602–611. [PubMed: 24940954]
45. Tillack K, Breiden P, Martin R, Sospedra M. T lymphocyte priming by neutrophil extracellular traps links innate and adaptive immune responses. *Journal of immunology*. 2012; 188:3150–3159.
46. Marko AJ, Miller RA, Kelman A, Frauwirth KA. Induction of glucose metabolism in stimulated T lymphocytes is regulated by mitogen-activated protein kinase signaling. *PloS one*. 2010; 5:e15425. [PubMed: 21085672]
47. Wieman HL, Wofford JA, Rathmell JC. Cytokine stimulation promotes glucose uptake via phosphatidylinositol-3 kinase/Akt regulation of Glut1 activity and trafficking. *Molecular biology of the cell*. 2007; 18:1437–1446. [PubMed: 17301289]
48. Macintyre AN, Rathmell JC. Activated lymphocytes as a metabolic model for carcinogenesis. *Cancer & metabolism*. 2013; 1:5. [PubMed: 24280044]
49. Jackson SH, Devadas S, Kwon J, Pinto LA, Williams MS. T cells express a phagocyte-type NADPH oxidase that is activated after T cell receptor stimulation. *Nature immunology*. 2004; 5:818–827. [PubMed: 15258578]

Summary statement

The lymphocyte metabolic switch, a transition from a mitochondrial to a more glycolytic metabolism, is prevented by neutrophil-derived hydrogen peroxide. We propose that NOX-2 is mediating the neutrophil-dependent suppression of lymphocyte effector function.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Figure 1. Stimulated neutrophil and lymphocyte immune function

Healthy donor CD15⁺ neutrophils were isolated from total Polymorphonuclear cells and lymphocytes by CD14⁺ monocyte, CD61⁺ platelet, and CD235⁺ RBC depletion. **A)** Neutrophils were plated on an extracellular Flux microplate and stimulated with 0, 10, 50, and 100ng/ml PMA for 100 min and the total oxygen consumption rate (OCR) measured as Area under the curve (AUC). **B)** Three hours after stimulation, cells were incubated with a final concentration of 1μM Sytoxgreen and fluorescence measured after 10 min. **C)** Lymphocytes stimulated with 50ng/ml PMA and 100ng/ml Ionomycin (P/I) for 4 days are positive for IFN-γ expression and **D)** undergo clonal expansion as seen by reduction of CFSE stain in subsequent daughter cells by FACS analysis. Values expressed as mean ± SEM with technical replicates of 3–6.

Figure 2. The mitochondrial and glycolytic function of lymphocytes and the oxidative burst of neutrophils with PMA stimulation

A) Lymphocytes (15×10^4 cells/well) and **B)** neutrophils (2.5×10^4 cells/well) isolated from freshly collected human blood were plated and stimulated with ± 100 ng/ml PMA and their OCR followed for 160 min before $1.0 \mu\text{g/ml}$ oligomycin (O), $0.6 \mu\text{M}$ FCCP (F), and $10 \mu\text{M}$ antimycin A (A) were injected for mitochondrial function profiling. **C)** Lymphocytes and **D)** neutrophil glycolytic function was monitored by Extracellular Acidification Rate (ECAR) concurrently with the OCR measurements. Values expressed as mean $\pm$ SEM with technical replicates of 3–6.

Figure 3. The neutrophil dose-dependent mitochondrial and glycolytic dysfunction of lymphocytes

Neutrophils were isolated and plated at 0, 5, 10, 25, 50, and 75 $\times 10^3$ (K) cells/well with 150 $\times 10^3$ (K) lymphocytes. **A**) OCR and **B**) ECAR were concurrently monitored for three basal measurements prior to 100ng/ml PMA injection in all co-culture groups and 160 min following injection. **C**) Following the oxidative burst of the neutrophils, 1.0 μ g/ml oligomycin (O), 0.6 μ M FCCP (F), and 10 μ M antimycin A (A) were injected for mitochondrial and glycolytic profiling. Values expressed as mean $\pm$ SEM with technical replicates of 3–6.

Figure 4. Lymphocyte mitochondrial and glycolytic profiling

Lymphocyte metabolic function was calculated using the OCR and ECAR measurements after oligomycin, FCCP, and antimycin A injection. **A)** Basal mitochondrial respiration (Basal OCR – Non-mitochondrial OCR), **B)** ATP-Linked respiration (oligo-sensitive OCR), **C)** Reserve Capacity (Maximal – Basal OCR), **D)** and Maximal mitochondrial respiration (Maximal OCR- Non-mitochondrial OCR) were calculated and plotted by neutrophil count (per 1,000 cells). Lymphocyte aerobic and oligomycin-stimulated glycolysis was measured using the ECAR values after **E)** 100ng/ml PMA injection (PMA-sensitive), **F)** and 1.0 µg/ml oligomycin injection (oligo-sensitive), respectively, and plotted by neutrophil count. Values expressed as mean ± SEM with technical replicates of 3–6. * = p < 0.05.

Figure 5. Lymphocyte metabolic dysfunction and prevention with catalase

Lymphocytes (15×10^4 cells/well) and neutrophils (2.5×10^4 cells/well) in co-culture were stimulated with PMA in the presence or absence of a 1000U/ml catalase 30 min pretreatment and **A**) OCR and **B**) ECAR followed until oligomycin, FCCP, and antimycin A injection. **C**) Basal mitochondrial, ATP-Linked, Reserve Capacity, and basal, PMA-sensitive, and oligomycin-sensitive ECAR were measured in lymphocytes $\pm$ PMA and in PMA stimulated lymphocyte and neutrophil $\pm$ catalase. Values expressed as mean $\pm$ SEM with technical replicates of 3–6. * = $p < 0.05$ compared to PMA stimulated lymphocytes.

Figure 6. Lymphocyte Glucose Stress Test in co-culture

Neutrophils were isolated and plated at 0, 5, 10, 25, 50, and 75 $\times 10^3$ (K) cells/well with 150 $\times 10^3$ (K) lymphocytes and were stimulated with PMA and no-glucose media exchanged after 160 min. **A**) ECAR was followed for three readings then 5mM glucose, 1.0 μ g/ml oligomycin, and 20mM 2-deoxy-glucose (2-DG) were injected. **B**) Glycolysis (Glucose ECAR – 2-DG ECAR) and glycolytic reserve (oligomycin-sensitive ECAR) were measured and plotted by neutrophil count (per 1,000 cells). **C**) These and non-glycolytic ECAR were measured in lymphocytes $\pm$ PMA and in PMA stimulated lymphocyte and neutrophil $\pm$ 1000U/ml catalase. Values expressed as mean $\pm$ SEM with technical replicates of 3–6. * = $p < 0.05$.

Figure 7. Metabolic regulation of activated lymphocyte by bolus hydrogen peroxide and DMNQ
 Isolated lymphocyte **A**) OCR and **B**) ECAR were assessed after a 30 min pretreatment of 100µM H₂O₂, 5µM DMNQ, or 25k neutrophils and the PMA sensitive changes measured for 160 min before the injection of oligomycin, FCCP, and antimycin A. **C**) The observed changes in Post-PMA Basal OCR/ECAR, ATP-linked OCR/Oligo-sensitive ECAR, Reserve capacity, and PMA sensitive ECAR were quantified and compared to lymphocyte + PMA control. OCR/ECAR traces were obtained from a single healthy donor lymphocytes and neutrophils and values expressed as mean ± SEM with technical replicates of 3–6. * = p 0.05.

Figure 8. Lymphocyte IFN- γ production in co-culture

CD3⁺ lymphocytes, from a representative healthy donor, are gated for IFN- γ secretion after 4 days of incubation with **A**) no stimulation: media control, **B**) co-culture with neutrophils, or **C**) Stimulation with 50ng/ml PMA and 100ng/ml Ionomycin (P/I). **D**) Co-culture of PMA/Ionomycin stimulated lymphocytes with neutrophils demonstrates reduced IFN- γ secretion. Catalase (1000U/ml) pretreated for 30 minutes on **E**) lymphocytes alone and **F**) lymphocyte and neutrophil co-culture, show similar IFN- γ expression compared to the lymphocytes with PMA and Ionomycin.

Figure 9. Lymphocyte clonal expansion in co-culture

CFSE analysis of lymphocyte proliferation from a representative healthy donor is demonstrated. Negative controls for **A**) media alone and **B**) neutrophils in co-culture show minimal proliferating cells in the absence of stimulation. **C**) The addition of 50ng/ml PMA and 100ng/ml Ionomycin (P/I) results in approximately 37% proliferation of lymphocytes. **D**) This proliferation is decreased when exposed to the neutrophil oxidative burst. **E**) Catalase (1000U/ml) pretreated on lymphocytes alone **F**) and in neutrophil co-culture, show loss of neutrophil-dependent suppression of proliferation as well as increased proliferative capacity compared to the lymphocytes with PMA and Ionomycin.