The use of nanoparticulates to treat breast cancer

Breast cancer is a major ongoing public health issue among women in both developing and developed countries. Significant progress has been made to improve the breast cancer treatment in the past decades. However, the current clinical approaches are invasive, of low specificity and can generate severe side effects. As a rapidly developing field, nanotechnology brings promising opportunities to human cancer diagnosis and treatment. The use of nanoparticulate-based platforms overcomes biological barriers and allows prolonged blood circulation time, simultaneous tumor targeting and enhanced accumulation of drugs in tumors. Currently available and clinically applicable innovative nanoparticulate-based systems for breast cancer nanotherapies are discussed in this review.

First draft submitted: 1 June 2017; Accepted for publication: 26 July 2017; Published online: 4 September 2017

Keywords: breast cancer • gene/drug delivery • nanoparticles

Breast cancer is the most common cancer for females both in the USA and other countries in the world. In the USA alone, approximately 61,000 new *in situ* breast cancer cases and 249,000 new invasive breast cancer cases have been estimated to occur in 2016, accounting for nearly one in three cancer cases [1]. Breast cancer is the leading cause of cancer death among women aged 20–59 years old and the second leading cause of cancer death among women aged 60 years or older behind lung and bronchus cancer [1]. In 2016, nearly 111 women per day in the USA have been projected to die from breast cancer [1]. The overall 5 years relative survival rate of women with breast cancer in the USA increased from 75% (1975–1977) to 91% (2005–2011), due to improved techniques for diagnosis and treatment [1]. However, the 5 years relative survival rate of breast cancer patients diagnosed at late stages is still as low as 26% [1] and a common cause of breast cancer mortality is metastasis to lymph nodes, lung, liver, bone and brain [2]. Therefore,

more efficacious treatment of breast cancer especially at late stages is crucial to the survival of breast cancer patients.

Current breast cancer treatment includes surgery (mastectomy and lumpectomy), radiotherapy, chemotherapy and hormone therapy [3]. A significant shortcoming associated with these therapies is the lack of specificity. Over the last decades, the development of cancer treatment has been significantly promoted by cancer nanomedicine, which is an interdisciplinary field focusing on the design and medical applications of materials and technologies at the nanoscale (typically up to 100 nm) [4]. Nanoparticulate-based delivery systems provide many potential benefits, including increased biocompatibility, multifunctional encapsulation of active agents, reduced degradation during blood circulation, passive or active targeting, effective delivery and reduced or eliminated side effects [5]. In this review, both advantages and limitations of nanomedicine formulations on breast cancer treatment were discussed using

Xiaomeng Tang^{1,2}, Welley S Loc^{1,2}, Cheng Dong³, Gail L Matters⁴, Peter J Butler³, Mark Kester⁵, Craig Meyers⁶, Yixing Jiang⁷ & James H Adair^{*,2}

¹Department of Chemistry, Pennsylvania State University, University Park, PA 16802, USA

²Department of Materials Science & Engineering, Pennsylvania State University, University Park, PA 16802, USA

³Department of Bioengineering, Pennsylvania State University, University Park, PA 16802, USA

⁴Department of Biochemistry & Molecular Biology, Pennsylvania State University College of Medicine, Hershey, PA 17033, USA

⁵Department of Pharmacology, University of Virginia School of Medicine, Charlottesville, VA 22908, USA

⁶Department of Microbiology & Immunology, Pennsylvania State University College of Medicine, Hershey, PA 17033, USA

⁷Marlene & Stewart Greenebaum Cancer Center, University of Maryland School of Medicine, Baltimore, MD 21201, USA

*Author for correspondence:

Tel.: +1 814 863 6047

jha3@ems.psu.edu

a material selection approach and biological system criteria rarely used in the medical community.

Selection of nanoparticulate-based systems for cancer treatment

Innumerable nanomaterials have been invented with the rapid development of nanotechnology over the past 30 years but only a few nanoparticulate-based systems can be used in biomedical field and even fewer of them can meet the requirements of the US FDA for clinical applications [5]. The physical and chemical characteristics of nanoscale materials that ensure effective drug delivery are summarized in Table 1 [5,6]. The primary requirement of a biomaterial is to be systemically nontoxic and biocompatible [5]. Furthermore, some nanomaterials are perceived as dysfunctional cellular components and initiate cell degradation through autophagy, a common result of material toxicity [7]. Autophagy can be induced by many common nanomaterials [8], including gold nanoparticles [9], quantum dots [10], titanium dioxide nanoparticles [11], zinc oxide nanoparticles [12], nano rare earth oxides [13], fullerene [14], fullerenol [15] and carbon nanotubes [16].

After screening over 240 nanotechnology strategies, researchers in the Nanotechnology Characterization Lab at the National Cancer Institute concluded that it was difficult to find a material satisfying all these criteria listed in Table 1. However, there are several nanoparticulate-based therapeutic systems that meet most if not all of these important criteria, which are considered to be promising for the diagnosis of cancers, and will be highlighted in this paper.

Clinical nanomedicine for breast cancer treatment

Systemic anticancer pharmaceuticals are usually hydrophobic, untargeted and toxic compounds, which can cause severe side effects [17]. Anthracyclines and taxanes are two mainstay chemotherapeutics used in advanced breast cancer chemotherapy over the last three decades [18]. The family of anthracyclines, especially doxorubicin formulations, has been reported to induce cardiotoxicity [19]. Also, taxanes (paclitaxel and docetaxel) have several side effects including bone marrow suppression [20], hypersensitivity reactions [21], cutaneous reactions [22] and dose-limiting neurotoxic-

Table 1. Selection criteria for a nanoparticulate-based therapeutic platform.

Desired characteristics	Comments
Inherently nontoxic and biocompatible materials	The initial material selection should be based on nontoxic and biocompatible materials especially with an aim toward human healthcare
Small size (10–200 nm)	Based on <i>in vivo</i> studies, there seems not a most efficacious particular size. This is the range of particle diameters that have proven most effective for a wide variety of delivery systems. There is also debate around the influence of particle shape [157]
Encapsulation of active molecules at therapeutic levels	The drugs should be encapsulated within the nanoparticles and be protected from unwanted degradation or clearance during blood circulation. To be efficacious, the nanoparticle vehicle must encapsulate a high percent of the agent (>50 mol%) to achieve therapeutic dosage
Colloidally stable in physiological environment	The nanoparticle and surface functionalization should be resistant to the agglomeration caused by solution pH values, ionic strength, macromolecular interactions and temperature in the physiological environment to avoid sequestration in organs or membranes
Reasonable circulation time	The nanoparticle and surface functionalization should be limited susceptible to photo-, thermal-, biological and chemical degradation, and thus there should be reasonable circulation time in the circulatory system for the nanoparticle to finish its task
Targeting to cell or tissue of choice	Targeting ensures the greatest uptake concentration of chemotherapeutics within the desired lesions and the least side effects with healthy tissues. Targeting could be active targeting via bioconjugation of receptor molecules on the surface or passive targeting, like through enhanced permeability and retention effect
Biologically or extrinsically controlled release of active molecules	There should be a trigger mechanism such as the acidic pH within the tumor or during endosome maturation to ensure the release of the encapsulated drug into the targeted tissues
Clearance	There must be clearance mechanisms for the nanoparticle vehicle after completing its task, to avoid the cumulative and/or long-term systemic side effects, such as possible interference with biological functions

Adapted with permission from [5].

ity [23]. Compared with conventional cancer therapies, nanomedicine has many benefits, such as less drug degradation during transport via protection from *in vivo* chemical or biological environments, minimized adverse effects via improved biocompatibility and targeting and increased dosage of chemotherapeutic delivered to the cancer tissue [5,24]. Nanomedicine also possesses great potential to effectively target and eliminate breast cancer stem cells, which is a possible important factor in the initiation, recurrence and chemo-/radiotherapy resistance of breast cancer [25]. Many nanoparticulate-based chemotherapeutic delivery platforms have been approved by the FDA or are in clinical trials for cancer treatment (Figure 1) and the nanomedicine formulations developed for breast cancer therapy are summarized in Table 2. The formulations based on liposomes, polymeric nanoparticles and nanoparticle albumin-bound (nabTM) paclitaxel to treat breast cancer are discussed in detail in this section.

Liposomes

Liposomes are a family of self-assembling closed spherical vesicles composed of an aqueous core and a membranous lipid bilayer, which were first described by Bangham *et al.* in 1964 [26]. The use of liposomes (typically 50–100 nm) to entrap drugs was reported by Gregoriadis *et al.* in 1971 [27]. Hydrophilic drugs, genes and siRNA (small interfering RNA) can be encapsulated within the core and hydrophobic drugs can also be incorporated within the lipid bilayer, which contribute to the reduced systemic toxicity and increased circulation time [28,29]. To achieve longer *in vivo* circulation of liposomes, biocompatible and inert polymers such as PEG are coated on the surface of liposomes, forming a protective layer, which decreases the recognition by the reticuloendothelial system and subsequent clearance [29]. Furthermore, the intratumoral localization of liposome-based drug delivery systems is improved via the enhanced permeability and retention effect or the bioconjugation of liposomes with a targeting ligand capable of recognizing and binding to cancer cells, including antibodies [30], peptides [31] and aptamers [32].

Doxil[®], PEGylated liposomes encapsulating doxorubicin was the first FDA-approved nanomedicine and the structure is shown in Figure 2 [33]. Doxorubicin intercalates between base pairs of DNA strands, inhibiting DNA synthesis and transcription and Doxil has been used to treat Kaposi's sarcoma, refractory breast and ovarian cancer [33]. The circulating half-life of Doxil (starting dose, 60 mg/m²) in breast cancer patients was about 74 h [34], compared with approximately 5 min for free doxorubicin [35]. The mean plasma clearance value of Doxil is 0.041 l/h/m² at a dose of 20 mg/m² in

contrast to doxorubicin, which displays a plasma clearance value ranging from 24 to 35 l/h/m² [36]. The treatment with liposomal doxorubicin was also more effective; a dose of 50 mg/m² given every 4 weeks was as effective as that of conventional doxorubicin 60 mg/m² given every 3 weeks [34]. More importantly, the risk of developing cardiotoxicity associated with peak concentration of free doxorubicin was significantly reduced and only 10 out of 254 patients treated with liposomal doxorubicin developed this side effect while 48 out of 255 breast cancer patients developed cardiotoxicity when treated with free doxorubicin [34].

Liposome-based nanoplatforams for breast cancer treatment are well developed, while there are still some limitations. One of the major shortcomings of liposomes is the physical and chemical instability. Physical instability can be caused by the aggregation or fusion of vesicles due to the poor colloid stability with the formed larger particles cleared rapidly [37]. Chemical instability may be caused by the oxidation of unsaturated acyl chains of lipids and/or the hydrolysis of ester bonds, leading to leakage of liposomes and release of drugs before the liposomes reach the cancer tissues [37]. Another problem with liposomes is the low encapsulation efficiency of hydrophobic drugs, such as

Figure 1. Nanoparticulate-based chemotherapeutic delivery platforms approved by the US FDA for cancer treatment. (A) Liposomes are spherical vesicle composed of an aqueous core and a membranous lipid bilayer and are often used for the encapsulation of hydrophilic chemotherapeutics. (B) Polymeric nanoparticles including polymeric micelles are self-assembled from amphiphilic and biodegradable polymers in aqueous solution and are effective to encapsulate hydrophobic drugs. (C) Nanoparticle albumin-bound paclitaxel is a colloid suspension of paclitaxel stabilized with human serum, which dissociates to albumin–paclitaxel complex in blood stream.

Table 2. Nanomedicines as breast cancer therapeutics approved by the US FDA or under different stages of development in clinical trials.

Product	Nanoplatfrom/agent	Manufacturer in USA	Status in USA
Doxil® [33]	PEGylated liposomal/doxorubicin hydrochloride	Janssen Products	Approved in 1995
Abraxane® [63]	Nanoparticle albumin-bound paclitaxel	Abraxis Bioscience	Approved in 2005
Myocet® [158]	Non-PEGylated liposomal/doxorubicin	Sopherion Therapeutics	Phase III
NK-105 [159]	PEG-polyaspartate/paclitaxel	Nippon Kayaku	Phase III
Genexol®-PM [49]	PEG-poly(D,L-lactide)/paclitaxel	Samyang Biopharmaceuticals	Phase III
LEP-ETU [160]	Liposome/Paclitaxel	INSYS Therapeutics	Phase II
NK-012 [161]	PEG-polyglutamic acid/SN-38	Nippon Kayaku	Phase II
Xyotax® [162]	Paclitaxel poliglumex	Dana-Farber Cancer Institute	Phase II
ThermoDox® [163]	Heat-activated liposomal/doxorubicin	Celsion	Phase I/II
Liposomal annamycin [164]	Liposome/semi-synthetic doxorubicin analogue annamycin	New York University, School of Medicine	Phase I/II
Rexin-G [165]	Targeting protein tagged phospholipid/microRNA-122	Epeius Biotechnologies	Phase I/II
SPI-077 [166]	Stealth liposomal cisplatin	LiPlasome Pharma	Phase I
S-CKD602 [167]	PEGylated liposomal/CKD602	Alza	Phase I
Nanoxel® [168]	PEG-poly(D,L-lactide)/docetaxel	Fresenius Kabi Oncology	Phase I
BIND-014 [169]	PEG-Polylactic-co-glycolic acid/docetaxel	BIND	Phase I

paclitaxel, due to the low affinity of hydrophobic drugs for the lipid bilayer [38]. These limitations should be fully studied and addressed in the development of new clinical nanomedicine based on liposomes.

Polymeric nanoparticles

Polymeric nanoparticles have been developed with the need of nanocarriers for hydrophobic chemotherapeutics and hormone regulators, such as taxanes, camptothecin, cisplatin and tamoxifen [39]. Polymeric nanoparticles including polymeric micelles, capsules, colloids and dendrimers are core-shell structures, self-assembled in aqueous solution by amphiphilic and biodegradable polymers, such as poly(D,L-lactic acid), poly(D,L-lactic-co-glycolic acid), poly(ϵ -caprolactone) and the copolymers with PEG, which is typically employed to prevent particle agglomeration [40]. Hydrophobic active agents can be encapsulated within the hydrophobic core while hydrophilic encapsulants can be attached to the outer shell or the corona of polymeric nanoparticles via covalent bonding, hydrogen bonding or electrostatic interaction [39]. Bioconjugation with targeting moieties, such as peptides and antibodies enhances delivery to cancer cells [41,42]. The most important property of polymeric nanoparticles is the controlled drug delivery not only via the diffusion and swelling of nanostructures but also as the response to changes in environments such as pH [43] and temperature [44] or to the external stimulus such as light [45] and ultrasound [46–48].

Genexol®-PM, paclitaxel-doped poly(D,L-lactic acid) is one of the polymeric nanomedicine in clinical trials for breast cancer treatment and the structure is shown in Figure 3 [49,50]. Paclitaxel is an antimitotic chemotherapeutic which inhibit the depolymerization of the microtubule and results in cell-cycle arrest and inhibition of mitosis [51]. It has been demonstrated clinically that Genexol-PM is able to deliver higher dose of paclitaxel (suggested dosage, 300 mg/m²) than the conventional chemotherapy (175 mg/m²) without additional dose-limiting toxicity [50,52]. The half-life of Genexol-PM is 18.8 $\pm$ 3.1 h, which is 1.8-times longer than 6.8 $\pm$ 1.4 h for free paclitaxel [53]. In a Phase II clinical trial of Genexol-PM for the treatment of metastatic breast cancer, 41 patients were administered intravenously with 300 mg/m² of Genexol-PM over a period of 3 h every 3 weeks [54]. The overall response rate was 43.5–73.7% (95% confidence interval) and the median time to progression for all patients was 9.0 months [54]. The lower systemic toxicity and fewer side effects could be contributed by the absence of Cremophor EL (polyoxyethylated castor oil), which was typically used to increase paclitaxel solubility and was associated with hypersensitivity reaction and neuropathy [55].

Although polymeric nanoparticles are effective drug carriers for hydrophobic chemotherapeutics or hor-

more regulators, there are still some shortcomings. The first concern is the stability of polymeric nanoparticles under harsh conditions such as high temperature and radiation during the sterilization, an essential step in nanomedicine manufacture [56,57]. Moreover, the presence of toxic residual solvents introduced during the manufacture of polymers and the exocytosis caused by undissolved polymeric structures cannot be ignored [5,58].

Nab paclitaxel

Paclitaxel is a complex compound first isolated from the bark of yew tree by Wall and Wani in 1967 [59], and is one of the primary drugs approved by the FDA for the treatment of breast, ovarian, prostate and non-small-cell lung cancer [60]. As a member of taxane family, paclitaxel binds to microtubule, promotes tubulin polymerization, stabilizes microtubule polymers and inhibits polymer disassembly during cell division [51]. Thus mitosis, motility and intracellular transportation within cancer cells are suppressed, leading to cancer cell apoptosis [60]. However, the clinical success of taxanes is limited by the poor water solubility. Cremophor EL is added to clinically available paclitaxel to improve the solubility of this hydrophobic drug, which may induce serious dose-dependent toxicities, including some fatal hypersensitivity reactions [61,62]. To reduce or eliminate the side effects caused by Cremophor EL, nab paclitaxel (Abraxane®) is designed and has been approved by the FDA in 2005 to treat metastatic breast cancer [63]. Abraxane is a Cremophor EL free colloid suspension of unmodified paclitaxel stabilized with

albumin with a particle size of approximately 130 nm, prepared by homogenization of paclitaxel and human serum albumin under high pressure [64,65]. Albumin is a natural carrier for hydrophobic molecules, such as most chemotherapeutics and hormones [65]. In nab, albumin and paclitaxel are linked via reversible non-covalent binding [66]. After the circulation in the blood stream, the administrated albumin–paclitaxel complex binds to the 60 kDa gp60 receptor on cell surface, and then the active agent, paclitaxel is transcytosed to the extravascular space through caveolae formed by the invagination of cell membranes [63].

In normal saline solution, the colloidal suspension of nanoparticles, nab paclitaxel can be reconstituted to 2–10 mg/ml, much higher than 0.3–1.2 mg/ml for Cremophor EL-paclitaxel [64]. Therefore, nab paclitaxel allows safe administration of paclitaxel with a much higher dose and a significant shorter injection time period without special tubing or premedications for hypersensitivity reactions, compared with conventional paclitaxel therapy [64,67]. In a Phase III FDA study of patients with metastatic breast cancer, an injection dose of 260 mg/m² Abraxane was administrated over 30 min, every 3 weeks, which is approximately 50% higher than the typical tolerated dose of Taxol® (175 mg/m², over 3 h every 3 weeks) [67]. The paclitaxel clearance rate and volume of distribution were higher for nab paclitaxel (21.13 l/h/m², 663.8 l/m²) than for Taxol (14.76 l/h/m², 433.4 l/m²) and the response rate for nab paclitaxel was also significantly higher (34 vs 19%) [67]. It should be noted that no hypersensitivity response was observed in 229 patients as a result of the

Figure 2. An illustration of Doxil®, PEGylated liposomes encapsulating doxorubicin (<100 nm). Anticancer drug, doxorubicin is encapsulated in the aqueous core and PEG is attached on the surface of liposomes for prolonged blood circulation and protection from unwanted rapid clearance. Reprinted with permission from [33].

absence of Cremophor EL [67]. More importantly, since paclitaxel in nab paclitaxel nanocomplex is delivered to tumors via the interaction between the complex and albumin receptors on cancer cell surface, a 9.9-fold increase in endothelial cell binding and a 4.2-fold increase in endothelial transcytosis for nab paclitaxel were observed *in vitro* and intratumor concentration of paclitaxel was 33% higher in MX-1 breast tumors of athymic mice, compared with standard paclitaxel [68].

Innovative nanotherapies for breast cancer HER2 targeted nanotherapies

As summarized in Table 3 for all innovative nanotherapies discussed in this section, HER2 is a transmembrane receptor with tyrosine kinase activity [69]. Unlike other members of the HER family (HER1, HER3 and HER4), there is no identifiable ligand for HER2, but HER2 can dimerize with other HER receptors [69,70]. The heterodimerization leads to the phosphorylation of the HER kinase domains. This activates several signal pathways related to transcription, translation and protein stability, including the MAPK, PI3K/Akt and phospholipase C γ [69,71]. Therefore, HER2 is important in the regulation of gene expression, cell mobility, growth, proliferation, apoptosis and other cellular responses, especially for cancerous cells [72]. HER2/HER3 is the most active and tumor promoting dimerized combination [73]. HER2 is overexpressed by ~20–30% of human breast tumors, leading to a higher recurrence rate and mortality rate for HER2-positive breast cancer [69,74,75]. However, the overexpression of HER2 can be utilized in the development of targeting nanomedicine to treat HER2-positive breast cancer. The first clinical HER2 targeting agent is the humanized monoclonal antibody, trastuzumab (Herceptin[®]) [76]. There are two antigen-specific sites in trastuzumab, which can bind to HER2, preventing the activation of tyrosine kinase and HER2 dimer-

ization [77]. In addition, trastuzumab activates the immune effector cells for antibody-dependent cell-mediated cytotoxicity to promote cancer cell death [78]. The presence of trastuzumab also enhances the therapeutic effects of other chemotherapeutics [79,80]. Therefore, trastuzumab is widely used as a single agent or combined with other chemotherapeutics in adjuvant therapy to reduce breast cancer recurrence rate and improve the overall survival of HER2-positive breast cancer patients clinically [81–83].

More importantly, the recognition and binding of the two antigen-specific sites in trastuzumab to HER2 make trastuzumab a targeting ligand to direct nanomedicine to breast cancer cells overexpressing HER2 [77,84]. Anhorn *et al.* have developed doxorubicin-doped human serum albumin nanoparticles (DOX-NP) surface functionalized with trastuzumab via covalent binding (DOX-NP–trastuzumab) with the structure shown in Figure 4 [85]. DOX-NP–trastuzumab bound to 73% of HER2 overexpressing SK-Br-3 breast cancer cells, while the DOX-NP bioconjugated with nonspecific IgG bound to less than 6% of SK-Br-3 cells [85]. This increase in cellular binding was dependent on the concentration of trastuzumab and was not observed with MCF7 breast cancer cells, a cell line that only weakly expresses HER2 [85]. Enhanced cellular uptake and intracellular distribution of DOX-NP–trastuzumab compared with the control DOX-NP PEGylated with Methoxy-PEG were also demonstrated by confocal laser scanning microscopy (Figure 4) [85]. This *in vitro* study displayed the potential of trastuzumab as a HER2-targeting ligand for breast cancer cells.

Unfortunately, many patients acquire trastuzumab resistance within a year after the initial response to trastuzumab formulations [69]. Therefore, other HER2-targeting strategies have been developed. Pertuzumab is another monoclonal antibody, which binds to HER2 and sterically blocks the homodimerization and heterodimerization of HER receptors [86]. Current clinical trials on pertuzumab focus on the evaluation of efficacy of pertuzumab with other drugs to treat breast cancer, especially plus trastuzumab and docetaxel [87–89]. Another type of HER2 inhibitors is tyrosine kinase antagonist lapatinib (trade name: Tykerb[®]) [69], which has been incorporated into lipid nanoparticles (LTNPs) by Gao *et al.* [90,91]. The recent *in vivo* study suggested that the area under the curve of LTNPs was approximately five- and threefold as that of free lapatinib and lapatinib suspension, respectively, demonstrating that LTNPs increased the bioavailability and blood circulation time of lapatinib [91]. Furthermore, the uptake of LTNPs by breast cancer cells was effective and led to apoptosis and about fivefold

Figure 3. Schematic illustration of the structure of Genexol[®]-PM, paclitaxel-doped PEGylated poly(D,L-lactic acid). Chemotherapeutic, paclitaxel is encapsulated within the hydrophobic poly(D,L-lactic acid) core and the surface of the nanoparticle is modified with polyethylene glycol to improve the colloidal stability.
PLA: Poly(D,L-lactic acid).

reduced tumor growth in a BT-474 xenograft-bearing mice model [90].

Nanotherapy for triple negative breast cancer

Triple-negative breast cancer (TNBC) is a subgroup of breast cancers, which is characterized by an absence or poor expression of estrogen receptor, progesterone receptor and HER2 [92]. Approximately 12–17% of female breast cancer patients are diagnosed with TNBC, which is aggressive and highly metastatic associated with poor prognosis [92]. Unlike patients with hormone or growth factor receptor-positive breast cancers who benefit from endocrine therapy or treatment targeting relative overexpressed receptors, conventional chemotherapy is the primary option for TNBC patients currently [93]. Chemotherapeutics for TNBC treatment include anthracyclines, taxanes and cisplatin [93], but the overall chemotherapy outcome for TNBC patients is worse than patients with other breast cancer subtypes [94]. The estimated 1-, 3- and 5-year overall survival rate were 90, 74 and 64% for TNBC and 97, 89 and 81% for non-TNBC, respectively [95]. Therefore, it is urgent to develop novel therapies for TNBC.

A novel therapy for TNBC was reported by Alam *et al.*, which induced necrotic death and inhibited of TNBC tumor growth *in vivo* using adeno-associated virus type 2 (AAV2) [96]. Triple-negative MDA-MB-435 cells derived from metastatic human breast carcinoma (5×10^6) were implanted subcutaneously in

the shoulder of mice and a significant suppression of tumor growth was recorded when the tumor received AAV2 (10^5 or 10^6 infectious units/200 μ l) compared with mock (not manipulated) and PBS-treated groups (Figure 5A) [96]. Dry and dark necrotic patches were observed on the skin over the tumors of mice in AAV2-treated groups and the hematoxylin and eosin (H&E) staining of the tumor sections also confirmed tumor necrosis treated with AAV2 and patterns of strong infiltrating macrophages in the mock and phosphate-buffered saline (PBS)-treated tumors (Figure 5B) [96]. This novel virus infection has shown the potential for the treatment of breast cancer, but safety is a concern with clinical virus involved applications. One solution to this issue is to encapsulate the virus in targeting nanocarriers and isolate the virus from healthy tissues.

The treatment of TNBC is a clinic challenge due to the high proliferative rates of TNBC cells, early recurrence and poor survival rate. Except for the AAV2 therapy, results of clinical neoadjuvant chemotherapy already demonstrated the improvement in breast conservation rates and response rates [97]. Obviously, there is a need for new, targeted therapeutic strategies for a better outcome for patients with TNBC.

RNA interference

RNA interference (RNAi) is able to silence tumor-associated genes that are difficult for antibodies or inhibitors to target and is expected to be a useful therapeutic approach for advanced breast cancer [98,99]. This pro-

Table 3. Summary of innovative nanotherapies for breast cancer.

Nanotherapy	Strategy	Active agents
HER2	Target overexpressed HER2 transmembrane receptor to inhibit tyrosine kinase activity and HER2 homo- and/or heterodimerization into tumor promoting combinations such as the HER2/HER3	Trastuzumab, pertuzuma and lapatinib
TNBC	Induce necrotic tumor cell death to inhibit tumor growth. Method can be further improved by encapsulation in targeting nanocarriers	AAV2
RNAi	Silence tumor-associated genes that cannot be targeted by antibodies or inhibitors. The result is to inhibit the growth and induce apoptosis of tumor cells. The active agent can be encapsulated in a targeting nanocarrier such as liposomes or form biocompatible complexes with atelocollagen to improve uptake	siRNA, siRNA–atelocollagen complex
Photothermal ablation (PTA)	Targeted nanoparticle-mediated PTA that can absorb NIR irradiation and release heat energy above the thermal ablation threshold (50°C) to damage tumor cells	Metallic-based nanomaterials (gold or silver) or materials with high photothermal conversion effects
CPSNPs	Nontoxic calcium phosphate-based nanoparticles doped with chemotherapeutic agents and bioconjugated with anti-CD71 to target breast cancer. CPSNPs dissolve at late stage endo-lysosome formation (pH 4.8–5.5) to release active agents. The particles remain intact at pH 7.4 for systemic delivery	Docetaxel, indocyanine green (photosensitizer for PINT)

AAV2: Adeno-associated virus type 2; CPSNP: Calcium phosphosilicate nanoparticle; NIR: Near infrared; PINT: Photoimmunonano therapy; PTA: Photothermal ablation; RNAi: RNA interference; TNBC: Triple-negative breast cancer.

cess is triggered by siRNA, the exogenously introduced double-strand RNA, which is able to bind to a group of protein complex, referred to RNA-induced silencing complex, leading to the degradation of target mRNA by a cleavage enzyme within RNA-induced silencing complex or suppression of mRNA translation [99–102]. However, there are several concerns that limit the clinical use of RNAi, including the rapid intravascular degradation of RNA and the safe and effective delivery of siRNA to target tissues [103,104]. Viral delivery systems widely used in gene therapy can be effective but carry safety concerns associated with the virus [105,106]. Non-viral delivery systems, such as liposomes, polymeric nanoparticles and polypeptides have been considered as potential alternatives [107]. Wang *et al.* reported a lipid nanoparticle (LNP) averaging 150 nm diameter loaded with a siRNA against cyclin-dependent kinase 4 (CDK4), an early G1 phase cell cycle factor associated with the progression of mammalian cells [108]. With the assistance of this lipid-based nanoparticle delivery system, a 16-fold increase in intracellular uptake of siRNA by MDA-MB-468 breast cancer cells was observed *in vitro* [108]. More importantly, the *in vitro* CDK4 suppression and G1 cell cycle arrest (13.8%) in MDA-MB-468 cells was effectively induced by LNP–siRNA while no CDK4 mRNA or G1 cell cycle down-regulation occurred after incubation with free CDK4 siRNA, indicating the potential of LNP–siRNA to inhibit breast cancer cell growth without adverse side effects [108].

In addition to silencing genes associated with cancer cell proliferation, RNAi can also be used to enhance the chemotherapeutic effects or to overcome the drug resistance. Honma *et al.* prepared a stable nano-sized RPN2 siRNA–atelocollagen complex and tested the effects on improving the *in vitro* and *in vivo* docetaxel sensitivity of MCF7-ADR cells, a human breast cancer cell line that is resistant to multiple drugs, such as doxorubicin and taxanes (Figure 6) [109]. In the *in vitro* study with MCF7-ADR cells, RPN2 siRNA–atelocollagen significantly inhibited cell growth and induced apoptosis compared with free siRNA in the presence of docetaxel [109]. MCF7-ADR cells treated with RPN2 siRNA–atelocollagen were 3.5- and 2.6-fold more sensitive to docetaxel and to paclitaxel than cells transfected with a nontargeting siRNA [109]. A 7-day treatment of siRNA–atelocollagen complex plus docetaxel effectively decreased the breast tumor size from 52 to 21 mm³ in mice orthotopically implanted with MCF7-ADR cells and controlled tumor growth (61–97 mm³) in mice orthotopically implanted with MDA-MB-231/MDR1 human breast cancer cell. There were no noticeable improvements when mice were treated with docetaxel, RPN2 siRNA or siRNA–atelocollagen

complex independently [109]. Enhanced chemosensitivities to doxorubicin, paclitaxel and cisplatin with nanoparticle-facilitated delivery of target siRNAs were also reported by Li *et al.* [110].

RNAi is a novel approach for breast cancer treatment with unique and critical potentials. However, there are still several challenges, such as intracellular targeting, immune-mediated toxicities [111] and oversaturation of extraneous RNAs [112], which need to be overcome before successful clinical applications [113].

Nanoparticle-mediated photothermal ablation

Photothermal ablation (PTA) is another innovative and noninvasive therapeutic approach which can be used for breast cancer treatment, especially for the patients who do not respond to conventional chemotherapies or radiation treatment. In nanoparticle-mediated PTA, nanoparticles delivered to cancer tissues absorb the energy from near infrared (NIR) irradiation and transform the energy into heat, known as the photothermal effect [114]. The generated heat can increase the temperature of the surrounding tissues above the thermal ablation threshold of 50°C [115]. When the tissue temperature is above 42°C, there is irreversible cellular damage [116], although the damaging mechanism of PTA is not yet fully understood. Possible explanations include membrane conformation and permeability change, cell deformation, cell cycle deregulation, which eventually lead to cell death [116–118]. *In vitro* studies have demonstrated that a few seconds of NIR laser irradiation are enough to disturb the plasma membrane integrity of tumor cells without noticeable damage to nonmalignant cells, making PTA a promising alternative for cancer treatment [119–121]. Metallic nanomaterials, especially gold or silver-based nanoparticles, are often used for PTA due to the strong absorption within NIR region and high photothermal conversion efficiency [122]. NIR resonant nanomaterials including gold nanoshells [123], gold nanorods [117], gold nanospheres [124], carbon nanotubes [125] and some other nanoparticles with NIR absorption were demonstrated to be potential photothermal sensitizers for PTA [115,126]. The nanoparticle surface can be modified with ligands to enhance targeting through hormone-receptor or antibody–antigen interaction [127–129]. Carpin *et al.* reported silica-gold nanoshells conjugated with anti-HER2 antibody for PTA of trastuzumab-resistance breast tumors [123]. Significantly more anti-HER2 nanoshells bound to HER2 overexpressed breast cancer cells SK-BR-3 (trastuzumab-sensitive) and BT474 AZ LR (trastuzumab-resistant) compared with HER2 negative breast cancer cells MCF10A [123]. After incubation with gold nanoshells, cells were exposed to 808 nm diode laser with 80 W/cm² and

Figure 4. Enhanced cellular uptake and intracellular distribution of DOX-NP-trastuzumab. (Top) Covalent linkage of thiolated antibody (trastuzumab or Immunoglobulins G) to cross-linker-activated drug-loaded human serum albumin nanoparticle (black square = doxorubicin molecules). (Bottom) Cellular uptake and intracellular distribution of the doxorubicin-nanoparticles studied by confocal laser scanning microscopy. SK-Br-3 cells are cultured on glass slides and treated with different doxorubicin-nanoparticle formulations for 4 h at 37°C. The green signal is the autofluorescence of the nanoparticles and the cell membranes are stained with a red fluorophore, Concanavalin A Alexa Fluor 594. Pictures are collected within inner sections of the cells. (A–C) Incubation of the cells with control Dox-NP-mPEG. (D–F) Incubation of the cells with the specific Dox-NP-trastuzumab. (A & D) display the green nanoparticles; (B & E) display the red cell membrane; (C & F) overlay both fluorescence channels. Adapted with permission from [85] © 2008 American Chemical Society.

Figure 5. Necrotic death and inhibition of TNBC tumor growth *in vivo* using AAV2. (A) AAV2 infection reduced MDA-MB-435 tumor growth and increased survival in nude mice. Four sets of five mice received a single dosage of AAV2 with 10⁵ and 10⁶ infectious units in 200 μ l PBS, 200 μ l PBS via intratumoral injections or not manipulated (Mock). On day 32, the mock and PBS-treated mice are sacrificed due to the grossly restricted ability to move and to feed arising from the tumor burden. The AAV2 treated group survived longer than the control mice and did not display similar restrictions. The mice are eventually euthanized on day 56. Statistical analyses are expressed as mean $\pm$ SD. (B) AAV2 infection of MDA-MB-435 xenografts induced dark patches of necrotic areas on the skin surrounding the tumors. Tumors are examined using hematoxylin and eosin staining (100 $\times$ magnification). Both mock and PBS-injected tumors showed areas of infiltrating macrophages as indicated with arrows, whereas AAV2-treated tumors showed large areas of tumor necrosis (N) that correlated with the appearance of dark patches on the external surface. AAV2: Adeno-associated virus type 2; IM: infiltrating macrophage; PBS: Phosphate-buffered saline.

Adapted with permission from [96].

1.5 mm as spot size for 5 min. The fluorescence microscope images confirmed the ablation of tumor cells mediated by gold nanoshells (Figure 7) [123]. Clearly localized regions of dead cells corresponding to the spot with laser exposure surrounded by live cells bounded with gold nanoshells without irradiation were observed in all HER2 positive breast cancer cells, regardless of the sensitive or resistant to trastuzumab, suggesting the potential of nanoshells-mediated ablation for the treatment of different types of breast cancer [123].

It should be noted that cancer cell death is caused by necrosis in PTA and the low metabolic activity of necrotic area [130] makes the antibiotics, antiseptics and disinfectants ineffective in killing present bacteria [131], leading to biofilm formation and chronic wounds [132]. Moreover, most nanomaterials used for PTA are metallic nanoparticles, which typically encounter ineffective clearance *in vivo* [133,134]. Therefore, some organic nanomaterials, such as conjugated polymeric nanoparticles also have been studied for PTA [115,135]. Future research on nanoparticle-mediated PTA should also focus on decreasing the dosage of photothermal coupling agents and the power of laser irradiation in order to minimize the potential damage to surrounding healthy cells.

Calcium phosphosilicate nanoparticles

As a prominent nanoparticulate-based drug delivery system, calcium phosphosilicate nanoparticles (CPSNPs) can be used to encapsulate many kinds of chemotherapeutics [5,6]. Calcium phosphate (CP) is a naturally occurring biomineral and CP-based nanoparticles are regarded as bioresorbable, innocuous, biocompatible and biodegradable [136–138]. Moreover, as a stable bioceramic, CP is relatively insoluble at physiological pH (pH 7.4), such as in the blood stream, but more soluble in acidic environments, such as solid tumor microenvironment (pH 5.8–7.8) and the intercellular late stage endo-lysosomes (pH 4.8–5.5), where the encapsulants are released after endocytosis [139–143]. Silicate decreases the solubility of the amorphous phase of CP, leaving CPSNPs without concerns of swelling or porosity [144]. Docetaxel-doped CPSNPs (Docetaxel-CPSNPs) are emerging as a useful nanomedicine in breast cancer treatment. Docetaxel, another member of the taxane family can be encapsulated within CPSNPs, which can be surface functionalized with either PEG for passive accumulation via enhanced permeability and retention effects or anti-CD71 for active targeting to breast cancer cells [145,146]. It has been demonstrated by Barth *et al.* that the targeting of CD71-CPSNPs to breast cancer cells are effective based on an *in vivo* study with an athymic nude mice model (Figure 8) [146]. CPSNPs are colloiddally stable

in the blood stream, protecting docetaxel from degradation and other unfavorable chemical or biological conditions during transport [140,141]. As a result, CD71-Docetaxel-CPSNPs can provide prolonged circulation time, enhanced accumulation and targeting, and improved drug efficiency with reduced or eliminated side effects. Future study should focus on improving the encapsulation of docetaxel and other chemotherapeutics for breast cancer treatment.

CPSNPs doped with a NIR fluorophore, indocyanine green (ICG-CPSNPs) used for NIR bioimaging can also act as a photosensitizer for the novel PhotoImmunoNano Therapy (PINT) as reported by Barth *et al.* (Figure 9) [147,148]. Athymic nude mice with MDA-MB-231 breast cancer tumors (T-cell deficient) and BALB/cJ mice with murine 410.4 breast cancer cells (T-cell competent) were injected with PEGylated ICG-CPSNPs and PBS, free ICG, PEGylated CPSNPs without encapsulant and citrate surface-functionalized ICG-CPSNPs as control followed 24 h later by NIR laser treatment (12.5 J/cm^2) at the tumor location [148]. The tumor growth in both models was effectively suppressed in the following 45 days after PINT mediated by PEGylated ICG-CPSNPs [148]. This relatively long-term suppression of tumor growth was caused by T-cell independent antitumor immune response as the result of the decrease in inflammation-expanded immature myeloid cells and the expansion of lymphocytes regulated by phosphorylated sphingolipid metabolites (Figure 10) [148]. Since PINT is a treatment based on the activation of the immune system, this therapy does not require continual injection of aggressive chemotherapeutics or exposure to high-dose radiation. ICG-mediated PINT is a revolutionary therapeutic approach for breast cancers which are inflammatory or immunosuppressed. More significantly, PINT was used to treat chronic myeloid leukemia [147], the first time a nonsolid tumor was treated with phototherapy, suggesting the application in the targeted treatment of metastatic breast cancer cells.

Continuing challenges in nanoparticulate developments

There are still many barriers to overcome before extensive clinical use of novel nanotherapies for human breast cancer can become standard treatment protocols. One is to improve the loading concentration and encapsulation efficiency of active agents. These two parameters used to evaluate nanomedicine can be the major drawbacks of some nanoparticle-based delivery platforms. A therapeutic concentration of drug or imaging contrast agents is essential to achieve anticancer effects or quality imaging of tumors. With a high drug loading capacity, the quantities of matrix

Figure 6. Delivery of RPN2 siRNA to orthotopic docetaxel-resistant breast tumors by RPN2 siRNA-ateolocollagen complex. (A) Reduction of MCF7-ADR breast tumor volume in mice given RPN2 siRNA or control nontargeting siRNA along with docetaxel (n = 6 per group, *p < 0.01). (B) siRNA-treated MCF7-ADR tumors in mice before and 7 days after docetaxel treatment. (C) Growth of MDA-MB-231/MDR1 breast tumor in mice administered RPN2 siRNA or nontargeting siRNA along with docetaxel (n = 6 per group, *p < 0.002). (D) MDA-MB-231/MDR1 tumors in mice 7 days after treatment with siRNA and docetaxel. (E) TUNEL staining of MCF7-ADR tumor tissues treated with RPN2 siRNAs or nontargeting siRNAs in the presence or absence of docetaxel. (F) TUNEL-positive cells are counted and are represented in the graph (n = 3 per group, *p < 0.01). (H) Expression of RPN2 mRNA in MCF7-ADR tumors treated with RPN2 siRNAs or nontargeting siRNAs (n = 3 per group, *p < 0.01). (H) Expression of RPN2 protein in MCF7-ADR tumors. H&E staining and RPN2 immunofluorescence staining (green, RPN2; blue, nuclei) of tissues treated with RPN2 siRNA or nontargeting siRNA. (I) Docetaxel retention in MCF7-ADR tumors in mice treated with RPN2 siRNAs or nontargeting siRNAs (n = 4 per group, *p < 0.001). Values are mean $\pm$ SD. ADR: Adriamycin resistant; TUNEL: TdT-mediated dUTP nick-end labeling. Adapted with permission from [109].

Figure 7. *In vitro* photothermal ablation of breast cancer cells mediated by silica-gold nanoshells. (A) Live/dead imaging demonstrates cell ablation in HER2-positive cell lines (SK-BR-3, JIMT-1, BT474 AZ LR) treated with anti-HER2 nanoshells and near infrared laser irradiation. (B) Silver staining shows nanoshells binding for each of the HER2-positive cell lines.

Reprinted with permission from [123].

nanomaterials to be administrated are reduced along with nanoparticle-related risks. Moreover, chemotherapeutics and other bioactive molecules, such as DNA, RNA and peptides, are typically costly or difficult to synthesize. Therefore, a high percentage (>50 mol%) of the encapsulant should be entrapped in the nanocarriers. The encapsulation of active agents is affected by several factors including the matrix composition and size, encapsulant size and structure, the interactions between encapsulants and nanovectors and synthetic methods. Various modifications on widely used lipid and polymeric nanoparticles have been exploited to promote the encapsulation [149–151]. However, improvements in loading capacity and entrapment efficiency remain as challenges for the translation of laboratory research to clinical trials for some novel nanocarriers.

Another obstacle in nanomedicine development is to select a proper targeting moiety. Nanoparticles that go through blood circulation should eventually clear from the body. A concern that stems from systemic circulation is the safety of targeting ligands, which must be nontoxic and biocompatible. Targeting of nanomedicine is based on the interaction between targeting agents and receptors on tumor cells with specificity, which is a primary selection criterion for targeting

agents. However, most relative receptors are expressed by both cancer cells and in other major organs. As a result, targeted nanoparticles that accumulate in both the target tumors and organs compromise the efficiency of nanomedicine and can potentially induce negative side effects to healthy tissue.

Targeting ligands can also colloiddally stabilize nanoparticles in physiological environments. Nanoparticle agglomeration imposes adverse effects on mechanical performance [152], affects the bioactivity [153] and leads to unwanted rapid clearance from the bloodstream [154]. Overall, the selection of a targeting ligand is a critical step in nanomedicine development that includes antibodies, aptamers, proteins, peptides and small molecules [155]. In addition to the scientific efficacy, the cost and accessibility of targeting ligands should also be evaluated and future attention should shift more toward early detection nanotechnology that target small tumor lesions.

Overall, the translation of discoveries from fundamental research to clinical trials is a highly selective process for evidence-based nanomedicine. Technical challenges such as the ability to scale-up for manufacturing and sterilization to produce clinical-grade nanoparticles are common barriers to this translation. The physicochemical features of nanoparticles (com-

Figure 8. Targeting transferrin receptors in an *in vivo* subcutaneous tumor model of breast cancer. (A) Schematic of calcium-phosphate nanoparticles (CPNPs). (B) transmission-electron micrograph of ~16 nm (mean diameter) indocyanine-green (ICG) CPNPs. (C) Human MDA-MB-231 metastatic breast cancer cells are analyzed via flow cytometry for the presence of the transferrin receptor (CD71). (D) MDA-MB-231 cells are xenografted subcutaneously into athymic nude mice. One week following engraftment, ICG-loaded CPNPs are administered systemically via tail vein injection and near-infrared images are taken 96 h post injection. From left to right, mice received (i) free ICG, (ii) ICG-loaded PEG-CPNPs, (iii) ICG-loaded anti-CD71-Avidin-CPNPs or (iv) ICG-loaded Human Holotransferrin-Avidin-CPNPs. (E) Excised tumors (mice ii, iii and iv from panel B) and spleen and stomach (mouse ii). All images are representative of four independent experiments.

Adapted with permission from [146,170].

position, structure, size, surface modification and colloidal stability) all contribute to the clinical outcome. Any small batch-to-batch variation in nanoparticle characterization can cause dramatic changes in material properties, such as toxicity, biocompatibility and *in vitro* and *in vivo* performances [156]. Therefore, the scale-up synthesis of nanoparticles must be reliable and consistent down to the nanoscale.

Conclusion & future perspective

As a rapidly developing interdisciplinary field, cancer nanotechnology brings various perspectives beyond conventional breast cancer treatment that are potentially safer and more efficacious. Several nanoparticle-based systems have been approved by FDA or are in clinical trials for the delivery of chemotherapeutics (such as liposomes, polymeric nanoparticles and nab

Figure 9. PhotoImmunoNanoTherapy decreases tumor burden and improves survival. (A) Tumor growth following PhotoImmunoNanoTherapy is monitored in athymic nude mice engrafted with human MDA-MB-231 breast cancer cells. ANOVA, $*p < 0.05$ compared with all, $n \geq 5$. (B) Tumor growth following PhotoImmunoNanoTherapy is monitored in BALB/cJ mice engrafted with murine 410.4 breast cancer cells. ANOVA, $*p < 0.05$ compared with all, $n \geq 8$. (C) Tumor growth following PhotoImmunoNanoTherapy is monitored in NOD.CB17-Prkdcscid/J mice engrafted with murine 410.4 breast cancer cells. ANOVA, $*p < 0.001$ compared with all, $n \geq 7$. (D) Tumor growth following PhotoImmunoNanoTherapy is monitored in C57BL/6J mice engrafted with murine Panc-02 pancreatic cancer cells. ANOVA, $*p < 0.05$ compared with all, $n \geq 6$. (E) Survival of athymic nude mice orthotopically implanted with human BxPC-3-GFP pancreatic cancer cells is monitored following PhotoImmunoNanoTherapy. Logrank test, $*p < 0.05$ compared with all, $n = 5$. (F) Survival of athymic nude mice with experimental lung metastases of human SAOS-2-LM7 osteosarcoma cells is monitored following PhotoImmunoNanoTherapy. Log-rank test, $*p < 0.05$, $n = 5$. (G) Survival of C3H/HeJ mice established with 32D-p210-GFP myeloid leukemia cells is monitored following PhotoImmunoNanoTherapy. Log-rank test, $*p < 0.05$ compared with all, $n = 7$. CD117: CD117 antibody bioconjugated; COOH: Citrate-functionalized; CPSNP: Calcium phosphosilicate nanoparticles; Ghost: Empty; ICG: Indocyanine green; PEG: PEGylated. Adapted with permission from [147,148].

Figure 10. PhotoImmunoNanoTherapy utilizing indocyanine green-alcium phosphosilicate nanoparticles-PEG exerts an antitumor effect by generating dihydrosphingosine-1-phosphate. Immunosuppressive cells that are often expanded as a function of tumor burden, namely, immature myeloid cells, are ablated by dihydrosphingosine-1-phosphate. This process allows the expansion of immune effector cells from hematopoietic progenitor cells and subsequently leads to an antitumor immune response. The mechanism is in addition, and independent from, traditional oxidative-mediated tumor cell death that is a hallmark of photodynamic therapy. Reprinted with permission from [148].

paclitaxel). To promote the treatment of breast cancer with nanotherapies, there are still many scientific contributions that can be made. For instance, there is a need to improve the treatment outcome of breast cancer types with higher recurrence rates and mortality rates such as HER2-positive or triple negative breast cancer. HER2 targeting nanotherapies based on HER2 targeting ligand such as trastuzumab, pertuzumab and lapatinib significantly increase the efficacy of HER2-

positive breast cancer treatments. While AAV2-induced necrosis of tumor tissues offers a novel solution to treat TNBC, safety regarding the use of virus vectors is generally the primary concern in clinical trial advancement.

Emerging anticancer mechanisms have also attracted attention in reference to RNAi and PTA. RNAi focuses on the treatment of cancer at gene level by degrading mRNA or suppressing RNA translation in cancerous cells when antibodies or inhibitors cannot target successfully. Immune-related side effects are the caveats of RNAi therapeutic research since hormone receptors are involved. With nanoparticle-mediated PTA, cancer cells are eradicated by the noninvasive treatment, which is beneficial for patients that are irresponsive to chemotherapeutic or radiation treatment.

Lastly, inventing novel nanocarriers is an alternate route to creating biological targets within the scope of perspective nanotherapies. Chemotherapeutics, hormones, RNA molecules or viruses can be encapsulated within nontoxic CPSNPs and be released in tumor cells in response to a pH trigger. Targeting ligands, such as antibodies, aptamers, peptides and proteins can be attached on the surface of CPSNPs via bioconjugation to direct the accumulation of CPNSPs within tumor tissues. The encapsulation of NIR fluorophore within CPSNPs also expands application to PTA. Overall, the CPSNP is a versatile nanocarrier for all kinds of active agents to encompass the many different types of therapies mentioned above, making CPSNPs-based nanotherapies and other personalized nanocarriers in general promising for cancer treatment.

In conclusion, the versatility of nanoparticulates allows for the delivery of multiple active agents with the ability to target various types of cancer. This leads

Executive summary

Background

- Cancer nanomedicine provides potential benefits to breast cancer treatment, including increased biocompatibility, prolonged blood circulation, targeted delivery and reduced side effects.
- Nanotechnology strategies for biomedical applications can be screened using a material selection criterion.

Nanotherapies for breast cancer

- Many nanoparticulate-based chemotherapeutic delivery platforms have been approved by the US FDA or are in clinical trials for breast cancer treatment, such as formulations based on liposomes, polymeric nanoparticles and nanoparticle albumin-bound paclitaxel.
- Innovative nanotherapies targeting HER2 breast cancer and triple negative breast cancer or nanotherapies based on RNA interference, nanoparticle-mediated photothermal ablation and calcium phosphosilicate nanoparticles bring various perspectives beyond conventional breast cancer treatment.
- Common obstacles include loading concentration and encapsulation efficiency of active agents, the selection of a proper targeting moiety and in manufacturing consistent clinical-grade nanoparticles at the industrial scale.

Conclusion & future perspective

- Novel nanocarriers such as CPSNPs are promising because the ability to target and treat various types of cancers depends on the design.
- The development of versatile nanomedicine is challenging, but significant in offering better alternatives to standard cancer procedures.

to improved efficacy and advancements in both diagnosis and treatment. These unique properties make nanomedicine challenging to study, but at the same time, appealing to the scientific community aiming to improve patient outcome.

Financial & competing interests disclosure

This review was funded partially by NIH grants R01CA167535 and R21CA170121 from the National Cancer Institute (NCI) for the CPSNP work. The project was also supported in part under a grant with the Pennsylvania Department of Health

using Tobacco CURE funds (SAP#4100072562 to GL Matters and JH Adair). Penn State Research Foundation has licensed CPSNP technology to Keystone Nano, Inc. (PA, USA). JH Adair and M Kester are co-founders of Keystone Nano and are CSO and CMO, respectively. The authors have no other relevant affiliations or financial involvement with any organization or entity with a financial interest in or financial conflict with the subject matter or materials discussed in the manuscript apart from those disclosed.

No writing assistance was utilized in the production of this manuscript.

References

Papers of special note have been highlighted as: • of interest; •• of considerable interest

- 1 Siegel RL, Miller KD, Jemal A. Cancer statistics, 2016. *CA Cancer J. Clin.* 66(1), 7–30 (2016).
- 2 Pogo BG, Holland JF. Possibilities of a viral etiology for human breast cancer. A review. *Biol. Trace Elem. Res.* 56(1), 131–142 (1997).
- 3 Hortobagyi GN. Treatment of breast cancer. *N. Engl. J. Med.* 339(14), 974–984 (1998).
- 4 Wang R, Billone PS, Mullett WM. Nanomedicine in action: an overview of cancer nanomedicine on the market and in clinical trials. *J. Nanomater.* 2013, 1–12 (2013).
- 5 Adair JH, Parette MP, Altinoglu EI, Kester M. Nanoparticulate alternatives for drug delivery. *ACS Nano* 4(9), 4967–4970 (2010).
- Provides a perspective for nanoparticulate design and criteria for drug delivery.
- 6 Altinoglu EI, Adair JH. Near infrared imaging with nanoparticles. *Wiley Interdiscip. Rev. Nanomed. Nanobiotechnol.* 2(5), 461–477 (2010).
- 7 Yoshimori T. Autophagy: a regulated bulk degradation process inside cells. *Biochem. Biophys. Res. Commun.* 313(2), 453–458 (2004).
- 8 Stern ST, Adiseshaiah PP, Crist RM. Autophagy and lysosomal dysfunction as emerging mechanisms of nanomaterial toxicity. *Part. Fibre Toxicol.* 9, 20 (2012).
- Covers a recently discovered and significant toxicity mechanism by nanomaterials in cells. Materials that have been long believed to be nontoxic (such as gold nanoparticles), can also induce toxicity by disrupting cellular function.
- 9 Li JJ, Hartono D, Ong CN, Bay BH, Yung LY. Autophagy and oxidative stress associated with gold nanoparticles. *Biomaterials* 31(23), 5996–6003 (2010).
- 10 Stern ST, Zolnik BS, Mcleland CB, Clogston J, Zheng J, Mcneil SE. Induction of autophagy in porcine kidney cells by quantum dots: a common cellular response to nanomaterials? *Toxicol. Sci.* 106(1), 140–152 (2008).
- 11 Zhao Y, Howe JL, Yu Z *et al.* Exposure to titanium dioxide nanoparticles induces autophagy in primary human keratinocytes. *Small* 9(3), 387–392 (2013).
- 12 Yu KN, Yoon TJ, Minai-Tehrani A *et al.* Zinc oxide nanoparticle induced autophagic cell death and mitochondrial damage via reactive oxygen species generation. *Toxicol. In Vitro* 27(4), 1187–1195 (2013).
- 13 Zhang Y, Yu C, Huang G, Wang C, Wen L. Nano rare-earth oxides induced size-dependent vacuolization: an independent pathway from autophagy. *Int. J. Nanomedicine* 5, 601–609 (2010).
- 14 Zhang Q, Yang W, Man N *et al.* Autophagy-mediated chemosensitization in cancer cells by fullerene C60 nanocrystal. *Autophagy* 5(8), 1107–1117 (2009).
- 15 Johnson-Lyles DN, Peifley K, Lockett S *et al.* Fullerenol cytotoxicity in kidney cells is associated with cytoskeleton disruption, autophagic vacuole accumulation, and mitochondrial dysfunction. *Toxicol. Appl. Pharmacol.* 248(3), 249–258 (2010).
- 16 Liu HL, Zhang YL, Yang N *et al.* A functionalized single-walled carbon nanotube-induced autophagic cell death in human lung cells through Akt-TSC2-mTOR signaling. *Cell Death Dis.* 2, e159 (2011).
- 17 Shapiro CL, Recht A. Side effects of adjuvant treatment of breast cancer. *N. Engl. J. Med.* 344(26), 1997–2008 (2001).
- 18 Group EBCTC. Comparisons between different polychemotherapy regimens for early breast cancer: meta-analyses of long-term outcome among 100,000 women in 123 randomised trials. *Lancet* 379(9814), 432–444 (2012).
- Highlights the standard protocols (anthracyclines and taxanes) for treating early breast cancer in the last three decades, regardless of known harmful side effects. Emphasizes the need to improve chemotherapy for targeting capabilities.
- 19 Smith LA, Cornelius VR, Plummer CJ *et al.* Cardiotoxicity of anthracycline agents for the treatment of cancer: systematic review and meta-analysis of randomised controlled trials. *BMC Cancer* 10, 337 (2010).
- 20 Nurgalieva Z, Liu CC, Du XL. Chemotherapy use and risk of bone marrow suppression in a large population-based cohort of older women with breast and ovarian cancer. *Med. Oncol.* 28(3), 716–725 (2011).
- 21 Lee C, Gianos M, Klaustermeyer WB. Diagnosis and management of hypersensitivity reactions related to common cancer chemotherapy agents. *Ann. Allergy Asthma Immunol.* 102(3), 179–187 (2009).

- 22 Wyatt AJ, Leonard GD, Sachs DL. Cutaneous reactions to chemotherapy and their management. *Am. J. Clin. Dermatol.* 7(1), 45–63 (2006).
- 23 Hagiwara H, Sunada Y. Mechanism of taxane neurotoxicity. *Breast Cancer* 11(1), 82–85 (2004).
- 24 Torchilin VP. *Nanoparticulates as Drug Carriers*. Imperial College Press, London, UK (2006).
- 25 He L, Gu J, Lim LY, Yuan ZX, Mo J. Nanomedicine-mediated therapies to target breast cancer stem cells. *Front. Pharmacol.* 7, 313 (2016).
- 26 Bangham AD, Horne RW. Negative staining of phospholipids and their structural modification by surface-active agents as observed in the electron microscope. *J. Mol. Biol.* 8(5), 660–668 (1964).
- 27 Gregoria G, Ryman BE. Liposomes as carriers of enzymes or drugs – new approach to treatment of storage diseases. *Biochem. J.* 124(5), P58 (1971).
- 28 Torchilin VP. Liposomes as targetable drug carriers. *Crit. Rev. Ther. Drug Carrier Syst.* 2(1), 65–115 (1985).
- 29 Torchilin VP. Recent advances with liposomes as pharmaceutical carriers. *Nat. Rev. Drug Discov.* 4(2), 145–160 (2005).
- 30 Sawant RR, Torchilin VP. Challenges in development of targeted liposomal therapeutics. *AAPS J.* 14(2), 303–315 (2012).
- 31 Chang DK, Li PC, Lu RM, Jane WN, Wu HC. Peptide-mediated liposomal doxorubicin enhances drug delivery efficiency and therapeutic efficacy in animal models. *PLoS ONE* 8(12), e83239 (2013).
- 32 Mann AP, Bhavane RC, Somasunderam A *et al.* Thioaptamer conjugated liposomes for tumor vasculature targeting. *Oncotarget* 2(4), 298–304 (2011).
- 33 Barenholz Y. Doxil(R) – the first FDA-approved nano-drug: lessons learned. *J. Control. Release* 160(2), 117–134 (2012).
- 34 O'Brien MER. Reduced cardiotoxicity and comparable efficacy in a Phase III trial of pegylated liposomal doxorubicin HCl (CAELYXTM/Doxil™) versus conventional doxorubicin for first-line treatment of metastatic breast cancer. *Ann. Oncol.* 15(3), 440–449 (2004).
- 35 Full prescribing information of doxorubicin hydrochloride. www.accessdata.fda.gov
- 36 Full prescribing information of Doxil(R). www.accessdata.fda.gov
- 37 Sharma A. Liposomes in drug delivery: progress and limitations. *Int. J. Pharm.* 154(2), 123–140 (1997).
- 38 Sharma A, Mayhew E, Bolcsak L *et al.* Activity of paclitaxel liposome formulations against human ovarian tumor xenografts. *Int. J. Cancer* 71(1), 103–107 (1997).
- 39 Vilar G, Tulla-Puche J, Albericio F. Polymers and drug delivery systems. *Curr. Drug Deliv.* 9(4), 367–394 (2012).
- 40 Chan JM, Valencia PM, Zhang L, Langer R, Farokhzad OC. Polymeric nanoparticles for drug delivery. *Methods Mol. Biol.* 624, 163–175 (2010).
- 41 Chandna P, Khandare JJ, Ber E, Rodriguez-Rodriguez L, Minko T. Multifunctional tumor-targeted polymer-peptide-drug delivery system for treatment of primary and metastatic cancers. *Pharm. Res.* 27(11), 2296–2306 (2010).
- 42 Vivek R, Thangam R, Nipunbabu V *et al.* Multifunctional HER2-antibody conjugated polymeric nanocarrier-based drug delivery system for multi-drug-resistant breast cancer therapy. *ACS Appl. Mater. Interfaces* 6(9), 6469–6480 (2014).
- 43 Du JZ, Du XJ, Mao CQ, Wang J. Tailor-made dual pH-sensitive polymer-doxorubicin nanoparticles for efficient anticancer drug delivery. *J. Am. Chem. Soc.* 133(44), 17560–17563 (2011).
- 44 Fitzpatrick SD, Fitzpatrick LE, Thakur A, Mazumder MA, Sheardown H. Temperature-sensitive polymers for drug delivery. *Expert Rev. Med. Devices* 9(4), 339–351 (2012).
- 45 Shamay Y, Adar L, Ashkenasy G, David A. Light induced drug delivery into cancer cells. *Biomaterials* 32(5), 1377–1386 (2011).
- 46 Lensen D, Gelderblom EC, Vriezema DM *et al.* Biodegradable polymeric microcapsules for selective ultrasound-triggered drug release. *Soft Matter* 7(11), 5417 (2011).
- 47 Liechty WB, Kryscio DR, Slaughter BV, Peppas NA. Polymers for drug delivery systems. *Annu. Rev. Chem. Biomol. Eng.* 1, 149–173 (2010).
- 48 Verma RK, Mishra B, Garg S. Osmotically controlled oral drug delivery. *Drug Dev. Ind. Pharm.* 26(7), 695–708 (2000).
- 49 Clinical Trials Database: NCT00876486. <https://clinicaltrials.gov/ct2/show/NCT00876486>
- 50 Kim TY, Kim DW, Chung JY *et al.* Phase I and pharmacokinetic study of Genexol-PM, a cremophor-free, polymeric micelle-formulated paclitaxel, in patients with advanced malignancies. *Clin. Cancer Res.* 10(11), 3708–3716 (2004).
- 51 Mackler NJ, Pienta KJ. Drug insight: use of docetaxel in prostate and urothelial cancers. *Nat. Clin. Pract. Urol.* 2(2), 92–100; quiz 101 p following 112 (2005).
- 52 Chu E, Devita VT. *Physicians' Cancer Chemotherapy Drug Manual 2013*. Jones & Bartlett Learning LLC, MA, USA (2012).
- 53 Dong Y, Feng SS. *In vitro* and *in vivo* evaluation of methoxy polyethylene glycol-poly(lactide) (MPEG-PLA) nanoparticles for small-molecule drug chemotherapy. *Biomaterials* 28(28), 4154–4160 (2007).
- 54 Lee KS, Chung HC, Im SA *et al.* Multicenter Phase II trial of Genexol-PM, a Cremophor-free, polymeric micelle formulation of paclitaxel, in patients with metastatic breast cancer. *Breast Cancer Res. Treat.* 108(2), 241–250 (2008).
- 55 Gelderblom H, Verweij J, Nooter K, Sparreboom A. Cremophor EL: the drawbacks and advantages of vehicle selection for drug formulation. *Eur. J. Cancer* 37(13), 1590–1598 (2001).
- 56 Ma Z, Moulton B. Recent advances of discrete coordination complexes and coordination polymers in drug delivery. *Coord. Chem. Rev.* 255(15–16), 1623–1641 (2011).
- 57 Gibson M. *Pharmaceutical Preformulation and Formulation: A Practical Guide from Candidate Drug Selection to Commercial Dosage Form*. Informa Healthcare USA (2009).

- 58 Han E-J, Chung A-H, Oh I-J. Analysis of residual solvents in poly(lactide-co-glycolide) nanoparticles. *J. Pharm. Invest.* 42(5), 251–256 (2012).
- 59 Wani MC, Taylor HL, Wall ME, Coggon P, Mcphail AT. Plant antitumor agents. VI. The isolation and structure of taxol, a novel antileukemic and antitumor agent from *Taxus brevifolia*. *J. Am. Chem. Soc.* 93(9), 2325–2327 (1971).
- 60 Jordan MA, Wilson L. Microtubules as a target for anticancer drugs. *Nat. Rev. Cancer* 4(4), 253–265 (2004).
- 61 Gelderblom H, Verweij J, Nooter K, Sparreboom A. Cremophor EL: the drawbacks and advantages of vehicle selection for drug formulation. *Eur. J. Cancer* 37(13), 1590–1598 (2001).
- 62 Desai N. Nanoparticle albumin-bound paclitaxel (abraxane®). In: *Albumin in Medicine: Pathological and Clinical Applications*. Otagiri M, Chuang VTG (Eds). Springer, Singapore, 101–119 (2016).
- 63 Hawkins MJ, Soon-Shiong P, Desai N. Protein nanoparticles as drug carriers in clinical medicine. *Adv. Drug Del. Rev.* 60(8), 876–885 (2008).
- 64 Ibrahim NK, Desai N, Legha S *et al.* Phase I and pharmacokinetic study of ABI-007, a Cremophor-free, protein-stabilized, nanoparticle formulation of paclitaxel. *Clin. Cancer Res.* 8(5), 1038–1044 (2002).
- 65 Miele E, Spinelli GP, Miele E, Tomao F, Tomao S. Albumin-bound formulation of paclitaxel (Abraxane ABI-007) in the treatment of breast cancer. *Int. J. Nanomedicine* 4, 99–105 (2009).
- 66 Paal K, Muller J, Hegedus L. High affinity binding of paclitaxel to human serum albumin. *Eur. J. Biochem.* 268(7), 2187–2191 (2001).
- 67 Sparreboom A, Scripture CD, Trieu V *et al.* Comparative preclinical and clinical pharmacokinetics of a cremophor-free, nanoparticle albumin-bound paclitaxel (ABI-007) and paclitaxel formulated in Cremophor (Taxol). *Clin. Cancer Res.* 11(11), 4136–4143 (2005).
- 68 Desai N, Trieu V, Yao Z *et al.* Increased antitumor activity, intratumor paclitaxel concentrations, and endothelial cell transport of cremophor-free, albumin-bound paclitaxel, ABI-007, compared with cremophor-based paclitaxel. *Clin. Cancer Res.* 12(4), 1317–1324 (2006).
- 69 Meric-Bernstam F, Hung MC. Advances in targeting human epidermal growth factor receptor-2 signaling for cancer therapy. *Clin. Cancer Res.* 12(21), 6326–6330 (2006).
- 70 Graus-Porta D, Beerli RR, Daly JM, Hynes NE. ErbB-2, the preferred heterodimerization partner of all ErbB receptors, is a mediator of lateral signaling. *EMBO J.* 16(7), 1647–1655 (1997).
- 71 Arteaga CL, Sliwkowski MX, Osborne CK, Perez EA, Puglisi F, Gianni L. Treatment of HER2-positive breast cancer: current status and future perspectives. *Nat. Rev. Clin. Oncol.* 9(1), 16–32 (2012).
- 72 Yarden Y. The EGFR family and its ligands in human cancer: signalling mechanisms and therapeutic opportunities. *Eur. J. Cancer* 37, 3–8 (2001).
- 73 Pinkas-Kramarski R, Lenferink AE, Bacus SS *et al.* The oncogenic ErbB-2/ErbB-3 heterodimer is a surrogate receptor of the epidermal growth factor and betacellulin. *Oncogene* 16(10), 1249–1258 (1998).
- 74 Tan M, Yao J, Yu DH. Overexpression of the c-erbB-2 gene enhanced intrinsic metastasis potential in human breast cancer cells without increasing their transformation abilities. *Cancer Res.* 57(6), 1199–1205 (1997).
- 75 Zhou BP, Hu MCT, Miller SA *et al.* HER-2/neu blocks tumor necrosis factor-induced apoptosis via the Akt/NF-kappa B pathway. *J. Biol. Chem.* 275(11), 8027–8031 (2000).
- 76 Carter P, Presta L, Gorman CM *et al.* Humanization of an anti-p185HER2 antibody for human cancer therapy. *Proc. Natl Acad. Sci. USA* 89(10), 4285–4289 (1992).
- 77 Hudis CA. Trastuzumab—mechanism of action and use in clinical practice. *N. Engl. J. Med.* 357(1), 39–51 (2007).
- 78 Park S, Jiang Z, Mortenson ED *et al.* The therapeutic effect of anti-HER2/neu antibody depends on both innate and adaptive immunity. *Cancer Cell* 18(2), 160–170 (2010).
- 79 Hudziak RM, Lewis GD, Winget M, Fendly BM, Shepard HM, Ullrich A. p185HER2 monoclonal antibody has antiproliferative effects *in vitro* and sensitizes human breast tumor cells to tumor necrosis factor. *Mol. Cell. Biol.* 9(3), 1165–1172 (1989).
- 80 Pegram M, Hsu S, Lewis G *et al.* Inhibitory effects of combinations of HER-2/neu antibody and chemotherapeutic agents used for treatment of human breast cancers. *Oncogene* 18(13), 2241–2251 (1999).
- 81 Romond EH, Perez EA, Bryant J *et al.* Trastuzumab plus adjuvant chemotherapy for operable HER2-positive breast cancer. *N. Engl. J. Med.* 353(16), 1673–1684 (2005).
- 82 Van Pelt AE, Mohsin S, Elledge RM *et al.* Neoadjuvant trastuzumab and docetaxel in patients with breast cancer: preliminary results. *Clin. Breast Cancer* 4(5), 348–353 (2003).
- 83 Ligibel JA, Winer EP. Trastuzumab/chemotherapy combinations in metastatic breast cancer. *Semin. Oncol.* 29(3 Suppl. 11), 38–43 (2002).
- 84 Austin CD, De Maziere AM, Pisacane PI *et al.* Endocytosis and sorting of ErbB2 and the site of action of cancer therapeutics trastuzumab and geldanamycin. *Mol. Biol. Cell* 15(12), 5268–5282 (2004).
- 85 Anhorn MG, Wagner S, Kreuter J, Langer K, Von Briesen H. Specific targeting of HER2 overexpressing breast cancer cells with doxorubicin-loaded trastuzumab-modified human serum albumin nanoparticles. *Bioconjug. Chem.* 19(12), 2321–2331 (2008).
- 86 Takai N, Jain A, Kawamata N *et al.* 2C4, a monoclonal antibody against HER2, disrupts the HER kinase signaling pathway and inhibits ovarian carcinoma cell growth. *Cancer* 104(12), 2701–2708 (2005).
- 87 Gianni L, Pienkowski T, Im YH *et al.* Efficacy and safety of neoadjuvant pertuzumab and trastuzumab in women with locally advanced, inflammatory, or early HER2-positive breast cancer (NeoSphere): a randomised multicentre, open-label, Phase II trial. *Lancet Oncol.* 13(1), 25–32 (2012).
- 88 Baselga J, Cortes J, Kim SB *et al.* Pertuzumab plus trastuzumab plus docetaxel for metastatic breast cancer. *N. Engl. J. Med.* 366(2), 109–119 (2012).

- 89 Baselga J, Gelmon KA, Verma S *et al.* Phase II trial of pertuzumab and trastuzumab in patients with human epidermal growth factor receptor 2-positive metastatic breast cancer that progressed during prior trastuzumab therapy. *J. Clin. Oncol.* 28(7), 1138–1144 (2010).
- 90 Gao H, Cao S, Chen C *et al.* Incorporation of lapatinib into lipoprotein-like nanoparticles with enhanced water solubility and anti-tumor effect in breast cancer. *Nanomedicine (Lond.)* 8(9), 1429–1442 (2013).
- 91 Gao H, Chen C, Xi Z *et al.* *In vivo* behavior and safety of lapatinib-incorporated lipid nanoparticles. *Curr. Pharm. Biotechnol.* 14(12), 1062–1071 (2014).
- 92 Foulkes WD, Smith IE, Reis-Filho JS. Triple-negative breast cancer. *N. Engl. J. Med.* 363(20), 1938–1948 (2010).
- 93 Hudis CA, Gianni L. Triple-negative breast cancer: an unmet medical need. *Oncologist* 16(Suppl. 1), 1–11 (2011).
- 94 Tan DS, Marchio C, Jones RL *et al.* Triple negative breast cancer: molecular profiling and prognostic impact in adjuvant anthracycline-treated patients. *Breast Cancer Res. Treat.* 111(1), 27–44 (2008).
- 95 Liedtke C, Mazouni C, Hess KR *et al.* Response to neoadjuvant therapy and long-term survival in patients with triple-negative breast cancer. *J. Clin. Oncol.* 26(8), 1275–1281 (2008).
- 96 Alam S, Bowser BS, Israr M, Conway MJ, Meyers C. Adeno-associated virus type 2 infection of nude mouse human breast cancer xenograft induces necrotic death and inhibits tumor growth. *Cancer Biol. Ther.* 15(8), 1013–1028 (2014).
- 97 Amos KD, Adamo B, Anders CK. Triple-negative breast cancer: an update on neoadjuvant clinical trials. *Int. J. Breast Cancer* 2012, 385978 (2012).
- 98 Rana TM. Illuminating the silence: understanding the structure and function of small RNAs. *Nat. Rev. Mol. Cell Biol.* 8(1), 23–36 (2007).
- 99 Pecot CV, Calin GA, Coleman RL, Lopez-Berestein G, Sood AK. RNA interference in the clinic: challenges and future directions. *Nat. Rev. Cancer* 11(1), 59–67 (2011).
- 100 Martinez J, Patkaniowska A, Urlaub H, Luhrmann R, Tuschl T. Single-stranded antisense siRNAs guide target RNA cleavage in RNAi. *Cell* 110(5), 563–574 (2002).
- 101 Liu J, Carmell MA, Rivas FV *et al.* Argonaute2 is the catalytic engine of mammalian RNAi. *Science* 305(5689), 1437–1441 (2004).
- 102 Meister G, Landthaler M, Patkaniowska A, Dorsett Y, Teng G, Tuschl T. Human Argonaute2 mediates RNA cleavage targeted by miRNAs and siRNAs. *Mol. Cell* 15(2), 185–197 (2004).
- 103 Haupenthal J, Baehr C, Kiermayer S, Zeuzem S, Piiper A. Inhibition of RNase A family enzymes prevents degradation and loss of silencing activity of siRNAs in serum. *Biochem. Pharmacol.* 71(5), 702–710 (2006).
- 104 Davidson BL, Mccray PB Jr. Current prospects for RNA interference-based therapies. *Nat. Rev. Genet.* 12(5), 329–340 (2011).
- 105 Katare DP, Aeri V. Progress in gene therapy: a review. *IJTPR* 1, 33–41 (2010).
- 106 Nayerossadat N, Maedeh T, Ali PA. Viral and nonviral delivery systems for gene delivery. *Adv. Biomed. Res.* 1, 27 (2012).
- 107 Zhang J, Li X, Huang L. Non-viral nanocarriers for siRNA delivery in breast cancer. *J. Control. Release* 190, 440–450 (2014).
- 108 Wang X, Yu B, Wu Y, Lee RJ, Lee LJ. Efficient down-regulation of CDK4 by novel lipid nanoparticle-mediated siRNA delivery. *Anticancer Res.* 31(5), 1619–1626 (2011).
- 109 Honma K, Iwao-Koizumi K, Takeshita F *et al.* RPN2 gene confers docetaxel resistance in breast cancer. *Nat. Med.* 14(9), 939–948 (2008).
- **Demonstrates the effectiveness of siRNA in silencing RPN2 gene to sensitize docetaxel-resistant breast cancer and increase drug impact on cancer cells.**
- 110 Li YT, Chua MJ, Kunnath AP, Chowdhury EH. Reversing multidrug resistance in breast cancer cells by silencing ABC transporter genes with nanoparticle-facilitated delivery of target siRNAs. *Int. J. Nanomedicine* 7, 2473–2481 (2012).
- 111 Judge AD, Sood V, Shaw JR, Fang D, McClintock K, MacLachlan I. Sequence-dependent stimulation of the mammalian innate immune response by synthetic siRNA. *Nat. Biotechnol.* 23(4), 457–462 (2005).
- 112 Grimm D, Streetz KL, Jopling CL *et al.* Fatality in mice due to oversaturation of cellular microRNA/short hairpin RNA pathways. *Nature* 441(7092), 537–541 (2006).
- 113 Guo S, Huang L. Nanoparticles escaping RES and endosome: challenges for siRNA delivery for cancer therapy. *J. Nanomater.* 2011, 1–12 (2011).
- 114 Guo Y, Zhang Z, Kim DH *et al.* Photothermal ablation of pancreatic cancer cells with hybrid iron-oxide core gold-shell nanoparticles. *Int. J. Nanomedicine* 8, 3437–3446 (2013).
- 115 Macneill CM, Coffin RC, Carroll DL, Levi-Polyachenko NH. Low band gap donor-acceptor conjugated polymer nanoparticles and their NIR-mediated thermal ablation of cancer cells. *Macromol. Biosci.* 13(1), 28–34 (2013).
- 116 Cobley CM, Au L, Chen J, Xia Y. Targeting gold nanocages to cancer cells for photothermal destruction and drug delivery. *Expert Opin. Drug Deliv.* 7(5), 577–587 (2010).
- 117 De Freitas LF, Zanelatto LC, Mantovani MS *et al.* *In vivo* photothermal tumour ablation using gold nanorods. *Laser Physics* 23(6), 066003 (2013).
- 118 Hildebrandt B, Wust P, Ahlers O *et al.* The cellular and molecular basis of hyperthermia. *Crit. Rev. Oncol. Hematol.* 43(1), 33–56 (2002).
- 119 Burke AR, Singh RN, Carroll DL *et al.* The resistance of breast cancer stem cells to conventional hyperthermia and their sensitivity to nanoparticle-mediated photothermal therapy. *Biomaterials* 33(10), 2961–2970 (2012).
- 120 Lapotko DO, Lukianova E, Oraevsky AA. Selective laser nano-thermolysis of human leukemia cells with microbubbles generated around clusters of gold nanoparticles. *Lasers Surg. Med.* 38(6), 631–642 (2006).
- 121 Lapotko D, Lukianova E, Potapnev M, Aleinikova O, Oraevsky A. Method of laser activated nano-thermolysis for elimination of tumor cells. *Cancer Lett.* 239(1), 36–45 (2006).

- 122 Huang X, Jain PK, El-Sayed IH, El-Sayed MA. Gold nanoparticles: interesting optical properties and recent applications in cancer diagnostics and therapy. *Nanomedicine (Lond.)* 2(5), 681–693 (2007).
- 123 Carpin LB, Bickford LR, Agollah G *et al.* Immunoconjugated gold nanoshell-mediated photothermal ablation of trastuzumab-resistant breast cancer cells. *Breast Cancer Res. Treat.* 125(1), 27–34 (2011).
- 124 Lu W, Xiong C, Zhang G *et al.* Targeted photothermal ablation of murine melanomas with melanocyte-stimulating hormone analog-conjugated hollow gold nanospheres. *Clin. Cancer Res.* 15(3), 876–886 (2009).
- 125 Moon HK, Lee SH, Choi HC. *In vivo* near-infrared mediated tumor destruction by photothermal effect of carbon nanotubes. *ACS Nano* 3(11), 3707–3713 (2009).
- 126 Rozanova N, Zhang J. Photothermal ablation therapy for cancer based on metal nanostructures. *Sci. China Ser. B* 52(10), 1559–1575 (2009).
- 127 Liu SY, Liang ZS, Gao F, Luo SF, Lu GQ. *In vitro* photothermal study of gold nanoshells functionalized with small targeting peptides to liver cancer cells. *J. Mater. Sci. Mater. Med.* 21(2), 665–674 (2010).
- 128 Bickford LR, Agollah G, Drezek R, Yu TK. Silica-gold nanoshells as potential intraoperative molecular probes for HER2-overexpression in *ex vivo* breast tissue using near-infrared reflectance confocal microscopy. *Breast Cancer Res. Treat.* 120(3), 547–555 (2010).
- 129 Loo C, Hirsch L, Lee MH *et al.* Gold nanoshell bioconjugates for molecular imaging in living cells. *Opt. Lett.* 30(9), 1012–1014 (2005).
- 130 Dobbs MR. *Clinical Neurotoxicology: Syndromes, Substances, Environments*. Saunders/Elsevier (2009).
- 131 Fisher J, Handel N. *Problems in Breast Surgery: A Repair Manual*. CRC Press (2014).
- 132 Edwards R, Harding KG. Bacteria and wound healing. *Curr. Opin. Infect. Dis.* 17(2), 91–96 (2004).
- 133 Choi HS, Liu W, Misra P *et al.* Renal clearance of quantum dots. *Nat. Biotechnol.* 25(10), 1165–1170 (2007).
- 134 Longmire M, Choyce PL, Kobayashi H. Clearance properties of nano-sized particles and molecules as imaging agents: considerations and caveats. *Nanomedicine (Lond.)* 3(5), 703–717 (2008).
- 135 Zha Z, Yue X, Ren Q, Dai Z. Uniform polypyrrole nanoparticles with high photothermal conversion efficiency for photothermal ablation of cancer cells. *Adv. Mater.* 25(5), 777–782 (2013).
- 136 Dorozhkin SV, Epple M. Biological and medical significance of calcium phosphates. *Angew. Chem. Int. Ed. Engl.* 41(17), 3130–3146 (2002).
- 137 He Q, Mitchell AR, Johnson SL, Wagner-Bartak C, Morcol T, Bell SJ. Calcium phosphate nanoparticle adjuvant. *Clin. Diagn. Lab. Immunol.* 7(6), 899–903 (2000).
- 138 Weber JN, White EW, Lebiedzki J. New porous biomaterials by replication of echinoderm skeletal microstructures. *Nature* 233(5318), 337–339 (1971).
- 139 Panyam J, Labhasetwar V. Biodegradable nanoparticles for drug and gene delivery to cells and tissue. *Adv. Drug Deliv. Rev.* 55(3), 329–347 (2003).
- 140 Pinto OA, Tabakovic A, Goff TM, Liu Y, Adair JH. Calcium phosphate and calcium phosphosilicate mediated drug delivery and imaging. *Fund. Biomed. Technol.* 5, 713–744 (2011).
- 141 Amjad Z. *Calcium Phosphates in Biological and Industrial Systems*. Kluwer Academic Publishers (1998).
- 142 Yamashiro DJ, Maxfield FR. Regulation of endocytic processes by pH. *Trends Pharmacol. Sci.* 9(6), 190–193 (1988).
- 143 Van Sluis R, Bhujwalla ZM, Raghunand N *et al.* *In vivo* imaging of extracellular pH using ¹H MRSI. *Magn. Reson. Med.* 41(4), 743–750 (1999).
- 144 Santra S, Dutta D. *Nanomaterials for Cancer Diagnosis*. Wiley-VCH Verlag GmbH & Co., Weinheim, 7 (2007).
- 145 Altinoglu EI, Russin TJ, Kaiser JM *et al.* Near-infrared emitting fluorophore-doped calcium phosphate nanoparticles for *in vivo* imaging of human breast cancer. *ACS Nano* 2(10), 2075–2084 (2008).
- 146 Barth BM, Sharma R, Altinoglu EI *et al.* Bioconjugation of calcium phosphosilicate composite nanoparticles for selective targeting of human breast and pancreatic cancers *in vivo*. *ACS Nano* 4(3), 1279–1287 (2010).
- 147 Barth BM, Altinoglu EI, Shanmugavelandy SS *et al.* Targeted indocyanine-green-loaded calcium phosphosilicate nanoparticles for *in vivo* photodynamic therapy of leukemia. *ACS Nano* 5(7), 5325–5337 (2011).
- Indocyanine-green-doped calcium phosphosilicate nanoparticles cannot only be used for early detection of tumor cells, but the dye also serves as near infrared photosensitizer with a second function in photodynamic therapy for treating the cancer. This is one of the first examples of treating a nonsolid tumor, chronic myeloid leukemia.
- 148 Barth BM, Shanmugavelandy SS, Kaiser JM *et al.* Photoimmunotherapy reveals an anticancer role for sphingosine kinase 2 and dihydrosphingosine-1-phosphate. *ACS Nano* 7(3), 2132–2144 (2013).
- After photodynamic treatment by indocyanine-green doped calcium phosphosilicate nanoparticle, it was discovered that long-term suppression (45 days following treatment) of tumor growth was due to, in part, by the expansion of lymphocytes regulated by phosphorylated sphingolipid metabolites. This potentially eliminates the need for aggressive chemotherapy and radiation protocols.
- 149 Mussi SV, Silva RC, Oliveira MC, Lucci CM, Azevedo RB, Ferreira LA. New approach to improve encapsulation and antitumor activity of doxorubicin loaded in solid lipid nanoparticles. *Eur. J. Pharm. Sci.* 48(1–2), 282–290 (2013).
- 150 Silva AL, Rosalia RA, Sazak A *et al.* Optimization of encapsulation of a synthetic long peptide in PLGA nanoparticles: low-burst release is crucial for efficient CD8(+) T cell activation. *Eur. J. Pharm. Biopharm.* 83(3), 338–345 (2013).

- 151 Zhang Z, Feng SS. The drug encapsulation efficiency, *in vitro* drug release, cellular uptake and cytotoxicity of paclitaxel-loaded poly(lactide)-tocopheryl polyethylene glycol succinate nanoparticles. *Biomaterials* 27(21), 4025–4033 (2006).
- 152 Bray DJ, Gilmour SG, Guild FJ, Hsieh TH, Masania K, Taylor AC. Quantifying nanoparticle dispersion: application of the Delaunay network for objective analysis of sample micrographs. *J. Mater. Sci.* 46(19), 6437–6452 (2011).
- 153 Sager T, Wolfarth M, Keane M, Porter D, Castranova V, Holian A. Effects of nickel-oxide nanoparticle pre-exposure dispersion status on bioactivity in the mouse lung. *Nanotoxicology* 10(2), 151–161 (2015).
- 154 Alexis F, Pridgen E, Molnar LK, Farokhzad OC. Factors affecting the clearance and biodistribution of polymeric nanoparticles.pdf. *Mol. Pharm.* 5(4), 505–515 (2008).
- 155 Bertrand N, Wu J, Xu X, Kamaly N, Farokhzad OC. Cancer nanotechnology: the impact of passive and active targeting in the era of modern cancer biology. *Adv. Drug Del. Rev.* 66, 2–25 (2014).
- 156 Wicki A, Witzigmann D, Balasubramanian V, Huwyler J. Nanomedicine in cancer therapy: challenges, opportunities, and clinical applications. *J. Control. Release* 200, 138–157 (2015).
- 157 Ferrari M. Nanogeometry: beyond drug delivery. *Nat. Nanotechnol.* 3(3), 131–132 (2008).
- 158 Clinical Trials database: NCT00294996. <http://clinicaltrials.gov/ct2/show/NCT00294996>
- 159 Clinical Trials database: NCT01644890. <https://clinicaltrials.gov/ct2/show/NCT01644890>
- 160 Caraglia M, De Rosa G, Salzano G *et al.* Nanotech revolution for the anti-cancer drug delivery through blood–brain barrier. *Curr. Cancer Drug Targets* 12(3), 186–196 (2012).
- 161 Clinical Trials database: NCT00951054. <http://clinicaltrials.gov/ct2/show/NCT00951054>
- 162 Clinical Trial database: NCT00148707. <http://clinicaltrials.gov/ct2/show/NCT00148707>
- 163 Clinical Trial database: NCT00826085. <http://clinicaltrials.gov/ct2/show/NCT00826085>
- 164 Clinical Trial database: NCT00012129. <http://clinicaltrials.gov/ct2/show/NCT00012129>
- 165 Clinical Trial database: NCT00505271. <http://clinicaltrials.gov/ct2/show/NCT00505271>
- 166 Clinical Trial database: NCT01861496. <http://clinicaltrials.gov/ct2/show/NCT01861496>
- 167 Clinical Trial database: NCT00177281. <http://clinicaltrials.gov/ct2/show/NCT00177281>
- 168 Clinical Trial database: NCT00915369. <http://clinicaltrials.gov/ct2/show/NCT00915369>
- 169 Clinical Trial database: NCT01300533. <http://clinicaltrials.gov/ct2/show/NCT01300533>
- 170 Mills W, Adair J, Altinoglu E. Novel approach to safe and highly efficient cancer imaging. *SPIE Newsroom* doi:10.1117/2.1200901.1459 (2009) (Epub ahead of print).