Published in final edited form as:

Science. 2007 July 27; 317(5837): 519–523. doi:10.1126/science.1139518.

The Near Eastern Origin of Cat Domestication

Carlos A. Driscoll^{1,2,*}, Marilyn Menotti-Raymond¹, Alfred L. Roca³, Karsten Hupe⁴, Warren E. Johnson¹, Eli Geffen⁵, Eric H. Harley⁶, Miguel Delibes⁷, Dominique Pontier⁸, Andrew C. Kitchener^{9,10}, Nobuyuki Yamaguchi², Stephen J. O'Brien^{1,*}, and David W. Macdonald^{2,*}

¹Laboratory of Genomic Diversity, National Cancer Institute, Frederick, MD 21702, USA

²Wildlife Conservation Research Unit, Department of Zoology, University of Oxford, Oxford OX1 3PS, UK

³Laboratory of Genomic Diversity, SAIC-Frederick Inc., NCI-Frederick, Frederick, MD 21702, USA

⁴JagdEinrichtungsbüro, Am Sahlbach 9a, 37170 Fürstentagen, Germany

⁵Department of Zoology, Tel Aviv University, Tel Aviv 69978, Israel

⁶Division of Chemical Pathology, University of Cape Town, Observatory 7925, Cape Town, South Africa

⁷Department of Applied Biology, Estación Biológica de Doñana, CSIC, Avda Maria Luisa s/n Pabellón del Perú, 41013 Sevilla, Spain

⁸UMR-CNRS 5558 Biométrie et Biologie Evolutive, Université Claude Bernard Lyon I, 43 boulevard du 11 novembre 1918, 69622 Villeurbanne, France

⁹Department of Natural Sciences, National Museums Scotland, Chambers Street, Edinburgh EH1 1JF, UK

¹⁰Institute of Geography, School of Geosciences, University of Edinburgh, Drummond Street, Edinburgh EH8 9XP, UK

Abstract

The world's domestic cats carry patterns of sequence variation in their genome that reflect a history of domestication and breed development. A genetic assessment of 979 domestic cats and their wild progenitors—*Felis silvestris silvestris* (European wildcat), *F. s. lybica* (Near Eastern wildcat), *F. s. ornata* (central Asian wildcat), *F. s. cafra* (southern African wildcat), and *F. s. bieti* (Chinese desert cat)—indicated that each wild group represents a distinctive subspecies of *Felis silvestris*. Further analysis revealed that cats were domesticated in the Near East, probably coincident with agricultural village development in the Fertile Crescent. Domestic cats derive from

*To whom correspondence should be addressed. obrien@ncifcrf.gov; driscoll@ncifcrf.gov; david.macdonald@zoology.oxford.ac.uk.

Supporting Online Material

www.sciencemag.org/cgi/content/full/1139518/DC1

SOM Text

Tables S1 to S6

Figs. S1 and S2

References

at least five founders from across this region, whose descendants were transported across the world by human assistance.

The domestic cat may be the world's most numerous pet, yet little is certain of the cat's origin (1–9). Archaeological remains and anthropological clues suggest that, unlike species domesticated for agriculture (e.g., cow, pig, and sheep) or transport (horse and donkey), the cat probably began its association with humans as a commensal, feeding on the rodent pests that infested the grain stores of the first farmers (1). The earliest evidence of cat-human association involves their co-occurrence in Cyprus deposits determined to be 9500 years old (6). Domestic cats are generally considered to have descended from the Old World wildcats, but they differ from these hypothesized progenitors in behavior, tameness, and coat color diversity (9, 10). Further, domestic cats appear to lack neotenous characteristics typical of other domesticated species (11).

Felis silvestris, from which domestic cats were derived, is classified as a polytypic wild species composed of three or more distinct interfertile subspecies: *F. s. silvestris* in Europe, *F. s. lybica* in Africa and the Near East, *F. s. ornata* in the Middle East and central Asia (1, 2, 12–15), and possibly the Chinese desert cat, *F. s. bieti* (Fig. 1A, inset). The domestic cat is sometimes considered an additional subspecies, *F. s. catus*, possibly derived from wildcats in the Middle East or Egypt (1, 12, 14, 15). The imprecise subspecific status of *F. silvestris* populations and of the relationship of the domestic cat within this assemblage stems from morphological similarities among these groups (1, 13). A feral domestic cat with a “wild-type” mackerel tabby pattern is difficult to distinguish visually from a “true” wildcat (15, 16), which is further confounded by ongoing admixture (16–19). Moreover, the relationship between *F. silvestris* and the Chinese desert cat—which may be a separate *Felis* species, *Felis bieti*, or a wildcat subspecies, *F. silvestris bieti* (9, 12)—is uncertain. The sand cat *F. margarita*, a distinct species of *Felis* that ranges across North Africa and the Middle East, is the closest outgroup of the *F. silvestris/bieti* complex on the basis of morphological and molecular data (12, 13, 20).

To investigate the relationships among domestic cats, their indigenous wild progenitors, and related species of the genus *Felis*, we collected tissue from 979 individuals (fig. S1; see table S1 for breakdown of number of cats tested for different genetic markers) including putative wildcats and feral domestic cats on three continents ($N = 629$), fancy-breed domestic cats ($N = 112$), sand cats (*F. margarita*, $N = 11$), and Chinese desert cats (*F. s. bieti*, $N = 5$). We extracted DNA and genotyped 851 cats for 36 short tandem repeat (STR) or microsatellite domestic cat loci (21) variable in *F. silvestris*, *F. s. bieti*, *F. margarita*, and domestic cats, and sequenced 2604 base pairs (bp) of mitochondrial DNA (mtDNA) genes *ND5* and *ND6* from 742 cats.

Neighbor-joining phylogenetic analyses for STR genotypes with kinship coefficient (*Dkf*) and proportion of shared alleles (*Dps*) genetic distance estimators provided concordant topologies that specified six clusters (Fig. 1B; referred to here as “clades” as also specified by mtDNA phylogenetic analyses; see below) corresponding to the following subspecies designations: (i) *F. s. silvestris*, wildcats from Europe (STR clade I, green in Fig. 1); (ii) *F. s. ornata*, wildcats from central Asia east of the Caspian Sea (STR clade III, purple); (iii) *F. s.*

lybica, wildcats from the Near East (STR clade IV, beige); (iv) *F. s. cafra*, wildcats from southern Africa (STR clade II, blue); (v) *F. s. bieti*, Chinese desert cats (STR clade V, red); and (vi) *F. margarita*, sand cat (STR clade VI, black). *Felis cafra* was first named in 1822 and renamed as *F. lybica cafra* subspecies in 1944 on the basis of a description of a wildcat specimen captured in “Kaffraria” (9), an area from whence our southern African wildcat samples derive.

The composite STR genotypes of all known domestic house cats, fancy-breed cats, and feral domestic cats occurring in the wild populations all fell within a large monophyletic group (clade IV) that also included wildcats from the Near East. The phylogenetic tree suggests that domestication occurred in the Near East, where STR clade IV wildcats live today. This inference was further explored by examining mtDNA variation, STR variation, and ongoing admixture hybridization in the study areas (17–19).

Phylogenetic analysis of *ND5* and *ND6* sequence reveals 245 parsimony-informative sites specifying 176 distinct mtDNA genotypes (Fig. 2A, fig. S2, and table S2). The mtDNA haplotypes were analyzed with Bayesian Markov chain Monte Carlo (MCMC), maximum parsimony, maximum likelihood, and distance-based methods (22, 23). All methods resulted in identical topologies for the principal groupings corresponding to both geographic origins and STR clade designations. The consensus mtDNA gene tree (Fig. 2A), rooted with *F. margarita*, shows *F. s. bieti* basal to *F. silvestris*, as inferred from morphology. However, the short branch lengths and relatively weak bootstrap support for the node separating *F. s. bieti* from *F. silvestris* (27 to 68% bootstrap) indicates a close genetic relationship between these two taxa, which supports the grouping of *F. s. bieti* and *F. silvestris* as a single species, *F. silvestris*.

The *F. silvestris* mtDNA haplotypes fall into specific geographic locales (Fig. 2A). A basal lineage [clade I, *F. s. silvestris* (European wildcat), green] is found in European populations from Scotland and Portugal in the west to Hungary and Serbia in the east and is sister to *F. silvestris* from Asia and Africa and to domestic cats. An early/basal European versus Africa-Asia divergence supported by recent morphological studies of fossil specimens of wildcats (15, 24) may reflect a postglacial repopulation of Europe from Iberian founders, as previously suggested (9, 15, 24). The basal position of an Iberian wildcat, Fsi-257, within mtDNA clade I also supports an Iberian refugium (Fig. 2A).

Beyond Europe, mtDNA clades II, III, and IV correspond with geography and STR analysis (Fig. 2A). Within mtDNA clade IV, we identified five principal lineages of mtDNA haplotypes (A to E, Fig. 2A) with no obvious phylo-geographic association among these lineages. Domestication appears to have occurred within the Near Eastern region where clade IV wildcats are currently extant (beige, Fig. 2A), because clade IV wildcats and domestic cats are monophyletic.

Because of hybridization between wildcats and feral domestic cats, domestic cat mtDNA haplotypes (clade IV in Fig. 2A) are commonly found in European, African, and central Asian populations along with indigenous wildcat haplotypes (Fig. 1A). The observed genetic admixture may be explained by the presence of feral domestic cats or by hybridization

between wildcats and domestic cats. Hybrid individuals carrying one mtDNA-clade genotype but a different STR-clade genotype can be identified. Such cytonuclear-discordant individuals were common in our data set (Figs. 1B and 2A). Of cats sampled for both STR and mtDNA genotypes, seven of the 472 cats in STR clade IV were discordant, with a wildcat mtDNA type (Fig. 1B). However, among 108 putative European wildcats (on the basis of STR genotype; Fig. 1B), 28 carried the clade IV (domestic) mtDNA type, as did 3 of 13 southern African (STR clade II) wildcats. The wildcats in central Asia (STR clade III) included the highest frequency of discordant individuals (mtDNA clades III and IV; Fig. 1B), perhaps as a result of incomplete lineage sorting or recent gene flow between adjacent populations (Fig. 2A).

We implemented the Bayesian population genetic analysis program STRUCTURE, which assesses population subdivision (25) and characterizes genomic evidence of recent hybridization. STRUCTURE analyses of the 851 STR genotypes placed cats into discrete population clusters corresponding to European, African, and central Asian wildcats and identified a subdivision of domestic cats from different regions (Fig. 2B). Interestingly, we identified a discrete population of wild and domestic cats from the Near East (brown group in Fig. 2B) distinct from the other *F. silvestris* subspecies, as well as three subgroupings of domestic cats. These 15 individuals had concordant mtDNA and STR phylogenies identical to those of domestic cats and were collected in remote deserts of Israel, United Arab Emirates, Bahrain, or Saudi Arabia. These data suggest that these Near Eastern wildcats may represent the ancestral founder population of domestic cats, supporting a domestication origin in the Near East.

Identification of hybrids (STRUCTURE $Q < 0.8$) revealed that some (~22%) of the identified cytonuclear-discordant cats in Figs. 1B and 2A showed evidence of recent hybridization. For this reason, we removed 81 hybrid cats defined by STRUCTURE and generated new phylogenies combining the STR genotypes of cats grouped within the distinct populations (Fig. 2C). This analysis reaffirms the recognition of the major *F. silvestris* subspecies groups illustrated in Fig. 1A and the distinctiveness of Near Eastern wildcats as the closest group to all domestic cats. The results also suggest a close affinity between *F. s. bieti* (Chinese desert cat) and the Asian wildcats, plus paraphyly of other *F. silvestris* subspecies with respect to *F. s. bieti*, in support of the recognition of *F. s. bieti* as a subspecies of *F. silvestris* (Fig. 2C).

The coalescence-based age of mtDNA ancestral nodes for domestic cats (clade IV) and all *F. silvestris* mtDNA lineages was estimated with the linearized tree method (26). After fulfilling the requirement for molecular clock rate homogeneity across all lineages (table S4), we constructed a neighbor-joining algorithm on the basis of the linearized tree with Kimura two-parameter distances. We adopted a sequence divergence rate specific for *ND5* and *ND6* genes of 2.24 bp per million years (27). This rate would predict one new variant, on average (range: 0 to 2), in the most recent 17,000-year period of domestic cat ancestry (28). Indeed, 90% of the domestic cats within the five lineages (A to E in Fig. 2A) share haplotypes that are 0 to 3 bp apart, reflecting modest mutation accumulation within lineages. By contrast, the estimated coalescent date on the basis of the mtDNA data for all *F. silvestris* (including *F. s. bieti*) subspecies is 230,000 years ago, whereas the estimated age for the

ancestor of *F. s. lybica* and domestic cats is 131,000 years. Other methods of date estimation suggested a range from 107,000 to 155,000 years (28). These estimates are all greater by an order of magnitude than the age implied by archaeological evidence for cat domestication (6). The persistence of five well-supported mtDNA lineages dating back 100,000 years before any archaeological record of domestication would suggest that domestic cats originated from at least five matrilineal mtDNA haplotypes.

The variation described here is important for the conservation and management of free-ranging wildcat populations (16, 29). In table S6 we present a full list of population-specific (private) STR alleles as well as mtDNA population-specific site genotypes suitable for assessment of a wildcat's population, subspecies of origin, and distinction from domestic cats. The domestication of wild species to complement human civilization stands as one of the more successful "biological experiments" ever undertaken. For cats, the process began more than 9000 years ago when the earliest farmers of the Fertile Crescent domesticated grains and cereals as well as livestock (1, 3, 4, 30–32). In parallel, the endemic wildcats of the region may have adapted by both regulating the rodents in the grain stores and abandoning their aggressive wild-born behaviors. The archaeological imprints left in the genomes of living cats here weigh into inferences about the timing, steps, and provenance of domestication—a dynamic exercise depicted in art, in history, and in human cultural development since recorded evidence began.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank M. W. Smith, A. Schmidt-Kuntzel, C. O'hUigen, and B. Gold for discussions and J. Bruksch, A. Brandt, S. Rosendale, and F. Hussain for technical assistance. We appreciate the efforts of all of our collaborators listed in fig. S1 who provided biological specimens used in this study. All tissues were collected in full compliance with federal fish and wildlife permits [Convention on International Trade in Endangered Species of Wild Fauna and Flora (CITES)] issued to the National Cancer Institute (NCI; principal officer, S.J.O.) by the Fish and Wildlife Service, U.S. Department of the Interior. Supported by NCI grant N01-CO-12400 and the Intramural Research Program of the NCI Center for Cancer Research. Sequences have been deposited in GenBank with accession numbers EF587016 to EF587179.

References and Notes

1. Clutton-Brock, JA. *Natural History of Domesticated Mammals*. Cambridge Univ Press; Cambridge: 1999.
2. Kitchener, A. *The Natural History of the Wild Cats*. Comstock Associates; Ithaca, NY: 1991.
3. Stephens, G., Yamazaki, T. *Legacy of the Cat: The Ultimate Illustrated Guide*. Chronicle; San Francisco: 2001.
4. Budiansky, S. *The Covenant of the Wild: Why Animals Chose Domestication*. Yale Univ Press; New Haven, CT: 1999.
5. Legay JM. *C R Acad Sci III*. 1986; 303:709. [PubMed: 3101986]
6. Vigne JD, Guilaine J, Debue K, Haye L, Gérard P. *Science*. 2004; 304:259. [PubMed: 15073370]
7. Donalson, MD. *The Domestic Cat in Roman Civilization*. Mellen; New York: 1999.
8. Kurten B. *Acta Zool Fenn*. 1965; 111:3.
9. Pocock, RI. *Catalogue of the Genus Felis*. British Museum–Natural History; London: 1951.

10. Vella, CM., Shelton, LM., McGonagle, JJ., Stanglein, TW. Robinson's Genetics for Cat Breeders & Veterinarians. Elsevier Science; Edinburgh: 2003.
11. Macdonald DW, et al. Adv Ethol. 1987; 28:1.
12. Nowell, K., Jackson, P. Status Survey and Conservation Action Plan, Wild Cats. International Union for Conservation of Nature and Natural Resources; Gland, Switzerland: 1996.
13. Sunquist, ME., Sunquist, F. Wild Cats of the World. Univ of Chicago Press; Chicago: 2002.
14. Pocock RI. Proc Zool Soc London. 1907; 1907:656.
15. Randi E, Ragni B. J Mammal. 1991; 72:79.
16. Macdonald, DW., Daniels, MJ., Driscoll, C., Kitchener, AC., Yamaguchi, N. The Scottish Wildcat: Analyses for Conservation and an Action Plan. Wildlife Conservation Research Unit; Oxford: 2004.
17. Beaumont M, et al. Mol Ecol. 2001; 10:319. [PubMed: 11298948]
18. Randi E, et al. Mol Biol Evol. 2001; 18:1679. [PubMed: 11504848]
19. Lecis R, et al. Mol Ecol. 2006; 15:119. [PubMed: 16367835]
20. Johnson WE, et al. Science. 2006; 311:73. [PubMed: 16400146]
21. Menotti-Raymond M, et al. Cytogenet Genome Res. 2003; 102:272. [PubMed: 14970716]
22. Swofford, DL. Phylogenetic Analysis Using Parsimony (PAUP). Natural History Survey; Champaign, IL: 1985.
23. Huelsenbeck JP, Ronquist F. Bioinformatics. 2001; 17:754. [PubMed: 11524383]
24. Yamaguchi N, et al. Anim Conserv. 2004; 7:339.
25. Pritchard JK, Donnelly P. Theor Popul Biol. 2001; 60:227. [PubMed: 11855957]
26. Takezaki N, et al. Mol Biol Evol. 1995; 12:828.
27. Lopez JV, Culver M, Stephens JC, Johnson WE, O'Brien SJ. Mol Biol Evol. 1997; 14:277. [PubMed: 9066795]
28. See supporting material on *Science* Online.
29. O'Brien SJ, Johnson WE. Annu Rev Genomics Hum Genet. 2005; 6:407. [PubMed: 16124868]
30. Tanno K, Willcox G. Science. 2006; 311:1886. [PubMed: 16574859]
31. Willcox, G. 13th IWGP Symposium. Buxo, R.Jacomet, S., Bitmann, F., editors. Springer; Girona, Spain: 2005. p. 534-541.
32. Morrell PL, Clegg MT. Proc Natl Acad Sci USA. 2007; 104:3289. [PubMed: 17360640]

Fig. 1.

(A) Current range of *F. silvestris* and areas of sample collection. Colored regions reflect the location of capture of individuals carrying different STR clade genotypes (defined at lower left). mtDNA haplotype frequencies are indicated in pie charts specifying the number of specimens carrying mtDNA haplotypes for each clade. Central Asian denotes Asian cats east of the Caspian Sea. Near Eastern denotes cats in Israel, Saudi Arabia, Bahrain, and the United Arab Emirates. European denotes specimens collected west of the Caspian Sea. Domestic cats (*F. s. catus*) are distributed worldwide and overwhelmingly carry clade IV

mtDNA haplotypes (beige). Inset: Current and historical range of *F. silvestris* subspecies on the basis of traditional morphology-based taxonomy (2, 12, 13). The Chinese desert cat is referred to throughout as a wildcat subspecies, *F. silvestris bieti* (9, 12), as supported by data presented here. **(B)** Phenogram of 851 domestic and wild specimens created on the basis of STRs, *Dps* genetic distance, and minimum evolution (neighbor-joining) algorithm. Color groups correspond to geographic locales specified in (A). Symbols indicate cytonuclear-discordant individuals that contain a STR composite clade of the indicated cluster but carry mtDNA of an alternative locale (see text); in parentheses are the numbers of cats in each STR clade that carry various mtDNA clade haplotypes.

Fig. 2.

(A) Phylogenetic tree of mitochondrial DNA sequence [minimum evolution (neighbor-joining) phylogram of 2604 bp of the *ND5* and *ND6* genes] of 176 haplotypes discerned from 742 cats sampled across the range of the domestic cat, European wildcat, Near Eastern wildcat, central Asian wildcat, southern African wildcat, Chinese desert cat, and sand cat. Trees created from Bayesian, maximum likelihood, and maximum parsimony methods result in identical topologies for principal clade groupings. Confidence/bootstrap values (from left to right: Bayes/maximum parsimony/maximum likelihood/minimum evolution) are based on 1000 iterations and are adjacent to nodes. The number of single-nucleotide differences is indicated in red below the corresponding branch. Clade designations and numbers of individuals are indicated in parentheses after the corresponding common name and taxonomic trinomial. A through E designate lineages within mtDNA clade IV. Confidence/bootstrap values for these nodes are as follows: A, 1.00/87/71/54; B, 1.00/82/80/80; C, 0.97/63/59/42; D, 1.00/98/99/88; E, 1.00/100/100/82. Purple and brown tree limbs within mtDNA clade IV reflect individuals from two locales that bear cytonuclear-discordant mtDNA versus STR genotypes (see text). Clade IV individuals bearing mtDNA haplotypes are found among domestic cats; in wild potentially admixed populations in Europe, Asia, or Africa (see Fig. 1A); and in Near Eastern wildcats (see text). (B) STRUCTURE-based populations resolved 851 cats into several wildcat groups, three domestic cat groups, and one group (brown) that included both domestic cats and Near Eastern wildcats. The *y* axis represents *Q*-value, the percent representation of resolved populations (colors) within each individual (listed on *x* axis). (C) Phylogenetic relationships among *F. silvestris* groups as defined by composite STR genotypes based on 36 STR loci. Tree is rooted at sand cat.

Bootstrap values at corresponding nodes are based on 1000 iterations with the following measures (from left to right): $Dps = 1 - (ps)/Dkf = 1 - (kf)/Dps = -\ln(ps)/Dkf = -\ln(kf)$. All methods resulted in identical topologies. Individuals were clustered into representative populations based on STRUCTURE Q -value of 0.80 or greater with the same loci (see text). All known domestic cats cluster into domestic-Asia, domestic-Europe, or Near Eastern wildcats, regardless of provenance, and these groups also cluster together.