CORRESPONDENCE

Lung Transplantation in Germany Since the Introduction of the Lung Allocation Score

A Retrospective Analysis

by Prof. Dr. med. Jens Gottlieb; Jacqueline Smits, MD, PhD;
PD Dr. med. Rene Schramm; PD Dr. med. Frank Langer; Prof. Dr. med. Roland
Buhl; Prof. Dr. med. Christian Witt; Prof. Dr. med. Martin Strueber; and
Prof. Dr. med. Hermann Reichenspurner in issue 11/2017

Acknowledgement

I gratefully acknowledge the help of Prof. Frank Detterbeck, MD, Thoracic Surgery at Yale University, and Margaret Cloud.

Thomas M. Egan, M.D., M.Sc.

University of North Carolina at Chapel Hill
thomas_egan@med.unc.edu

Imitation is the Sincerest Form of Flattery

Gottlieb et al. reported encouraging results after three years' use of the Lung Allocation Score (LAS) in Germany (1). I congratulate the German lung transplant community for embracing a radically different organ allocation system to provide some objectivity to lung allocation when numbers are inadequate to meet the need. For six years, I chaired the subcommittee that, with help from many people, developed the LAS for the US Organ Procurement and Transplant Network (OPTN). Until recently, LAS was the only US organ allocation system that considered post-transplant survival. I am excited that Eurotransplant uses the LAS to distribute donor lungs (2). Gottlieb reports that in Germany, the LAS was associated with more lung transplants performed (without concomitant increase in donors), a significant reduction in waitlist deaths, with no increase in post-transplant mortality, and improved distribution of recipient diagnoses. This is similar to the impact in the US (3). LAS adoption appears to have been less contentious in Germany than in the US. Now Germany has enough data to make its LAS a *German* entity, instead of relying on old US waitlist and transplant data like us (4). Our Subcommittee intended to use new data to re-analyze risks of waitlist death and post-transplant survival every six months, and modify variable coefficients that factored into LAS calculation. Variables could be eliminated or added as the population of waitlisted patients changed, to keep the model current. Sadly, the OPTN took years to make substantive changes. The *Deutsche Stiftung Organtransplantation* may be more nimble. DOI: 10.3238/arztebl.2017.0543a

REFERENCES

1. Gottlieb J, Smits J, Schramm R, et al.: Lung transplantation in Germany since the introduction of the Lung Allocation Score—a retrospective analysis. *Dtsch Arztebl Int* 2017; 114: 179–85.
2. Eurotransplant: Introduction of the LAS in the Netherlands. 2014. www.eurotransplant.org/cms/index.php?page=newsitems&oid=458&action=viewitem&id=4505 (last accessed on 23 May 2017).
3. Egan TM, Edwards LB: Effect of the lung allocation score on lung transplantation in the United States. *J Heart Lung Transplant* 2016; 35: 433–9.
4. Egan TM, Murray S, Bustami RT: Development of the new lung allocation system in the United States. *Am J Transplant* 2006; 6: 1212–27.

In Reply:

We thank Dr. Egan for his interest in our publication (1) and for his endorsement of the German experience with the Lung Allocation Score (LAS). He pioneered the important concept of transplant benefit in donor lung allocation within the US, balancing the patient mortality risk after transplantation against that of remaining on the wait list. Translating this into clinical practice required great courage in 2005, given the fundamental shift needed to prioritize outcome benefit rather than current clinical need. The results remain unequivocal, demonstrating improved wait list mortality and transplant outcomes subsequently in the US (2).

Germany adopted the LAS in 2011, with the Netherlands following in 2014, primarily for national allocation in contrast to the primarily local allocation in the US. From the outset both countries have introduced special business rules, adapted to the needs specific to the European population. One of the reasons was that the LAS does not factor in the need for extracorporeal life support (ECLS). In the 5 years that the LAS has been in operation in Germany, more than a dozen special rules have been designed and introduced to provide clarity to physicians and allow parity among candidates.

On February 19th, 2015 the US adopted a modified LAS, entailing adjustments in of some existing coefficients, alongside introducing new parameters such as cardiac index, bilirubin and creatinine. The European countries await with anticipation the upcoming publications outlining experience and the impact of these amendments.

We agree wholeheartedly with Dr. Egan that the European model needs to be kept current by analyzing waitlist outcomes and survival after transplantation in Europe as well as additional parameters continuously. We firmly believe that data quality is of extreme importance to avoid “garbage in, garbage out” in statistical modeling (3). Eurotransplant has introduced practical business rules for oxygen titration to minimize this problem and national audits in Germany have improved data quality since their introduction in 2014.

DOI: 10.3238/arztebl.2017.0543b

REFERENCES

1. Gottlieb J, Smits J, Schramm R, et al.: Lung transplantation in Germany since the introduction of the Lung Allocation Score—a retrospective analysis. *Dtsch Arztebl Int* 2017; 114: 179–85.

2. Egan TM, Edwards LB: Effect of the lung allocation score on lung transplantation in the United States. *J Heart Lung Transplant*. 2016; 35: 433–9.
3. N.N.: Work with new electronic 'brains' opens field for army math experts. *The Hammond Times* 1957; 65 (retrieved on 20 March 2016 via Newspapers.com).

On behalf of the authors

Prof. Dr. med. Jens Gottlieb

Medizinische Hochschule Hannover, Biomedical Research in Endstage and Obstructive Lung Disease Hannover (BREATH), Mitglied des Deutschen Forschungszentrums für Lungenforschung
Gottlieb.Jens@mh-hannover.de

Jacqueline Smits, MD, PhD

Eurotransplant International Foundation, Leiden, Niederlande

Conflict of interest statement

The authors of both contributions declare that no conflict of interest exists.

CLINICAL SNAPSHOT

A 76-Year-Old Woman With Chest Pain and Tarry Stool

Computerized tomography on admission (a + b) reveals the perforation (arrow) of the intrathoracic stomach (asterisk) into the pleural cavity. An extensive seropneumothorax is seen (triangles).

and partial pleurectomy. A persistent pleural fluid collection was drained under CT guidance. The patient was discharged in stable condition after three weeks in the hospital.

Dr. med. Lars-Arne Schaafs, PD Dr. med. Zarko Grozdanovic, Klinik und Hochschulambulanz für Radiologie, Campus Benjamin Franklin, Charité-Universitätsmedizin Berlin, lars-arne.schaafs@charite.de

Conflict of interest statement

The authors state that they have no conflict of interest.

Cite this as:

Schaafs L-A, Grozdanovic Z: A 76-year old woman with chest pain and tarry stool. *Dtsch Arztebl Int* 2017; 114: 544. DOI: 10.3238/arztebl.2017.0544

Translated from the original German by Ethan Taub, M.D.

A 76-year-old woman was taken to the emergency room because of chest pain and tarry stool. She stated that she frequently used a combined preparation containing acetylsalicylic acid, acetaminophen, and caffeine.

Esophagogastroduodenoscopy was performed because of the suspected diagnosis of upper gastrointestinal bleeding. This revealed a paraesophageal hernia and an ulcer with covered perforation in the area of the gastric cardia and fundus. Computerized tomography (CT) of the chest and abdomen yielded the diagnosis of an intrathoracic stomach with perforation into the left pleural cavity (*Figure*). In an emergency procedure, the proximal portion of the stomach and the neighboring distal portion of the esophagus were resected, with ensuing gastric reconstruction, esophago-gastrostomy, and hiato-plasty. Pleural empyema arose as a postoperative complication and was treated with irrigation