Phosphoserine Phosphatase Is Required for Serine and One-Carbon Unit Synthesis in *Hydrogenobacter thermophilus*

Keugtae Kim,^a Yoko Chiba,^b Azusa Kobayashi,^a Hiroyuki Arai,^a Masaharu Ishii^a

Department of Biotechnology, Graduate School of Agricultural and Life Sciences, The University of Tokyo, Tokyo, Japan^a; Department of Subsurface Geobiological Analysis and Research, Japan Agency for Marine-Earth Science and Technology, Kanagawa, Japan^b

ABSTRACT *Hydrogenobacter thermophilus* is an obligate chemolithoautotrophic bacterium of the phylum *Aquificae* and is capable of fixing carbon dioxide through the reductive tricarboxylic acid (TCA) cycle. The recent discovery of two novel-type phosphoserine phosphatases (PSPs) in *H. thermophilus* suggests the presence of a phosphorylated serine biosynthesis pathway; however, the physiological role of these novel-type metal-independent PSPs (iPSPs) in *H. thermophilus* has not been confirmed. In the present study, a mutant strain with a deletion of *pspA*, the catalytic subunit of iPSPs, was constructed and characterized. The generated mutant was a serine auxotroph, suggesting that the novel-type PSPs and phosphorylated serine synthesis pathway are essential for serine anabolism in *H. thermophilus*. As an autotrophic medium supplemented with glycine did not support the growth of the mutant, the reversible enzyme serine hydroxymethyltransferase does not appear to synthesize serine from glycine and may therefore generate glycine and 5,10-CH₂-tetrahydrofolate (5,10-CH₂-THF) from serine. This speculation is supported by the lack of glycine cleavage activity, which is needed to generate 5,10-CH₂-THF, in *H. thermophilus*. Determining the mechanism of 5,10-CH₂-THF synthesis is important for understanding the fundamental anabolic pathways of organisms, because 5,10-CH₂-THF is a major one-carbon donor that is used for the synthesis of various essential compounds, including nucleic and amino acids. The findings from the present experiments using a *pspA* deletion mutant have confirmed the physiological role of iPSPs as serine producers and show that serine is a major donor of one-carbon units in *H. thermophilus*.

IMPORTANCE Serine biosynthesis and catabolism pathways are intimately related to the metabolism of 5,10-CH₂-THF, a one-carbon donor that is utilized for the biosynthesis of various essential compounds. For this reason, determining the mechanism of serine synthesis is important for understanding the fundamental anabolic pathways of microorganisms. In the present study, we experimentally confirmed that a novel phosphoserine phosphatase in the obligate chemolithoautotrophic bacterium *Hydrogenobacter thermophilus* is essential for serine biosynthesis. This finding indicates that serine is synthesized from an intermediate of gluconeogenesis in *H. thermophilus*. In addition, because glycine cleavage system activity and genes encoding an enzyme capable of producing 5,10-CH₂-THF were not detected, serine appears to be the major one-carbon donor to tetrahydrofolate (THF) in *H. thermophilus*.

KEYWORDS amino acid biosynthesis, carbon metabolism, thermophiles

Hydrogenobacter thermophilus, which belongs to the order *Aquificales*, is a strictly thermophilic hydrogen-oxidizing chemolithoautotroph that fixes carbon dioxide through the reductive tricarboxylic acid (TCA) cycle (1, 2). Recently, two novel-type phosphoserine phosphatases (PSPs; EC 3.1.3.3), metal-independent PSP iPSP1 and iPSP2, were identified and characterized following the purification of proteins with

Received 26 June 2017 Accepted 31 July 2017

Accepted manuscript posted online 7 August 2017

Citation Kim K, Chiba Y, Kobayashi A, Arai H, Ishii M. 2017. Phosphoserine phosphatase is required for serine and one-carbon unit synthesis in *Hydrogenobacter thermophilus*. *J Bacteriol* 199:e00409-17. <https://doi.org/10.1128/JB.00409-17>.

Editor William W. Metcalf, University of Illinois at Urbana-Champaign

Copyright © 2017 American Society for Microbiology. All Rights Reserved.

Address correspondence to Yoko Chiba, ychiba@jamstec.go.jp.

K.K. and Y.C. contributed equally to this work.

FIG 1 Metabolic pathway related to serine biosynthesis in *H. thermophilus*. The unbroken arrows indicate the biochemically confirmed reactions, and dashed arrows indicated predicted reactions based on genomic analyses. CoA, coenzyme A; rTCA, reductive TCA.

phosphoserine phosphatase activity from *H. thermophilus* cells (3–5). iPSP1 is a homodimer of PspA subunits, whereas iPSP2 is a heterodimer of PspA and PspB. Although PspA and PspB share 37% identity and exhibit phosphatase activity, only the PspA subunit recognizes phosphoserine as a substrate (4). The discovery of iPSP1 and iPSP2 in *H. thermophilus* strongly suggests the presence of a phosphorylated serine biosynthesis pathway, in which serine is produced from the gluconeogenesis intermediate 3-phosphoglycerate in a three-step conversion process (Fig. 1) (6). Despite this speculation, the synthesis of serine from phosphoserine by the catalysis of iPSP in *H. thermophilus* has not been experimentally confirmed, and it is unclear whether other serine synthesis pathways are active in this bacterium.

In addition to iPSPs, *H. thermophilus* also possesses the enzyme serine hydroxymethyltransferase (SHMT), which has the potential to synthesize serine by catalyzing the reversible conversion of glycine using tetrahydrofolate (THF) as a one-carbon carrier (Fig. 1) (7). Methanotrophic and methylotrophic bacteria use SHMT for serine synthesis (8). Higher plants also synthesize a portion of serine from glycine (6). A one-carbon donor, 5,10-CH₂-tetrahydrofolate (5,10-CH₂-THF), which is used for this direct conversion reaction, is provided by the glycine cleavage system (9) or through the direct fixation of a one-carbon compound to THF (8, 10). Numerous organisms use SHMT for glycine synthesis from serine (11, 12). The one-carbon units produced in this direct reaction are used for the synthesis of several important compounds, including methionine, purine, and pyrimidine (13, 14). Therefore, identifying the components of the serine biosynthesis pathway is necessary to determine how an organism provides the one-carbon donor 5,10-CH₂-THF for the biosynthesis of various fundamental compounds, including serine, methionine, purines, and pyrimidines.

Here, we analyzed the serine biosynthesis pathway of *H. thermophilus* using a *pspA* deletion mutant (Δ *pspA*). A component involved in the uptake of serine and the main reaction by which 5,10-CH₂-THF is produced in *H. thermophilus* are also discussed.

RESULTS

Physiological characterization of *pspA* and *pspB* deletion mutants. The constructed *pspA* deletion mutant of *H. thermophilus* was cultivated in a medium supple-

FIG 2 Growth profiles of the $\Delta pspA$ mutant cultured in autotrophic medium supplemented with various serine concentrations. Growth of the $\Delta pspA$ mutant was monitored in an autotrophic medium not supplemented with serine ($\circ$) or supplemented with 10 μ M ($\square$), 25 μ M ($\diamond$), 50 μ M ($\triangle$), 100 μ M ($\blacklozenge$), 300 μ M ($\bullet$), 400 μ M ($\blacksquare$), or 500 μ M ($\blacktriangle$) serine. The data are expressed as the mean $\pm$ standard error (SE) ($n = 3$).

mented with various concentrations of L-serine, and the cell densities at stationary phase were compared (Fig. 2). Clear growth was observed in medium supplemented with serine at concentrations of 50 μ M or higher, confirming that the $\Delta pspA$ mutant is a serine auxotroph. The final cell yields were comparable between the cultures supplemented with 300, 400, and 500 μ M L-serine; however, the $\Delta pspA$ mutant cells cultured in medium supplemented with 300 μ M serine grew faster than those in the medium containing 400 or 500 μ M serine (Fig. 2). Therefore, 300 μ M serine was selected as the optimum concentration for the growth of the $\Delta pspA$ mutant. The serine concentration in the medium decreased with increasing growth of the $\Delta pspA$ mutant (Fig. 3A), indicating that serine was incorporated into the cells. In contrast, the supplementation of medium with serine had no detectable effects on cell growth of a $pspB$ deletion ($\Delta pspB$) mutant of *H. thermophilus* (Fig. 3B) or the wild type (Fig. 3C). No clear serine uptake by wild-type cells was observed either (Fig. 3C).

To confirm whether compounds derived from serine or 5,10-CH₂-THF (15–17) could support the growth of the $\Delta pspA$ mutant, $\Delta pspA$ cells were cultivated in inorganic medium supplemented with 300 μ M glycine, L-tryptophan, methionine, histidine, or formate (Fig. 3A). No growth was detected in the first 24 h of culture (Fig. 3A). In contrast, the growth of wild-type cells was not inhibited, indicating that these compounds were not incorporated into cells, or that they were incorporated into cells but could not compensate for the inability of the $\Delta pspA$ mutant to synthesize serine.

Evaluation of serine transport. The uptake of serine by the $\Delta pspA$ mutant indicated the existence of a serine transporter in *H. thermophilus*. However, homologs of known serine transporters in *Escherichia coli* (18, 19, 34) (Na⁺/serine threonine symporter [accession no. [P0AGE4.1](#)], H⁺/threonine serine symporter [accession no.

FIG 3 Growth profiles and serine uptake by the $\Delta pspA$ mutant (A), $\Delta pspB$ mutant (B), and wild-type (C) strains. Growth was monitored in an autotrophic medium not supplemented ($\circ$) or supplemented with 300 μ M Ser ($\square$), Gly ($\triangle$), Trp ($\diamond$), Met ($\circ$), His ($\dagger$), or formate ($\ast$). Serine and glycine concentrations (con.) within the medium supplemented with serine are shown with closed squares and closed circles, respectively. The data are expressed as the mean $\pm$ SE ($n = 3$).

FIG 4 Inhibitory effect of Leu, Ile, and Val on the growth of $\Delta pspA$ (A) and wild-type (B) cells. A 300 μ M concentration of Val (●), Ile (◇), or Leu (△) or nothing (control, □) was added to autotrophic medium not supplemented or supplemented with 300 μ M Ser for the wild-type and $\Delta pspA$ mutant strains, respectively. The data are expressed as the mean $\pm$ SE ($n = 3$).

CDH66785], and serine transporter [accession no. ADX49536.1]) were not found in the *H. thermophilus* genome. In contrast, homologs of all five proteins that comprise the leucine-isoleucine-valine (LIV) transport system, which has been proposed to function as a serine transporter (18, 20), were detected in the genome. The candidates of the *H. thermophilus* LIV system components showed amino acid identities ranging from 26% to 38% and E values of 10^{-13} to 10^{-48} with those of *E. coli* (see Table S1 in the supplemental material).

Inhibition tests using leucine, isoleucine, and valine were performed to explore the possibility that the LIV system of *H. thermophilus* functions as a serine transporter. Growth inhibition of the $\Delta pspA$ mutant was detected when 300 μ M leucine, isoleucine, or valine was added to medium containing 300 μ M serine (Fig. 4A). In contrast, no growth inhibition was observed when leucine, isoleucine, or valine was added to the autotrophic medium of the wild-type strain (Fig. 4B). These data suggest that serine uptake is competitively inhibited by leucine, isoleucine, or valine and therefore indicate that the LIV system may function as a serine transporter in *H. thermophilus*.

Detection of the glycine cleavage system. A homology search was performed for *E. coli* homologs of the glycine cleavage system in *H. thermophilus*. Homologs of all protein components (P, H, T, and L proteins) of the *E. coli* glycine cleavage system were found. Notably, the N- and C-terminal domains of the *E. coli* P protein exist as two separate proteins in *H. thermophilus*. The candidate proteins shared 30% to 48% identity and had E values of 10^{-32} to 10^{-70} with those in *E. coli* (Table S2). Notably, the glycine cleavage system proteins were not highly conserved in the phylum *Aquificae*. For example, organisms in the families *Desulfurobacteriaceae* and *Hydrogenothermaceae* and several species in *Aquificaceae* lack at least one component of the glycine cleavage system (Table 1).

To confirm whether the glycine cleavage system is functional in *H. thermophilus*, the glycine cleavage activity in cell extract was assayed by measuring both the THF and NAD⁺- or NADP⁺-dependent decrease of glycine and the THF-dependent reduction of benzyl viologen (BV), an artificial electron acceptor alternative to NAD(P)⁺. However, no activity (less than 20 nmol glycine cleaved by 1 mg of protein per min) was detected under the present experimental conditions (data not shown).

DISCUSSION

In this study, the physiological role of novel-type PSPs in *H. thermophilus* was investigated using *pspA* and *pspB* deletion strains. The $\Delta pspB$ mutant did not show apparent phenotypic change under our standard autotrophic cultivation conditions. A physiological role of PspB is unclear so far, because previous biochemical analyses failed to identify a physiological substrate of PspB, while it seems to have relatively low phosphatase activity against *p*-nitrophenyl phosphate (3). Further studies are required to understand the reason why iPSP2, a heteromer of PspA and PspB, exists at the same level as iPSP1, a homodimer of PspAs in *H. thermophilus* (3). In contrast, a deletion

TABLE 1 Homolog search of glycine cleavage system proteins in *Aquificae*^a

Family	Organism	P protein N-terminal domain		P protein C-terminal domain		T protein		H protein		L protein	
		% identity	Accession no.	% identity	Accession no.	% identity	Accession no.	% identity	Accession no.	% identity	Accession no.
Aquificaceae	<i>Aquifex aeolicus</i> VF5	33	WP_010880695	39	WP_010881244	38	O67441.2	49	WP_010880694	28	WP_010880444
	<i>Hydrogenobacter thermophilus</i> TK-6	35	WP_012964165	38	WP_012963212	37	WP_012963253	48	WP_012964164	30	WP_012963843
	<i>Hydrogenobaculum</i> sp. strain Y04AAS1	ND		ND		ND		37	WP_012514115	27	WP_012513374
	<i>Thermocrinis albus</i> DSM 14484	37	WP_012991979	38	WP_012992260	37	WP_012991224	45	WP_012991980	30	WP_012992228
Hydrogenothermaceae	<i>Persephonella marina</i> EX-H1	ND		ND		ND		44	WP_012675418	35	WP_012676035
	<i>Sulfurihydrogenibium</i> sp. strain YO3AOP1	ND		ND		ND		48	WP_012459599	30	WP_012459600
	<i>Desulfurobacterium thermolithotrophum</i>	ND		ND		ND		ND		ND	
Desulfurobacteriaceae	<i>Thermovibrio ammonificans</i> HB-1	ND		ND		ND		ND		ND	

^aProteins which showed an E value of $<10^{-10}$ against queries are listed as possible candidates of glycine cleavage (GCV) proteins. *E. coli* genes (GenBank accession numbers NP_417379, NP_417381, NP_417380, and NP_414658 for GCV P, T, H, and L proteins, respectively) were used as queries (33). ND, not determined.

mutant of the PSPs' catalytic subunit, the $\Delta pspA$ mutant, was revealed to require serine for growth. Based on this finding, PspA and, most likely, the phosphorylated serine synthesis pathway, appear to be essential for serine synthesis in *H. thermophilus*.

The observed decrease in serine in the culture medium of the $\Delta pspA$ mutant suggests that this obligate autotroph possesses a serine uptake system. *H. thermophilus* lacks homologs of all three serine uptake systems in *E. coli* but encodes homologs of all *E. coli* LIV transport system components. In addition, growth impairment of $\Delta pspA$ cells was observed in cultures supplemented with branched-chained amino acids. The transport of serine by the *E. coli* LIV system has not been conclusively determined (18, 20, 21); however, the present results suggest that this system is involved in the uptake of serine. In contrast to the case of the $\Delta pspA$ mutant, the wild-type strain did not show clear uptake of serine. Therefore, quantitative real-time reverse transcription-PCR analysis was performed to see if the presence or absence of serine uptake is due to the presence or absence of the LIV transporter using *livK*, the first gene of the operon of the LIV transporter genes. Statistically significant differences were observed neither between the expression levels in the wild-type strains cultivated with and without serine nor the wild-type strain and $\Delta pspA$ mutant cultivated with serine (data not shown). These results suggest that the uptake of serine is regulated not by the expression level of the LIV transporter but by the intracellular serine concentration, although further studies are required to confirm this speculation.

H. thermophilus possesses SHMT, a reversible enzyme with the potential to synthesize serine from glycine and 5,10-CH₂-THF (7, 22). If *H. thermophilus* has the ability to synthesize glycine from compounds other than serine, such as glyoxylate, and can produce 5,10-CH₂-THF without an SHMT reaction, we speculated that the $\Delta pspA$ mutant would be able to grow without exogenously added serine. However, the $\Delta pspA$ mutant required supplemented serine to grow. This result seems to be reasonable because *H. thermophilus* has neither genes encoding the enzymes for the glyoxylate cycle nor enzymatic activities for the metabolism (23). Furthermore, if *H. thermophilus* possesses an active glycine cleavage system, supplementation of the culture medium with glycine may support the growth of the $\Delta pspA$ mutant, even if cells cannot produce sufficient glycine for growth without functional iPSPs. However, neither the growth of the $\Delta pspA$ mutant in autotrophic medium supplemented with glycine nor glycine cleavage system activity was observed. These results suggest that SHMT catalyzes the generation of glycine and 5,10-CH₂-THF in *H. thermophilus*.

THF-based one-carbon units are essential for fundamental metabolic processes, including nucleic acid biosynthesis and amino acid and vitamin metabolism (24). In the case of prokaryotes, serine cleavage by SHMT, glycine cleavage by the glycine cleavage system, and the direct fixation of formate to THF by formate-THF ligase (EC 6.3.4.3; also called formyltetrahydrofolate synthetase) are the major reactions that produce THF-based one-carbon donors (10, 25, 26). Although several glycine cleavage system candidates were detected in the genome of *H. thermophilus*, this system does not appear to be operative. As the components of this system are not highly conserved among members of the order *Aquificales*, and because truncated homologs of P protein and low-identity L proteins were detected in *H. thermophilus*, these proteins might have other functions in *H. thermophilus*. Formate-THF ligase is used for CO₂ fixation in the Wood-Ljungdahl pathway (10, 27) or one-carbon compounds by methylotrophs (28). However, *H. thermophilus*, which fixes CO₂ using the reductive TCA cycle, lacks a homolog of this enzyme. Taken together, these findings indicate that SHMT is the major provider of one-carbon units in *H. thermophilus*.

If SHMT is the major provider of one-carbon units and a glycine cleavage system is absent, the amount of glycine produced and one-carbon units in biosynthesis are stoichiometrically coupled. Therefore, excess glycine needs to be excreted or further converted. Since no clear accumulation of glycine in the medium was observed when the $\Delta pspA$ mutant and the wild type were cultivated with supplemented serine (Fig. 3A and C), excess glycine should be further catabolized *in vivo*. We are expecting that glycine is converted into glyoxylate and then used as a carbon source, because *H.*

thermophilus possesses at least 13 transaminase genes and a glycine oxidase (EC 1.4.3.19) candidate gene.

The present results demonstrate that iPSPs are essential for the synthesis of serine in *H. thermophilus*. This finding, together with the lack of detectable glycine cleavage system activity in this autotrophic bacterium, strongly suggests that serine is the major THF-based one-carbon donor in *H. thermophilus*.

MATERIALS AND METHODS

Bacterial strains and culture conditions. *H. thermophilus* TK-6 (IAM 12695, DSMZ 6534) was cultivated at 70°C under a gas phase of H₂-O₂-CO₂ (75:10:15 [vol/vol/vol]) in a 100-ml vial containing 10 ml of autotrophic medium, as described previously (2). A *pspA* deletion mutant of strain TK-6 (Δ *pspA*) was maintained in the autotrophic medium supplemented with 300 μ M L-serine. To monitor the growth of *H. thermophilus* and the Δ *pspA* mutant, cells in early stationary phase were inoculated into fresh medium to give an optical density at 540 nm (OD₅₄₀) of approximately 0.1. For solid culture, autotrophic medium containing 300 μ M L-serine and 10 μ g · ml⁻¹ CuSO₄·5H₂O was solidified with 1.0% (wt/vol) Gelrite (Wako Pure Chemical Industries, Osaka, Japan). When required, 100 μ g · ml⁻¹ ampicillin and 500 μ g · ml⁻¹ kanamycin were added to the medium for *Escherichia coli* and *H. thermophilus*, respectively.

Deletion of *pspA* or *pspB* in *H. thermophilus*. The *pspA* and *pspB* genes were deleted in *H. thermophilus*, using a previously established homologous recombination method (29), with pEX18Ap used as a vector in place of pUC19. To construct plasmids pEX18Ap-d*pspA* and pEX18Ap-d*pspB*, approximately 1.1-kbp fragments of the *pspA* and *pspB* genes were PCR amplified from *H. thermophilus* chromosomal DNA using the primer pairs *pspA1_F* (5'-TAT Gga att cGT AAG CCT GGA GAG-3') and *pspA1_R* (5'-Gag gta cCA AGA AAA AGA CAC TGG-3'), and *pspA2_F* (5'-CAA ggt acc TAG CCT GCT TTT TAC-3') and *pspA2_R* (5'-TAG tct aga GGA AGC TCT GGG AAG-3') (lowercase letters indicate EcoRI, KpnI, or XbaI restriction endonuclease sites) for pEX18Ap-d*pspA*; and *pspB1_F* (5'-GCT gag ctc GTT TTG CCT ACA CC-3') and *pspB1_R* (5'-TAA ggt acc TTT GCC ACA CAA GAC-3'), and *pspB2_F* (5'-GAG ggt acc GCA TAT CAC TTA TGG-3') and *pspB2_R* (5'-AGC gga tcc TTT ACA GAA GAG GAG-3') (lowercase letters indicate SacI, KpnI, or BamHI restriction endonuclease site) for pEX18Ap-d*pspB*. A thermostable kanamycin nucleotidyltransferase gene (*htk*) (30) was amplified using the primers *htk_F* (5'-AAA ggt acc ACA TTC GGT GAG AAG CTA CAG G-3') and *htk_R* (5'-AAA ggt acc GGT CAT CGT TCA AAA TGG TAT G-3') (lowercase letters indicate KpnI sites). These three fragments were digested with the appropriate restriction enzymes and cloned into pEX18Ap as a fused sequence.

After transfecting the constructed pEX18Ap-d*pspA* or pEX18Ap-d*pspB* into *H. thermophilus* by electroporation, positive transformants (designated Δ *pspA* and Δ *pspB* mutants) were selected on a solid medium supplemented with kanamycin. Disruption of the genes was confirmed by checking the length and sequence of PCR-amplified DNA fragments containing the *pspA* or *pspB* and *htk* genes.

Quantification of serine and glycine. Serine and glycine were detected and quantified using high-performance liquid chromatography with a reverse-phase column after phenylthiocarbamyl derivatization, as described previously (31, 32). Alanine was used as an internal standard.

Preparation of cell extract. *H. thermophilus* cells were harvested during the early stationary phase by centrifugation (10,000 × *g* for 15 min). The collected cells were resuspended in a 4-fold volume of 20 mM Tris-HCl (pH 8.0) and disrupted by sonication (Ultrasonic disruptor UD-201; Tomy, Japan). After centrifugation at 100,000 × *g* and 4°C for 1 h, the supernatant was collected and used as the cell extract.

Glycine cleavage system assay. Glycine cleavage system activity was assayed using two methods. In the first method, the decrease in glycine was monitored in 150 μ l of a reaction mixture consisting of 20 mM Tris-HCl (pH 7.5 at 70°C), 5 mM glycine, 3.5 mM THF, 5 mM NAD⁺ or NADP⁺, 10 mM 2-mercaptoethanol, and cell extract (80 μ g of protein) during incubation at 70°C. The second method utilized an artificial electron acceptor, benzyl viologen (BV), in place of NAD(P)⁺ to monitor the reduction of BV. Four hundred microliters of the reaction mixture, which contained 20 mM Tris-HCl (pH 7.5 at 70°C), 5 mM glycine, 3.5 mM THF, 5 mM BV, 0.125 mM dithiothreitol, and cell extract (200 μ g of protein), was incubated at 70°C in argon gas. The reduction of BV was monitored photometrically at 578 nm.

SUPPLEMENTAL MATERIAL

Supplemental material for this article may be found at <https://doi.org/10.1128/JB.00409-17>.

SUPPLEMENTAL FILE 1, PDF file, 0.1 MB.

ACKNOWLEDGMENTS

We gratefully acknowledge Mami Kato for providing technical assistance with the maker gene (*htk*) construction and Maiko Yamamoto and Gaku Kono for helping with qRT-PCR experiments.

This work was supported in part by Grant-in-aid for Scientific Research (A) 21248010 (to M.I.) and Grant-in-Aid for Scientific Research on Innovative Areas (research in a proposed research area) 17H05242 (to Y.C.) from the Japan Society for the Promotion of Science.

REFERENCES

- Shiba H, Kawasumi T, Igarashi Y, Kodama T, Minoda Y. 1985. The CO₂ assimilation via the reductive tricarboxylic acid cycle in an obligately autotrophic, aerobic hydrogen-oxidizing bacterium, *Hydrogenobacter thermophilus*. Arch Microbiol 141:198–203. <https://doi.org/10.1007/BF00408058>.
- Shiba H, Kawasumi T, Igarashi Y, Kodama T, Minoda Y. 1982. The deficient carbohydrate metabolic pathways and the incomplete tricarboxylic acid cycle in an obligately autotrophic hydrogen-oxidizing bacterium. Agric Biol Chem 46:2341–2345. <https://doi.org/10.1080/00021369.1982.10865433>.
- Chiba Y, Oshima K, Arai H, Ishii M, Igarashi Y. 2012. Discovery and analysis of cofactor-dependent phosphoglycerate mutase homologs as novel phosphoserine phosphatases in *Hydrogenobacter thermophilus*. J Biol Chem 287:11934–11941. <https://doi.org/10.1074/jbc.M111.330621>.
- Chiba Y, Horita S, Ohtsuka J, Arai H, Nagata K, Igarashi Y, Tanokura M, Ishii M. 2013. Structural units important for activity of a novel-type phosphoserine phosphatase from *Hydrogenobacter thermophilus* TK-6 revealed by crystal structure analysis. J Biol Chem 288:11448–11458. <https://doi.org/10.1074/jbc.M112.449561>.
- Kim KT, Chiba Y, Arai H, Ishii M. 2015. Discovery of an intermolecular disulfide bond required for the thermostability of a heterodimeric protein from the thermophile *Hydrogenobacter thermophilus*. Biosci Biotechnol Biochem 80:232–240. <https://doi.org/10.1080/09168451.2015.1079476>.
- Ho C-L, Saito K. 2001. Molecular biology of the plastidic phosphorylated serine biosynthetic pathway in *Arabidopsis thaliana*. Amino Acids 20:243–259. <https://doi.org/10.1007/s007260170042>.
- Chiba Y, Terada T, Kameya M, Shimizu K, Arai H, Ishii M, Igarashi Y. 2012. Mechanism for folate-independent aldolase reaction catalyzed by serine hydroxymethyltransferase. FEBS J 279:504–514. <https://doi.org/10.1111/j.1742-4658.2011.08443.x>.
- Šmejkalová H, Erb TJ, Fuchs G. 2010. Methanol assimilation in *Methylobacterium extorquens* AM1: demonstration of all enzymes and their regulation. PLoS One 5:e13001. <https://doi.org/10.1371/journal.pone.0013001>.
- Kikuchi G. 1973. The glycine cleavage system: composition, reaction mechanism, and physiological significance. Mol Cell Biochem 1:169–187. <https://doi.org/10.1007/BF01659328>.
- Zhuang W-Q, Yi S, Bill M, Brisson VL, Feng X, Men Y, Conrad ME, Tang YJ, Alvarez-Cohen L. 2014. Incomplete Wood-Ljungdahl pathway facilitates one-carbon metabolism in organohalide-respiring *Dehalococcoides mccartyi*. Proc Natl Acad Sci U S A 111:6419–6424. <https://doi.org/10.1073/pnas.1321542111>.
- Sato T, Kochi H, Motokawa Y, Kawasaki H, Kikuchi G. 1969. Glycine metabolism by rat liver mitochondria: I. Synthesis of two molecules of glycine from one molecule each of serine, bicarbonate and ammonia. J Biochem 65:63–70.
- Harder W, Quayle J. 1971. The biosynthesis of serine and glycine in *Pseudomonas* AM1 with special reference to growth on carbon sources other than C₁ compounds. Biochem J 121:753–762. <https://doi.org/10.1042/bj1210753>.
- Locasale JW. 2013. Serine, glycine and one-carbon units: cancer metabolism in full circle. Nat Rev Cancer 13:572–583. <https://doi.org/10.1038/nrc3557>.
- Bailey LB, Gregory JF, III. 1999. Folate metabolism and requirements. J Nutr 129:779–782.
- Phillips RS, Miles EW, Cohen LA. 1984. Interactions of tryptophan synthase, tryptophanase, and pyridoxal phosphate with oxindolyl-L-alanine and 2,3-dihydro-L-tryptophan: support for an indolenine intermediate in tryptophan metabolism. Biochem J 23:6228–6234. <https://doi.org/10.1021/bi00320a052>.
- Tsoi BM, Beckhouse AG, Gelling CL, Rafferty MJ, Chiu J, Tsoi AM, Lauterbach L, Rogers PJ, Higgins VJ, Dawes IW. 2009. Essential role of one-carbon metabolism and Gcn4p and Bas1p transcriptional regulators during adaptation to anaerobic growth of *Saccharomyces cerevisiae*. J Biol Chem 284:11205–11215. <https://doi.org/10.1074/jbc.M809225200>.
- Fox JT, Stover PJ. 2008. Folate-mediated one-carbon metabolism. Vitam Horm 79:1–44.
- Hama H, Shimamoto T, Tsuda M, Tsuchiya T. 1988. Characterization of a novel L-serine transport system in *Escherichia coli*. J Bacteriol 170:2236–2239. <https://doi.org/10.1128/jb.170.5.2236-2239.1988>.
- Yamamoto T, Nishizaki I, Furuya S, Hirabayashi Y, Takahashi K, Okuyama S, Yamamoto H. 2003. Characterization of rapid and high-affinity uptake of L-serine in neurons and astrocytes in primary culture. FEBS Lett 548:69–73. [https://doi.org/10.1016/S0014-5793\(03\)00742-7](https://doi.org/10.1016/S0014-5793(03)00742-7).
- Robbins JC, Oxender DL. 1973. Transport systems for alanine, serine, and glycine in *Escherichia coli* K-12. J Bacteriol 116:12–18.
- Velayudhan J, Jones MA, Barrow PA, Kelly DJ. 2004. L-Serine catabolism via an oxygen-labile L-serine dehydratase is essential for colonization of the avian gut by *Campylobacter jejuni*. Infect Immun 72:260–268. <https://doi.org/10.1128/IAI.72.1.260-268.2004>.
- Florio R, di Salvo ML, Vivoli M, Contestabile R. 2011. Serine hydroxymethyltransferase: a model enzyme for mechanistic, structural, and evolutionary studies. Biochim Biophys Acta 1814:1489–1496. <https://doi.org/10.1016/j.bbapap.2010.10.010>.
- Kameya M, Arai H, Ishii M, Igarashi Y. 2010. Purification of three aminotransferases from *Hydrogenobacter thermophilus* TK-6—novel types of alanine or glycine aminotransferase: enzymes and catalysis. FEBS J 277:1876–1885. <https://doi.org/10.1111/j.1742-4658.2010.07604.x>.
- Appling DR. 1991. Compartmentation of folate-mediated one-carbon metabolism in eukaryotes. FASEB J 5:2645–2651.
- Braakman R, Smith E. 2012. The emergence and early evolution of biological carbon-fixation. PLoS Comput Biol 8:e1002455. <https://doi.org/10.1371/journal.pcbi.1002455>.
- Dev IK, Harvey RJ. 1982. Sources of one-carbon units in the folate pathway of *Escherichia coli*. J Biol Chem 257:1980–1986.
- Poehlein A, Schmidt S, Kaster A-K, Goenrich M, Vollmers J, Thürmer A, Bertsch J, Schuchmann K, Voigt B, Hecker M, Daniel R, Thauer RK, Gottschalk G, Müller V. 2012. An ancient pathway combining carbon dioxide fixation with the generation and utilization of a sodium ion gradient for ATP synthesis. PLoS One 7:e33439. <https://doi.org/10.1371/journal.pone.0033439>.
- Marx CJ, Laukel M, Vorholt JA, Lidstrom ME. 2003. Purification of the formate-tetrahydrofolate ligase from *Methylobacterium extorquens* AM1 and demonstration of its requirement for methylotrophic growth. J Bacteriol 185:7169–7175. <https://doi.org/10.1128/JB.185.24.7169-7175.2003>.
- Yamamoto M, Arai H, Ishii M, Igarashi Y. 2006. Role of two 2-oxoglutarate: ferredoxin oxidoreductases in *Hydrogenobacter thermophilus* under aerobic and anaerobic conditions. FEMS Microbiol Lett 263:189–193. <https://doi.org/10.1111/j.1574-6968.2006.00415.x>.
- Hoseki J, Yano T, Koyama Y, Kuramitsu S, Kagamiyama H. 1999. Directed evolution of thermostable kanamycin-resistance gene: a convenient selection marker for *Thermus thermophilus*. J Biochem 126:951–956. <https://doi.org/10.1093/oxfordjournals.jbchem.a022539>.
- Heinrikson RL, Meredith SC. 1984. Amino acid analysis by reverse-phase high-performance liquid chromatography: precolumn derivatization with phenylisothiocyanate. Anal Biochem 136:65–74. [https://doi.org/10.1016/0003-2697\(84\)90307-5](https://doi.org/10.1016/0003-2697(84)90307-5).
- Kameya M, Ikeda T, Nakamura M, Arai H, Ishii M, Igarashi Y. 2007. A novel ferredoxin-dependent glutamate synthase from the hydrogen-oxidizing chemoautotrophic bacterium *Hydrogenobacter thermophilus* TK-6. J Bacteriol 189:2805–2812. <https://doi.org/10.1128/JB.01360-06>.
- Okamura-Ikeda K, Ohmura Y, Fujiwara K, Motokawa Y. 1993. Cloning and nucleotide sequence of the *gcv* operon encoding the *Escherichia coli* glycine-cleavage system. Eur J Biochem 216:539–548. <https://doi.org/10.1111/j.1432-1033.1993.tb18172.x>.
- Hama H, Shimamoto T, Tsuda M, Tsuchiya T. 1987. Properties of a Na⁺-coupled serine-threonine transport system in *Escherichia coli*. Biochim Biophys Acta 905:231–239. [https://doi.org/10.1016/0005-2736\(87\)90451-2](https://doi.org/10.1016/0005-2736(87)90451-2).