Published in final edited form as:

AIDS Behav. 2017 July ; 21(7): 1914–1925. doi:10.1007/s10461-017-1740-7.

Prevalence and factors associated with hazardous alcohol use among persons living with HIV across the US in the current era of antiretroviral treatment

Heidi M Crane¹, Mary E McCaul², Geetanjali Chander³, Heidi Hutton², Robin M Nance¹, Joseph AC Delaney⁴, Joseph O Merrill¹, Bryan Lau⁵, Kenneth H Mayer⁶, Michael J Mugavero⁷, Matthew Mimiaga⁸, James H Willig⁷, Greer A. Burkholder⁷, Daniel R Drozd¹, Rob J Fredericksen¹, Karen Cropsey⁹, Richard D Moore³, Jane M. Simoni¹⁰, W Christopher Mathews¹¹, Joseph J Eron¹², Sonia Napravnik¹², Katerina Christopoulos¹³, Elvin Geng¹³, Michael S Saag⁷, and Mari M Kitahata¹

¹Department of Medicine, University of Washington, Seattle, WA

²Department of Psychiatry and Behavioral Sciences, Johns Hopkins University, Baltimore, MD

³Department of Medicine, Johns Hopkins University, Baltimore, MD

⁴Department of Epidemiology, University of Washington

⁵Department of Epidemiology, Johns Hopkins University, Baltimore, MD

⁶Fenway Institute and Department of Medicine, Harvard Medical School, Boston, MA

⁷Department of Medicine, University of Alabama at Birmingham, Birmingham, AL

⁸Fenway Institute and Division of Psychiatry, Harvard Medical School, Boston MA

⁹Department of Psychiatry, University of Alabama at Birmingham, Birmingham, AL

¹⁰Department of Psychology, University of Washington, Seattle, WA

¹¹Department of Medicine, University of California at San Diego, San Diego, CA

¹²Department of Medicine, University of North Carolina, Chapel Hill, NC

¹³Department of Medicine, University of California at San Francisco, San Francisco, CA

Abstract

Hazardous alcohol use is associated with detrimental health outcomes among persons living with HIV (PLWH). We examined the prevalence and factors associated with hazardous alcohol use in

Corresponding author: Heidi M Crane, MD, MPH, University of Washington, 325 9th Ave, Box 359931, Seattle, WA, 206-744-6649 206-744-3693 fax, hcrane@uw.edu.

Data presented in part at the 38th Annual Research Society on Alcoholism Scientific Meeting, San Antonio, Texas, 6/2015.

Conflicts of interest: There are no conflicts of interest for this paper.

Compliance with ethical standards

The authors declare they have no conflicts of interest. All sites have approval from Human Subjects committees at their local institution and participants signed informed consent to participate in CNICS. All data submitted to the CNICS Data Repository are deidentified. For this retrospective data analysis of existing deidentified data in the CNICS Data Repository, no additional formal consent is required. This article does not contain any studies with animals performed by any of the authors.

the current era using several hazardous drinking definitions and binge drinking defined as 5 drinks for men vs. 4 for women. We included 8567 PLWH from 7 U.S. sites from 2013–2015. Current hazardous alcohol use was reported by 27% and 34% reported binge drinking. In adjusted analyses, current and past cocaine/crack (odd ratio [OR] 4.1:3.3–5.1, $p<0.001$ and OR 1.3:1.1–1.5, $p<0.001$ respectively), marijuana (OR 2.5:2.2–2.9, $p<0.001$ and OR 1.4:1.2–1.6, $p<0.001$), and cigarette use (OR 1.4:1.2–1.6, $p<0.001$ and OR 1.3:1.2–1.5, $p<0.001$) were associated with increased hazardous alcohol use. The prevalence of hazardous alcohol use remains high in the current era, particularly among younger men. Routine screening and targeted interventions for hazardous alcohol use, potentially bundled with interventions for other drugs, remain a key aspect of HIV care.

Keywords

HIV; alcohol use; hepatitis C; substance use

INTRODUCTION

Hazardous alcohol use is common among persons living with HIV (PLWH)[1] with studies such as the HIV Cost and Service Utilization survey and the HIV Research Network (HIVRN) previously reporting rates of 8 and 11%[1,2]. Hazardous drinking rates among PLWH have previously been found to be higher than the general population[1]. Predictors of hazardous alcohol use among PLWH include illicit drug use use[1,2] and depression or other mental health disorders[3] while the impact of other factors such as hepatitis C virus (HCV) is less clear.

Hazardous alcohol use is associated with detrimental health outcomes including multiple steps along the HIV treatment cascade such as retention in care[4,5]; lower CD4 counts among those not on antiretroviral therapy (ART)[6], delayed initiation of and decreased adherence to ART[7,8], and poorer survival[9]. Hazardous alcohol use potentially increases the risk of HIV transmission by an increased likelihood of risky sexual behavior including unprotected sex with multiple partners[10,11].

With potent ART, survival has increased dramatically[12,13]. Consequently, the demographic and clinical characteristics of PLWH in care in the US are changing; for example there is an increasingly disproportionate share of both new HIV diagnoses and individuals living with HIV among Blacks and Latinos and an increasingly larger proportion of those living with HIV among those age 50 or older [14,15]. However, many earlier studies of alcohol use particularly hazardous alcohol use in PLWH were conducted before the current ART treatment era[1,2,8,16,17]; had small sample sizes[17,18]; and unrepresentative patient groups, including few women or other restrictive patient characteristics[16,19] yielding results with limited generalizability to PLWH currently in care.

The purpose of this study was to determine the prevalence and factors associated with hazardous alcohol use among PLWH in care across the US in the current era. We hypothesized that in the current ART era substance use and depression would be associated

with hazardous alcohol use and HCV co-infection would be associated with no/low alcohol use.

METHODS

Centers for AIDS Research Network of Integrated Clinical Systems (CNICS) cohort

CNICS is a longitudinal observational study of PLWH receiving primary care from 8 clinical sites from 1/1/1995 to the present[20]. CNICS was created to better define the relationship between patient and treatment factors and long-term clinical outcomes among PLWH in the ART era and to investigate questions related to HIV disease management that cannot be readily addressed through traditional randomized controlled trials and other cohort studies[20]. PLWH from 7 CNICS sites with comprehensive collection of the CNICS assessment including hazardous alcohol use during the study time period were included in this study to ensure geographic and racial/ethnic diversity: University of Alabama at Birmingham; University of California, San Francisco; University of Washington; University of California, San Diego; Fenway Community Health Center; University of North Carolina, Chapel Hill; and Johns Hopkins University. Substance use, particularly hazardous alcohol use has been an area of particular interest for CNICS including studies examining how patients and providers prioritize its assessment in clinical care[21] and its impact along with other factors on outcomes such as liver fibrosis, adherence and sexual transmission risk[22–24].

Study Subjects—All PLWH eighteen years of age or older who completed a clinical assessment of patient reported behaviors and outcomes as part of a clinical care visit at least once between 2013–2015 were eligible. We did not include assessments from before 1/2013 to ensure that this was representative of the current treatment era. The clinical assessment is completed approximately every 4–6 months. For those who completed multiple assessments during the study period, the most recent assessment was used. CNICS data collection was approved by Institutional Review Boards at each site.

Data Sources—The CNICS data repository integrates longitudinal data including comprehensive clinical information from outpatient and inpatient encounters, demographic, clinical, medication, laboratory, and socioeconomic data obtained from each site's electronic health record and other institutional data sources[20].

The CNICS data repository integrates clinical assessment data. Patients used touch-screen tablets to complete the ~10–12 minute clinical assessment including measures of alcohol use (Alcohol Use Disorders Identification Test (AUDIT-C)[25,26], substance use (modified Alcohol, Smoking, and Substance Involvement Screening Test [ASSIST])[27,28], depressive symptoms (Patient Health Questionnaire [PHQ-9])[29] and other domains.

Instrument Scoring—AUDIT-C scores for current alcohol consumption measured over the prior year were calculated by summing scores for each question[30]. AUDIT-C asks three questions (0–4 points each) about alcohol use during the past year; how often a person has a drink containing alcohol, the usual quantity of drinks consumed, and the frequency of drinking a large number of drinks at one time. We categorized scores as a tertiary variable of

No use, Low alcohol use, and Hazardous alcohol use using a cut-off score of 4 for men and 3 for women to define Hazardous alcohol use[31]. We repeated analyses using a score of 5 for men and 4 for women to define Hazardous alcohol use[32]. We also created a binary alcohol use variable of No/low use (No use + Low alcohol use) and Hazardous alcohol use. Finally, we used binge drinking as an outcome comparing recent binge drinking (less than monthly, monthly, weekly, daily or almost daily) vs. no binge drinking. Binge drinking was defined as 5 drinks for men vs. 4 drinks for women on one occasion.

ASSIST categorizes drug use as current use (past 3 months), prior use, or never used[27,28]. We examined illicit substance use by 1) specific type of drug used (marijuana, crack/cocaine, methamphetamines/crystal, or illicit opioids/heroin); 2) any drug use; and 3) any drug use excluding marijuana given its evolving legal status. Cigarette use was categorized as current, prior, or never used.

Depressive symptom scores using the PHQ-9 range from 0–27 and were categorized as: none (0–4), mild (5–9), moderate (10–19), or severe (≥ 20 points)[29].

Statistical Analyses—We performed bivariate analysis using chi-squared tests. We compared participants who completed the clinical assessment and were therefore included in the study to those who did not complete an assessment during the study period. We compared characteristics among those with no use, low use, and hazardous alcohol use. We examined demographic (age, race/ethnicity, gender, HIV transmission risk factor), and clinical characteristics including CD4⁺ cell count nadir (< 350, 350–499, and ≥ 500 cells/mm³), current CD4⁺ cell count (< 350, 350–499, and ≥ 500 cells/mm³), current HIV-1 RNA viral load level (detectable vs. undetectable), current ART use, HCV infection indicated by either the presence of HCV antibody or HCV RNA, depression category, and substance use (overall and by individual drug class). We examined the percentage of PLWH in different demographic and clinical categories with hazardous alcohol use.

We used multivariate logistic regression to examine factors associated with hazardous versus no/low alcohol use. Inclusion in models was based on hypotheses, known associations, potential confounders, and bivariate results. In addition to including demographic characteristics, we also included HIV treatment related characteristics and substance use. Adjusted models included age, race/ethnicity, gender, HIV transmission risk factor, clinical site, HCV status, CD4 cell count nadir, current viral load level, depression symptoms, substance use, and smoking status. Partial correlations were assessed and too low to create bias due to collinearity. Analyses were repeated excluding those who reported no alcohol use as this is a heterogeneous group including both current non-drinkers who never were hazardous alcohol users and prior hazardous alcohol user who due to illness, consequences of alcohol use, or other reasons became non-drinkers[33,34]. We repeated analyses using higher cut-offs (AUDIT-C scores of 5 for men and 4 for women) to define hazardous alcohol use, and using binge drinking. As odds ratios from logistic regression analyses for common outcomes can be different from underlying prevalence ratios, we conducted secondary analyses using generalized linear models with relative risks rather than odds ratios. We repeated models stratified by individual sites to look for differences across site. Lastly, we conducted sensitivity analyses stratified by gender to examine findings of models

specifically for women vs. men. Stratified models included similar covariates as the main models except gender and HIV transmission risk factor were excluded from the models. Two-tailed p values <0.05 were considered significant.

Results

Between 2013–2015 the clinical assessment was completed by 8567 PLWH. Demographic and clinical characteristics are presented in Table I categorized by current alcohol use. Mean age of participants was 46 years with 3441 (41%) aged 50 or older, 1305 (15%) women, and 1298 with HCV (15%). Demographic and clinical characteristics of participants were similar to those of CNICS patients who did not complete an assessment during the study period and to 345 individuals who started the assessment but did not complete it (data not shown).

Hazardous alcohol use

No current alcohol use was reported by 33%, 41% reported low alcohol use and 27% reported hazardous alcohol use (AUDIT-C scores of 4 for men and 3 for women). Binge drinking was reported by 34%. Rates of hazardous alcohol use were higher among men, those with a detectable viral load, and those not receiving ART (Table I). Hazardous alcohol use was less common with older ages. Hazardous alcohol use was less frequently reported by PLWH with HCV (21%) compared to no HCV (27%).

Hazardous alcohol use was more prevalent among PLWH who reported current illicit drug use (38%) compared with past (22%) or no substance use (15%) (Table II). This pattern was consistent across all drugs except methamphetamine use where hazardous alcohol use was most prevalent among those with past not current use (Table II), with particularly prevalent hazardous alcohol use with cocaine/crack use.

Factors associated with hazardous alcohol use

In bivariate analyses (Table III, column 1), factors associated with hazardous alcohol use included male gender, younger age, white race, not having HCV, and having a detectable VL. Past methamphetamine/crystal use, past and current cocaine/crack and marijuana use, and current opioid/heroin use were associated with hazardous alcohol use.

In adjusted logistic regression analyses of hazardous alcohol use (AUDIT-C scores of 4 for men and 3 for women), compared to those reporting no/low alcohol use, current illicit drug use *including marijuana* were associated with hazardous alcohol use (OR 3.21: 2.75–3.73, $p<0.001$) and past (OR 1.60: 1.37–1.87, $p<0.001$). In adjusted models, current substance use *excluding marijuana* were also associated with hazardous alcohol use (OR 1.74: 1.48–2.06, $p<0.001$) and past (OR 1.24: 1.09–1.42, $p<0.001$).

We repeated analyses using individual illicit drugs. Factors associated with no/low alcohol use included older age, black race, and HCV co-infection while past and present cocaine/crack use and marijuana use were associated with hazardous alcohol use as was past and current cigarette use (Table III, column 2).

We conducted sensitivity analyses comparing those with hazardous alcohol use to those with low use excluding those reporting that they did not drink at all (Table III, column 3). This was to exclude those with prior hazardous alcohol use who had become non-drinkers[33–35]. Findings were similar to results including nondrinkers except HCV was no longer associated with low alcohol use. We also repeated analyses using a higher cutoff to define hazardous alcohol use (Table III, Column 4) with a similar pattern of results.

In adjusted analyses using binge drinking to characterize drinking patterns (Table III, column 5), female gender was associated with a lower odds of binge drinking than male gender as was older age. Past and current cocaine/crack and marijuana use, and past and current cigarette smoking were all associated with increased odds of binge drinking. In contrast to several other alcohol measures, HCV was not associated with significantly less binge drinking. In addition, we examined those with hazardous alcohol use vs. no/low alcohol use using relative risk rather than odds ratios and found similar findings (Table III, column 6).

Because there are differences in CNICS sites in terms of participants (e.g. California sites have a higher proportion of Latino PLWH than the site in Birmingham, Alabama), we repeated models looking at individual sites. The pattern of findings in the adjusted models were similar by site although not all findings that were statistically significant in the overall model were significant in the individual site models with smaller sample sizes.

Lastly we conducted analyses stratified by gender (Table IV). Results were similar to overall findings, however a few differences were notable. The stratified analyses highlighted that associations between black race and no/low alcohol use were driven by men and associations between CD4 nadir >350 cells/ml³ and hazardous alcohol use were driven by women. In contrast, past and current cocaine/crack and marijuana use were associated with hazardous alcohol use among both men and women.

Discussion

This study describes alcohol use among a population of >8000 PLWH in care across the U.S. in the current treatment era. It demonstrates the high prevalence of hazardous alcohol use (27%) which has important implications on outcomes among PLWH such as adherence to ART and survival. There was a lower prevalence of hazardous alcohol use among older individuals and those with HCV co-infection, though alcohol use was still substantial among these groups. Past and present cocaine/crack use and marijuana use were associated with increased odds of hazardous alcohol use. The results of this study have important implications for the care of PLWH including highlighting the crucial need for on-going screening and improved treatment delivery for alcohol use in clinical care systems.

Prevalence of hazardous alcohol use

The prevalence of hazardous alcohol use was higher in this study compared to the HIV Cost and Services Utilization survey or the HIV Research Network (HIVRN) where 8% or 11% reported heavy/hazardous alcohol use[1,2]. These differences may be explained in part by definitions. For example, HIVRN defined hazardous drinking as >14 drinks/week or 5

drinks/occasion for men and >7 drinks/week or 4 drinks/occasion for women. Of note, even when using the more stringent AUDIT-C definition of 5 for men and 4 for women, 18% in the current study reported hazardous alcohol use. Differences in how alcohol use was collected may also contribute to the higher prevalence in CNICS. CNICS uses touch-screen tablets to collect assessments, lessening some of the social desirability bias and underreporting associated with interviewer-based collection. Several studies have indicated patient preference for self-administered electronic over interviewer-based data collection, and electronic administration has yielded more disclosure of sensitive behaviors[36–39]. It may be that study differences can also be explained in part by changes in demographic and clinical characteristics of PLWH in the US in recent years. Only 33% in CNICS reported no alcohol use which is notable given increasing evidence that perhaps no level of alcohol consumption is ‘safe’ among PLWH[40]. The high prevalence of hazardous alcohol use found in this study has clinical implications in terms of ensuring clinical care settings have enough resources to ensure adequate identification and treatment of this complex and yet crucial issue among PLWH.

Aging

Effective ART has resulted in many PLWH living into middle and old age. Of note, among older PLWH, hazardous alcohol use was less likely for those 60, who were less than half as likely to report hazardous alcohol use compared with those <30. This is consistent with general population studies which have found an association between older age and less alcohol use[41,42]. Despite this, 17% of PLWH 60 reported hazardous alcohol use suggesting that there is still need for additional interventions targeting all ages of PLWH, and for additional research to better understand the effectiveness of interventions across the age spectrum for PLWH, particularly given the associations between alcohol and poor outcomes among the elderly such as falls, fatal injuries and adverse drug reactions[43,44].

Substance use

We found hazardous alcohol use was more common among current substance users and that this association varied by individual drug. Both current and past cocaine/crack and marijuana use were associated with a higher odds of hazardous alcohol use. These findings highlight the importance of research that evaluates associations between individual substances and alcohol use as it can vary across drugs and may differentially affect outcomes. Further research is needed to identify whether certain patterns of substance use, for instance alcohol and cocaine/crack, are particularly detrimental to HIV outcomes and warrant specific interventions and to better identify the role of joint interventions that simultaneously target both alcohol and drug use.

HCV

HCV co-infection is common among PLWH and liver disease is a leading cause of non-AIDs mortality[45,46]. Hazardous alcohol use exacerbates HCV-related liver disease through increased HCV viral replication, toxic effects on the liver, and indirectly via effects on ART adherence and decreased HCV treatment eligibility[47–51]. In addition, alcohol use among PLWH co-infected with HCV is associated with faster liver fibrosis progression, while sustained virological response to HCV therapy is associated with slower liver fibrosis

progression[52]. HCV treatment is rapidly evolving with higher treatment success rates and easier treatment regimens, however alcohol use remains a potential contraindication to HCV treatment among PLWH[53]. This is an important target population for interventions to reduce preventable morbidity and mortality.

We found 56% of PLWH co-infected with HCV reported alcohol use with 21% reporting hazardous alcohol use. This rate is high given the negative consequences. HCV was associated with a lower odds of hazardous alcohol use. One possible explanation for this could be selection effects with the combination of HCV and hazardous alcohol use making it less likely for a PLWH to survive and thereby be part of the cohort. However, in a sensitivity analysis that excluded non-drinkers, there was not a decreased odds of hazardous alcohol use among those co-infected with HCV. These findings raises the possibility that those with HCV are to some extent making different decisions about drinking than those without HCV, possibly heeding advise to reduce their risk for liver toxicity from alcohol.

Strengths

A study strength is the large, diverse cohort of PLWH. This cohort represents the changing epidemiology of HIV across the US with substantial numbers of women, racial and ethnic diversity, and a population increasing in mean age. A second strength is the focus on 2013 and after. Finally, the comprehensive measurement of not just alcohol but other drug use allows assessment of the role of individual drugs among PLWH who may have high rates of co-occurring substance use.

Limitations

Several limitations are worth noting. We evaluated associations with alcohol use, but associations do not necessarily indicate causation. Alcohol and drug use were collected in the clinical assessment which could lead to underestimates. However, use of electronic collection allows patient burden to be reduced permitting integration into care and decreasing underreporting of risk behaviors due to social desirability bias[54]. While the clinical assessment has expanded to include additional languages such as Amharic, this study included only English and Spanish-speaking PLWH which may reduce generalizability to PLWH who do not speak English or Spanish and/or are not in care. Studies are needed that focus more broadly, including additional socioeconomic and contextual factors to inform public policies and interventions, with effective interventions that can be routinely incorporated into clinical settings being of particular value.

Conclusions

We describe prevalence and factors associated with hazardous alcohol use among PLWH in care across the U.S in the current era. The purpose is to better understand alcohol use not to stigmatize PLWH but instead to ensure that risk behaviors are identified and addressed to prevent negative consequences. Rates of alcohol use remain high. Factors associated with hazardous alcohol use included younger age and past and present cocaine use and marijuana use. Strengths include inclusion of women, the large sample size and examining individual substances and patterns of substance use while accounting for overlapping use. While rates of hazardous alcohol use were lower among PLWH with HCV, the rates remained

substantial given the negative consequences. Routine screening and targeted interventions for hazardous alcohol and other substance use remain a key aspect of HIV care.

Acknowledgments

We thank the patients and providers throughout the CNICS network.

This work was supported by the National Institutes of Alcohol Abuse and Alcoholism (NIAAA) at the National Institutes of Health [U24AA020801, U01AA020793 and U01AA020802]. Additional support came from the National Institute of Allergy and Infectious Diseases (NIAID) at the National Institutes of Health [CNICS R24 AI067039, UW CFAR NIAID Grant P30 AI027757, UNC CFAR grant P30 AI50410, and UAB CFAR grant P30 AI027767].

Funding: This work was supported by the National Institutes of Alcohol Abuse and Alcoholism (NIAAA) at the National Institutes of Health [U24AA020801, U01AA020793 and U01AA020802]. Additional support came from the National Institute of Allergy and Infectious Diseases (NIAID) at the National Institutes of Health [CNICS R24 AI067039, UW CFAR NIAID Grant P30 AI027757; UNC CFAR grant P30 AI50410, JHU CFAR grant P30 AI094189, and UAB CFAR grant P30 AI027767].

References

- Galvan FH, Bing EG, Fleishman JA, et al. The prevalence of alcohol consumption and heavy drinking among people with HIV in the United States: results from the HIV Cost and Services Utilization Study. *Journal of Studies on Alcohol*. Mar; 2002 63(2):179–186. [PubMed: 12033694]
- Chander G, Josephs J, Fleishman JA, et al. Alcohol use among HIV-infected persons in care: results of a multi-site survey. *HIV Med*. Apr; 2008 9(4):196–202. [PubMed: 18366443]
- Galvan FH, Burnam MA, Bing EG. Co-occurring psychiatric symptoms and drug dependence or heavy drinking among HIV-positive people. *Journal of Psychoactive Drugs*. May; 2003 35(Suppl 1): 153–160. [PubMed: 12825758]
- Vagenas P, Azar MM, Copenhaver MM, Springer SA, Molina PE, Altice FL. The impact of alcohol use and related disorders on the HIV continuum of care: a systematic review : alcohol and the HIV continuum of care. *Curr HIV/AIDS Rep*. Dec; 2015 12(4):421–436. [PubMed: 26412084]
- Monroe AK, Lau B, Mugavero MJ, et al. Heavy alcohol use is associated with worse retention in HIV care. *J Acquir Immune Defic Syndr*. May 26.2016
- Samet JH, Cheng DM, Libman H, Nunes DP, Alperen JK, Saitz R. Alcohol consumption and HIV disease progression. *Journal of Acquired Immune Deficiency Syndromes*. Oct 1; 2007 46(2):194–199. [PubMed: 17667330]
- Chander G, Lau B, Moore R. Hazardous alcohol use: a risk factor for non-adherence and lack of suppression in HIV infection. *J Acquir Immune Defic Syndr*. 2006; 43(4):411–417. [PubMed: 17099312]
- Cook RL, Sereika SM, Hunt SC, Woodward WC, Erlen JA, Conigliaro J. Problem drinking and medication adherence among persons with HIV infection. *J Gen Intern Med*. Feb; 2001 16(2):83–88. [PubMed: 11251758]
- Braithwaite RS, Conigliaro J, Roberts MS, et al. Estimating the impact of alcohol consumption on survival for HIV+ individuals. *AIDS Care*. Apr; 2007 19(4):459–466. [PubMed: 17453583]
- Stein M, Herman DS, Trisvan E, Pirraglia P, Engler P, Anderson BJ. Alcohol use and sexual risk behavior among human immunodeficiency virus-positive persons. *Alcohol Clin Exp Res*. May; 2005 29(5):837–843. [PubMed: 15897729]
- Rehm J, Shield KD, Joharchi N, Shuper PA. Alcohol consumption and the intention to engage in unprotected sex: systematic review and meta-analysis of experimental studies. *Addiction*. Jan; 2012 107(1):51–59. [PubMed: 22151318]
- Palella FJ Jr, Delaney KM, Moorman AC, et al. Declining morbidity and mortality among patients with advanced human immunodeficiency virus infection. *HIV Outpatient Study Investigators*. *N Engl J Med*. 1998; 338(13):853–860. [PubMed: 9516219]
- Hogg RS, Heath KV, Yip B, et al. Improved survival among HIV-infected individuals following initiation of antiretroviral therapy. *JAMA*. Feb 11; 1998 279(6):450–454. [PubMed: 9466638]

14. The Henry J Kasier Foundation. The HIV/AIDS epidemic in the United States. 2017. <http://kff.org/hivaids/fact-sheet/the-hivaids-epidemic-in-the-united-states/>
15. Centers for Disease Control and Prevention. HIV among people aged 50 and over. 2016. <https://www.cdc.gov/hiv/group/age/olderamericans/>
16. Cook RL, Zhu F, Belnap BH, et al. Longitudinal trends in hazardous alcohol consumption among women with human immunodeficiency virus infection, 1995–2006. *Am J Epidemiol*. Apr 15; 2009 169(8):1025–1032. [PubMed: 19270052]
17. Lake-Bakaar G, Grimson R. Alcohol abuse and stage of HIV disease in intravenous drug abusers. *Journal of the Royal Society of Medicine*. Jul; 1996 89(7):389–392. [PubMed: 8774537]
18. Welch KJ. Correlates of alcohol and/or drug use among HIV-infected individuals. *AIDS Patient Care STDS*. Jun; 2000 14(6):317–323. [PubMed: 10897504]
19. Bertholet N, Cheng DM, Samet JH, Quinn E, Saitz R. Alcohol consumption patterns in HIV-infected adults with alcohol problems. *Drug Alcohol Depend*. Nov 1; 2010 112(1–2):160–163. [PubMed: 20579819]
20. Kitahata MM, Rodriguez BG, Haubrich R, et al. Cohort profile: the Centers for AIDS Research (CFAR) Network of Integrated Clinical Systems (CNICS). *Int J Epidemiol*. 2008; 37(5):948–955. [PubMed: 18263650]
21. Fredericksen RJ, Edwards TC, Merlin JS, et al. Patient and provider priorities for self-reported domains of HIV clinical care. *AIDS Care*. Oct; 2015 27(10):1255–1264. [PubMed: 26304263]
22. Kim HN, Nance R, Van Rompaey S, et al. Poorly controlled HIV infection: an independent risk factor for liver fibrosis. *J Acquir Immune Defic Syndr*. Aug 01; 2016 72(4):437–443. [PubMed: 26990826]
23. Mimiaga MJ, Biello K, Reisner SL, et al. Latent class profiles of internalizing and externalizing psychosocial health indicators are differentially associated with sexual transmission risk: findings from the CFAR Network of Integrated Clinical Systems (CNICS) cohort study of HIV-infected men engaged in primary care in the United States. *Health Psychol*. Feb 2.2015
24. Mimiaga MJ, Reisner SL, Grasso C, et al. Substance use among HIV-infected patients engaged in primary care in the United States: findings from the Centers for AIDS Research Network of Integrated Clinical Systems cohort. *Am J Public Health*. Aug; 2013 103(8):1457–1467. [PubMed: 23763417]
25. Bradley KA, McDonell MB, Bush K, Kivlahan DR, Diehr P, Fihn SD. The AUDIT alcohol consumption questions: reliability, validity, and responsiveness to change in older male primary care patients. *Alcohol Clin Exp Res*. Nov; 1998 22(8):1842–1849. [PubMed: 9835306]
26. Bradley KA, Bush KR, Epler AJ, et al. Two brief alcohol-screening tests From the Alcohol Use Disorders Identification Test (AUDIT): validation in a female Veterans Affairs patient population. *Arch Intern Med*. Apr 14; 2003 163(7):821–829. [PubMed: 12695273]
27. Newcombe DA, Humeniuk RE, Ali R. Validation of the World Health Organization Alcohol, Smoking and Substance Involvement Screening Test (ASSIST): report of results from the Australian site. *Drug and Alcohol Review*. May; 2005 24(3):217–226. [PubMed: 16096125]
28. WHO Assist Working Group. The Alcohol, Smoking and Substance Involvement Screening Test (ASSIST): development, reliability and feasibility. *Addiction*. Sep; 2002 97(9):1183–1194. [PubMed: 12199834]
29. Kroenke K, Spitzer RL, Williams JB. The PHQ-9: validity of a brief depression severity measure. *J Gen Intern Med*. Sep; 2001 16(9):606–613. [PubMed: 11556941]
30. Bush K, Kivlahan DR, McDonell MB, Fihn SD, Bradley KA. The AUDIT alcohol consumption questions (AUDIT-C): an effective brief screening test for problem drinking. Ambulatory Care Quality Improvement Project (ACQUIP). Alcohol Use Disorders Identification Test. *Arch Intern Med*. Sep 14; 1998 158(16):1789–1795. [PubMed: 9738608]
31. Bradley KA, DeBenedetti AF, Volk RJ, Williams EC, Frank D, Kivlahan DR. AUDIT-C as a brief screen for alcohol misuse in primary care. *Alcohol Clin Exp Res*. Jul; 2007 31(7):1208–1217. [PubMed: 17451397]
32. Gual A, Segura L, Contel M, Heather N, Colom J. AUDIT-3 and AUDIT-4: effectiveness of two short forms of the alcohol use disorders identification test. *Alcohol and Alcoholism*. Nov-Dec; 2002 37(6):591–596. [PubMed: 12414553]

33. Lucas N, Windsor TD, Caldwell TM, Rodgers B. Psychological distress in non-drinkers: associations with previous heavy drinking and current social relationships. *Alcohol and Alcoholism*. Jan-Feb;2010 45(1):95–102. [PubMed: 19917637]
34. Liang W, Chikritzhs T. The association between alcohol exposure and self-reported health status: the effect of separating former and current drinkers. *PLoS One*. 2013; 8(2):e55881. [PubMed: 23405228]
35. Crane HM, Nance RM, Merrill JO, et al. Not all non-drinkers with HIV are equal: demographic and clinical comparisons among current non-drinkers with and without a history of prior alcohol use disorders. *AIDS Care*. Feb; 2017 29(2):177–184. [PubMed: 27482893]
36. Newman JC, Des Jarlais DC, Turner CF, Gribble J, Cooley P, Paone D. The differential effects of face-to-face and computer interview modes. *Am J Public Health*. Feb; 2002 92(2):294–297. [PubMed: 11818309]
37. Perlis TE, Des Jarlais DC, Friedman SR, Arasteh K, Turner CF. Audio-computerized self-interviewing versus face-to-face interviewing for research data collection at drug abuse treatment programs. *Addiction*. Jul; 2004 99(7):885–896. [PubMed: 15200584]
38. Kurth AE, Martin DP, Golden MR, et al. A comparison between audio computer-assisted self-interviews and clinician interviews for obtaining the sexual history. *Sex Transm Dis*. Dec; 2004 31(12):719–726. [PubMed: 15608586]
39. Dolezal C, Marhefka SL, Santamaria EK, Leu CS, Brackis-Cott E, Mellins CA. A comparison of audio computer-assisted self-interviews to face-to-face interviews of sexual behavior among perinatally HIV-exposed youth. *Archives of Sexual Behavior*. Apr; 2012 41(2):401–410. [PubMed: 21604065]
40. Bryant KJ. Expanding research on the role of alcohol consumption and related risks in the prevention and treatment of HIV/AIDS. *Subst Use Misuse*. Jan 1; 2006 41(10–12):1465–1507. [PubMed: 17002990]
41. Eigenbrodt ML, Mosley TH Jr, Hutchinson RG, Watson RL, Chambless LE, Szklo M. Alcohol consumption with age: a cross-sectional and longitudinal study of the Atherosclerosis Risk in Communities (ARIC) study, 1987–1995. *Am J Epidemiol*. Jun 1; 2001 153(11):1102–1111. [PubMed: 11390330]
42. Merrick EL, Horgan CM, Hodgkin D, et al. Unhealthy drinking patterns in older adults: prevalence and associated characteristics. *J Am Geriatr Soc*. Feb; 2008 56(2):214–223. [PubMed: 18086124]
43. Mukamal KJ, Mittleman MA, Longstreth WT Jr, Newman AB, Fried LP, Siscovick DS. Self-reported alcohol consumption and falls in older adults: cross-sectional and longitudinal analyses of the cardiovascular health study. *J Am Geriatr Soc*. Jul; 2004 52(7):1174–1179. [PubMed: 15209658]
44. Sorock GS, Chen LH, Gonzalgo SR, Baker SP. Alcohol-drinking history and fatal injury in older adults. *Alcohol*. Nov; 2006 40(3):193–199. [PubMed: 17418699]
45. Smith C. Data Collection on Adverse Events of Anti-HIV drugs (D:A:D) Study Group. Factors associated with specific causes of death amongst HIV-positive individuals in the D:A:D Study. *AIDS (London, England)*. Jun 19; 2010 24(10):1537–1548.
46. Sherman KE, Rouster SD, Chung RT, Rajicic N. Hepatitis C Virus prevalence among patients infected with Human Immunodeficiency Virus: a cross-sectional analysis of the US adult AIDS Clinical Trials Group. *Clin Infect Dis*. Mar 15; 2002 34(6):831–837. [PubMed: 11833007]
47. Cheng DM, Nunes D, Libman H, et al. Impact of hepatitis C on HIV progression in adults with alcohol problems. *Alcohol Clin Exp Res*. May 1; 2007 31(5):829–836. [PubMed: 17403066]
48. Cooper CL, Cameron DW. Effect of alcohol use and highly active antiretroviral therapy on plasma levels of hepatitis C virus (HCV) in patients coinfecting with HIV and HCV. *Clin Infect Dis*. Jul 1; 2005 41(Suppl 1):S105–109. [PubMed: 16265607]
49. Fishbein DA, Lo Y, Netski D, Thomas DL, Klein RS. Predictors of hepatitis C virus RNA levels in a prospective cohort study of drug users. *Journal of acquired immune deficiency syndromes (1999)*. Apr 1; 2006 41(4):471–476. [PubMed: 16652056]
50. National Institute of Diabetes and Digestive and Kidney Diseases. [Accessed 1/3/2010] Chronic Hepatitis C: Current Disease Management. 2010. <http://digestive.niddk.nih.gov/ddiseases/pubs/chronichepc/>

51. Weis N, Lindhardt BO, Kronborg G, et al. Impact of hepatitis C virus coinfection on response to highly active antiretroviral therapy and outcome in HIV-infected individuals: a nationwide cohort study. *Clin Infect Dis*. May 15; 2006 42(10):1481–1487. [PubMed: 16619163]
52. Loko MA, Bani-Sadr F, Valantin MA, et al. Antiretroviral therapy and sustained virological response to HCV therapy are associated with slower liver fibrosis progression in HIV-HCV-coinfected patients: study from the ANRS CO 13 HEPAVIH cohort. *Antivir Ther*. 2012; 17(7): 1335–1343. [PubMed: 23052829]
53. Maier MM, He H, Schafer SD, Ward TT, Zaman A. Hepatitis C treatment eligibility among HIV-hepatitis C virus coinfecting patients in Oregon: a population-based sample. *AIDS Care*. 2014; 26(9):1178–1185. [PubMed: 24601687]
54. Fairley CK, Sze JK, Vodstrcil LA, Chen MY. Computer-assisted self interviewing in sexual health clinics. *Sex Transm Dis*. Nov; 2010 37(11):665–668. [PubMed: 20975481]

Author Manuscript Author Manuscript Author Manuscript Author Manuscript

Table 1

Clinical and demographic characteristics by alcohol use among persons living with HIV in clinical care at 7 CNICS sites across the U.S. in 2013–2015

N=8567

	Total N=8567	No alcohol use N=2804	Low alcohol use N=3493	Hazardous alcohol use N=2270	P value
	N	N	N	N	
Gender					
Male	7262	2211	3068	1983	27%
Female	1305	593	425	287	22%
					<0.001
Age (years)					
<30	793	145	313	335	42%
30–39	1768	427	763	578	33%
40–49	2565	854	1096	615	24%
50–59	2615	995	1018	602	23%
60	826	383	303	140	17%
					<0.001
Race/ethnicity					
White	4164	1257	1736	1171	28%
Black	2677	970	1102	605	23%
Hispanic	1268	436	466	366	29%
Other	458	141	189	128	28%
					<0.001
HIV transmission risk factor					
MSM	5392	1433	2392	1567	29%
IDU*	1082	500	374	208	19%
Heterosexual	1854	788	639	427	23%
Other	239	83	88	68	28%
					<0.001
CD4+ cell count (nadir) (N=8556)					
350	5033	1746	2073	1214	24%
351–500	1285	400	511	374	29%
500	2238	654	904	680	30%
					<0.001

	Total N=8567 N		100%		No alcohol use N=2804 N		33%		Low alcohol use N=3493 N		41%		Hazardous alcohol use N=2270 N		27%		P value
CD4+ cell count (current) (N=8556)																	
350	1638	19%			602	37%			641	39%			395	24%			
351–500	1382	16%			435	31%			579	42%			368	27%			
500	5536	65%			1763	32%			2268	41%			1505	27%			0.003
Currently receiving ART (N=8465)																	
No	742	9%			229	31%			286	39%			227	31%			
Yes	7723	91%			2534	33%			3169	41%			2020	26%			0.03
Current viral load (N=8539)																	
Detectable	1113	13%			326	29%			455	41%			332	30%			
Undetectable	7426	87%			2471	33%			3024	41%			1931	26%			0.007
Hepatitis C virus																	
No	7269	85%			2228	31%			3049	42%			1992	27%			
Yes	1298	15%			576	44%			444	34%			278	21%			<0.001
Depression symptoms (N=8548)																	
None	4718	55%			1521	32%			1973	42%			1224	26%			
Mild	1969	23%			624	32%			810	41%			535	27%			
Moderate	1517	18%			525	35%			578	38%			414	27%			
Severe	344	4%			125	36%			125	36%			94	27%			0.09

IDU: injection drug use; MSM: men who have sex with men; ART: antiretroviral therapy

* IDU includes patients who report being both MSM and IDU

Substance use by alcohol use among persons living with HIV in clinical care at 7 CNICS sites across the U.S. in 2013–2015 N=8567

Table II

	No alcohol use N=2804 N %	Lower risk alcohol use N=3493 N %	Hazardous alcohol use N=2270 N %	P value
Illicit drug use (including marijuana)				
None	996	987	349	15%
Past	1131	1154	658	22%
Current	677	1352	1263	38%
				<0.001
Illicit drug use* (excluding marijuana)				
None	1378	1778	848	21%
Past	1096	1200	907	28%
Current	330	515	515	38%
				<0.001
Methamphetamine/crystal use				
None	1803	2293	1347	29%
Past	746	827	682	30%
Current	255	373	241	28%
				<0.001
Cocaine/crack use				
None	1614	2049	1017	22%
Past	1107	1268	941	28%
Current	83	176	312	55%
				<0.001
Opiate/heroin use				
None	2233	2957	1809	26%
Past	513	455	383	28%
Current	58	81	78	36%
				<0.001
Marijuana use				
None	1224	1163	477	17%
Past	1086	1199	713	24%
Current	494	1131	1080	40%
				<0.001

	No alcohol use N=2804 N %	Lower risk alcohol use N=3493 N %	Hazardous alcohol use N=2270 N %	P value
Cigarette use				
Never	1148 35%	1469 45%	662 20%	<0.001
Past	740 31%	961 40%	707 29%	
Current	916 32%	1063 37%	901 31%	

Table III

Factors associated with at-risk alcohol use among persons living with HIV in clinical care at 7 CNICS sites across the U.S. in 2013–2015 in bivariate and multivariate analyses

	Bivariate	Multivariate				
	Hazardous vs. no/low* alcohol use OR: 95% CI, p value	Hazardous vs. no/low* alcohol use OR: 95% CI, p value	Hazardous vs. low risk** alcohol use OR: 95% CI, p value	Hazardous vs. no/ low*** alcohol use using higher cut-off OR: 95% CI, p value	Binge drinking vs. not binge drinking OR: 95% CI, p value	Hazardous vs. no/low* alcohol use with relative risks rather than odds ratios RR: 95% CI, p value
Sex						
Male	Ref	Ref	Ref	Ref	Ref	Ref
Female	0.75: 0.65–0.86, <0.001	1.03: 0.85–1.25, 0.8	1.16: 0.93–1.45, 0.2	0.94: 0.75–1.19, 0.6	0.77: 0.64–0.93, 0.007	0.96: 0.83–1.11, 0.6
Age						
<30	Ref	Ref	Ref	Ref	Ref	Ref
30–39	0.66: 0.56–0.79, <0.001	0.70: 0.58–0.84, <0.001	0.66: 0.54–0.81, <0.001	0.78: 0.63–0.96, 0.02	0.75: 0.62–0.90, 0.002	0.86: 0.78–0.94, 0.002
40–49	0.43: 0.36–0.51, <0.001	0.47: 0.39–0.57, <0.001	0.48: 0.39–0.59, <0.001	0.54: 0.44–0.66, <0.001	0.44: 0.37–0.53, <0.001	0.69: 0.62–0.77, <0.001
50–59	0.41: 0.35–0.48, <0.001	0.44: 0.37–0.53, <0.001	0.51: 0.42–0.63, <0.001	0.50: 0.41–0.62, <0.001	0.34: 0.28–0.41, <0.001	0.65: 0.58–0.72, <0.001
60	0.28: 0.22–0.35, <0.001	0.33: 0.26–0.42, <0.001	0.42: 0.32–0.55, <0.001	0.34: 0.25–0.46, <0.001	0.21: 0.16–0.26, <0.001	0.52: 0.44–0.62, <0.001
Race/ethnicity						
White	Ref	Ref	Ref	Ref	Ref	Ref
Black	0.75: 0.67–0.84, <0.001	0.74: 0.64–0.85, <0.001	0.77: 0.66–0.90, 0.001	0.86: 0.73–1.01, 0.07	0.81: 0.71–0.92, 0.002	0.85: 0.77–0.93, 0.001
Hispanic	1.04: 0.90–1.19, 0.6	1.05: 0.89–1.22, 0.6	1.13: 0.95–1.34, 0.2	1.03: 0.86–1.23, 0.7	1.24: 1.07–1.43, 0.005	1.03: 0.94–1.14, 0.5
Other	0.99: 0.80–1.23, 0.9	0.88: 0.70–1.12, 0.3	0.89: 0.69–1.15, 0.4	0.88: 0.67–1.15, 0.4	0.96: 0.77–1.20, 0.7	0.96: 0.84–1.10, 0.6
HIV transmission risk factor						
MSM	Ref	Ref	Ref	Ref	Ref	Ref
IDU*	0.58: 0.49–0.68, <0.001	0.56: 0.46–0.68, <0.001	0.69: 0.56–0.86, 0.001	0.65: 0.52–0.81, <0.001	0.58: 0.49–0.70, <0.001	0.68: 0.59–0.77, <0.001
Heterosexual	0.73: 0.65–0.83, <0.001	1.08: 0.90–1.29, 0.4	1.34: 1.10–1.63, 0.004	1.14: 0.93–1.39, 0.2	0.96: 0.81–1.13, 0.6	1.06: 0.93–1.20, 0.4
Other	0.97: 0.73–1.29, 0.8	1.06: 0.77–1.45, 0.7	1.15: 0.81–1.63, 0.4	0.81: 0.55–1.19, 0.3	1.16: 0.87–1.56, 0.3	1.05: 0.86–1.28, 0.6
CD4+ cell count nadir						

	Bivariate	Multivariate					
	Hazardous vs. no/low [*] alcohol use	Hazardous vs. no/low [*] alcohol use	Hazardous vs. low ^{**} risk alcohol use	Hazardous vs. no/ low ^{***} alcohol use using higher cut-off	Binge drinking vs. not binge drinking	Hazardous vs. no/low [*] alcohol use with relative risks rather than odds ratios	
	OR: 95% CI, p value	OR: 95% CI, p value	OR: 95% CI, p value	OR: 95% CI, p value	OR: 95% CI, p value	RR: 95% CI, p value	
350	Ref	Ref	Ref	Ref	Ref	Ref	
351–499	1.29: 1.13–1.48, <0.001	1.14: 0.98–1.32, 0.09	1.18 1.01–1.39, 0.04	1.13: 0.95–1.34, 0.2	1.11: 0.97–1.28, 0.1	1.08: 0.99–1.19, 0.09	
500	1.37: 1.23–1.53, <0.001	1.16: 0.99–1.34, 0.06	1.17: 0.99–1.38, 0.06	1.22: 1.03–1.45, 0.02	1.09: 0.95–1.26, 0.2	1.10: 1.00–1.21, 0.04	
Current viral load							
Detectable	Ref	Ref	Ref	Ref	Ref	Ref	
Undetectable	0.83: 0.72–0.95, 0.007	0.95: 0.81–1.10, 0.5	0.98: 0.83–1.15, 0.8	0.92: 0.77–1.09, 0.3	0.90: 0.78–1.04, 0.2	0.97: 0.88–1.07, 0.6	
Hepatitis C virus							
No	Ref	Ref	Ref	Ref	Ref	Ref	
Yes	0.72: 0.63–0.83, <0.001	0.83: 0.70–0.98, 0.03	0.97: 0.80–1.17, 0.7	0.75: 0.62–0.92, 0.004	0.88: 0.75–1.03, 0.1	0.84: 0.75–0.95, 0.004	
Depression symptoms							
None	Ref	Ref	Ref	Ref	Ref	Ref	
Mild	1.06: 0.95–1.20, 0.3	0.91: 0.80–1.04, 0.2	0.95: 0.82–1.09, 0.4	0.97: 0.84–1.12, 0.7	0.98: 0.87–1.22, 0.8	0.93: 0.86–1.01, 0.1	
Moderate	1.07: 0.94–1.22, 0.3	0.90: 0.78–1.03, 0.1	0.97: 0.83–1.14, 0.7	1.03: 0.88–1.21, 0.7	0.99: 0.87–1.13, 0.9	0.89: 0.81–0.97, 0.01	
Severe	1.07: 0.84–1.37, 0.6	0.88: 0.68–1.15, 0.4	0.98: 0.73–1.31, 0.9	1.06: 0.79–1.41, 0.7	0.86: 0.66–1.10, 0.2	0.85: 0.71–1.03, 0.09	
Methamphetamine/crystal use							
None	Ref	Ref	Ref	Ref	Ref	Ref	
Past	1.32: 1.18–1.47, <0.001	0.88: 0.75–1.02, 0.09	0.95: 0.80–1.11, 0.5	0.90: 0.76–1.07, 0.2	0.94: 0.81–1.09, 0.4	0.93: 0.85–1.02, 0.1	
Current	1.17: 0.99–1.37, 0.06	0.58: 0.47–0.71, <0.001	0.60: 0.48–0.75, <0.001	0.62: 0.50–0.78, <0.001	0.70: 0.58–0.85, <0.001	0.70: 0.61–0.80, <0.001	
Cocaine/crack use							
None	Ref	Ref	Ref	Ref	Ref	Ref	
Past	1.43: 1.29–1.58, <0.001	1.31: 1.14–1.51, <0.001	1.33: 1.14–1.55, <0.001	1.28: 1.08–1.50, 0.003	1.28: 1.12–1.46, <0.001	1.16: 1.05–1.28, 0.003	
Current	4.34: 3.63–5.19, <0.001	4.08: 3.29–5.05, <0.001	3.34: 2.65–4.21, <0.001	4.05: 3.25–5.06, <0.001	4.09: 3.29–5.08, <0.001	1.92: 1.72–2.14, <0.001	
Opiate/heroin use							
None	Ref	Ref	Ref	Ref	Ref	Ref	

Bivariate		Multivariate										
	Hazardous vs. no/low* alcohol use	OR: 95% CI, p value	Hazardous vs. no/low* alcohol use	OR: 95% CI, p value	Hazardous vs. low risk** alcohol use	OR: 95% CI, p value	Hazardous vs. no/ low*** alcohol use using higher cut-off	OR: 95% CI, p value	Binge drinking vs. not binge drinking	OR: 95% CI, p value	Hazardous vs. no/low* alcohol use with relative risks rather than odds ratios	RR: 95% CI, p value
Past		1.14; 1.00–1.29, 0.06		0.95; 0.81–1.11, 0.5		1.10; 0.92–1.31, 0.3		1.12; 0.94–1.33, 0.2		0.92; 0.80–1.07, 0.3		1.00; 0.91–1.09, 1.0
Current		1.73; 1.21–2.14, 0.001		1.05; 0.76–1.45, 0.8		1.09; 0.76–1.56, 0.6		1.14; 0.81–1.63, 0.4		0.97; 0.71–1.34, 0.9		1.12; 0.94–1.33, 0.2
Marijuana use												
None	Ref		Ref		Ref		Ref		Ref		Ref	
Past		1.56; 1.37–1.78, <0.001		1.37; 1.17–1.59, <0.001		1.21; 1.03–1.42, 0.02		1.23; 1.03–1.48, 0.02		1.22; 1.06–1.40, 0.005		1.28; 1.12–1.45, <0.001
Current		3.33; 2.94–3.77, <0.001		2.50; 2.15–2.91, <0.001		1.69; 1.44–1.99, <0.001		2.17; 1.82–2.58, <0.001		2.20; 1.91–2.53, <0.001		1.84; 1.63–2.09, <0.001
Cigarette use												
Never	Ref		Ref		Ref		Ref		Ref		Ref	
Past		1.64; 1.45–1.86, <0.001		1.33; 1.16–1.52, <0.001		1.37; 1.18–1.59, <0.001		1.39; 1.19–1.64, <0.001		1.40; 1.23–1.59, <0.001		1.18; 1.07–1.29, 0.001
Current		1.80; 1.60–2.02, <0.001		1.38; 1.20–1.58, <0.001		1.55; 1.34–1.80, <0.001		1.60; 1.37–1.87, <0.001		1.39; 1.23–1.58, <0.001		1.18; 1.07–1.29, 0.001

* No/low alcohol use includes no alcohol use and lower risk alcohol use defined as AUDIT-C scores of less than 4 for men and less than 3 for women

** Low risk alcohol use defined as AUDIT-C scores of less than 4 for men and less than 3 for women, excluding those with no alcohol use

*** No/low alcohol use using the higher cut-off includes no alcohol use and lower risk alcohol use defined as AUDIT-C scores of less than 5 for men and less than 4 for women

OR: odds ratio; 95% CI: 95% confidence interval; RR: relative risk

All models also adjusted for site

Table IV

Factors associated with hazardous alcohol use compared with no/low alcohol use among persons living with HIV in clinical care at 7 CNICS sites across the U.S. in 2013–2015 in multivariate analyses stratified by sex

	Multivariate OR Females	Multivariate OR Males
Age		
<30	1 Ref	1 Ref
30–39	0.74: 0.40–1.38, 0.3	0.68: 0.56–0.82, <0.001
40–49	0.66: 0.36–1.23, 0.2	0.44: 0.36–0.53, <0.001
50–59	0.65: 0.34–1.21, 0.2	0.41: 0.34–0.50, <0.001
60	0.34: 0.16–0.74, 0.006	0.31: 0.24–0.41, <0.001
Race/ethnicity		
White	1 Ref	1 Ref
Black	0.94: 0.65–1.35, 0.7	0.71: 0.61–0.83, <0.001
Hispanic	1.11: 0.64–1.94, 0.7	1.05: 0.89–1.24, 0.5
Other	0.40: 0.17–0.97, 0.04	0.96: 0.75–1.23, 0.8
CD4+ cell count (nadir)		
350	1 Ref	1 Ref
351–500	1.50: 0.99–2.25, 0.05	1.09: 0.93–1.27, 0.3
>500	1.23: 0.82–1.87, 0.3	1.14: 0.96–1.34, 0.1
Current viral load		
Detectable	1 Ref	1 Ref
Undetectable	0.84: 0.57–1.24, 0.4	0.97: 0.82–1.14, 0.7
Hepatitis C virus		
No	1 Ref	1 Ref
Yes	0.81: 0.53–1.25, 0.3	0.69: 0.58–0.82, <0.001
Depression symptoms		
None	1 Ref	1 Ref
Mild	0.96: 0.67–1.37, 0.8	0.90: 0.79–1.03, 0.1
Moderate	1.35: 0.94–1.94, 0.1	0.83: 0.71–0.97, 0.02
Severe	1.71: 0.93–3.13, 0.09	0.76: 0.57–1.03, 0.08
Methamphetamine/crystal use		
None	1 Ref	
Past	0.58: 0.35–0.97, 0.04	0.88: 0.75–1.04, 0.1
Current	1.09: 0.51–2.33, 0.8	0.53: 0.42–0.65, <0.001
Cocaine/crack use		
None	1 Ref	1 Ref
Past	1.55: 1.03–2.34, 0.04	1.26: 1.08–1.47, 0.003
Current	3.66: 2.10–6.37, <0.001	4.11: 3.25–5.19, <0.001

	Multivariate OR Females	Multivariate OR Males
Opiate/heroin use		
None	1 Ref	1 Ref
Past	0.88; 0.55–1.41, 0.6	0.87; 0.74–1.02, 0.1
Current	0.65; 0.27–1.53, 0.3	0.99; 0.70–1.41, 1.0
Marijuana use		
None	1 Ref	1 Ref
Past	1.60; 1.07–2.40, 0.02	1.30; 1.10–1.53, 0.002
Current	2.79; 1.85–4.21, <0.001	2.47; 2.11–2.91, <0.001
Cigarette use		
Never	1 Ref	1 Ref
Past	2.09; 1.40–3.12, <0.001	1.25; 1.08–1.44, 0.003
Current	1.10; 0.74–1.63, 0.7	1.38; 1.19–1.59, <0.001