Published in final edited form as:

Biochemistry. 2017 October 10; 56(40): 5229–5244. doi:10.1021/acs.biochem.7b00771.

Radical SAM enzymes involved in RiPP biosynthesis

Nilkamal Mahanta^{†,‡}, Graham A. Hudson[†], and Douglas A. Mitchell^{†,‡,*}

[†]Department of Chemistry, University of Illinois at Urbana-Champaign, 600 South Mathews Avenue, Urbana, Illinois 61801, USA

[‡]Carl R. Woese Institute for Genomic Biology, University of Illinois at Urbana-Champaign, 1206 West Gregory Drive, Urbana, Illinois 61801, USA

Abstract

Ribosomally synthesized and post-translationally modified peptides (RiPPs) display a diverse range of structures and continue to expand as a natural product class. Accordingly, RiPPs exhibit a wide array of bioactivities, such as broad and narrow spectrum growth suppressors, antidiabetics, as well as antinociception and anticancer agents. Owing to these properties, and the complex repertoire of post-translational modifications (PTMs) that give rise to these molecules, RiPP biosynthesis has been intensely studied. RiPP biosynthesis often involves enzymes that perform unique chemistry with intriguing reaction mechanisms, which attract chemists and biochemists alike to study and re-engineer these pathways. One particular type of RiPP biosynthetic enzyme is the so-called radical S-adenosylmethionine (rSAM) enzyme, which utilize radical-based chemistry to install several distinct PTMs. Here, we describe the rSAM enzymes characterized over the past decade that catalyze six reactions types from several RiPP biosynthetic pathways. We present the current state of mechanistic understanding and conclude with possible directions for future characterization of this enzyme family.

Graphical Abstract

Introduction

Ribosomally synthesized and post-translationally modified peptides (RiPPs) are a functionally and structurally diverse family of natural product that are divided into several

*Corresponding Author: douglasm@illinois.edu.

ORCID

Nilkamal Mahanta: 0000-0001-8901-2531

Graham A Hudson: 0000-0002-3715-4279

Douglas A. Mitchell: 0000-0002-9564-0953

subclasses based on common structural motifs installed by conserved biosynthetic enzymes.¹ Recent genome sequencing efforts have revealed that RiPPs form a large natural product class produced by all domains of life.^{2,3} In general, RiPP biosynthesis commences with the post-translational modification (PTM) of a precursor peptide, typically consisting of *N*-terminal leader and *C*-terminal core regions.¹⁻⁴ The RiPP precursor peptide gene is often encoded adjacent to other genes responsible for regulation, biosynthesis, and export. Collectively, these genes comprise the biosynthetic gene cluster (BGC) for the natural product. Recognition sequences in the leader peptide are bound by key biosynthetic proteins with many relying on a conserved structural domain, referred to as the RiPP precursor peptide Recognition Element (RRE) while others bind the peptide substrate using alternative strategies.⁵ Upon substrate engagement, the RiPP biosynthetic enzymes carry out modifications to the core region. The leader peptide is subsequently cleaved from the modified core, frequently by a peptidase encoded within the BGC.^{1,2} Other tailoring enzymes⁶, if present, can further modify the core peptide in a leader peptide-independent fashion. Once all of the PTMs are installed, the mature RiPP is exported, sometimes by a locally encoded ATP-binding cassette (ABC) transporter. Since RiPP precursor peptides are direct gene products (i.e. ribosomally synthesized), RiPP biosynthetic pathways tend to exhibit significant malleability for biosynthetic re-engineering and hold potential for numerous biotechnological applications.⁷

Radical S-adenosylmethionine (rSAM) enzymes are known to catalyze chemically difficult transformations involved in both primary metabolism and natural product biosynthesis.^{8,9} rSAM enzymes invariably contain an active site [4Fe-4S] cluster, coordinated by a canonical cysteine-rich CX₃CX₂C motif that transfers an electron to a bound SAM cofactor. This transfer usually leads to the reductive cleavage of the 5'-C-S bond, generating a highly reactive 5'-deoxyadenosyl (5'-DA) radical and methionine. The 5'-DA radical then abstracts a hydrogen atom from the substrate to form 5'-deoxyadenosine and a substrate radical which undergoes further reaction.⁸ In some cases, the 5'-DA radical first forms a protein active site radical (i.e. glycyl or thiyl radicals), which subsequently generates the substrate radical.⁸ In the last decade, a plethora of studies have been reported on rSAM enzymes that carry out fascinating transformations in natural product biosynthesis. In this review, we discuss developments in the discovery and characterization of rSAM enzymes that are involved in the biosynthesis of several classes of RiPPs. We have not attempted to be comprehensive. Instead, we provide representative examples from several RiPP classes that underscore the diversity of the chemistry along with the current mechanistic understanding of the enzymology driving these reactions.

RiPP biosynthetic reactions catalyzed by rSAM enzymes

For organizational purposes, we have grouped rSAM enzymes involved in RiPP biosynthesis into the following six reaction types: (A) *C*-methylation, (B) thioether (Cα-S) bond formation, (C) C-C bond formation, (D) epimerization, (E) decarboxylation/C-C bond formation, and (F) rearrangements. Each group features rSAMs with unique domain architectures that give rise to the mechanistic diversity observed within this catalytically gifted family of enzymes (Figure 1). A few illustrative examples of each reaction type are discussed.

A) C-methylation reactions

Methylation is a widespread chemical modification that often plays a pivotal role in major biological processes. Nucleophilic positions in biomolecules are usually methylated via a SAM-dependent S_N2 reaction.¹² However, recent studies have also revealed rSAM-dependent methylation of non-nucleophilic centers in natural product biosynthesis.^{13–15} At present, there are four classes of rSAM methyltransferases (MTs). Class A includes RNA base-modifying enzymes that use active site cysteines (non-cluster binding cysteines, such as Cys118 and Cys355 in RlmN) during methylation. Class B is characterized by the presence of an *N*-terminal cobalamin-binding domain and a *C*-terminal rSAM domain. In contrast, class C are cobalamin-independent and use two active site SAM molecules to achieve methyl transfer while class D use methylenetetrahydrofolate as the methyl donor.^{8,15} However, we note that with the exception of class B, which transfers a complete methyl group from SAM, the other three classes actually transfer a methylene group and hence might be called methylases rather than formal methyltransferases. Below, we highlight a few examples of rSAM enzymes that methylate unactivated sp^2 or sp^3 carbons in RiPP biosynthetic pathways.¹⁵

A.1. TsrM in thiostrepton A biosynthesis—The thiopeptide antibiotic thiostrepton A (**3**, Figure 2) produced by several *Streptomyces sp.* (such as *S. azureus*, *S. laurentii*, etc.) targets the 50S ribosome and has growth-suppressive activity against Gram-positive bacteria.^{16–19} Thiostrepton biosynthesis involves a complex array of PTMs on a ribosomal precursor peptide.^{18,19} One of these modifications appends an L-tryptophan (**1**)-derived quinaldic acid moiety to the main thiostrepton scaffold. It has been proposed that TsrM, a class B rSAM MT, catalyzes the first step of this unusual ring expansion by methylation of **1** at C-2 to form 2-methyltryptophan (**2**).¹⁹ This reaction proceeds with net retention of stereochemistry¹⁹ and with concomitant production of S-adenosylhomocysteine (SAH, **5**), a SAM (**4**) byproduct.^{20,21} Labeling studies have shown the methyl group of SAM undergoes transfer to **1**. However, no net generation of 5'-deoxyadenosine (**10**) was observed, suggesting that reductive cleavage of SAM was not required. This conclusion is also supported by the observation that a TsrM variant lacking the cysteines that coordinate the [4Fe-4S] cluster can still catalyze the conversion of **4** to **5**. However, this variant was unable to catalyze the methyl transfer to **1**, suggesting the [4Fe-4S] cluster plays a critical role in this step.^{20,21}

Mössbauer and hyperfine sublevel correlation (HYSCORE) spectroscopy studies have confirmed the presence of the [4Fe-4S] cluster and revealed that **4** and **1** do not directly coordinate to the cluster.²² UV-Vis spectroscopy studies detected a cobalamin (II) intermediate¹⁹ and electron paramagnetic resonance (EPR) experiments have indicated the presence of a five-coordinate Co(II) species, which changes dramatically upon addition of **4**.²² The native cofactor for TsrM has recently been identified as methylcob(III)alamin (MeCbl).²¹ Based on these studies²¹, it has been proposed that Co-C bond MeCbl undergoes homolytic cleavage to generate a putative methyl radical, which attacks the sp^2 C-2 of **1** (Figure 2C). The resulting Trp-based radical loses an electron back to the [4Fe-4S] cluster and undergoes deprotonation to yield **2**. The [4Fe-4S] cluster can then reduce the cobalamin (II) intermediate, forming the supernucleophile cobalamin (I), which would then attack the

electrophilic methyl group of **4** to generate MeCbl and **5**. This proposal is consistent with known aspects of how methionine synthase and corrinoid/iron-sulfur protein function.^{21,22} However, a very recent study has cast doubt on a radical-based mechanism for TsrM. Using analogs of **1**, it was revealed that the N1-amine group is critical for the reaction rate and suggests that the methyl transfer from methylcobalamin to **1** may follow S_N2 chemistry.²³ The postulated mechanism that involves nucleophilic attack from C-2 of **1** onto the methyl group of methyl cob(III)alamin would rule out a radical-based mechanism involving the homolytic cleavage of the Co-C bond. Homolytic cleavage would generate cob(II)alamin as an intermediate, which was not detected.²³ Further characterization of TsrM and its homologs is in order to resolve these mechanistic ambiguities.

A.2. PoyC in polytheonamide biosynthesis—Polytheonamides (**6**, Figure 3) are cytotoxic, pore-forming RiPPs that undergo extensive modification. Polytheonamides were isolated from a microbial symbiont (*Candidatus entothionella*) of *Theonella swinhoei*, a marine sponge.^{24,25} Out of the total 48 PTMs, 17 are C-methylations on *sp*³ carbon centers (13 are located on C_β, 3 are on C_γ, and 1 is on C_δ) that are proposed to be formed by the cobalamin-dependent class B rSAM MTs, PoyB and PoyC (Figure 4). Also, 18 of the PTMs are epimerizations which are proposed to be catalyzed by the rSAM epimerase, PoyD (discussed later in this review).

Using a rhizobial heterologous host that naturally produces cobalamin, PoyA (precursor peptide) was co-expressed with PoyB and/or PoyC. From these co-expression studies, 13 of the 17 expected C-methylations on the PoyA peptide were detected and localized.²⁶ It was gleaned that the C-methyltransferase activity was dependent on the PoyA core region with PoyC acting on residues 1–21 and PoyB on residues 23–49 (Figure 3).²⁶ In vitro experiments using a synthetic PoyA₁₋₁₅ peptide led to the formation of **8** (Figure 4) that contained only one methylation at L-Val₁₄. **5** and **10** were detected as byproducts.²⁷ It was also shown that the peptide containing D-Val at position 14 was not modified, suggesting that C-methylation likely occurs prior to epimerization at that position.²⁷ When the Cys in the CX₇CX₂C motif in PoyC were substituted with Ala, the variant was unable to produce **5** or **10** in addition to being incapable of transferring a methyl group to the substrate. This observation suggests different functions for the [4Fe-4S] cluster in two class B rSAM MTs, PoyC and TsrM.^{21,22} It was observed that purified PoyC was loaded with MeCbl and adenosylcobalamin; however, the triple Cys variant lacked MeCbl, suggesting a critical role of the [4Fe-4S] cluster in cobalamin methylation. When the reaction was performed with CD₃-SAM, MeCbl was converted to CD₃-Cbl, establishing that the enzyme-bound cobalamin is recycled from SAM during catalysis (Figure 4C).²⁷ Such class B rSAM MTs are also involved in bottromycin biosynthetic pathway; however, in vitro characterization has not yet been reported for those enzymes. While these are interesting targets for future enzymatic characterization, they are not considered further for this review.^{1,28,29}

Based on the available data it was proposed that, unlike TsrM, PoyC catalyzes the reductive cleavage of **4** (SAM 1) to form the 5'-DA radical (**9**, also referred to as "dAdo radical"), which then abstracts the C_β-H atom of Val₁₄ (**7**) forming a carbon-centered radical intermediate (**11**).²⁷ It remains unclear how methyl transfer to form **8** takes place, but

potentially, **11** could undergo a radical recombination reaction with the methyl radical generated from the homolytic cleavage of the Co-C bond. Cobalamin (II), after reduction to cobalamin (I), could react with **4** (SAM 2) to regenerate MeCbl. Based on TsrM and PoyB/C, it appears that the hybridization of the target carbon atom (sp^2 vs. sp^3) dictates whether class B rSAM enzymes require the reductive cleavage of SAM for catalysis.

A.3. NosN in nosiheptide biosynthesis—Nosiheptide (**14**, Figure 5) is a thiopeptide antibiotic produced by *Streptomyces actuosus* that, similar to **3** (Figure 2B), contains a secondary macrocycle derived from **1** with a similar antibacterial mode of action.³⁰ NosN belongs to the class C rSAM MTs¹³ and is responsible for appending a methyl group onto the C-4 of an indole moiety during the maturation of **14**.³⁰

In vitro studies on NosN using a thioester analog of 3-methyl-2-indolic acid (**12**, Figure 5) demonstrated that it catalyzes the transfer of the methyl group onto the C-4 of the indolyl moiety of **12** to form **13**.³¹ It was observed that two hydrogen atoms from the methyl group of **4** were transferred to the substrate³¹, similar to that observed for class A rSAM MTs.¹⁵ However, upon substituting the active site cysteines Cys118 and Cys355 of RlmN, a class A enzyme, with alanine, catalysis was abolished. Importantly, Cys118 and Cys355 of RlmN are not involved in [4Fe-4S] cluster binding. In contrast, substitution of non-cluster binding cysteines of NosN with alanine did not abolish catalysis (hence its designation as class C). It has also been reported that NosN produces methylthioadenosine (MTA, **15**, Figure 5C) from SAM. MTA is proposed to be the direct methyl donor; accordingly, 5'-thioadenosine (**19**) is produced as a byproduct instead of SAH (**5**).³¹

Based on the proposed mechanism,³¹ the first SAM molecule (SAM 1) is reductively cleaved to generate **9** (Figure 5C). Subsequently, **9** abstracts a hydrogen from the methyl group of **15** (generated from SAM 2) and the resulting methylene thioadenosyl radical (**16**) adds to the C-4 of the indolyl substrate **12** to form intermediate **17**. Radical **17** is proposed to be deprotonated to produce the radical anion **18** which would undergo C-S bond cleavage to eliminate the thioadenosine anion (**19**) and generate the methylene radical (**20**). An electron transfer to **20**, likely from the [4Fe-4S] cluster, followed by protonation would yield the methylated indole product (**13**). Density functional theory (DFT) calculations support a heterolytic C-S bond cleavage to release **19** and **20**. NosN assays using SAM homologs (i.e. *S*-guanosylmethionine and *S*-cytidinylmethionine) have provided indirect support for this mechanistic proposal.³² The work on NosN presents a new paradigm for cobalamin-independent rSAM MTs. Whether this proposed mechanism holds for other class C rSAM MTs remains an area of active investigation.

A.4. TbtI in thiomuracin biosynthesis—Thiomurcin (**24**, Figure 6) is also a thiopeptide, but unlike **3** and **14**, its antibiotic activity derives from binding to elongation factor thermo unstable (EF-Tu) involved in ribosomal peptide synthesis.³³ Through in vitro reconstitution experiments using genes from *Thermobispora bispora*, it was revealed that the precursor peptide, TbtA, undergoes an array of PTMs including: six ATP-dependent cyclodehydrations (TbtF/G), six FMN-dependent dehydrogenations (TbtE), four tRNA-dependent glutamylations with subsequent elimination (TbtB/C), and finally, a formal [4+2]

cycloaddition (TbtD) to form a trithiazole-substituted pyridine with concomitant ejection of the leader peptide as a carboxamide to yield thiomuracin GZ, an in vitro biosynthetic product. Compared to thiomuracin A1, the GZ variant lacks additional peripheral tailoring modifications including epoxidation, β -hydroxylation of Phe5, C-terminal trimming/amidation, and C-methylation of one of six thiazole moieties.^{34,35} The latter is the relevant modification for this review. Recently, the activity of the responsible enzyme, a class C rSAM MT (TbtI), was reconstituted in vitro.³⁶ This report demonstrated that a linear intermediate bearing six thiazoles was a substrate for TbtI, while the macrocyclized peptide was not, suggesting that C-methylation occurs prior to the [4+2] cycloaddition reaction (Figure 6B). As a class C rSAM MT, TbtI shows homology to HemN, involved in heme biosynthesis,¹³ and acts in a leader peptide-independent manner. This observation was also supported by that fact that it does not have a bioinformatically identifiable RRE. It was also demonstrated that TbtI requires a preceding Asn (residue 3 of the core peptide) and at least one downstream thiazole to regiospecifically install the methyl group at thiazole 4 (Thz4).³⁶ Homologous genes (*pbtM2* and *pbtM3*) are encoded by the *Planobispora rosea* BGC that produces GE2270A (**23**), a structurally related thiopeptide.³⁷ However, the in vitro activity and the substrate scope for these enzymes have not yet been reported. Future work will elucidate the enzymatic mechanism of TbtI to determine if the observations noted in the NosN mechanism are applicable to other class C rSAM MTs.

B) Thioether bond (S to C α) formation reaction

Sulfur-to- α carbon thioether cross-linked peptides (sactipeptides)³⁸ are RiPPs that show antibacterial, spermicidal, and hemolytic properties.³⁹ Sactipeptides are characterized by a rSAM-catalyzed, intramolecular thioether bond that crosslinks the sulfur atom of Cys to the α -carbon of an acceptor amino acid. This “sactionine” linkage is distinct from the C-S bond found in lanthipeptides, which link Cys residues to the β -carbon of dehydrated Ser/Thr residues via a Michael-like nucleophilic addition reaction (Figure 7E).⁴⁰ Five sactipeptides have been reported³⁹ including: subtilisin A, **27** from *Bacillus subtilis* 168;⁴¹ thurincin H, **28** from *B. thuringiensis* SF361;⁴² sporulation killing factor (SKF, **29**) from various *B. subtilis*;⁴³ and the two-component thuricin CD (Trn- α , **30** and Trn- β , **31**) from *Bacillus thuringiensis* DPC 6431.^{44,45} Herein, we discuss recent advances in our understanding of rSAM-mediated thioether bond formation during sactipeptide biosynthesis. While there are reports of rSAM enzymes that install thioether linkages within protein substrates (e.g. QhpD acting on QhpC),⁴⁶ these are not considered RiPPs; thus, they will not be mentioned further.

B.1. AlbA in subtilisin A biosynthesis—Subtilisin A (**27**) is 35-residue, head-to-tail macrocyclized peptide encoded by the *sbo-alb* locus of *B. subtilis* 168 that displays antibiotic activity (Figure 7).⁴⁷ The overall conformation of **27** is constrained by three C-S bonds between Cys4, Cys7, and Cys13 and the α -carbons of Phe31, Thr28, and Phe22, respectively. The *sbo-alb* locus consists of the precursor gene *sboA* and processing genes *albA-G*, which are responsible for the PTMs as well as for the export of **27**.

In vitro studies on the rSAM enzyme AlbA demonstrated that it contains two [4Fe-4S] clusters and catalyzes sactionine formation on the linear precursor peptide SboA (**25** to **26**). This reaction was found to be leader peptide-dependent, which was also supported by the

presence of the RRE in AlbA.⁴⁷ Thioether formation is proposed to be followed by AlbE/F (Zn-dependent protease)-catalyzed leader peptide cleavage and macrocyclization.⁴⁷ Using biochemical, mutational, and biophysical techniques, it was elucidated that both the [4Fe-4S] clusters are essential for AlbA activity. The first [4Fe-4S] cluster is within the rSAM domain and is bound by Cys129, Cys133, and Cys136. This cluster binds **4** and carries out reductive cleavage to form **9**. The second [4Fe-4S] cluster is located within the “SPASM” domain (subtilisin A/pyrroloquinoline quinone/anaerobic sulfatase/mycofactacin maturation enzymes)¹¹ and is bound by Cys408, Cys414, and Cys417. Although not required for **9** formation, the [4Fe-4S] cluster of the SPASM domain is believed to bind and orient the substrate, SboA, via its cysteine residues for thioether formation (Figure 7D).⁴⁷ Deprotonation forms **25** and subsequently **9**, coordinated by the rSAM [4Fe-4S] cluster, then abstracts hydrogen from the α -carbon of the acceptor amino acid (Thr or Phe) of **25** forming a substrate radical (**33**). This radical can then attack the coordinated SboA sulfur atom by forming the sactonine bridge in **26** with the SPASM cluster accepting one electron. The reduced form of the SPASM [4Fe-4S] cluster may transfer an electron to the rSAM cluster, converting both into their active forms to prime the enzyme for a second turnover.⁴⁷ A recent study reported that AlbA contains three [4Fe-4S] clusters, two of which are located in the SPASM domain. It has been proposed that both SPASM clusters could be involved in electron transfer although one may simply play a structural role.⁴⁸ As with AlbA, additional rSAM enzymes previously reported to contain two [4Fe-4S] clusters might actually contain a total of three (two auxiliary clusters being bound by the SPASM domain). However, there are known “SPASM” domains that only contain one [4Fe-4S] cluster, which have been dubbed “twitch” domains.¹¹ Additional studies are required clarify the roles of these auxiliary clusters in AlbA and other SPASM-domain containing rSAM proteins.

B.2. SkfB in sporulation killing factor biosynthesis—The sporulation killing factor (**29**, Figure 7) is a 26-residue RiPP that plays a critical role during the sporulation of certain *Bacillus* strains.^{43,49} **29** contains a sactonine linkage between Cys4 and Met12, a disulfide between Cys1 and Cys16, and a head-to-tail cyclic architecture. The *skf* gene cluster is responsible for **29** biosynthesis and is comprised of a precursor gene, *skfA*, tailoring genes *skfB* (rSAM enzyme), *skfC* (putative protease), *skfH* (putative thioredoxin), and *skfE/skfF* for export/immunity. The function of *skfG* remains unknown.⁴³

SkfB is a rSAM enzyme that contains two [4Fe-4S] clusters and is responsible for sactonine formation.⁵⁰ From mutational and biochemical studies, it was evident that the positions of the donor and acceptor amino acids play important roles during the SkfB-catalyzed reaction, as they were not interchangeable. Although SkfB was specific towards the donor Cys4 site, it showed broader tolerance towards the acceptor site, suggesting that radical generation at the α -carbon is independent of the acceptor amino acid. It was observed for the acceptor Met12 site that hydrophobic amino acids were tolerated most efficiently. Hydrophilic amino acids were either moderately tolerated or not at all. Like AlbA, SkfB also showed leader peptide dependence. Both the rSAM (coordinated by Cys117, Cys121, and Cys124) and SPASM [4Fe-4S] (coordinated by Cys380, Cys385, and Cys387) clusters were found to be catalytically essential and the formation of **9** was detected.⁵⁰ These data, along with a separate study that demonstrated that SkfB abstracts hydrogen from the α -carbon of Met12

on SkfA,⁵¹ have permitted a mechanistic proposal to be put forth that is akin to AlbA. It remains to be determined if SkfB harbors a divergent RRE domain like other sactonine-forming rSAM enzymes (Figure 1). Also, while SkfB is reported to contain one SPASM-bound [4Fe-4S] cluster (which would classify it as a twitch domain), multiple sequence alignment shows that there are cysteines in SkfB that align with AlbA residues that bind the second auxiliary cluster. This suggests that SkfB may contain a total of three [Fe-S] clusters.

B.3. ThnB in thurincin H biosynthesis and TrnC/D in thuricin CD biosynthesis

—Thurincin H (**28**, Figure 7) is a 31-residue antibacterial RiPP biosynthesized by the *thn* gene cluster of *B. thuringiensis* SF361.⁴² Detailed structural characterization of thurincin H revealed that it contains four sactonine linkages between Cys4, Cys7, Cys10, and Cys13 to the α -carbons of Ser28, Thr25, Thr22, and Asn19, respectively.⁴² The BGC is comprised of three identical copies of the 40-residue precursor peptide *thnA*, *thnB* (rSAM enzyme), *thnP* (protease), *thnD/E/T* (three-component ABC transporter), *thnR* (transcriptional regulator), and *thnI* (unknown function). Studies using purified ThnB demonstrated that it was responsible for sactonine formation.⁵² Like SkfB, ThnB was shown to contain two catalytically essential [4Fe-4S] clusters, the second contained within the SPASM domain, although a third, undiscovered [4Fe-4S] cluster may be present.⁵³ Mutational studies revealed that the rSAM [4Fe-4S] cluster is coordinated by Cys165, Cys169 and Cys172 and is required for **9** formation while a SPASM domain cluster is putatively coordinated by Cys449, Cys455 and Cys458. A variant of ThnB which lacked the *N*-terminal RRE domain was incapable of thioether bond formation but still capable of **9** formation, suggesting the RRE participates in mediating the interaction with ThnA.^{5,52} The proposed mechanism for ThnB is similar to that proposed for AlbA and SkfB (Figure 7D).

Thuricin CD (Figure 7) is a two-component RiPP produced by *Bacillus thuringiensis* DPC 6431.⁴⁴ Two distinct peptides, Trn α , **30** and Trn β , **31**, act synergistically to kill *Clostridium difficile*. A recent mode of action study has shown that thuricin CD inserts into the membrane of the target cell, forms pores, and causes ion leakage, membrane depolarization, and finally, cell lysis.⁵⁴ Structural elucidation studies of thuricin CD have shown that each of the two components contains three sactonines. The linkages in Trn α are from Cys5, Cys9, and Cys13 to Thr28, Thr25, and Ser21, respectively. The sactonines in Trn β are from Cys5, Cys9, and Cys13 to Tyr28, Ala25, and Thr21, respectively.⁴⁵ In addition to the precursor peptides, the thurincin CD BGC encodes five proteins: TrnC/D (rSAM enzymes), TrnE (unknown), TrnF/G (ABC transporter).⁴⁴ Future studies will clarify the roles of these proteins in thuricin CD biosynthesis.

B.4. Tte1186 and CteB in SCIFF peptide maturation

—A bioinformatics study identified a gene encoding a putative rSAM enzyme adjacent to a gene encoding for a peptide in several clostridial genomes.⁵⁵ It was proposed that the rSAM enzyme might modify the peptide to serve a housekeeping role based on the conservation of local genes. Sequence analysis of the peptides showed that nearly all contained variable *N*-terminal and conserved *C*-terminal regions distinguished by the presence of six cysteines in a forty-five-residue (SCIFF) peptide. Tte1186 from *Caldanaerobacter subterraneus* subsp. *tengcongensis* MB4 represents the first characterized SCIFF-related rSAM enzyme, which catalyzes

thioether formation on the associated peptide, Tte1186a (Figure 7).³⁸ This SPASM domain-containing rSAM enzyme contains three essential [4Fe-4S] clusters. Biochemical studies revealed that the rSAM cluster, coordinated by Cys104, Cys108, and Cys111, was responsible for **9** formation. The two auxiliary SPASM clusters are coordinated by Cys344, Cys362, and Cys413 for one auxiliary cluster and Cys400, Cys403, and Cys432 for the second auxiliary cluster. Although multiple cysteines were present in the SCIFF precursor peptide, only one thioether crosslink between Cys32 and Thr37 (**32a**) was formed by Tte1186 during in vitro reconstitution experiments.³⁸

A new mechanism was postulated for Tte1186 (Figure 8).³⁸ Instead of direct coupling between the α -carbon-centered radical and the thiol of **35**, the former could transform into a resonance-stabilized ketyl radical (**36**) with subsequent oxidation to a ketoimine intermediate (**37**). This planar intermediate could then be trapped by the thiol of the respective cysteine residue of the substrate on either face, leading to stereoisomeric *re* and *si* adducts (**38** or **39**). Such dual stereochemical outcomes have been observed in other sactipeptides (e.g. subtilisin A)⁴⁷ and the authors propose this mechanism better explains this phenomenon than one that involves direct thiol attack of the α -carbon-centered radical (Figure 7).³⁸ Additional studies will be required to delineate the mechanistic details and the role of auxiliary clusters in rSAM enzymes that contain SPASM domains.

A recent report describes yet another SPASM domain-containing rSAM enzyme, CteB from *Clostridium thermocellum* ATCC 27405 (Figure 7A). CteB installs a thioether linkage on CteA, a putative SCIFF peptide (**32b**), and contain three [4Fe-4S] clusters.⁵⁶ This report marks the first crystal structure of a thioether bond-forming enzyme that has both **4** and an *N*-terminal fragment of CteA bound at the active site. CteB has a TIM barrel fold that is characteristic of rSAM enzymes and represents the SAM-activating domain, as well as a *C*-terminal SPASM domain that contains two auxiliary [4Fe-4S] clusters. Based on sequence alignments of CteB with other thioether-forming rSAMs, it was postulated that there are variations on the number of the auxiliary clusters in SPASM domains (i.e. twitch domains are annotated as SPASM domains but contain only a single cluster). As mentioned above, additional studies are required on previously characterized enzymes to confirm the number and the roles of these additional [4Fe-4S] clusters. With CteB, it was also found that one [4Fe-4S] cluster in the SPASM domain exhibited an open coordination site in the absence of the peptide substrate. In the presence of substrate, the previously open coordination site was occupied by a cysteine of the peptide substrate.⁵⁶ This structure also revealed that an *N*-terminal RRE domain⁵ that aided substrate binding.⁵⁶ Collectively, these studies provided significant insight into the mechanism of thioether formation and has opened up the area for further investigations.

C) Epimerization reactions

Radical SAM epimerases are responsible for the regioselective introduction of D-amino acids into RiPPs. Herein, we discuss two rSAM epimerases that have been described recently in RiPP biosynthetic pathways.

C.1. PoyD in polytheonamide biosynthesis—The pore-forming polytheonamides (**6**, Figure 3) are the best-known proteusins.^{24,25} Among other structural features, **6** possesses 18 D-configured residues positioned in near-perfect alternation with L-amino acids.²⁴ PoyD is the responsible rSAM epimerase which was shown to install numerous D-stereocenters (7/10 D-Asn and 1/5 D-Val) in PoyA, as determined from in vivo co-expression studies.²⁴ The majority of the D-Asn reside on the C-terminal half of the polytheonamide peptide, while most D-Val are found in the N-terminal region. In a subsequent study, using in vivo co-expression with PoyA variants, it was established that PoyD actually installs all 18 of the D-amino acids appearing in PoyA.²⁶ Another study reported the existence of a larger rSAM enzyme subfamily catalyzing peptide epimerization, which are architecturally distinct and occur in various bacteria.⁵⁷ Three PoyD homologs from cyanobacteria, namely PlpD (*Pleurocapsa* sp. PCC 7319), AvpD (*Anabaena variabilis* ATCC 29413) and OspD (*Oscillatoria* sp. PCC 6506) were shown to convert their precursor peptides to a single product containing D-amino acids. After co-expression of the rSAM epimerases with the cognate precursor peptides, the epimerized residues were localized, revealing that the enzymes exhibited high site-selectivity and tended to epimerize Val, Ile, Ala, and Met residues within the substrate.⁵⁷ The authors proposed that the epimerization is initiated by hydrogen abstraction by **9** to generate a resonance-stabilized substrate radical, which could then obtain hydrogen from an external source at the opposite face to complete the epimerization. Labeling studies showed that deuterium was lost after epimerization when a precursor peptide containing deuterated amino acids was used, consistent with the proposed mechanism. This also indicated that **9** would not be recycled during the catalytic cycle and one molecule of **4** will be required per epimerization.⁵⁷ Further studies are need to further clarify the mechanistic details.

C.2. YydG in epipeptide biosynthesis—Epipeptide (**40**, Figure 9) is a recently discovered class of RiPP from *B. subtilis* which also features epimerization. While the molecular target of epipeptide remains unknown, like vancomycin, **40** induces the bacterial cell envelope stress-response system and inhibits bacterial growth.⁵⁸ The *yyd* gene cluster of *B. subtilis* encodes a precursor peptide (YydF), a rSAM enzyme (YydG), a protease (YydH) and an ABC transporter (YydI/J).⁵⁹ Bioinformatics analysis identified *yydF* and *yydG* homologs in human-associated Gram-positive bacteria, including *Enterococcus faecalis*, *Streptococcus agalactiae*, and *Staphylococcus epidermidis*.⁶⁰

Studies on purified YydG suggested it contains two [4Fe-4S] clusters and forms **9**.⁶⁰ Sequence similarity analysis revealed a closer relationship to sactonine-forming rSAM enzymes over other rSAM epimerases.⁶⁰ Mutational analysis indicated that the one [4Fe-4S] cluster was coordinated by a canonical rSAM domain while the second cluster was probably bound by a C-terminal SPASM domain. Both of these [4Fe-4S] clusters were catalytically essential. Biochemical studies on a truncated peptide substrate (YydF₁₈₋₄₉) revealed Val36, Ile44, or both were epimerized (**40**).⁶⁰ According to the proposed mechanism, **9** abstracts the C α hydrogen from Val36 and/or Ile44 of **41**, resulting in radical **43** with the loss of C α -atom stereochemistry. **43** is then quenched by the thiol hydrogen of Cys223, a mode of reactivity observed in ribonucleotide reductase and DNA spore photoproduct lyase.^{8,60} The authors implicated Cys223 from a mutational study that showed it was not involved in [Fe-S] cluster

formation but was catalytically important. After radical quenching by Cys223, an epimerized product (**42**) and a thiyl radical (**44**) are formed. This proposal is further supported by the stable incorporation of a solvent-derived hydrogen atom into the peptide backbone. Finally, **44** is proposed to coordinate one Fe atom of the SPASM [4Fe-4S] cluster, followed by reduction and protonation to regenerate the Cys223 thiol for the next catalytic cycle.⁶⁰ These mechanistic details remain speculative and require further experimental confirmation.

D) C-C crosslinking reactions

Several biosynthetic strategies have emerged to generate a diverse array of peptide crosslinks, including thioethers, disulfides, isopeptides, esters, thiolactones, and P450-mediated cross-couplings.¹ In this section, we discuss two rSAM-mediated C-C crosslinking reactions reported in RiPP biosynthesis.

D.1. StrB in streptide biosynthesis—Streptide (**45**, Figure 10) is a cyclic peptide produced by *Streptococcus thermophilus* and bears a covalent linkage between two unactivated carbons of lysine and tryptophan.^{61,62} *S. thermophilus* harbors a quorum-sensing system common to pathogenic streptococci species.⁶² This quorum-sensing network, comprised of a short hydrophobic peptide (SHP) and the Rgg transcriptional regulator, upregulates the *str* gene cluster leading to streptide formation. The *str* gene cluster contains a 30-residue precursor peptide (*strA*), a rSAM enzyme (*strB*), and a putative transporter (*strC*).^{61,62}

The structure of streptide was elucidated using NMR and high-resolution mass spectrometry techniques and established that the Lys-Trp crosslink identified eight years earlier⁶² connects the C β of Lys2 and the C-7 of Trp6 (Figure 10B).⁶¹ Deletion studies in *S. thermophilus* confirmed that StrB was essential for streptide production. The authors hypothesized that StrB installed the Lys-to-Trp crosslink, which would be followed by cleavage and export by StrC to yield mature streptide.⁶¹ This proposal was supported by in vitro assays of StrB that showed the presence of two [4Fe-4S] clusters, with the canonical rSAM domain containing one [4Fe-4S] cluster. Like the thioether-forming rSAMs,⁴⁷ the C-terminal SPASM domain⁵³ coordinated the second [4Fe-4S] cluster. The production of the crosslinked peptide correlated with the formation of **9**.⁶¹ A potential mechanism was proposed (Figure 10D) in which **9** abstracts the hydrogen atom from β -carbon of Lys2 of **46**. The resulting lysyl radical (**48**) was supported by studies using a deuterated StrA substrate. Compound **48** then undergoes an intramolecular radical addition on to the C-7 of indolyl side chain of Trp6, forming the crosslinked intermediate (**49**). Deprotonation and re-aromatization with concomitant reduction of the auxiliary [4Fe-4S] cluster yields **47**.⁶¹ A redox role for the auxiliary SPASM cluster was proposed based on EPR data, which suggested that after one turnover, intramolecular electron transfer from the SPASM cluster to the rSAM cluster would return both to their catalytic active states. This proposal is consistent with other rSAM enzymes, such as DesII.⁶³

D.2. PqqE in pyrroloquinoline quinone (PQQ) biosynthesis—Pyrroloquinoline quinone (PQQ, **52**, Figure 11) is a redox active cofactor predominantly utilized by alcohol

and sugar dehydrogenases localized in the periplasm of Gram-negative bacteria to produce ATP via an alternate, non-glycolytic pathway.⁶⁴ The conserved *pqq* gene cluster possesses six genes (*pqqA-F*) and is found in many pathogens, making this pathway a potential target for future antibiotics.⁶⁵ PQQ is formed from the fusion of evolutionarily conserved Glu and Tyr residues located within a peptidic substrate in a complex and poorly understood biosynthetic pathway.⁶⁵ Gene deletion studies in *Klebsiella pneumoniae* have shown that four out of six gene products (PqqA, PqqC, PqqD, and PqqE) are required for PQQ production.⁶⁶ PqqA serves as the precursor peptide. PqqB shares homology to β -lactamases and has an unknown function. PqqC is a cofactor-independent, oxygen-activating enzyme that catalyzes ring closure through an 8-electron oxidation of the final biosynthetic intermediate, 3a-(2-amino-2-carboxyethyl)-4,5-dioxo-4,5,6,7,8,9-hexahydroquinoline-7,9-dicarboxylic acid. PqqD is a small protein that binds PqqA and potentiates its interaction with PqqE, a rSAM enzyme. Lastly, PqqF is the putative protease that releases mature PQQ from the processed precursor peptide.⁶⁵

Extensive biochemical studies^{67–69} have shown that PqqE catalyzes the C-C linkage between Glu15 and Tyr19 of the PqqA substrate (**50** to **51**, Figure 11B) in a PqqD-dependent manner. Recently, PqqD was shown to associate with PqqA with sub-micromolar affinity and acts as a RRE to deliver PqqA to PqqE.^{70,71} PqqE is a founding member of the SPASM domain-containing rSAM enzymes,⁵³ which contain one or more important auxiliary [4Fe-4S] clusters in their C-terminal regions, as described at numerous points in this review.⁷⁰

Using *Methylobacterium extorquens* as a model,⁷² it was shown that PqqE contains three [Fe-S] centers, of which at least two are [4Fe-4S].⁷¹ Reconstituted PqqE was capable of crosslinking Glu15 and Tyr19 of *M. extorquens* PqqA.⁷¹ A mechanistic proposal was postulated (Figure 11C) in which **9** abstracts the γ -hydrogen from Glu15 of **50**, forming a carbon-based radical (**53**). This then undergoes an intramolecular radical addition on to the 3-position of Tyr19 to form a bridged radical intermediate (**54**). **54** then would lose a proton and an electron, potentially to the auxiliary cluster, to form the rearomatized, crosslinked product (**51**), which will be further processed to **52**. This study provided new insights regarding the role of PqqD⁵ and clarified several questions regarding PQQ biosynthesis.⁷¹ Further studies will be required to delineate the role of the auxiliary clusters and the precise details of the crosslinking reaction.

E) Decarboxylation and C-C bond forming crosslinking reaction

E.1. MftC in mycofactocin biosynthesis—Mycofactocin (**58**, Figure 12) is a putative RiPP that is primarily associated with mycobacteria.⁷³ Mycofactocin BGCs are comprised of six genes (*mftA-F*) and often co-occur with dehydrogenases, suggesting that the cluster transforms the precursor peptide MftA to a unique redox-active cofactor used by the dehydrogenases.⁷⁴ MftA is the precursor peptide and ~30 residues in length. MftB is an RRE-containing protein while MftC is a SPASM domain-containing rSAM enzyme that catalyzes the oxidative decarboxylation of the C-terminal tyrosine of MftA in a MftB-dependent manner (**56** to **57**). Mature mycofactocin has yet to be isolated.^{73,75}

Characterization of the MftC reaction identified two isomeric products which provided some mechanistic insight.⁷⁶ The minor product was the previously identified **57**. The major product (**58**, Figure 12C) contained a crosslink between C β of Val29 and either the C α or C β of the decarboxylated Tyr30 (tyramine). NMR simulations on model compounds, along with other spectroscopic support, favored the C α position as the crosslinked site. It was further established that MftC uses two equivalents of **4** to catalyze **58** formation and that the phenolic group of Tyr30 was critical for the reaction. In the proposed mechanism, **9** abstracts a C β hydrogen from Tyr30 of MftA (**56**). Radical intermediate (**59**) would then undergo decarboxylation to form **57**. In the second turnover, **9** from a second equivalent of **4** would abstract a C β hydrogen from Val29 of **57** to yield radical intermediate **60**. Subsequently, **60** would intramolecularly attack the C α of the tyramine double bond of Tyr30 forming a C-C crosslink. The resulting radical intermediate (**61**) would be resonance-stabilized, which upon addition of an electron and a proton, would yield the bridged product (**58**, major compound). Recently, a creatininase homolog, MftE, was reported to catalyze the cleavage of **58** to liberate the final two residues (VY) in a MftB-independent manner. It was proposed that this dipeptide would undergo further processing to the mature mycofactocin.⁷⁷ While additional studies are required to validate this hypothesis, these advances mark, to our knowledge, the first time a rSAM-SPASM enzyme has been shown to catalyze two different chemistries on the same substrate. Future studies will surely aim to delineate the structure and function of mature mycofactocin.

F) Rearrangement reaction

Radical SAM enzymes catalyze rearrangement reactions for the biosynthesis of primary and secondary metabolites. In known examples, the substrate radicals generated undergo remarkable diversity of rearrangement reactions. Here, we describe one example of this reaction type.

F.1. NosL in nosiheptide biosynthesis—Nosiheptide (**14**, Figure 5), produced by *Streptomyces actuosus*, is a thiopeptide that consists of a central tetra-substituted pyridine, five thiazoles, one dehydroalanine, one dehydrobutyrine, and an indolic acid moiety derived from L-Trp (**1**).³⁰ Gene deletion experiments have revealed that formation of the indolic acid moiety requires the previously described NosN, as well as NosL, which is the second rSAM enzyme encoded in the *nos* gene cluster. In vivo complementation and in vitro studies on purified NosL⁷⁸ established that it catalyzes the conversion of **1** to 3-methyl-2-indolic acid (MIA, **62**, Figure 13). **62** is a key biosynthetic intermediate processed by NosN³¹ to complete the attachment of the indolic moiety to the main nosiheptide framework. NosL has one [4Fe-4S] cluster bound by the canonical rSAM motif.⁷⁸ NosL belongs to the aromatic amino acid lyase family, which includes ThiH (thiamin biosynthesis), HydG (Fe-Fe hydrogenase maturation) and CofH (F₄₂₀ cofactor biosynthesis).⁹ A highly similar enzyme, NocL, from the nocathiacin biosynthetic pathway has also been characterized.⁷⁹

Extensive biochemical and structural studies^{30,78,80–88} have demonstrated that the NosL-catalyzed reaction (Figure 13) involves an unprecedented, intramolecular migration of the carboxylate of **1** to the C-2 of the indole ring. This reaction also transforms the methylene carbon to a methyl group with elimination of the α -carbon as formaldehyde (**63**) and the

amino group as ammonia (**64**). 3-methylindole (**72**) and glyoxalate (**73**) have been identified as shunt products.^{78,86} According to one mechanistic proposal,⁸⁷ **9** abstracts an amino hydrogen from Trp to generate amino radical **65** which undergoes C α -C β bond scission to yield methylene radical **66** and dehydroglycine (**67**). These two intermediates then recombine to yield ketyl radical **68** which can then undergo a second β scission to form **69** which can isomerize to MIA (**62**). The byproduct imine radical (**70**) may abstract hydrogen to give imine **71** which then undergoes hydrolysis to yield **63** and **64**. This mechanism is supported by labeling experiments that identified the origin of key atoms in the product, by trapping **66** as 3-methylindole (**72**), and by hydrolysis of **67** to yield glyoxalate (**73**).^{78,81,87} The most compelling evidence for amino hydrogen abstraction comes from the crystal structure of the NosL-**1** complex⁸⁵ which is further supported by studies employing analogs of **1**.^{81,87} Active site residues suspected to play a catalytic role were probed by mutagenesis.^{85,87} Arg323 was apparently hydrogen bonded to the amino and carboxyl groups of **1** and was suspected to control the regiochemistry of the β scission reaction and in binding **67** for subsequent reaction at C-2 of the indole. In support of this function, the replacement of Arg323 with Lys produced indole-3-pyruvic acid and no products arising from C α -C β bond scission were detected.

A recent study provided further insight to the NosL mechanism.⁸⁶ EPR experiments using multiple isotopologues of **1** were consistent with a C-3-centered radical (**74**, Figure 13C) which could be explained by the cleavage of the C α -C bond from **65** with concomitant migration of the carboxyl radical fragment from C α to C-2. According to the proposed mechanism, radical **74** could undergo a second β scission forming intermediate **69** which can then tautomerize to form the product (**62**) while radical **70** can form **63** and **64**, as previously proposed.⁸⁶ In this regard, NosL differs from other tyrosine lyases (e.g. ThiH),⁹ although all start with amino hydrogen abstraction to form **65**. It was proposed that subtle substrate motions in the active site of NosL are responsible for a fine tuned radical chemistry, leading to cleavage of the C α -C bond.⁸⁶ However, NosL could also catalyze C α -C β cleavage in vitro, producing **72** and **73** as shunt products.⁷⁸ This in vitro NosL promiscuity is modulated by the amount and the nature of the reductant employed in the reaction. Presumably, these alternative products would not form under physiological conditions.

Summary and outlook

With the advent of new genome-mining tools^{89,90} and the accessibility of vast genomic databases, natural products research has been revitalized. The diversity and malleability of RiPP scaffolds underscores the versatility of the biosynthetic enzymes which create these novel architectures. This review aimed to summarize recent insights into rSAM enzymes involved in RiPP biosynthesis. The enzymes highlighted in this review perform several chemically difficult transformations with a high degree of mechanistic novelty. These rSAM-installed PTMs may also be used in the future to re-engineer RiPP biosynthetic pathways and to generate new-to-nature RiPP analogs that display enhanced bioactivity and improved pharmacological properties.⁹¹ With genome-mining studies predicting that many new rSAM-dependent RiPP classes await discovery, we believe that future efforts will

characterize the requisite enzymes and the mature RiPP products. Undoubtedly, these pathways will display paradigm-shifting enzyme chemistry but also, the products themselves may hold value as chemical probes or drug leads.

Acknowledgments

This work was supported by the National Institutes of Health (GM097142 to D.A.M.) and the Seemon Pines Fellowship from the Department of Chemistry at the University of Illinois at Urbana-Champaign (to G.A.H.).

References

1. Arnison PG, Bibb MJ, Bierbaum G, Bowers AA, Bugni TS, Bulaj G, Camarero JA, Campopiano DJ, Challis GL, Clardy J, Cotter PD, Craik DJ, Dawson M, Dittmann E, Donadio S, Dorrestein PC, Entian KD, Fischbach MA, Garavelli JS, Goeransson U, Gruber CW, Haft DH, Hemscheidt TK, Hertweck C, Hill C, Horswill AR, Jaspars M, Kelly WL, Klinman JP, Kuipers OP, Link AJ, Liu W, Marahiel MA, Mitchell DA, Moll GN, Moore BS, Mueller R, Nair SK, Nes IF, Norris GE, Olivera BM, Onaka H, Patchett ML, Piel J, Reaney MJT, Rebuffat S, Ross RP, Sahl HG, Schmidt EW, Selsted ME, Severinov K, Shen B, Sivonen K, Smith L, Stein T, Suessmuth RD, Tagg JR, Tang GL, Truman AW, Vederas JC, Walsh CT, Walton JD, Wenzel SC, Willey JM, van der Donk WA. Ribosomally synthesized and post-translationally modified peptide natural products: overview and recommendations for a universal nomenclature. *Nat Prod Rep*. 2013; 30:108–160. [PubMed: 23165928]
2. Ortega MA, van der Donk WA. New insights into the biosynthetic logic of ribosomally synthesized and post-translationally modified peptide natural products. *Cell Chem Biol*. 2016; 23:31–44. [PubMed: 26933734]
3. Hetrick KJ, van der Donk WA. Ribosomally synthesized and post-translationally modified peptide natural product discovery in the genomic era. *Curr Opin Chem Biol*. 2017; 38:36–44. [PubMed: 28260651]
4. Burkhardt BJ, Schwalen CJ, Mann G, Naismith JH, Mitchell DA. YcaO-dependent posttranslational amide activation: biosynthesis, structure, and function. *Chem Rev*. 2017; 117:5389–5456. [PubMed: 28256131]
5. Burkhardt BJ, Hudson GA, Dunbar KL, Mitchell DA. A prevalent peptide-binding domain guides ribosomal natural product biosynthesis. *Nat Chem Biol*. 2015; 11:564–570. [PubMed: 26167873]
6. Funk MA, van der Donk WA. Ribosomal natural products, tailored to fit. *Acc Chem Res*. 2017; 50:1577–1586. [PubMed: 28682627]
7. Sardar D, Lin Z, Schmidt EW. Modularity of RiPP enzymes enables designed synthesis of decorated peptides. *Chem Biol*. 2015; 22:907–916. [PubMed: 26165156]
8. Broderick JB, Duffus BR, Duschene KS, Shepard EM. Radical S-adenosylmethionine enzymes. *Chem Rev*. 2014; 114:4229–4317. [PubMed: 24476342]
9. Mehta AP, Abdelwahed SH, Mahanta N, Fedoseyenko D, Philmus B, Cooper LE, Liu Y, Jhulki I, Ealick SE, Begley TP. Radical S-adenosylmethionine (SAM) enzymes in cofactor biosynthesis: a treasure trove of complex organic radical rearrangement reactions. *J Biol Chem*. 2015; 290:3980–3986. [PubMed: 25477515]
10. Söding J, Biegert A, Lupas AN. The HHpred interactive server for protein homology detection and structure prediction. *Nucleic Acids Res*. 2005; 33:W244–W248. [PubMed: 15980461]
11. Grell TAJ, Goldman PJ, Drennan CL. SPASM and twitch domains in S-adenosylmethionine (SAM) radical enzymes. *J Biol Chem*. 2015; 290:3964–3971. [PubMed: 25477505]
12. Struck AW, Thompson ML, Wong LS, Micklefield J. S-adenosyl-methionine-dependent methyltransferases: highly versatile enzymes in biocatalysis, biosynthesis and other biotechnological applications. *ChemBioChem*. 2012; 13:2642–2655. [PubMed: 23180741]
13. Zhang Q, van der Donk WA, Liu W. Radical-mediated enzymatic methylation: A tale of two SAMS. *Acc Chem Res*. 2012; 45:555–564. [PubMed: 22097883]
14. Fujimori DG. Radical SAM-mediated methylation reactions. *Curr Opin Chem Biol*. 2013; 17:597–604. [PubMed: 23835516]

15. Bauerle MR, Schwalm EL, Booker SJ. Mechanistic diversity of radical S-adenosylmethionine (SAM)-dependent methylation. *J Biol Chem.* 2015; 290:3995–4002. [PubMed: 25477520]
16. Walsh CT, Acker MG, Bowers AA. Thiazolyl peptide antibiotic biosynthesis: a cascade of post-translational modifications on ribosomal nascent proteins. *J Biol Chem.* 2010; 285:27525–27531. [PubMed: 20522549]
17. Bagley MC, Dale JW, Merritt EA, Xiong X. Thiopeptide antibiotics. *Chem Rev.* 2005; 105:685–714. [PubMed: 15700961]
18. Just-Baringo X, Albericio F, Alvarez M. Thiopeptide antibiotics: retrospective and recent advances. *Mar Drugs.* 2014; 12:317–351. [PubMed: 24445304]
19. Kelly WL, Pan L, Li C. Thiostrepton biosynthesis: prototype for a new family of bacteriocins. *J Am Chem Soc.* 2009; 131:4327–4334. [PubMed: 19265401]
20. Pierre S, Guillot A, Benjdia A, Sandstroem C, Langella P, Berteau O. Thiostrepton tryptophan methyltransferase expands the chemistry of radical SAM enzymes. *Nat Chem Biol.* 2012; 8:957–959. [PubMed: 23064318]
21. Benjdia A, Pierre S, Gherasim C, Guillot A, Carmona M, Amara P, Banerjee R, Berteau O. The thiostrepton A tryptophan methyltransferase TsrM catalyzes a cob(II)alamin-dependent methyl transfer reaction. *Nat Commun.* 2015; 6:8377. [PubMed: 26456915]
22. Blaszczyk AJ, Silakov A, Zhang B, Maiocco SJ, Lanz ND, Kelly WL, Elliott SJ, Krebs C, Booker SJ. Spectroscopic and electrochemical characterization of the iron-sulfur and cobalamin cofactors of TsrM, an unusual radical S-adenosylmethionine methylase. *J Am Chem Soc.* 2016; 138:3416–3426. [PubMed: 26841310]
23. Blaszczyk AJ, Wang B, Silakov A, Ho JV, Booker SJ. Efficient methylation of C2 in L-tryptophan by the cobalamin-dependent radical S-adenosylmethionine methylase TsrM requires an unmodified N1 amine. *J Biol Chem.* 2017; doi: 10.1074/jbc.M117.778548
24. Freeman MF, Gurgui C, Helf MJ, Morinaka BI, Uria AR, Oldham NJ, Sahl HG, Matsunaga S, Piel J. Metagenome mining reveals polytheonamides as posttranslationally modified ribosomal peptides. *Science.* 2012; 338:387–390. [PubMed: 22983711]
25. Freeman MF, Vagstad AL, Piel J. Polytheonamide biosynthesis showcasing the metabolic potential of sponge-associated uncultivated ‘Entotheonella’ bacteria. *Curr Opin Chem Biol.* 2016; 31:8–14. [PubMed: 26625171]
26. Freeman MF, Helf MJ, Bhushan A, Morinaka BI, Piel J. Seven enzymes create extraordinary molecular complexity in an uncultivated bacterium. *Nat Chem.* 2017; 9:387–395. [PubMed: 28338684]
27. Parent A, Guillot A, Benjdia A, Chartier G, Leprince J, Berteau O. The B12-radical SAM enzyme PoyC catalyzes valine C β -methylation during polytheonamide biosynthesis. *J Am Chem Soc.* 2016; 138:15515–15518. [PubMed: 27934015]
28. Gomez-Escribano JP, Song L, Bibb MJ, Challis GL. Posttranslational β -methylation and macrolactamidation in the biosynthesis of the bottromycin complex of ribosomal peptide antibiotics. *Chem Sci.* 2012; 3:3522–3525.
29. Crone WJK, Vior NM, Santos-Aberturas J, Schmitz LG, Leeper FJ, Truman AW. Dissecting bottromycin biosynthesis using comparative untargeted metabolomics. *Angew Chem, Int Ed.* 2016; 55:9639–9643.
30. Yu Y, Duan L, Zhang Q, Liao R, Ding Y, Pan H, Wendt-Pienkowski E, Tang G, Shen B, Liu W. Nosiheptide biosynthesis featuring a unique indole side ring formation on the characteristic thiopeptide framework. *ACS Chem Biol.* 2009; 4:855–864. [PubMed: 19678698]
31. Ding W, Li Y, Zhao J, Ji X, Mo T, Qianzhu H, Tu T, Deng Z, Yu Y, Chen F, Zhang Q. The catalytic mechanism of the class C radical S-adenosylmethionine methyltransferase NosN. *Angew Chem, Int Ed.* 2017; 56:3857–3861.
32. Ding W, Wu Y, Ji X, Qianzhu H, Chen F, Deng Z, Yu Y, Zhang Q. Nucleoside-linked shunt products in the reaction catalyzed by the class C radical S-adenosylmethionine methyltransferase NosN. *Chem Commun.* 2017; 53:5235–5238.
33. Morris RP, Leeds JA, Naegeli HU, Oberer L, Memmert K, Weber E, LaMarche MJ, Parker CN, Burrer N, Esterow S, Hein AE, Schmitt EK, Krastel P. Ribosomally synthesized thiopeptide

- antibiotics targeting elongation factor Tu. J Am Chem Soc. 2009; 131:5946–5955. [PubMed: 19338336]
34. Hudson GA, Zhang Z, Tietz JI, Mitchell DA, van der Donk WA. In vitro biosynthesis of the core scaffold of the thiopeptide thiomuracin. J Am Chem Soc. 2015; 137:16012–16015. [PubMed: 26675417]
35. Zhang Z, Hudson GA, Mahanta N, Tietz JI, van der Donk WA, Mitchell DA. Biosynthetic timing and substrate specificity for the thiopeptide thiomuracin. J Am Chem Soc. 2016; 138:15511–15514. [PubMed: 27700071]
36. Mahanta N, Zhang Z, Hudson GA, van der Donk WA, Mitchell DA. Reconstitution and substrate specificity of the radical S-adenosylmethionine thiazole C-methyltransferase in thiomuracin biosynthesis. J Am Chem Soc. 2017; 139:4310–4313. [PubMed: 28301141]
37. Tocchetti A, Maffioli S, Iorio M, Alt S, Mazzei E, Brunati C, Sosio M, Donadio S. Capturing linear intermediates and C-terminal variants during maturation of the thiopeptide GE2270. Chem Biol. 2013; 20:1067–1077. [PubMed: 23932526]
38. Bruender NA, Wilcoxon J, Britt RD, Bandarian V. Biochemical and spectroscopic characterization of a radical S-adenosyl-L-methionine enzyme involved in the formation of a peptide thioether cross-Link. Biochemistry. 2016; 55:2122–2134. [PubMed: 27007615]
39. Fluehe L, Marahiel MA. Radical S-adenosylmethionine enzyme catalyzed thioether bond formation in sactipeptide biosynthesis. Curr Opin Chem Biol. 2013; 17:605–612. [PubMed: 23891473]
40. Repka LM, Chekan JR, Nair SK, van der Donk WA. Mechanistic understanding of lanthipeptide biosynthetic enzymes. Chem Rev. 2017; 117:5457–5520. [PubMed: 28135077]
41. Kawulka K, Sprules T, McKay RT, Mercier P, Diaper CM, Zuber P, Vederas JC. Structure of subtilosin A, an antimicrobial peptide from *Bacillus subtilis* with unusual posttranslational modifications linking cysteine sulfurs to α -carbons of phenylalanine and threonine. J Am Chem Soc. 2003; 125:4726–4727. [PubMed: 12696888]
42. Sit CS, van Belkum MJ, McKay RT, Worobo RW, Vederas JC. The 3D solution structure of thurincin H, a bacteriocin with four sulfur to α -carbon crosslinks. Angew Chem, Int Ed. 2011; 50:8718–8721.
43. Gonzalez-Pastor JE, Hobbs EC, Losick R. Cannibalism by sporulating bacteria. Science. 2003; 301:510–513. [PubMed: 12817086]
44. Rea MC, Sit CS, Clayton E, O'Connor PM, Whittall RM, Zheng J, Vederas JC, Ross RP, Hill C. Thuricin CD, a posttranslationally modified bacteriocin with a narrow spectrum of activity against *Clostridium difficile*. Proc Natl Acad Sci U S A. 2010; 107:9352–9357. [PubMed: 20435915]
45. Sit CS, McKay RT, Hill C, Ross RP, Vederas JC. The 3D structure of thuricin CD, a two-component bacteriocin with cysteine sulfur to α -carbon cross-links. J Am Chem Soc. 2011; 133:7680–7683. [PubMed: 21526839]
46. Nakai T, Ito H, Kobayashi K, Takahashi Y, Hori H, Tsubaki M, Tanizawa K, Okajima T. The radical S-adenosyl-L-methionine enzyme QhpD catalyzes sequential formation of intra-protein sulfur-to-methylene carbon thioether bonds. J Biol Chem. 2015; 290:11144–11166. [PubMed: 25778402]
47. Fluehe L, Knappe TA, Gattner MJ, Schaefer A, Burghaus O, Linne U, Marahiel MA. The radical SAM enzyme AlbA catalyzes thioether bond formation in subtilosin A. Nat Chem Biol. 2012; 8:350–357. [PubMed: 22366720]
48. Benjdia A, Guillot A, Lefranc B, Vaudry H, Leprince J, Berteau O. Thioether bond formation by SPASM domain radical SAM enzymes: C α H-atom abstraction in subtilosin A biosynthesis. Chem Commun. 2016; 52:6249–6252.
49. Liu W-T, Yang Y-L, Xu Y, Lamsa A, Haste NM, Yang JY, Ng J, Gonzalez D, Ellmermeier CD, Straight PD, Pevzner PA, Pogliano J, Nizet V, Pogliano K, Dorrestein PC. Imaging mass spectrometry of intraspecies metabolic exchange revealed the cannibalistic factors of *Bacillus subtilis*. Proc Natl Acad Sci U S A. 2010; 107:16286–16290. [PubMed: 20805502]
50. Fluehe L, Burghaus O, Wieckowski BM, Giessen TW, Linne U, Marahiel MA. Two [4Fe-4S] clusters containing radical SAM enzyme SkfB catalyze thioether bond formation during the

maturation of the sporulation killing factor. *J Am Chem Soc.* 2013; 135:959–962. [PubMed: 23282011]

51. Bruender NA, Bandarian V. SkfB abstracts a hydrogen atom from C α on SkfA to initiate thioether cross-link formation. *Biochemistry.* 2016; 55:4131–4134. [PubMed: 27410522]
52. Wieckowski BM, Hegemann JD, Mielcarek A, Boss L, Burghaus O, Marahiel MA. The PqqD homologous domain of the radical SAM enzyme ThnB is required for thioether bond formation during thurincin H maturation. *FEBS Lett.* 2015; 589:1802–1806. [PubMed: 26026269]
53. Lanz ND, Booker SJ. Auxiliary iron-sulfur cofactors in radical SAM enzymes. *Biochim Biophys Acta, Mol Cell Res.* 2015; 1853:1316–1334.
54. Mathur H, Fallico V, O'Connor PM, Rea MC, Cotter PD, Hill C, Ross RP. Insights into the mode of action of the sactibiotic thuricin CD. *Front Microbiol.* 2017; 8:696. [PubMed: 28473822]
55. Haft DH, Basu MK. Biological systems discovery in silico: radical S-adenosylmethionine protein families and their target peptides for posttranslational modification. *J Bacteriol.* 2011; 193:2745–2755. [PubMed: 21478363]
56. Grove TL, Himes PM, Hwang S, Yumerefendi H, Bonanno JB, Kuhlman B, Almo SC, Bowers AA. Structural insights into thioether bond formation in the biosynthesis of sactipeptides. *J Am Chem Soc.* 2017; 139:11734–11744. [PubMed: 28704043]
57. Morinaka BI, Vagstad AL, Helf MJ, Gugger M, Kegler C, Freeman MF, Bode HB, Piel J. Radical S-adenosylmethionine epimerases: regioselective introduction of diverse D-amino acid patterns into peptide natural products. *Angew Chem, Int Ed.* 2014; 53:8503–8507.
58. Jordan S, Junker A, Helmann JD, Mascher T. Regulation of LiaRS-dependent gene expression in *Bacillus subtilis*: identification of inhibitor proteins, regulator binding sites, and target genes of a conserved cell envelope stress-sensing two-component system. *J Bacteriol.* 2006; 188:5153–5166. [PubMed: 16816187]
59. Butcher BG, Lin YP, Helmann JD. The yydFGHIJ operon of *Bacillus subtilis* encodes a peptide that induces the LiaRS two-component system. *J Bacteriol.* 2007; 189:8616–8625. [PubMed: 17921301]
60. Benjdia A, Guillot A, Ruffie P, Leprince J, Berteau O. Post-translational modification of ribosomally synthesized peptides by a radical SAM epimerase in *Bacillus subtilis*. *Nat Chem.* 2017; 9:698–707. [PubMed: 28644475]
61. Schramma KR, Bushin LB, Seyedsayamdost MR. Structure and biosynthesis of a macrocyclic peptide containing an unprecedented lysine-to-tryptophan crosslink. *Nat Chem.* 2015; 7:431–437. [PubMed: 25901822]
62. Ibrahim M, Guillot A, Wessner F, Algaron F, Besset C, Courtin P, Gardan R, Monnet V. Control of the transcription of a short gene encoding a cyclic peptide in *Streptococcus thermophilus*: a new quorum-sensing system? *J Bacteriol.* 2007; 189:8844–8854. [PubMed: 17921293]
63. Ruszczycky MW, Choi S-h, Liu H-w. Stoichiometry of the redox neutral deamination and oxidative dehydrogenation reactions catalyzed by the radical SAM enzyme DesII. *J Am Chem Soc.* 2010; 132:2359–2369. [PubMed: 20121093]
64. Anthony C. Pyrroloquinoline quinone (PQQ) and quinoprotein enzymes. *Antioxid Redox Signaling.* 2001; 3:757–774.
65. Shen YQ, Bonnot F, Imsand EM, RoseFigura JM, Sjolander K, Klinman JP. Distribution and properties of the genes encoding the biosynthesis of the bacterial cofactor, pyrroloquinoline quinone. *Biochemistry.* 2012; 51:2265–2275. [PubMed: 22324760]
66. Meulenberg JJM, Sellink E, Riegman NH, Postma PW. Nucleotide sequence and structure of the *Klebsiella pneumoniae* pqq operon. *Mol Gen Genet.* 1992; 232:284–294. [PubMed: 1313537]
67. Wecksler SR, Stoll S, Tran H, Magnusson OT, Wu S-p, King D, Britt RD, Klinman JP. Pyrroloquinoline quinone biogenesis: demonstration that PqqE from *Klebsiella pneumoniae* is a radical S-adenosyl-L-methionine enzyme. *Biochemistry.* 2009; 48:10151–10161. [PubMed: 19746930]
68. Wecksler SR, Stoll S, Iavarone AT, Imsand EM, Tran H, Britt RD, Klinman JP. Interaction of PqqE and PqqD in the pyrroloquinoline quinone (PQQ) biosynthetic pathway links PqqD to the radical SAM superfamily. *Chem Commun.* 2010; 46:7031–7033.

69. Klinman JP, Bonnot F. Intrigues and intricacies of the biosynthetic pathways for the enzymatic quinocofactors: PQQ, TTQ, CTQ, TPQ, and LTQ. *Chem Rev.* 2014; 114:4343–4365. [PubMed: 24350630]
70. Latham JA, Iavarone AT, Barr I, Juthani PV, Klinman JP. PqqD is a novel peptide chaperone that forms a ternary complex with the radical S-adenosylmethionine protein PqqE in the pyrroloquinoline quinone biosynthetic pathway. *J Biol Chem.* 2015; 290:12908–12918. [PubMed: 25817994]
71. Barr I, Latham JA, Iavarone AT, Chantarojsiri T, Hwang JD, Klinman JP. Demonstration that the radical S-adenosylmethionine (SAM) enzyme PqqE catalyzes de novo carbon-carbon cross-linking within a peptide substrate PqqA in the presence of the peptide chaperone PqqD. *J Biol Chem.* 2016; 291:8877–8884. [PubMed: 26961875]
72. Saichana N, Tanizawa K, Pechousek J, Novak P, Yakushi T, Toyama H, Frebortova J. PqqE from *Methylobacterium extorquens* AM1: a radical S-adenosyl-L-methionine enzyme with an unusual tolerance to oxygen. *J Biochem.* 2016; 159:87–99. [PubMed: 26188050]
73. Khaliullin B, Aggarwal P, Bubas M, Eaton GR, Eaton SS, Latham JA. Mycofactocin biosynthesis: modification of the peptide MftA by the radical S-adenosylmethionine protein MftC. *FEBS Lett.* 2016; 590:2538–2548. [PubMed: 27312813]
74. Haft DH. Bioinformatic evidence for a widely distributed, ribosomally produced electron carrier precursor, its maturation proteins, and its nicotinoprotein redox partners. *BMC Genomics.* 2011; 12:21. [PubMed: 21223593]
75. Bruender NA, Bandarian V. The radical S-adenosyl-L-methionine enzyme MftC catalyzes an oxidative decarboxylation of the C-terminus of the MftA peptide. *Biochemistry.* 2016; 55:2813–2816. [PubMed: 27158836]
76. Khaliullin B, Ayikpoe R, Tuttle M, Latham JA. Mechanistic elucidation of the mycofactocin-biosynthetic radical-S-adenosylmethionine protein, MftC. *J Biol Chem.* 2017; doi: 10.1074/jbc.M117.795682
77. Bruender NA, Bandarian V. The creatininase homolog MftE from *Mycobacterium smegmatis* catalyzes a peptide cleavage reaction in the biosynthesis of a novel ribosomally synthesized post-translationally modified peptide (RiPP). *J Biol Chem.* 2017; 292:4371–4381. [PubMed: 28077628]
78. Zhang Q, Li YX, Chen DD, Yu Y, Duan L, Shen B, Liu W. Radical-mediated enzymatic carbon chain fragmentation-recombination. *Nat Chem Biol.* 2011; 7:154–160. [PubMed: 21240261]
79. Zhang Q, Chen D, Lin J, Liao R, Tong W, Xu Z, Liu W. Characterization of NocL involved in thiopeptide nocathiacin I biosynthesis: a [4Fe-4S] cluster and the catalysis of a radical S-adenosylmethionine enzyme. *J Biol Chem.* 2011; 286:21287–21294. [PubMed: 21454624]
80. Ji X, Li Y, Ding W, Zhang Q. Substrate-tuned catalysis of the radical S-adenosyl-L-methionine enzyme NosL involved in nosiheptide biosynthesis. *Angew Chem, Int Ed.* 2015; 54:9021–9024.
81. Ji X, Li Y, Jia Y, Ding W, Zhang Q. Mechanistic insights into the radical S-adenosyl-L-methionine enzyme NosL from a substrate analogue and the shunt products. *Angew Chem, Int Ed.* 2016; 55:3334–3337.
82. Ji X, Li Y, Xie L, Lu H, Ding W, Zhang Q. Expanding radical SAM chemistry by using radical addition reactions and SAM analogues. *Angew Chem, Int Ed.* 2016; 55:11845–11848.
83. Ji X, Liu WQ, Yuan S, Yin Y, Ding W, Zhang Q. Mechanistic study of the radical SAM-dependent amine dehydrogenation reactions. *Chem Commun.* 2016; 52:10555–10558.
84. Qianzhu H, Ji W, Ji X, Chu L, Guo C, Lu W, Ding W, Gao J, Zhang Q. Reactivity of the nitrogen-centered tryptophanyl radical in the catalysis by the radical SAM enzyme NosL. *Chem Commun.* 2017; 53:344–347.
85. Nicolet Y, Zeppieri L, Amara P, Fontecilla-Camps JC. Crystal structure of tryptophan lyase (NosL): evidence for radical formation at the amino group of tryptophan. *Angew Chem, Int Ed.* 2014; 53:11840–11844.
86. Sicoli G, Mouesca JM, Zeppieri L, Amara P, Martin L, Barra AL, Fontecilla-Camps JC, Gambarelli S, Nicolet Y. Fine-tuning of a radical-based reaction by radical S-adenosyl-L-methionine tryptophan lyase. *Science.* 2016; 351:1320–1323. [PubMed: 26989252]

87. Bhandari DM, Xu H, Nicolet Y, Fontecilla-Camps JC, Begley TP. Tryptophan lyase (NosL): mechanistic insights from substrate analogues and mutagenesis. *Biochemistry*. 2015; 54:4767–4769. [PubMed: 26204056]
88. Bhandari DM, Fedoseyenko D, Begley TP. Tryptophan lyase (NosL): a cornucopia of 5'-deoxyadenosyl radical mediated transformations. *J Am Chem Soc*. 2016; 138:16184–16187. [PubMed: 27998091]
89. Mohimani H, Kersten RD, Liu WT, Wang M, Purvine SO, Wu S, Brewer HM, Pasa-Tolic L, Bandeira N, Moore BS, Pevzner PA, Dorrestein PC. Automated genome mining of ribosomal peptide natural products. *ACS Chem Biol*. 2014; 9:1545–1551. [PubMed: 24802639]
90. Tietz JI, Schwalen CJ, Patel PS, Maxson T, Blair PM, Tai HC, Zakai UI, Mitchell DA. A new genome-mining tool redefines the lasso peptide biosynthetic landscape. *Nat Chem Biol*. 2017; 13:470–478. [PubMed: 28244986]
91. Latham JA, Barr I, Klinman JP. At the confluence of ribosomally synthesized peptide modification and radical S-adenosylmethionine (SAM) enzymology. *J Biol Chem*. 2017; doi: 10.1074/jbc.R117.797399

Figure 1.

rSAM enzymes catalyze a wide variety of chemical complex transformations in RiPP biosynthesis. The variety of domain architectures contained within this family echoes the observed mechanistic diversity. Class C methyltransferases are homologous to HemN (orange) and the rSAM (green) domain. It is possible that additional enzymes may actually harbor a RiPP precursor peptide Recognition Element (RRE) domain (e.g. SkfB); however, the sequence hypervariability of the RRE may preclude bioinformatic identification by tools such as HHpred.¹⁰ The SPASM domain of YydG (*) features only one auxiliary [4Fe-4S] cluster. SPASM domains that harbor a single auxiliary [4Fe-4S] cluster have recently been referred to as “Twitch” domains.¹¹

Figure 2.

The biosynthetic gene cluster (A) for thiostrepton (**3**) features TsrM, an atypical rSAM MT which methylates C2 of Trp (B). Catalysis of this reaction is mediated by a cobalamin-based cofactor. Competing radical (C) and polar (D) mechanisms are presented.

Figure 3.
Structure of polytheonamide A/B. Sites of sp^3 C-methylation are indicated in red while the side chains of Ca epimerized sites are shown in blue. Both reactions are mediated by rSAM enzymes.

Figure 4.

The biosynthetic gene cluster for polytheonamide features three rSAM enzymes: two MTs (PoyB and PoyC) as well as an epimerase (PoyD) (A). PoyB/C catalyze the C-methylation of specific residues on the PoyA precursor peptide (B) using a cobalamin-dependent reaction mechanism (C).

Figure 5.

Nosiheptide (A) biosynthesis requires two rSAM enzymes, NosL and NosN. NosL converts Trp (**1**) to 3-methyl-2-indolic acid (**62**) while NosN *C*-methylates the indolyl moiety of **12** (a derivative of **62**) prior to its incorporation into **14** (B) through a mechanism that utilizes MTA (**15**) as a methyl donor (C). The indolyl moiety is shown in blue for structural clarity.

Figure 6.

Structurally related thiopeptides GE2270A (**23**) and thiomuracin (**24**) are encoded by similar BGCs which both feature class C rSAM MTs (A). These enzymes *C*-methylate the 5 position of thiazoles on a linear intermediate (B) in a regioselective manner. Both compounds feature methylation at Thz4 while GE2270A has an additional modification at Thz6 (C).

Figure 7.

Several RiPP BGCs have been identified that feature a rSAM responsible for sactionine installation (A). Structure of a sactionine linkage (B). Sactipeptides exhibit variation in the number and location of the sactionine linkages (C). * denotes sactionines only observed through in vitro reconstitution (i.e. the natural product has not been characterized). Proposed mechanism for sactionine biosynthesis (D). In contrast to the radical-based mechanism of sactionine formation, lanthionines, which feature a C_β thioether linkage, are formed through a Michael-type addition (E).

Figure 8.

An alternative mechanism proposed for Tte1186-dependent thioether formation.

Figure 9.

Epipeptide BGC (A). Using residues 18–49 of the precursor peptide YydF (B), the installation of up to two D-amino acids was observed (C). Proposed enzymatic mechanism (D).

Figure 10.

The streptide BGC from *S. thermophilus* encodes one rSAM enzyme (A). Structure of streptide (B). The Lys-Trp crosslinking is formed by StrB (C). Proposed StrB mechanism (D).

Figure 11.

PQQ BGC from *Klebsiella pneumoniae* (A). PqqE catalyzes an intramolecular crosslink in PqqA (B). Proposed PqqE mechanism (C).

Figure 12.

Mycofactocin is encoded by a BGC from *Mycobacterium smegmatis*(A) MftA biosynthetic intermediate produced (B), decarboxylation and cross-linking reactions catalyzed by MftC (C). Proposed mechanism for MftC (D).

Figure 13.

NosL catalyzed transformation of **1** to **62**, **63**, and **64**; colored atoms show the fate of the carbon atoms during rearrangement (A). Two mechanisms have been proposed for NosL (B, C).