Erratum for Salma et al., “Tfe3 and Tfeb Transcriptionally Regulate Peroxisome Proliferator-Activated Receptor γ 2 Expression in Adipocytes and Mediate Adiponectin and Glucose Levels in Mice”

Nunciada Salma,^a Jun S. Song,^{b,c,d} Akinori Kawakami,^a Suprabha P. Devi,^a Mehdi Khaled,^e José M. Cacicedo,^f David E. Fisher^a

Department of Dermatology, Cutaneous Biology Research Center, Massachusetts General Hospital, Harvard Medical School, Charlestown, Massachusetts, USA^a; Institute for Genomic Biology, University of Illinois, Urbana-Champaign, Urbana, Illinois, USA^b; Department of Bioengineering, University of Illinois, Urbana-Champaign, Urbana, Illinois, USA^c; Department of Physics, University of Illinois, Urbana-Champaign, Urbana, Illinois, USA^d; Gustave Roussy, INSERM U753, Villejuif, France^e; Diabetes and Metabolism Unit, Department of Medicine Section of Endocrinology, Boston University School of Medicine, Boston, Massachusetts, USA^f

Volume 37, no. 15, e00608-16, 2017, <https://doi.org/10.1128/MCB.00608-16>. Page 13, Fig. 11 legend: Line 5 should read “. . .preimmune IgG (white bars), anti-Tfeb (gray bars), anti-Tfe3 (black bars), anti-Ppary (diagonal line bars), and anti-RNA polymerase II (phospho S2) (dotted bars) antibodies.”

Citation Salma N, Song JS, Kawakami A, Devi SP, Khaled M, Cacicedo JM, Fisher DE. 2017. Erratum for Salma et al., “Tfe3 and Tfeb transcriptionally regulate peroxisome proliferator-activated receptor γ 2 expression in adipocytes and mediate adiponectin and glucose levels in mice.” *Mol Cell Biol* 37:e00466-17. <https://doi.org/10.1128/MCB.00466-17>.

Copyright © 2017 American Society for Microbiology. All Rights Reserved.