Figure 1. Tubulysin ADCs containing (A) carbamate or (B) quaternary ammonium salt linker connections are cleaved and eliminated to release tubulysin drugs containing either a 2° amine (1) or a 3° amine (2), respectively.

pipecolic acid N-terminus of the peptide drug could generate a secondary amine for linker attachment.

We employed this strategy with an easily accessible, potent analogue tubulysin M^{10,22} (2) to generate NH-tubulysin M (1) and its carbamate-linked ADC using a cysteine-engineered antibody at site LC-K149C (Figure 1A). Unfortunately, consistent with other recent reports,^{15,18} removal of the methyl group to enable linker attachment resulted in a dramatic loss in potency as 1 was 18–177-fold less potent than 2 (Table 1). As might be expected, this loss of free drug potency abolished ADC activity, as the anti-CD22 carbamate linked-ADC 4 was inactive *in vitro* against lymphoma cell lines.

To prevent this loss of activity, we needed another method for linking tubulysin. We recently described a novel quaternary ammonium salt peptide linker that is amenable to the stable connection and traceless release of tertiary amines.^{23,24} Utilizing this concise, protecting-group-free synthesis, we generated

tubulysin M ADCs (Figure 1B) in just three steps from commercially available starting materials. The resulting anti-CD22 tubulysin ADC 5 was potent *in vitro*, in a range similar to MMAE ADC 8 in the CD22-expressing sensitive cell lines BJAB and WSU-DLCL2 (Table 1). Significantly, tubulysin ADC 5 retained activity in a Pgp-overexpressing cell line (BJAB.Luc/Pgp),²⁵ while MMAE ADC 8 was completely inactive. The ADC activity was antigen-specific as evidenced by the lack of activity of 5 against CD22-negative Jurkat cells as well as non-targeting control anti-NaPi ADC 6 being inactive. Furthermore, the requirement for linker cleavage was supported by the inactivity of tubulysin ADC 7, which is epimeric at the citrulline amino acid required for proteolytic linker cleavage.

Due to its potent *in vitro* activity against sensitive and resistant cell lines, we selected quaternary ammonium salt-linked tubulysin ADC 5 to evaluate in a human lymphoma xenograft in mice. Unexpectedly, 5 showed very modest tumor growth inhibition (TGI) after a single IV dose of 1 mg/kg (57% TGI), whereas MMAE ADC 8 afforded complete tumor regression through day 28 at a matched dose (Figure 2A). This *in vitro*–*in vivo* disconnect led us to investigate the *in vivo*

Table 1. *In Vitro* Cytotoxicity of Tubulysin and MMAE Free Drugs and ADCs

#	free drug or ADC	BJAB IC ₅₀ (nM)	BJAB.Luc/Pgp IC ₅₀ (nM)	WSU IC ₅₀ (nM)	Jurkat IC ₅₀ (nM)
1	NH-tubulysin M	2.1	23	2.0	5.0
2	tubulysin M	0.12	0.13	0.11	0.10
3	MMAE	0.42	>30	0.19	0.13
4	anti-CD22-MC-VC-PABC-NH-tubulysin M	>190		>190	>190
5	anti-CD22-MC-VC(S)-PABQ-tubulysin M	6.8	25	0.27	236
6	anti-NaPi-MC-VC(S)-PABQ-tubulysin M	>253		185	>253
7	anti-CD22-MC-VC(R)-PABQ-tubulysin M	>253		>253	>253
8	anti-CD22-MC-VC-PABC-MMAE	3.3	>380	0.95	>253
9	tubulysin Pr	0.14	0.34	0.19	0.26
10	anti-CD22-MC-VC-PABQ-tubulysin Pr	5.3	19	0.28	
11	anti-NaPi-MC-VC-PABQ-tubulysin Pr	>253		200	>253

Figure 2. *In vivo* efficacy and stability of a first generation tubulysin ADC. (A) Efficacy of quaternary ammonium salt-linked tubulysin M ADC 5 (K149C) compared to MMAE ADC 8 (K149C) in a BJAB.Luc human lymphoma xenograft model in mice. ADCs were given as a single IV dose of 1 mg/kg. (B) Acetate cleavage was observed *in vivo* in mice. (C) *In vivo* stability (expressed as drug-to-antibody ratio or DAR) of tubulysin M ADC 5 over time in mice compared to MMAE ADC 8.

stability of the two ADCs. Both ADCs comprised maleimides conjugated to the same site (LC-K149C) of a cysteine-engineered antibody with a drug-to-antibody ratio (DAR) of 2. This antibody site was selected because it forms a highly stable connection to maleimide-containing linker drugs (Figure S1).²⁶ Utilizing affinity-capture LC–MS to determine *in vivo* stability,²⁷ we were able to observe a small loss in mass (−43 Da) of the tubulysin ADC 5 over time (Figure S2). This loss is consistent with cleavage of the acetate (Figure 2B), which is known to cause a significant reduction in potency for tubulysins²⁸ and has been recently described with tubulysin ADCs.¹⁴ Due to the expected loss of ADC activity upon acetate cleavage, we assigned this drug modification as a loss in “drug” in DAR calculations. With this in mind, we observed a significant loss of DAR over time for tubulysin ADC 5 (Figure 2C) with no active “drug” left after 4 days. In contrast, MMAE ADC 8 had no loss in DAR over 4 days. The cleavage of the acetate and corresponding loss in drug activity helps explain the limited efficacy of tubulysin ADC 5.

We explored two approaches to improve the *in vivo* stability of tubulysin ADCs, specifically aiming to minimize or prevent the putatively enzyme-mediated deacetylation. One approach is to utilize cysteine engineering to identify an antibody site to protect the conjugated small molecule from metabolism during circulation.¹⁴ Toward this effort, we identified an antibody site that decreased tubulysin acetate cleavage and subsequently improved efficacy (manuscript in preparation).²⁹ Yet we preferred a more general solution not requiring antibody engineering. Thus, the second approach was to replace the acetate with a stable isostere while retaining potency. Tubulysin analogues with a propyl ether and/or an *N*-propyl amide afforded low nanomolar potency.³⁰ We therefore aimed to replace the acetate and modify the amide of tubulysin M (2) to generate tubulysin Pr (9) (Figure 3).

Figure 3. Stabilizing tubulysin M (2) by replacing the acetate with a propyl ether. This modification combined with installation of an *N*-propyl amide in place of the *N*-Me amide led to the design of target tubulysin Pr (9).

Preparation of 9 required an 11-step synthesis (Figure S4), and step 7 required an amide coupling involving a now increasingly sterically hindered amine (12) with a hindered acid chloride 13 (Scheme 1). The resulting amide was only obtained in a 5% yield and necessitated a new synthetic route. We found that swapping the Fmoc-amino acid chloride 13 for azido acid chloride 15 resulted in a 14-fold increase in yield of the desired amide and, upon quantitative reduction of the azide, provided amine 14 to intersect the original route.

With the new tubulysin Pr analogue 9, we were able to remove the metabolic liability, while retaining the low double digit picomolar potency found with tubulysin M (2) (Table 1). This retained potency was also observed with the quaternary ammonium linked anti-CD22 tubulysin Pr ADC 10, which killed lymphoma cells in a target-dependent manner.

An *in vivo* evaluation of the conjugate was necessary to fully understand how changing the tubulysin structure would impact both ADC stability and efficacy. We evaluated the antitumor

Scheme 1. Improving Tubulysin Amide Coupling with Sterically Hindered Substrates

effects of the anti-CD22 tubulysin Pr ADC 10 in a human lymphoma xenograft model in mice (Figure 4).

Figure 4. *In vivo* efficacy of a second generation tubulysin ADC. (A) Structure of tubulysin Pr ADCs. (B) Efficacy of quaternary ammonium salt linked-tubulysin Pr ADC 10 (K149C) compared to MMAE ADC 8 (K149C) in a BJAB.Luc or (C) BJAB.Luc/Pgp human lymphoma model in mice. ADCs were given as a single IV dose.

The stabilized tubulysin Pr ADC 10 resulted in tumor stasis for 21 days when administered at a single dose of 1 mg/kg (Figure 4B). The response was dose-dependent with modest tumor growth inhibition (77% TGI, day 11) observed at 0.5 mg/kg, and target-specific, as anti-NaPi tubulysin Pr ADC 11 gave no activity (identical to the vehicle control). While the tubulysin Pr ADC 10 was not quite as efficacious as MMAE ADC 8 at a matched 0.5 mg/kg dose (80% TGI, day 11), it had a much-improved efficacy compared to the unstable tubulysin ADC 5 (Figure 2A). Stability was also dramatically improved with no loss of DAR observed for 7 days *in vivo* (Figure S3).

One of the most compelling reasons to utilize tubulysins as ADC payloads is their potential to overcome multidrug resistance. As previously reported, we derived lymphoma xenograft models resistant to MMAE ADCs.²⁵ These models were characterized to have an upregulation of Pgp while maintaining equivalent levels of target (CD22). We used this model here as it represents acquired resistance to the previous generation of ADCs targeting tubulin. We tested tubulysin and MMAE ADCs head-to-head in the BJAB.Luc-Pgp lymphoma model in mice (Figure 4C). MMAE ADC 8 was dosed up to 8

mg/kg and gave minimal tumor growth inhibition (41% TGI, day 13), which reflected more than 16-fold reduction compared to activity observed in the parental BJAB model. Gratifyingly, the stabilized tubulysin Pr ADC **10** gave clear dose-dependent tumor growth inhibition, resulting in 74% TGI (day 13) at 2 mg/kg, which reflected approximately only 4-fold reduction compared to activity in the parental BJAB model. This improved activity of the tubulysin ADC relative to the MMAE ADC in an MDR model is likely a result of the released tubulysin payload not being a substrate for efflux by Pgp. Finally, the tubulysin ADCs were tolerated at all doses with mouse body weights increasing over the duration of the study (data not shown).

New classes of highly potent cytotoxic drugs with distinct mechanisms and activity are a source of excitement to synthetic and medicinal chemists. While their incorporation into conjugates for targeted therapy can improve their tolerability and therefore therapeutic potential, new challenges arise. Among these are ways to stably and reversibly connect the drug and, particularly with ADCs, the increased risk for metabolic instability accompanying prolonged half-life. We have employed a peptide linker combined with a quaternary ammonium salt linker connection to provide a stable and bioreversible connection of tubulysins to antibodies. Furthermore, we identified and addressed the metabolism of an acetate ester critical to activity through its replacement with a propyl ether. The resulting conjugate was highly stable and efficacious against both sensitive and multidrug-resistant tumor models. The latter activity substantially differentiates this tubulysin ADC from the clinically validated MMAE ADC and generates excitement for its further investigation.

■ ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: [10.1021/acsmedchemlett.7b00243](https://doi.org/10.1021/acsmedchemlett.7b00243).

In vivo stability data, methods for conjugate preparation, *in vitro* cytotoxicity, *in vivo* efficacy and stability, experimental procedures for the synthesis of new compounds, and the corresponding characterization data (PDF)

■ AUTHOR INFORMATION

Corresponding Author

*E-mail: thomashp@gene.com.

ORCID

Thomas H. Pillow: [0000-0001-7300-1002](https://orcid.org/0000-0001-7300-1002)

Present Address

[§]23andMe, Mountain View, California 94041, United States.

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

We thank Yanzhou Liu and Kewei Xu for analytical support and Terry Crawford and Phil Bergeron for assistance during manuscript preparation.

■ ABBREVIATIONS

ADC, antibody–drug conjugate; DAR, drug-to-antibody ratio; MMAE, monomethyl auristatin E; Pgp, p-glycoprotein

■ REFERENCES

- (1) Polakis, P. Antibody Drug Conjugates for Cancer Therapy. *Pharmacol. Rev.* **2016**, 68 (1), 3–19.
- (2) Chari, R. V. J. Expanding the Reach of Antibody-Drug Conjugates. *ACS Med. Chem. Lett.* **2016**, 7 (11), 974–976.
- (3) Doronina, S. O.; Toki, B. E.; Torgov, M. Y.; Mendelsohn, B. A.; Cervený, C. G.; Chace, D. F.; DeBlanc, R. L.; Gearing, R. P.; Bovee, T. D.; Siegall, C. B.; Francisco, J. A.; Wahl, A. F.; Meyer, D. L.; Senter, P. D. Development of Potent Monoclonal Antibody Auristatin Conjugates for Cancer Therapy. *Nat. Biotechnol.* **2003**, 21 (7), 778–784.
- (4) Widdison, W. C.; Wilhelm, S. D.; Cavanagh, E. E.; Whiteman, K. R.; Leece, B. A.; Kovtun, Y.; Goldmacher, V. S.; Xie, H.; Steeves, R. M.; Lutz, R. J.; Zhao, R.; Wang, L.; Blättler, W. A.; Chari, R. V. J. Semisynthetic Maytansine Analogues for the Targeted Treatment of Cancer. *J. Med. Chem.* **2006**, 49 (14), 4392–4408.
- (5) Schumacher, D.; Hackenberger, C. P. R.; Leonhardt, H.; Helma, J. Current Status: Site-Specific Antibody Drug Conjugates. *J. Clin. Immunol.* **2016**, 36 (Suppl 1), 100–107.
- (6) Chudasama, V.; Maruani, A.; Caddick, S. Recent Advances in the Construction of Antibody-Drug Conjugates. *Nat. Chem.* **2016**, 8 (2), 114–119.
- (7) Sasse, F.; Steinmetz, H.; Heil, J.; Höfle, G.; Reichenbach, H. Tubulysins, New Cytostatic Peptides From Myxobacteria Acting on Microtubuli. Production, Isolation, Physico-Chemical and Biological Properties. *J. Antibiot.* **2000**, 53 (9), 879–885.
- (8) Steinmetz, H.; Glaser, N.; Herdtweck, E.; Sasse, F.; Reichenbach, H.; Höfle, G. Isolation, Crystal and Solution Structure Determination, and Biosynthesis of Tubulysins—Powerful Inhibitors of Tubulin Polymerization From Myxobacteria. *Angew. Chem., Int. Ed.* **2004**, 43 (37), 4888–4892.
- (9) Khalil, M. W.; Sasse, F.; Lünsdorf, H.; Elnakady, Y. A.; Reichenbach, H. Mechanism of Action of Tubulysin, an Antimitotic Peptide From Myxobacteria. *ChemBioChem* **2006**, 7 (4), 678–683.
- (10) Wang, Y.; Benz, F. W.; Wu, Y.; Wang, Q.; Chen, Y.; Chen, X.; Li, H.; Zhang, Y.; Zhang, R.; Yang, J. Structural Insights Into the Pharmacophore of Vinca Domain Inhibitors of Microtubules. *Mol. Pharmacol.* **2016**, 89 (2), 233–242.
- (11) Kaur, G.; Hollingshead, M.; Holbeck, S.; Schauer-Vukasinović, V.; Camalier, R. F.; Dömling, A.; Agarwal, S. Biological Evaluation of Tubulysin a: A Potential Anticancer and Antiangiogenic Natural Product. *Biochem. J.* **2006**, 396 (2), 235–242.
- (12) Giddens, A. C.; Lee, H. H.; Lu, G.-L.; Miller, C. K.; Guo, J.; Lewis Phillips, G. D.; Pillow, T. H.; Terce, M. Analogues of DNA Minor Groove Cross-Linking Agents Incorporating aminoCBI, an Amino Derivative of the Duocarmycins: Synthesis, Cytotoxicity, and Potential as Payloads for Antibody-Drug Conjugates. *Bioorg. Med. Chem.* **2016**, 24 (22), 6075–6081.
- (13) Murray, B. C.; Peterson, M. T.; Fecik, R. A. Chemistry and Biology of Tubulysins: Antimitotic Tetrapeptides with Activity Against Drug Resistant Cancers. *Nat. Prod. Rep.* **2015**, 32 (5), 654–662.
- (14) Tumey, L. N.; Leverett, C. A.; Vetelino, B.; Li, F.; Rago, B.; Han, X.; Loganzo, F.; Musto, S.; Bai, G.; Sukuru, S. C. K.; Graziani, E. I.; Puthenveetil, S.; Casavant, J.; Ratnayake, A.; Marquette, K.; Hudson, S.; Doppalapudi, V. R.; Stock, J.; Tchistiakova, L.; Bessire, A. J.; Clark, T.; Lucas, J.; Hosselet, C.; O'Donnell, C. J.; Subramanyam, C. Optimization of Tubulysin Antibody-Drug Conjugates: a Case Study in Addressing ADC Metabolism. *ACS Med. Chem. Lett.* **2016**, 7 (11), 977–982.
- (15) Leverett, C. A.; Sukuru, S. C. K.; Vetelino, B. C.; Musto, S.; Parris, K.; Pandit, J.; Loganzo, F.; Varghese, A. H.; Bai, G.; Liu, B.; Liu, D.; Hudson, S.; Doppalapudi, V. R.; Stock, J.; O'Donnell, C. J.; Subramanyam, C. Design, Synthesis, and Cytotoxic Evaluation of Novel Tubulysin Analogues as ADC Payloads. *ACS Med. Chem. Lett.* **2016**, 7 (11), 999–1004.
- (16) Li, J. Y.; Perry, S. R.; Muniz-Medina, V.; Wang, X.; Wetzel, L. K.; Rebelatto, M. C.; Hinrichs, M. J. M.; Bezabeh, B. Z.; Fleming, R. L.; Dimasi, N.; Feng, H.; Toader, D.; Yuan, A. Q.; Xu, L.; Lin, J.; Gao, C.; Wu, H.; Dixit, R.; Osbourn, J. K.; Coats, S. R. A Biparatopic HER2-

Targeting Antibody-Drug Conjugate Induces Tumor Regression in Primary Models Refractory to or Ineligible for HER2-Targeted Therapy. *Cancer Cell* **2016**, 29 (1), 117–129.

(17) Thompson, P.; Fleming, R.; Bezabeh, B.; Huang, F.; Mao, S.; Chen, C.; Harper, J.; Zhong, H.; Gao, X.; Yu, X.-Q.; Hinrichs, M. J.; Reed, M.; Kamal, A.; Strout, P.; Cho, S.; Woods, R.; Hollingsworth, R. E.; Dixit, R.; Wu, H.; Gao, C.; Dimasi, N. Rational Design, Biophysical and Biological Characterization of Site-Specific Antibody-Tubulysin Conjugates with Improved Stability, Efficacy and Pharmacokinetics. *J. Controlled Release* **2016**, 236, 100–116.

(18) Burke, P. J.; Hamilton, J. Z.; Pires, T. A.; Setter, J. R.; Hunter, J. H.; Cochran, J. H.; Waight, A. B.; Gordon, K. A.; Toki, B. E.; Emmerton, K. K.; Zeng, W.; Stone, I. J.; Senter, P. D.; Lyon, R. P.; Jeffrey, S. C. Development of Novel Quaternary Ammonium Linkers for Antibody-Drug Conjugates. *Mol. Cancer Ther.* **2016**, 15 (5), 938–945.

(19) Cohen, R.; Vugts, D. J.; Visser, G. W. M.; Stigter-van Walsum, M.; Bolijn, M.; Spiga, M.; Lazzari, P.; Shankar, S.; Sani, M.; Zanda, M.; van Dongen, G. A. M. S. Development of Novel ADCs: Conjugation of Tubulysin Analogues to Trastuzumab Monitored by Dual Radiolabeling. *Cancer Res.* **2014**, 74 (20), 5700–5710.

(20) Shen, B.-Q.; Xu, K.; Liu, L.; Raab, H.; Bhakta, S.; Kenrick, M.; Parsons-Reponte, K. L.; Tien, J.; Yu, S.-F.; Mai, E.; Li, D.; Tibbitts, J.; Baudys, J.; Saad, O. M.; Scales, S. J.; McDonald, P. J.; Hass, P. E.; Eigenbrot, C.; Nguyen, T.; Solis, W. A.; Fuji, R. N.; Flagella, K. M.; Patel, D.; Spencer, S. D.; Khawli, L. A.; Ebens, A.; Wong, W. L.; Vandlen, R.; Kaur, S.; Sliwkowski, M. X.; Scheller, R. H.; Polakis, P.; Junutula, J. R. Conjugation Site Modulates the in Vivo Stability and Therapeutic Activity of Antibody-Drug Conjugates. *Nat. Biotechnol.* **2012**, 30 (2), 184–189.

(21) Dubowchik, G. M.; Firestone, R. A.; Padilla, L.; Willner, D.; Hofstead, S. J.; Mosure, K.; Knipe, J. O.; Lasch, S. J.; Trail, P. A. Cathepsin B-Labile Dipeptide Linkers for Lysosomal Release of Doxorubicin From Internalizing Immunoconjugates: Model Studies of Enzymatic Drug Release and Antigen-Specific in Vitro Anticancer Activity. *Bioconjugate Chem.* **2002**, 13 (4), 855–869.

(22) Wang, Z.; McPherson, P. A.; Raccor, B. S.; Balachandran, R.; Zhu, G.; Day, B. W.; Vogt, A.; Wipf, P. Structure-Activity and High-Content Imaging Analyses of Novel Tubulysins. *Chem. Biol. Drug Des.* **2007**, 70 (2), 75–86.

(23) Lehar, S. M.; Pillow, T. H.; Xu, M.; Staben, L.; Kajihara, K. K.; Vandlen, R.; DePalatis, L.; Raab, H.; Hazenbos, W. L.; Morisaki, J. H.; Kim, J.; Park, S.; Darwish, M.; Lee, B.-C.; Hernandez, H.; Loyet, K. M.; Lupardus, P.; Fong, R.; Yan, D.; Chalouni, C.; Luis, E.; Khalfin, Y.; Plise, E.; Cheong, J.; Lyssikatos, J. P.; Strandh, M.; Koefoed, K.; Andersen, P. S.; Flygare, J. A.; Wah Tan, M.; Brown, E. J.; Mariathasan, S. Novel Antibody-Antibiotic Conjugate Eliminates Intracellular *S. Aureus*. *Nature* **2015**, 527 (7578), 323–328.

(24) Staben, L. R.; Koenig, S. G.; Lehar, S. M.; Vandlen, R.; Zhang, D.; Chuh, J.; Yu, S.-F.; Ng, C.; Guo, J.; Liu, Y.; Fourie-O'Donohue, A.; Go, M.; Linghu, X.; Segraves, N. L.; Wang, T.; Chen, J.; Wei, B.; Phillips, G. D. L.; Xu, K.; Kozak, K. R.; Mariathasan, S.; Flygare, J. A.; Pillow, T. H. Targeted Drug Delivery Through the Traceless Release of Tertiary and Heteroaryl Amines From Antibody-Drug Conjugates. *Nat. Chem.* **2016**, 8 (12), 1112–1119.

(25) Yu, S.-F.; Zheng, B.; Go, M.; Lau, J.; Spencer, S.; Raab, H.; Soriano, R.; Jhunjhunwala, S.; Cohen, R.; Caruso, M.; Polakis, P.; Flygare, J.; Polson, A. G. A Novel Anti-CD22 Anthracycline-Based Antibody-Drug Conjugate (ADC) That Overcomes Resistance to Auristatin-Based ADCs. *Clin. Cancer Res.* **2015**, 21 (14), 3298–3306.

(26) Pillow, T. H.; Sadowsky, J. D.; Zhang, D.; Yu, S.-F.; Del Rosario, G.; Xu, K.; He, J.; Bhakta, S.; Ohri, R.; Kozak, K. R.; Ha, E.; Junutula, J. R.; Flygare, J. A. Decoupling stability and release in disulfide bonds with antibody-small molecule conjugates. *Chem. Sci.* **2017**, 8, 366–370.

(27) Xu, K.; Liu, L.; Saad, O. M.; Baudys, J.; Williams, L.; Leipold, D.; Shen, B.; Raab, H.; Junutula, J. R.; Kim, A.; Kaur, S. Characterization of Intact Antibody-Drug Conjugates From Plasma/Serum in Vivo by Affinity Capture Capillary Liquid Chromatography-Mass Spectrometry. *Anal. Biochem.* **2011**, 412 (1), 56–66.

(28) Patterson, A. W.; Peltier, H. M.; Sasse, F.; Ellman, J. A. Design, Synthesis, and Biological Properties of Highly Potent Tubulysin D Analogues. *Chem. - Eur. J.* **2007**, 13 (34), 9534–9541.

(29) Genentech. Cysteine Engineered Antibodies and Conjugates. WO 2016/040856 A2, 2016.

(30) Richter, W. WO 2011/057805 A1, 2011.