Published in final edited form as:

Tetrahedron Lett. 2017 May 24; 58(21): 2031–2033. doi:10.1016/j.tetlet.2017.04.026.

Asymmetric solution-phase mixture aldol reaction using oligomeric ethylene glycol tagged chiral oxazolidinones

Serhan Turkeyilmaz^{a,b} and Craig S. Wilcox^{a,*}

^aDepartment of Chemistry, University of Pittsburgh, Pittsburgh, PA 15260, USA

^bDepartment of Pharmaceutical Chemistry, Faculty of Pharmacy, Istanbul University, 34116 Beyazit, Istanbul, Turkey

Abstract

Sorting tags are oligomeric structures that can be used as protecting groups or chiral auxiliaries enabling solution-phase mixture syntheses of multiple tagged compounds in one pot and allowing for facile and predictable chromatographic separation of products at the end of synthetic sequences. Perfluorinated hydrocarbon and oligomeric ethylene glycol (OEG) derivatives are known classes of sorting tags. Herein we describe the preparation of OEGylated chiral oxazolidinones and their use in asymmetric solution-phase mixture aldol reactions. Through the use of such oxazolidinones based on tyrosine four different individually tagged aldol adducts were obtained as a mixture, chromatographically demixed, detagged, and it was shown that these processes gave the desired aldol products in good yield and enantioselectivity.

Graphical abstract

Keywords

Solution-Phase Mixture Synthesis; Oligomeric Ethylene Glycol; Evans Aldol Reaction; Oxazolidinone; Sorting Tag

*Corresponding author. Tel.: +1 4126248269; fax: +1 4126248611; daylite@pitt.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Supplementary Information: Supplementary data (synthetic procedures, NMR data, and HPLC data) associated with this article can be found, in the online version, at ...

Introduction

Drawbacks associated with solid-phase organic synthesis have prompted a search for alternative methods for the generation of libraries of structurally diverse compounds. Such methods include liquid phase organic synthesis^{1a,b} along with solution-phase strategies such as the generation of indexed combinatorial libraries^{2a-c} and approaches involving the use of phase tags such as precipitons,^{3a-c} boronic acid derivatives,⁴ and fluororous synthesis.^{5a-g} Among these alternative methods fluororous synthesis is particularly important as it provides for a reliable strategy, termed Fluororous Mixture Synthesis (FMS),^{6a-i} allowing for the mixture synthesis of target structures and their separation into individual products at the end of the synthetic sequence. This is accomplished through labelling of each substrate with a fluorocarbon sorting tag of unique chain length, taking mixtures of these tagged substrates through a number of synthetic steps, and separation (i.e. demixing) of these tagged substrates by chromatography with a fluororous stationary phase where substrate elution order is dictated by the chain length of the sorting tag.

We have discovered that oligomeric ethylene glycol (OEG) derivatives can also be used as sorting tags.^{7a,b} OEG tags allow for the orderly and predictable demixing of a mixture of tagged substrates through chromatography on regular silica gel where elution times are directly proportional to OEG chain length (Figure 1).^{7a} Demixing efficiency can be further enhanced by addition of lithium salts to silica gel (i.e. complexation chromatography).^{7a} OEG tags are particularly noteworthy because they are separable under orthogonal chromatographic conditions with respect to fluororous tags. This chromatographic orthogonality allows for double tagging of substrates through which a larger number of parallel solution-phase reactions can be carried out in the same reaction vessel compared to the use of a single class of sorting tag.^{7c-d}

The aldol reaction is a very important method for C-C bond formation. Use of chiral oxazolidinones (i.e. Evans auxiliaries) as chiral auxiliaries for asymmetric aldol reactions has been a seminal contribution to this area.^{8a-c} The high synthetic value of Evans auxiliaries made the preparation and application of OEG tagged chiral oxazolidinones an attractive target for us. Chiral oxazolidinones derived from tyrosine were of particular interest as they allow for straightforward derivatization with OEGs through etherification. In this communication we would like to report on the preparation and application of OEG tagged chiral oxazolidinones derived from tyrosine.

The objectives of this study were: *i*. The preparation of acylated-OEGylated chiral oxazolidinones **2a–d** starting from **1a–b**; *ii*. Execution of mixture solution-phase aldol reactions of **2a–d** with benzaldehyde; *iii*. Demonstration of demixing of the products using chromatography on normal phase silica; *iv*. Stereochemical analysis of the aldol products thus formed (**3a–d**) to establish the efficacy and suitability of these OEGylated chiral oxazolidinones for asymmetric mixture aldol reactions. Structures **2a–d** were targeted because they would maximize the chromatographic resolution of products **3a–d**. Aldol reactions with benzaldehyde were selected since the resulting aldol adducts are known and thus would allow for facile assessment of the utility of OEGylated chiral oxazolidinones (Scheme 1).

Results and Discussion

While a number of routes could be considered for the preparation of tyrosine based chiral oxazolidinones, we found the method developed by Green *et al.* to be particularly convenient and high yielding.⁹ Boc protection of *R*- and *S*-tyrosine (**1a–b**) followed by benzylation afforded the globally protected amino acids (**4a–b**) in good yields. LiAlH₄ reduction of **4a–b**, NaH mediated oxazolidinone formation, and removal of the benzyl protecting group through hydrogenation afforded chiral oxazolidinones **5a–b** in good yields as well (Scheme 1). The optical rotation of **5b** (–11.2°) matched the values reported in the literature. The optical rotation of **5a** (+11.2°) was, as expected, the opposite of **5b** (Scheme 2).

OEGylation of oxazolidinones **5a–b** was accomplished with acceptable yields through Cs₂CO₃/KI mediated etherification with OEG-chlorides of varying chain lengths. Thus the *R*-isomers were tagged with OEG-1 and OEG-3 (**6a**, **6c**) and the *S*-isomers were tagged with OEG-2 and OEG-4 (**6b**, **6d**). **6a–b** and **6c–d** were acylated in good yields with, respectively, butyryl and propionyl chloride using *n*-BuLi to give **2a–d** (Scheme 3).⁸

With **2a–d** in hand, mixture aldol reactions of these OEG-tagged substrates with benzaldehyde were carried out. Enolization of a mixture of **2a–d** using Bu₂BOTf (1.8 eq.) and NEt₃ (1.9 eq.), and subsequent addition of benzaldehyde afforded a mixture of OEG-tagged aldol products **3a–d**. This mixture was demixed using silica gel flash column chromatography (Scheme 4). It is remarkable that it was possible to separate 9 compounds (4 products, 4 starting materials, and excess benzaldehyde) with good purity in a single run. Demixing was also demonstrated using normal phase HPLC and a chromatogram for this separation is provided in Figure 2 (for chromatographic parameters see Table SI-1, Supporting Information). Product elution times in either case were-as expected-directly proportional to bound OEG chain length.

Aldol products **7a–d** were liberated from the OEG-tagged chiral auxiliaries using LiOOH (LiOH mixed with H₂O₂).¹⁰ To facilitate the stereochemical analysis of these products they were converted to the corresponding methyl esters **7a–d** through reaction with freshly prepared diazomethane (CH₂N₂). Both steps proceeded with good to excellent yields (Scheme 5).

Relative and absolute stereochemical configurations, and enantiomeric purities of **7a–d** and **8a–d** were determined, respectively, using ¹H-NMR, optical rotation measurements, and chiral HPLC. α-β hydrogen spin-spin coupling constants (i.e. J_{AB}) as determined through ¹H-NMR indicated gauche conformations in all cases (i.e. J_{AB} < 6 Hz), thus it was inferred that *syn*-aldol products were obtained (Table SI-2, Supporting Information).¹¹ Optical rotation (i.e. [α]_D) values for **7c–d** and **8a–d** matched those previously reported (Table SI-2, Supporting Information),^{12a–e} and it was concluded that products with the expected absolute configuration were obtained. Chiral HPLC analysis of **8a–d** revealed enantiomeric excess (i.e. % ee) values of 99% for **8a**, 98% for **8b**, 94% for **8c**, and 93% for **8d** (Table SI-4, Supporting Information).

In this study we have demonstrated the preparation of OEG-tagged chiral oxazolidinones and have used them in solution-phase mixture aldol reactions. The products obtained were easily demixed using silica gel flash column chromatography. As the chromatogram in Figure 2 shows, even using an extremely steep gradient excellent baseline resolution was observed between **3a–d**. Thus it is conceivable that it would be possible to demix more than 4 products using just 4 OEG tags. We have observed that OEG tags were compatible with all reaction conditions required in this work. Furthermore, aldol products were obtained in good yields and their stereochemical configurations were identical to what would be expected of the corresponding products of single component reactions. We believe that OEG-tagged protecting groups and chiral auxiliaries like **6a–d** will be important tools in the arsenal of chemists seeking to generate libraries of compounds through solution-phase mixture synthesis.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This work was supported by the National Institutes of Health (5R01GM068005).

References

1. (a) Gravet DJ, Janda KD. *Chem Rev.* 1997; 97:489–509. [PubMed: 11848880] (b) Toy PH, Janda KD. *Acc Chem Res.* 2000; 33:546–554. [PubMed: 10955985]
2. (a) Pirrung MC, Chen J. *J Am Chem Soc.* 1995; 117:1240–1245. (b) Andrus MB, Asgari D, Wenke L. *J Org Chem.* 1999; 64:2978–2979. [PubMed: 11674382] (c) Andrus MB, Turner TM, Sauna ZE, Ambudkar SV. *J Org Chem.* 2000; 65:4973–4983. [PubMed: 10956480]
3. (a) Bosanac T, Yang J, Wilcox CS. *Angew Chem Int Ed.* 2001; 40:1875–1879. (b) Bosanac T, Wilcox CS. *J Am Chem Soc.* 2002; 124:4194–4195. [PubMed: 11960432] (c) Bosanac T, Wilcox CS. *Org Lett.* 2004; 6:2321–2324. [PubMed: 15228269]
4. Mothana S, Grassot JM, Hall DG. *Angew Chem Int Ed.* 2010; 49:2883–2887.
5. Reviews: (a) Curran DP. *Angew Chem Int Ed.* 1998; 37:1174–1196. (b) Gladysz JA, Curran DP. *Tetrahedron.* 2002; 58:3823–3825. (c) Zhang W, Curran DP. *Tetrahedron.* 2006; 62:11837–11865. [PubMed: 18509513] (d) Curran DP. *J Fluor Chem.* 2008; 129:898–902. [PubMed: 19802321] (e) Fustero S, Acena JL, Catalán S. *Top Curr Chem.* 2012; 308:45–68. [PubMed: 21952845] (f) Jiang Z-X, Zhi S, Zhang W. *Mol Divers.* 2014; 18:203–218. [PubMed: 24443173] (g) Mukherjee T, Gladysz JA. *Aldrichimica Acta.* 2015; 48:25–28.
6. (a) Luo Z, Zhang Q, Oderatashi Y, Curran DP. *Science.* 2001; 291:1766–1769. [PubMed: 11230688] (b) Zhang W, Luo Z, Chen CH-T, Curran DP. *J Am Chem Soc.* 2002; 124:10443–10450. [PubMed: 12197746] (c) Zhang Q, Rivkin A, Curran DP. *J Am Chem Soc.* 2002; 124:5774–5781. [PubMed: 12010052] (d) Curran DP, Moura-Letts G, Pohlman M. *Angew Chem Int Ed Eng.* 2006; 45:2423–2436. (e) Yang F, Newsome JJ, Curran DP. *J Am Chem Soc.* 2006; 128:14200–14205. [PubMed: 17061905] (f) Curran DP, Sinha MK, Zhang K, Sabatini JJ, Cho D-H. *Nature Chem.* 2012; 4:124–129. [PubMed: 22270645] (g) Moretti JD, Wang X, Curran DP. *J Am Chem Soc.* 2012; 134:7963–7970. [PubMed: 22515682] (h) Yeh EA-H, Kumli E, Damodaran K, Curran DP. *J Am Chem Soc.* 2013; 135:1577–1584. [PubMed: 23297872] (i) Takahashi H, Endo N, Takanose H, Sugiyama Y, Eguchi F, Oguri K, Hamamoto H, Shioiri T, Matsugi M. *Eur J Org Chem.* 2015:3971–3982.
7. (a) Wilcox CS, Turkyilmaz S. *Tetrahedron Lett.* 2005; 46:1827–1829. (b) Gudipati V, Curran DP, Wilcox CS. *J Org Chem.* 2006; 71:3599–3607. [PubMed: 16626147] (c) Wilcox CS, Gudipati V, Lu H, Turkyilmaz S, Curran DP. *Angew Chem Int Ed.* 2005; 44:6938–6940. (d) Curran DP, Zhang Q,

- Richard C, Lu H, Gudipati V, Wilcox CS. J Am Chem Soc. 2006; 128:9561–9573. [PubMed: 16848495]
8. (a) Evans DA, Nelson JV, Vogel E, Taner TR. J Am Chem Soc. 1981; 103:3099–3111.(b) Evans DA, Bartoli J, Shih TL. J Am Chem Soc. 1981; 103:2127–2129.(c) Evans DA, Takacs JM, McGee LR, Ennis MD, Mathre DJ. Pure & Appl Chem. 1981; 53:1109–1127.
9. Green R, Taylor PJM, Bull SD, James TD, Mahon MF, Merritt AT. Tetrahedron Asymm. 2003; 14:2619–2623.
10. Evans DA, Ellman JA, Dow RL. Tetrahedron Lett. 1987; 28:1123–1126.
11. Heathcock, CH. Asymmetric Synthesis. Morrison, JD., editor. Vol. 3. Academic Press; New York, NY: 1984. p. 111-212.
12. (a) Kim M-S, Jun J-G. Synth Comm. 2002; 32:3851–3864.(b) Casper DM, Burgeson JR, Esken JM, Ferrence GM, Hitchcock SR. Org Lett. 2002; 4:3739–3742. [PubMed: 12375932] (c) Vicario JL, Badía D, Domínguez E, Rodríguez M, Carrillo L. J Org Chem. 2000; 65:3754–3760. [PubMed: 10864760] (d) Yan T-H, Tan C-W, Lee H-C, Lo H-C, Huang T-Y. J Am Chem Soc. 1993; 115:2613–2621.(e) Hsiao C-N, Liu L, Miller MJ. J Org Chem. 1987; 52:2201–2206.

Highlights

- Oligomeric ethylene glycol (OEG) derivatives act as chromatographic sorting tags.
- Chiral oxazolidinones tagged with OEGs of varying length were synthesized.
- 4 aldol products were synthesized in one pot using tagged chiral oxazolidinones.
- Products were easily separated using silica gel flash column chromatography.
- Aldol products were obtained with good yield and enantioselectivity.

Figure 1.
Principle of solution-phase mixture synthesis using OEG-based sorting tags.

Figure 2.

Panel A shows a normal phase HPLC chromatogram for a mixture of aldol adducts **3a–d** (Conditions: 250 × 4.6 mm Supelco Supelcosil silica column, gradient elution (1:1 EtOAc:hexanes for 5 min., then EtOAc to 5% IPA in EtOAc in 3 min.), 1 mL/min flow rate, UV-Vis detection). Panel B shows 2D-UV/Vis spectra demonstrating identical absorption profiles for the chromatographic peaks corresponding to **3a–d** (right to left).

Scheme 1.

Objectives of this study.

Scheme 2.

i. Boc_2O , dioxane: H_2O (1:1), TEA, 20 °C, 20 h; **ii.** BnBr, K_2CO_3 , Bu_4NI , DMF, rt, 2 d; **iii.** LiAlH_4 , THF, 0 °C to rt, 2 h; **iv.** NaH, THF, rt, 24 h; **v.** Pd/C, H_2 , rt, 24 h.

Scheme 3.

i. Cs₂CO₃, KI, Cl-OEG_nMe, DMF, 60 °C, 24 h; **ii.** *n*-BuLi, THF, -78 °C, 20 min.; **iii.** RCH₂CH₂COCl, THF, -78 °C, 30 min.

Scheme 4.

i. 1.8 eq. Bu₂BOTf, 1.9 eq. NEt₃, CH₂Cl₂, 0 °C, 10 min; ii. PhCHO, CH₂Cl₂, -78 °C to 0 °C, 2 h; iii. Demixing using flash column chromatography on silica.

Scheme 5.

i. LiOH, H₂O₂, THF:H₂O (3:1), 0 °C, 2 h; ii. CH₂N₂, Et₂O, 0 °C, 1 h.