Published in final edited form as:

Childs Nerv Syst. 2017 November ; 33(11): 1917–1926. doi:10.1007/s00381-017-3588-6.

Time-to-event Analysis of Surgically Treated Posthemorrhagic Hydrocephalus in Preterm Infants: A Single-Institution Retrospective Study

Rowland H. Han^{1,*}, Daniel Berger^{1,*}, Mohamed Gabir¹, Brandon S. Baksh¹, Diego M. Morales¹, Amit M. Mathur², Christopher D. Smyser^{2,3,4}, Jennifer M. Strahle^{1,2}, and David D. Limbrick Jr.^{1,2}

¹Department of Neurological Surgery, Washington University School of Medicine, St. Louis, MO, USA

²Department of Pediatrics, Washington University School of Medicine, St. Louis, MO, USA

³Department of Neurology, Washington University School of Medicine, St. Louis, MO, USA

⁴Mallinckrodt Institute of Radiology, Washington University School of Medicine, St. Louis, MO, USA

Abstract

Purpose—The purpose of this study is to report time-points relevant to the neurosurgical management of posthemorrhagic hydrocephalus (PHH).

Methods—Data were collected retrospectively on 104 preterm infants with intraventricular hemorrhage (IVH) who received neurosurgical intervention for PHH at St. Louis Children's Hospital from 1994 to 2016. Kaplan-Meier curves were constructed for various endpoints.

Results—IVH grade on head ultrasound obtained through routine clinical care was II, III, and IV in 5 (4.8%), 33 (31.7%), and 66 (63.5%) of the patients, respectively. Neither IVH size or location appeared to affect development of PHH. Days from birth to IVH, ventriculomegaly, temporizing

To whom correspondence should be addressed: Rowland H. Han, Washington University School of Medicine, One Children's Place, Suite 4S20, St. Louis Children's Hospital, St. Louis, MO 63110, Telephone: 314-397-9116, rowland.han@wustl.edu, Fax: 314-454-2818.

*These authors contributed equally to this work.

Authors' Contributions

R.H.H. and D.B. performed literature review, study design, collected data, and wrote the manuscript. R.H.H. performed data analysis. B.S.B. wrote and critically revised the manuscript. M.G. and D.M.M. collected data and critically revised the manuscript. A.M.M., C.D.S., J.M.S., and D.D.L. designed the study and critically revised the manuscript.

Compliance with Ethical Standards

Conflict of interest

Dr. Limbrick receives research funds and/or research equipment for unrelated projects from Medtronic, Inc., Karl Storz, Inc., and Microbot Medical, Inc. Dr. Limbrick has received philanthropic equipment contributions for humanitarian relief work from Karl Storz, Inc. and Aesculap, Inc. The authors have no personal, financial, or institutional interest in any of the drugs, materials, or devices described in this article.

Research involving human participants and/or animals

All procedures performed were in accordance with the ethical standards of the Washington University in St. Louis Institutional Review Board and with the 1964 Helsinki declaration and its later amendments. For this type of study formal consent is not required. This article does not contain any studies with animals performed by any of the authors.

Data availability

The datasets during and/or analysed during the current study are available from the corresponding author on reasonable request.

neurosurgical procedure (TNP), and permanent neurosurgical intervention were 2.0 (95% CI 1.7–2.3), 3.0 (2.5–3.5), 24.0 (22.2–25.8), and 101.0 (90.4–111.6), respectively. Grades III and IV IVH did not differ in age at IVH diagnosis (X^2 (1 d.f.) = 1.32, p = 0.25), ventriculomegaly (X^2 = 0.73, p = 0.40), TNP (X^2 = 0.61, p = 0.43), or permanent intervention (X^2 = 2.48, p = 0.17). Ventricular reservoirs and ventriculosubgaleal shunts were used in 71 (68.3%) and 30 (28.8%), respectively. Eighty (76.9%) of the patients ultimately received a VPS. Five (4.8%) underwent a primary endoscopic third ventriculostomy (ETV), and 2 (1.9%) had ETV for a revision procedure. Four of the 7 ETVs had choroid plexus cauterization.

Conclusions—Although most infants who develop IVH and ventriculomegaly will do so within a few days of birth, at-risk infants should be observed for at least four weeks with serial head ultrasounds to monitor for PHH requiring surgery.

Keywords

intraventricular hemorrhage; ventriculomegaly; ventriculoperitoneal shunt; temporizing neurosurgical procedure

Introduction

Intraventricular hemorrhage (IVH) remains a serious challenge in the care of preterm infants and is associated with significant morbidities, including cognitive deficits, cerebral palsy, and vision and hearing impairment [1]. The incidence of IVH is well-documented, and 25–28% of infants with grade III and IV IVH develop posthemorrhagic hydrocephalus (PHH) during their initial admission [6, 27, 35, 36]. Even with appropriate intervention, children with PHH have chronic neurological deficits and poor developmental outcomes [3, 32].

Initial management of PHH frequently utilizes temporizing neurosurgical procedures (TNP) such as ventricular reservoirs (VRs) or ventriculosubgaleal shunts (VSGS) [11, 16, 26]. Permanent long-term surgical options for PHH include placement of a ventriculoperitoneal shunt (VPS) or endoscopic third ventriculostomy (ETV) with or without choroid plexus cauterization. These surgical procedures and their implementation have been well documented by a number of groups over the years [15, 39, 41].

While a number of studies have examined the timing of IVH development in preterm neonates [2, 4, 8–10, 17–19, 23, 33, 38], the few that have focused on the time-course of PHH included only a small number of subjects who developed the condition [13, 21, 25, 37]. Understanding the temporal profile of this disease may elucidate the optimal diagnosis and treatment timeframe and help improve clinical management. This study examined the timing of IVH, PHH, and surgical treatment in affected preterm infants.

Methods

Patient Population

This study was approved by the Washington University in St. Louis Human Research Protection Office and is in accordance with the 1964 Helsinki declaration and its later amendments.

Data were collected retrospectively on all preterm infants (estimated gestational age at birth [EGA] 34 weeks) admitted to St. Louis Children's Hospital (SLCH) from 1994 to 2016 with a diagnosis of IVH, and who required a neurosurgical procedure to treat PHH either during or after their NICU course. Standard neurosurgical care of preterm infants with IVH and PHH at SLCH has been detailed in a previous article [15]; though the current study and the previous report investigate different aspects of PHH management, there is some overlap in the subjects analyzed (54 of 104 subjects). Among these infants, head ultrasound (HUS) examinations were obtained in the setting of abrupt clinical change or acute event requiring resuscitation (e.g. chest compressions or epinephrine use). In infants without acute clinical status changes, HUS was performed routinely on day of life 3 to screen for IVH. Of the 104 subjects in the study, 23 (22.1%) were scanned on day 0, 45 (43.3%) on day 1, 23 (22.1%) on day 2, and 13 (12.5%) on day 3. Those found to have IVH then had HUS at least 1–2 times per week to monitor for PHH. No HUS was performed beyond what was required for clinical purposes.

All study subjects developed IVH, ventriculomegaly, and required surgical treatment for PHH. Ventriculomegaly was defined by semi-quantitative assessment of ventricular size and frontal-occipital horn ratio (FOR) ≥ 0.5 [41]. PHH was defined by clinical criteria including progressive increase in occipitofrontal circumference, full fontanel (palpated above the level of surrounding bone), and splaying of the sagittal suture ≥ 2 mm in the mid-parietal region [20, 22, 40, 41]. All patients displaying these parameters underwent neurosurgical intervention at earliest possible date; therefore, TNP was used as a proxy for PHH diagnosis. Initial neurosurgical intervention was with VR or VSGS, depending on the surgeon's training or preference.

Data Collection

Primary time-points of interest included the dates of initial IVH diagnosis, initial ventriculomegaly diagnosis, PHH diagnosis or TNP, permanent neurosurgical intervention (VPS or ETV), and first revision. The Papile system was used to assign IVH grade [24]. FOR was calculated at three time points (first imaging study, ventriculomegaly diagnosis, and first procedure) for the 75 subjects with imaging available in the electronic medical record. Rates of known risk factors for shunt complications including placement of a tracheostomy or gastrostomy tube, surgery for necrotizing enterocolitis or intestinal perforation, and surgery for patent ductus arteriosus (PDA) were collected [30, 31, 34]. Additional variables queried included sex, race, birthweight, EGA at birth, and initial procedure type (VR or VSGS). Volume of cerebrospinal fluid (CSF) removed during the temporization phase was not routinely recorded in the medical records and was thus not included in the current study.

Statistical Analysis

Statistical analyses were performed using SPSS version 24 (IBM Corporation, NY, USA). Baseline characteristics were summarized as percentage of the total cohort, mean with standard deviation for normally distributed variables, and median with interquartile range for other numerical variables. Kaplan-Meier (KM) survival curves were constructed for time from birth to IVH, ventriculomegaly, TNP, VPS or ETV, and first revision. Significant

differences between curves were detected using the log rank test. Cox proportional hazards regression was used to test effects of continuous predictors on time to revision. Differences between FOR at the three time points were determined with repeated measures ANOVA and post-hoc pairwise comparisons with Bonferroni adjustment for multiple comparisons. All tests of significance were conducted at a predetermined alpha level of 0.05.

Results

Baseline Characteristics

All 104 preterm infants who met the inclusion criteria were included in the analysis. Baseline characteristics for the subjects are included in Table 1. Duration of follow-up ranged from 11.1 to 202.2 months. Highest documented IVH grade was II for 5 patients (4.8%), III for 33 (31.7%), and IV for 66 (63.5%). Representative HUS for each of the IVH grades are shown in Figure 1, and characteristics of the IVH on HUS including size and location are summarized in Table 2. All subjects developed ventriculomegaly and underwent neurosurgical intervention for PHH. Seventy-one (68.3%) subjects were treated initially with a VR and 30 (28.8%) with a VSGS. Eighty (76.9%) of the patients ultimately received a VPS during the entire follow-up duration. Two (2.5%) of the shunted patients experienced an infection of their primary shunt (described in Table 3). Thirty-six (45.0%) of the shunted patients required at least one revision within six months of shunt insertion, and 53 (66.3%) required a revision during the entire follow-up duration.

Changes in Frontal-Occipital Horn Ratio

FOR measured at the time of each subject's first ever imaging study (mean $\pm$ standard deviation, 0.456 ± 0.058), ventriculomegaly diagnosis (0.517 ± 0.054), and TNP (0.647 ± 0.061) are summarized in Figure 2. The FOR values were significantly different on repeated measures ANOVA ($F(2 \text{ d.f.}) = 235.52$, $p < 0.001$), and they increased significantly between each of the three time-points (adjusted $p < 0.001$ for all post-hoc comparisons).

Time-course for Intraventricular Hemorrhage

KM survival curves showing time from birth to development of IVH are displayed in the top row of Figure 3. Curves are shown for the entire pooled cohort (left), as well as broken down by grade for those with grades III and IV IVH (right). Median time to IVH for the pooled cohort was 2.0 (95% CI 1.7–2.3) days. No significant difference was found in age at IVH ($\chi^2 (1 \text{ d.f.}) = 1.32$, $p = 0.25$) between patients with grades III (median 2.0 [95% CI 1.4–2.6] days) or IV IVH (2.0 [1.7–2.3] days).

Time-course for Ventriculomegaly

KM survival curves showing time from birth to development of ventriculomegaly are displayed in the middle row of Figure 3. Median time to ventriculomegaly for the pooled cohort was 3.0 (95% CI 2.5–3.5) days. No significant difference was found in age at ventriculomegaly ($\chi^2 (1 \text{ d.f.}) = 0.73$, $p = 0.40$) between patients with grades III (median 3.0 [95% CI 1.6–4.4] days) or IV IVH (2.0 [1.5–2.5] days).

Time-course for Temporizing Neurosurgical Procedure

Median postmenstrual age (PMA) at TNP was 29.6 (interquartile range, IQR 28.1–31.6) weeks for the pooled cohort. KM survival curves showing time from birth to TNP are displayed in the bottom row of Figure 3. Median time to intervention for the pooled cohort was 24.0 (95% CI 22.2–25.8) days. No significant difference was found in age at intervention (χ^2 (1 d.f.) = 0.61, $p = 0.43$) between patients with grades III (median 23.0 [95% CI 20.2–25.8] days) or IV IVH (24.0 [22.0–26.0] days).

Time-course for Permanent Neurosurgical Procedure

Among subjects with grade III IVH, 23 of 33 (69.7%) required a permanent neurosurgical intervention (VPS or ETV); among subjects with grade IV IVH, 54 of 66 (81.8%) required permanent intervention. Rates of permanent interventions were not significantly different between grades III and IV IVH (χ^2 (1 d.f.) = 1.87, $p = 0.17$). Five (4.8%) of the patients underwent an ETV for their first permanent neurosurgical intervention, and 2 (1.9%) had ETV for a revision procedure. Four of the 7 ETVs also had choroid plexus cauterization.

Median PMA at permanent neurosurgical intervention was 38.4 (IQR 36.7–42.9) weeks for the 80 patients who received a VPS or ETV. KM survival curves showing time from birth to permanent intervention are displayed in Figure 4. Curves are shown for all 104 subjects (left), as well as broken down by grade for those with grades III and IV IVH (right). Censoring is represented by tick marks for patients who had not yet received a VPS or ETV by the end of the study period, and for 1 patient who died from non-neurological causes. Median time from birth to permanent intervention for all 104 subjects was 101.0 (95% CI 90.4–111.6) days. There was no significant difference in age at permanent intervention (χ^2 (1 d.f.) = 2.48, $p = 0.17$) between patients with grades III (median 118.0 [95% CI 62.9–173.1] days) or IV IVH (95.0 [81.5–108.5] days). None of the patient characteristics shown in Table 1 were significantly predictive of time to VPS or ETV by log rank test or unadjusted Cox regression.

Time-course for First Revision

Median PMA at the time of first revision procedure (VPS or ETV) was 52.8 (IQR 43.7–95.3) weeks for the 54 patients who required revisions. KM survival curves showing time from initial permanent intervention to first revision are displayed in Figure 5. Curves are shown for all 80 patients who required a VPS or ETV (left), as well as broken down by grade for those with grades III and IV IVH (right). Censoring is represented by tick marks for patients who had not yet had a revision by the end of the study period. Median time from initial permanent intervention to revision for the 80 patients was 50.4 (95% CI 17.9–82.9) weeks. There was no significant difference in time to first revision (χ^2 (1 d.f.) = 0.58, $p = 0.45$) between patients with grades III (median 57.6 [95% CI 5.5–109.6] weeks) or IV IVH (22.9 [0.0–53.7] weeks). None of the patient characteristics or risk factors shown in Table 1 were significantly predictive of time to first shunt revision by log rank test or unadjusted Cox regression.

Discussion

The goal of this study was to report the time-course of development of IVH, ventriculomegaly, and PHH requiring surgery in preterm infants who ultimately underwent surgical intervention, as well as our institution's experience with timing of temporary and permanent CSF diversion. While a number of studies have examined the timing of IVH development in preterm neonates, few have detailed the other endpoints in a sample of this size. In general, we found that patients who received subsequent surgical intervention developed IVH, ventriculomegaly, and PHH by day of life 2, 3, and 24, respectively. There were no differences in IVH or PHH timing between IVH grades III and IV in our study population. These findings suggest that although most infants who develop IVH and ventriculomegaly will do so within a few days of birth, at-risk infants should be observed for at least four weeks with serial HUS to monitor for development of PHH requiring surgery.

Our findings are in agreement with existing literature on IVH and PHH timing. A 2014 meta-analysis by Al-Abdi and Al-Aamri incorporated 5 prospective studies and found two major modes of IVH timing in 279 cases—0–6 hours (48% of cases) and after 24 hours of life (38%) [2, 4, 8, 10, 17, 18]. Although our retrospective review was unable to provide more resolution within the first day of life due to the hours when routine HUS were available, we showed that the majority of IVH occurs within the first 2 days, a finding that is consistent with previous studies [38]. Two studies have touched on timing of ventriculomegaly associated with IVH. In 1983, Partridge et al. reported PHH in 18 patients, with 6% diagnosed by day 1, 39% by day 4, 56% by day 7, 83% by day 14, and 94% by day 28 [25]. In 1984, Szymonowicz and Yu performed HUS every 12 hours until 72 hours of age in infants with IVH, and weekly thereafter until discharge or death. They reported the optimal timing for diagnosis of ventriculomegaly at two to three weeks, although they only had 5 cases of ventricular dilatation in 30 patients with IVH [37]. In our cohort of infants ultimately receiving surgical intervention, we found a more rapid course of ventriculomegaly with median time-to-event of 3 days. We agree that diagnosis of ventriculomegaly is likely to be made within the first weeks of life.

We used TNP as a proxy for PHH, and found that this develops three weeks after IVH. Few papers have focused on timing of temporary or permanent CSF diversion in infants with PHH. Wang et al. reported VR insertion at mean 33 days of life (31.8 weeks EGA, $n = 44$) and VSGS insertion at mean 29 days of life (30.1 weeks EGA, $n = 46$) [39]. They also described VPS placement at mean 81.7 and 109.5 days of life for patients with VR and VSGS, respectively. Likewise, Karas et al. reported VSGS at median 25 days of life (30.3 weeks EGA, $n = 10$) and VPS placement at 61 days of life ($n = 9$) [12]. Christian et al. found in 91 patients that those receiving a VR or VSGS as initial procedure did so at a mean age of 29 days, and initial permanent shunt by 56 days [7]. Our results are consistent with these studies (median 24 days for VR and VSGS). Time from initial permanent intervention to first revision varies across the literature from 59 to 156 weeks [5, 29]. We report a median time of 50.4 (95% CI 17.9–82.9) weeks, which is consistent with previous studies.

FOR was first described as a ratio of linear measures of ventricular size that accounts for disproportionate occipital horn enlargement in pediatric hydrocephalus [22] and has been

used in a number of recent studies due to its ability to approximate ventricular volumes [14, 28, 30]. Compared to FOR at time of first ever image, FOR is significantly larger at time of ventriculomegaly diagnosis and is even larger at TNP, which provides evidence of the concurrent validity of this measure for distinguishing between patients at various stages of the PHH disease course.

A primary limitation of this study is that our database only contained patients who developed IVH and PHH requiring neurosurgical intervention. Therefore, our results are not generalizable to the population of all preterm infants with IVH or ventriculomegaly, but only to the more severe subset of this group that is managed by neurosurgeons. Patients who have lower grade IVH that does not progress to PHH may be expected to have a more prolonged or 'arrested' time-course. Additionally, because the timing and frequency of HUS were not standardized in this retrospective study, the time resolution of endpoints is limited. A larger, prospective study examining the incidence of ventriculomegaly and PHH in the broader population of all IVH patients (grades I–IV) would facilitate clinical decision making at the onset of disease. However, the timings described in this study may be used to inform expectations regarding the time-course of high grade IVH and PHH. Finally, this single institution series only captures the experience at SLCH, and a multi-institutional study on PHH timing would produce more generalizable results.

In conclusion, we report the typical time-course of development of IVH, ventriculomegaly, PHH, VPS or ETV, and first revision in preterm infants who ultimately require neurosurgery. We also show a significantly increasing trend in FOR between first imaging study, ventriculomegaly diagnosis, and TNP. Our data suggest that at-risk infants should be observed for at least four weeks with serial HUS to monitor for development of PHH. Future studies should aim to generalize our results to all preterm infants who develop IVH, as well as validate the findings at other institutions.

Acknowledgments

Funding

This publication was made possible by the Washington University Institute of Clinical and Translational Sciences (National Institutes of Health/National Center for Advancing Translational Sciences UL1 TR000448, TL1 TR000449, and UL1 RR024992), the Washington University Intellectual and Developmental Disabilities Research Center (National Institutes of Health/*Eunice Kennedy Shriver* National Institute of Child Health and Human Development U54 HD087011), and St. Louis Children's Hospital.

References

1. Adams-Chapman I, Hansen NI, Stoll BJ, Higgins R. Neurodevelopmental outcome of extremely low birth weight infants with posthemorrhagic hydrocephalus requiring shunt insertion. *Pediatrics*. 2008; 121:e1167–77. DOI: 10.1542/peds.2007-0423 [PubMed: 18390958]
2. Al-Abdi SY, Al-Aamri MA. A Systematic Review and Meta-analysis of the Timing of Early Intraventricular Hemorrhage in Preterm Neonates: Clinical and Research Implications. *J Clin Neonatol*. 2014; 3:76–88. DOI: 10.4103/2249-4847.134674 [PubMed: 25024973]
3. Badhiwala JH, Hong CJ, Nassiri F, Hong BY, Riva-Cambrin J, Kulkarni AV. Treatment of posthemorrhagic ventricular dilation in preterm infants: a systematic review and meta-analysis of outcomes and complications. *J Neurosurg Pediatr*. 2015; 16:545–555. DOI: 10.3171/2015.3.PEDS14630

4. Beverley DW, Chance GW, Coates CF. Intraventricular haemorrhage--timing of occurrence and relationship to perinatal events. *Br J Obstet Gynaecol.* 1984; 91:1007–1013. [PubMed: 6435667]
5. Chittiboina P, Pasieka H, Sonig A, Bollam P, Notarianni C, Willis BK, Nanda A. Posthemorrhagic hydrocephalus and shunts: what are the predictors of multiple revision surgeries? *J Neurosurg Pediatr.* 2013; 11:37–42. DOI: 10.3171/2012.8.PEDS11296 [PubMed: 23101558]
6. Christian EA, Jin DL, Attenello F, Wen T, Cen S, Mack WJ, Krieger MD, McComb JG. Trends in hospitalization of preterm infants with intraventricular hemorrhage and hydrocephalus in the United States, 2000–2010. *J Neurosurg Pediatr.* 2016; 17:260–269. DOI: 10.3171/2015.7.PEDS15140 [PubMed: 26544084]
7. Christian EA, Melamed EF, Peck E, Krieger MD, McComb JG. Surgical management of hydrocephalus secondary to intraventricular hemorrhage in the preterm infant. *J Neurosurg Pediatr.* 2016; 17:278–84. DOI: 10.3171/2015.6.PEDS15132 [PubMed: 26565942]
8. de Crespigny LC, Mackay R, Murton LJ, Roy RN, Robinson PH. Timing of neonatal cerebroventricular haemorrhage with ultrasound. *Arch Dis Child.* 1982; 57:231–233. DOI: 10.1136/adsc.57.10.811-a [PubMed: 7073304]
9. Dani C, Bertini G, Pezzati M, Poggi C, Guerrini P, Martano C, Rubaltelli FF. Prophylactic ibuprofen for the prevention of intraventricular hemorrhage among preterm infants: a multicenter, randomized study. *Pediatrics.* 2005; 115:1529–35. DOI: 10.1542/peds.2004-1178 [PubMed: 15930213]
10. Dolfin T, Skidmore MB, Fong KW, Hoskins EM, Shennan AT. Incidence, severity, and timing of subependymal and intraventricular hemorrhages in preterm infants born in a perinatal unit as detected by serial real-time ultrasound. *Pediatrics.* 1983; 71:541–546. [PubMed: 6835737]
11. Fulmer BB, Grabb PA, Oakes WJ, Mapstone TB, Rekate HL. Neonatal ventriculosubgaleal shunts. *Neurosurgery.* 2000; 47:80–84. DOI: 10.1097/00006123-200007000-00018 [PubMed: 10917350]
12. Karas CS, Baig MN, Elton SW. Ventriculosubgaleal shunts at Columbus Children's Hospital: Neurosurgical implant placement in the neonatal intensive care unit. *J Neurosurg.* 2007; 107:220–223. DOI: 10.3171/PED-07/09/220
13. Korobkin R. The relationship between head circumference and the development of communicating hydrocephalus in infants following intraventricular hemorrhage. *Pediatrics.* 1975; 56:74–77. [PubMed: 1080558]
14. Limbrick DD, Baksh B, Morgan CD, Habiyaemye G, McAllister JP, Inder TE, Mercer D, Holtzman DM, Strahle J, Wallendorf MJ, Morales DM. Cerebrospinal fluid biomarkers of infantile congenital hydrocephalus. *PLoS One.* 2017; 12:1–20. DOI: 10.1371/journal.pone.0172353
15. Limbrick DD, Mathur A, Johnston JM, Munro R, Sagar J, Inder T, Park TS, Leonard JL, Smyth MD. Neurosurgical treatment of progressive posthemorrhagic ventricular dilation in preterm infants: a 10-year single-institution study. *J Neurosurg Pediatr.* 2010; 6:224–30. DOI: 10.3171/2010.5.PEDS1010 [PubMed: 20809705]
16. McComb JG, Ramos AD, Platzker CG, Henderson DJ, Segall HD. Management of hydrocephalus secondary to intraventricular hemorrhage in the preterm infant with a subcutaneous ventricular catheter reservoir. *Neurosurgery.* 1983; 13:295–300. DOI: 10.1227/00006123-198309000-00014 [PubMed: 6621842]
17. McDonald MM, Koops BL, Johnson ML, Guggenheim MA, Rumack CM, Mitchell SA, Hathaway WE. Timing and antecedents of intracranial hemorrhage in the newborn. *Pediatrics.* 1984; 74:32–36. [PubMed: 6739217]
18. Meidell R, Marinelli P, Pettett G. Perinatal factors associated with early-onset intracranial hemorrhage in premature infants. A prospective study. *Am J Dis Child.* 1985; 139:160–3. [PubMed: 3883746]
19. Ment LR, Oh W, Ehrenkranz RA, Philip AGS, Duncan CC, Makuch RW. Antenatal steroids, delivery mode, and intraventricular hemorrhage in preterm infants. *Am J Obstet Gynecol.* 1995; 172:795–800. DOI: 10.1016/0002-9378(95)90001-2 [PubMed: 7892866]
20. Morales DM, Holubkov R, Inder TE, Ahn HC, Mercer D, Rao R, McAllister JP, Holtzman DM, Limbrick DD. Cerebrospinal fluid levels of amyloid precursor protein are associated with ventricular size in post-hemorrhagic hydrocephalus of prematurity. *PLoS One.* 2015; 10:e0115045.doi: 10.1371/journal.pone.0115045 [PubMed: 25738507]

21. Murphy BP, Inder TE, Rooks V, Taylor GA, Anderson NJ, Mogridge N, Horwood LJ, Volpe JJ. Posthaemorrhagic ventricular dilatation in the premature infant: natural history and predictors of outcome. *Arch Dis Child Fetal Neonatal Ed.* 2002; 87:37–41. DOI: 10.1136/fn.87.1.F37
22. O'Hayon BB, Drake JM, Ossip MG, Tuli S, Clarke M. Frontal and occipital horn ratio: A linear estimate of ventricular size for multiple imaging modalities in pediatric hydrocephalus. *Pediatr Neurosurg.* 1998; 29:245–249. DOI: 10.1159/000028730 [PubMed: 9917541]
23. Paneth N, Pinto-Martin J, Gardiner J, Wallenstein S, Katsikiotis V, Hegyi T, Hiatt IM, Susser M. Incidence and timing of germinal matrix/intraventricular hemorrhage in low birth weight infants. *Am J Epidemiol.* 1993; 137:1167–76. [PubMed: 8322758]
24. Papile LA, Burstein J, Burstein R, Koffler H. Incidence and evolution of subependymal and intraventricular hemorrhage: A study of infants with birth weights less than 1,500 gm. *J Pediatr.* 1978; 92:529–534. DOI: 10.1016/S0022-3476(78)80282-0 [PubMed: 305471]
25. Partridge JC, Babcock DS, Steichen JJ, Han BK. Optimal timing for diagnostic cranial ultrasound in low-birth-weight infants: Detection of intracranial hemorrhage and ventricular dilation. *J Pediatr.* 1983; 102:281–287. DOI: 10.1016/S0022-3476(83)80544-7 [PubMed: 6822939]
26. Perret GE, Graf CJ. Subgaleal shunt for temporary ventricle decompression and subdural drainage. *J Neurosurg.* 1977; 47:590–595. DOI: 10.3171/jns.1977.47.4.0590 [PubMed: 903811]
27. du Plessis AJ. The role of systemic hemodynamic disturbances in prematurity-related brain injury. *J Child Neurol.* 2009; 24:1127–40. DOI: 10.1177/0883073809339361 [PubMed: 19745087]
28. Ragan DK, Cerqua J, Nash T, McKinstry RC, Shimony JS, Jones BV, Mangano FT, Holland SK, Yuan W, Limbrick DD. The accuracy of linear indices of ventricular volume in pediatric hydrocephalus: technical note. *J Neurosurg Pediatr.* 2015; 15:547–51. DOI: 10.3171/2014.10.PEDS14209 [PubMed: 25745953]
29. Reinprecht A, Dietrich W, Berger A, Bavinszki G, Weninger M, Czech T. Posthemorrhagic hydrocephalus in preterm infants: Long-term follow-up and shunt-related complications. *Child's Nerv Syst.* 2001; 17:663–669. DOI: 10.1007/s00381-001-0519-2 [PubMed: 11734984]
30. Riva-Cambrin J, Kestle JRW, Holubkov R, Butler J, Kulkarni AV, Drake J, Whitehead WE, Wellons JC, Shannon CN, Tamber MS, Limbrick DD, Rozzelle C, Browd SR, Simon TD. Risk factors for shunt malfunction in pediatric hydrocephalus: a multicenter prospective cohort study. *J Neurosurg Pediatr.* 2015; :1–9. DOI: 10.3171/2015.6.PEDS14670
31. Riva-Cambrin J, Shannon CN, Holubkov R, Whitehead WE, Kulkarni AV, Drake J, Simon TD, Browd SR, Kestle JRW, Wellons JC. Center effect and other factors influencing temporization and shunting of cerebrospinal fluid in preterm infants with intraventricular hemorrhage. *J Neurosurg Pediatr.* 2012; 9:473–81. DOI: 10.3171/2012.1.PEDS11292 [PubMed: 22546024]
32. Robinson S. Neonatal posthemorrhagic hydrocephalus from prematurity: pathophysiology and current treatment concepts. *J Neurosurg Pediatr.* 2012; 9:242–258. DOI: 10.3171/2011.12.PEDS11136 [PubMed: 22380952]
33. Sarkar S, Kaplan C, Wiswell TE, Spitzer AR. Histological chorioamnionitis and the risk of early intraventricular hemorrhage in infants born < or =28 weeks gestation. *J Perinatol.* 2005; 25:749–52. DOI: 10.1038/sj.jp.7211399 [PubMed: 16237461]
34. Simon TD, Butler J, Whitlock KB, Browd SR, Holubkov R, Kestle JRW, Kulkarni AV, Langley M, Limbrick DD, Mayer-Hamblett N, Tamber M, Wellons JC, Whitehead WE, Riva-Cambrin J. Risk factors for first cerebrospinal fluid shunt infection: findings from a multi-center prospective cohort study. *J Pediatr.* 2014; 164:1462–8. e2. DOI: 10.1016/j.jpeds.2014.02.013 [PubMed: 24661340]
35. Stoll BJ, Hansen NI, Bell EF, Shankaran S, Laptook AR, Walsh MC, Hale EC, Newman NS, Schibler K, Carlo WA, Kennedy KA, Poindexter BB, Finer NN, Ehrenkranz RA, Duara S, Sánchez PJ, O'Shea TM, Goldberg RN, Van Meurs KP, Faix RG, Phelps DL, Frantz ID, Watterberg KL, Saha S, Das A, Higgins RD, Hack M, Horbar J, Malloy M, Tyson J, Wright E, Wright L, Hack M, Wright L, Shankaran S, Fanaroff A, Wright L, Stevenson D, Stevenson D, Wright L, Lemons J, Lemons J, Bauer C, Oh W, Fanaroff A, Stoll B, Wright L, Muglia L, Katz M, Iams J, Romero R, Culhane J, Goldenberg R, Alexander G, Himes J, Kaufman R, Mor J, Kogan M, Stoll B, Hansen N, Fanaroff A, Stoll B, Hansen N, Higgins R, Jobe A, Bancalari E, Walsh M, Yao Q, Gettner P, Zou G, Horbar J, Badger G, Carpenter J, Chan K, Ohlsson A, Synnes A, Meadow W, Lee G, Lin K, Lantos J, Group E, Vohr B, Wright L, Dusick A, Lee S, Chan K, Ohlsson A, Walsh M, Laptook A, Kazzi S. Eunice Kennedy Shriver National Institute of Child Health and Human Development

Neonatal Research Network for the EKSNI of CH and HDNR, EpicS Group, Network NI of CH and HDNR, Network CN. Neonatal outcomes of extremely preterm infants from the NICHD Neonatal Research Network. *Pediatrics*. 2010; 126:443–56. DOI: 10.1542/peds.2009-2959 [PubMed: 20732945]

36. Stoll BJ, Hansen NI, Bell EF, Walsh MC, Carlo WA, Shankaran S, Laptook AR, Sánchez PJ, Van Meurs KP, Wyckoff M, Das A, Hale EC, Ball MB, Newman NS, Schibler K, Poindexter BB, Kennedy KA, Cotten CM, Watterberg KL, D'Angio CT, DeMauro SB, Truog WE, Devaskar U, Higgins RD. Network EKSNI of CH and HDNR. Trends in Care Practices, Morbidity, and Mortality of Extremely Preterm Neonates, 1993–2012. *JAMA*. 2015; 314:1039–1051. DOI: 10.1001/jama.2015.10244 [PubMed: 26348753]
37. Szymonowicz W, Yu VY, Wilson FE. Antecedents of periventricular haemorrhage in infants weighing 1250 g or less at birth. *Arch Dis Child*. 1984; 59:13–17. DOI: 10.1136/adc.59.1.13 [PubMed: 6696488]
38. Tsiantos A, Victorin L, Relier JP, Dyer N, Sundell H, Brill AB, Stahlman M. Intracranial hemorrhage in the prematurely born infant. Timing of clots and evaluation of clinical signs and symptoms. *J Pediatr*. 1974; 85:854–859. DOI: 10.1016/S0022-3476(74)80360-4 [PubMed: 4417550]
39. Wang JY, Amin AG, Jallo GI, Ahn ES. Ventricular reservoir versus ventriculosubgaleal shunt for posthemorrhagic hydrocephalus in preterm infants: infection risks and ventriculoperitoneal shunt rate. *J Neurosurg Pediatr*. 2014; 14:447–454. DOI: 10.3171/2014.7.PEDS13552 [PubMed: 25148212]
40. Wellons JC, Holubkov R, Browd SR, Riva-Cambrin J, Whitehead W, Kestle J, Kulkarni AV. The assessment of bulging fontanel and splitting of sutures in premature infants: an interrater reliability study by the Hydrocephalus Clinical Research Network. *J Neurosurg Pediatr*. 2013; 11:12–4. DOI: 10.3171/2012.10.PEDS12329 [PubMed: 23121114]
41. Wellons JC, Shannon CN, Holubkov R, Riva-Cambrin J, Kulkarni AV, Limbrick DD, Whitehead W, Browd S, Rozzelle C, Simon TD, Tamber MS, Oakes WJ, Drake J, Luerssen TG, Kestle J. Hydrocephalus Clinical Research Network. Shunting outcomes in posthemorrhagic hydrocephalus: results of a Hydrocephalus Clinical Research Network prospective cohort study. *J Neurosurg Pediatr*. 2017; :1–11. DOI: 10.3171/2017.1.PEDS16496

Fig. 1. Coronal (left) and sagittal (right) head ultrasound screen captures from patients with grade II (top row), III (middle row), and IV (bottom row) IVH, respectively.

Fig. 2. Boxplots of FOR measured at the time of each subject's first ever imaging study, ventriculomegaly diagnosis, and TNP. Boxes indicate interquartile ranges; lines inside boxes represent medians; whiskers at 5 to 95 percentiles

Fig. 3.

KM survival curves showing time from birth to IVH diagnosis (top row), ventriculomegaly diagnosis (middle row), and TNP (bottom row). Data are shown for all 104 preterm infants (left column) and separated by IVH grade for 33 infants with grade III and 66 infants with grade IV IVH (right column). All subjects experienced the event, and none were censored

Fig. 4.

KM survival curves showing time from birth to permanent neurosurgical intervention. Data are shown for all 104 preterm infants (left) and separated by IVH grade for 33 infants with grade III and 66 infants with grade IV IVH (right). Tick marks indicate censoring due to end of study period or death (1 subject)

Fig. 5.

KM survival curves showing time from initial permanent intervention to first revision. Data are shown for all 80 preterm infants who required a VPS or ETV (left) and separated by IVH grade for 23 infants with grade III and 54 infants with grade IV IVH (right). Tick marks indicate censoring due to end of study period or death

Table 1Baseline characteristics, all subjects ($N = 104$)

Characteristic	Count (%)	Values ^a
Sex		
Male	58 (55.8)	
Female	46 (44.2)	
Race		
Caucasian	49 (47.1)	
African American	50 (48.1)	
Other	5 (4.8)	
Birthweight (g)	104 (100)	860 (700–1166)
EGA at birth (weeks)	104 (100)	26 (24–28)
Highest IVH grade		
II	5 (4.8)	
III	33 (31.7)	
IV	66 (63.5)	
Temporizing procedure type		
Ventricular reservoir	71 (68.3)	
Ventriculosubgaleal shunt	30 (28.8)	
PMA at temporizing procedure (weeks)	101 (97.1)	30 (28–32)
All ventriculoperitoneal shunts	80 (76.9)	
Shunt revisions within six months ($N = 80$ shunts)	36 (45.0 ^b)	
Shunt infection ^c ($N = 80$ shunts)	2 (2.5 ^b)	
PMA at initial shunt insertion (weeks)	80 (76.9)	39 (37–43)
Risk factors		
Patent ductus arteriosus	39 (37.5)	
Tracheostomy	4 (3.8)	
Gastrostomy tube	24 (23.1)	
Intestinal perforation	13 (12.5)	
Necrotizing enterocolitis	7 (6.7)	
Frontal-occipital horn ratio		
First imaging study	74 (71.2)	0.456 ± 0.058
Ventriculomegaly	75 (72.1)	0.517 ± 0.054
Temporizing procedure	75 (72.1)	0.647 ± 0.061

^aSummary statistics are median (interquartile range) for birthweight and EGA, and mean ± standard deviation for frontal-occipital horn ratio measurements.

^bPercentage out of 80 shunts.

^cInfections of first ever shunt prior to any revisions.

EGA, estimated gestational age; IVH, intraventricular hemorrhage; g, grams

Table 2IVH characteristics on neonatal head ultrasound ($N = 74$)^a

Characteristic	Left hemisphere	Right hemisphere
IVH grade ^b		
I	1 (1.4)	3 (4.1)
II	9 (12.2)	11 (14.9)
III	30 (40.5)	31 (41.9)
IV	34 (45.9)	26 (35.1)
Location		
Frontal horn	40 (54.1)	32 (43.2)
Body	65 (87.8)	59 (79.7)
Atrium	62 (83.8)	56 (75.7)
Occipital horn	35 (47.3)	34 (45.9)
Temporal horn	17 (23.0)	14 (18.9)
Maximum coronal dimension (cm)	0.95 ± 0.41	0.86 ± 0.45
Maximum sagittal dimension (cm)	1.05 ± 0.34	0.95 ± 0.43

^aValues are count (%) for IVH grade and location, and mean ± standard deviation for dimensions.^bHighest IVH grade was II or above for all subjects.

IVH, intraventricular hemorrhage; cm, centimeters

Table 3

Characteristics of the two patients who had infections of their primary shunt

Characteristic	Patient 1	Patient 2
Sex	Male	Male
Race	Caucasian	Caucasian
Birthweight (g)	1275	1230
EGA at birth (weeks)	29	28
Highest IVH grade	3	4
Initial procedure type	VR	VSGS
EGA at initial procedure (weeks)	33	31
Risk factors		
Patent ductus arteriosus	Yes	Yes
Tracheostomy	No	No
Gastrostomy tube	Yes	No
Intestinal perforation	No	No
Necrotizing enterocolitis	No	No
Loculation	No	No
Birth to IVH (days)	2	2
Birth to ventriculomegaly (days)	2	2
Birth to first procedure (days)	25	20
Birth to initial shunt insertion (days)	104	67
EGA at initial shunt insertion (weeks)	44	38
Initial shunt insertion to infection (days)	63	28
Organism cultured	Coagulase- negative staphylococcus	<i>S. aureus</i>

EGA, estimated gestational age; IVH, intraventricular hemorrhage; g, grams