CASE REPORT

Interrupted aortic arch complicated with takotsubo cardiomyopathy mimicking aortic dissection

Farhala Mari Baloch,¹ Javed Majid Tai,¹ Aamir Hameed Khan,² Abdul Baqi²

¹Department of Cardiology,
Aga Khan University Hospital,
Karachi, Pakistan

²Department of Cardiology,
Aga Khan University Hospital,
Karachi, Pakistan

Correspondence to
Farhala Mari Baloch,
farhala.baloch@aku.edu

Accepted 26 September 2017

SUMMARY

A 50-year-old man presented to the emergency department with interscapular pain, diaphoresis and restlessness. Initial examination raised the possibility of aortic dissection; however, the CT scan did not concur with the diagnosis. An ECG showed ST segment elevation in leads V1–V6 and echocardiography showed severe left ventricular systolic dysfunction. Coronary angiography through the right femoral artery was attempted but the diagnostic catheter could not be advanced to the ascending aorta. Radiocontrast injection showed complete obstruction of the descending aorta. Coronary angiography through right radial approach showed mild left anterior descending disease. The aortogram showed complete interruption of the ascending aorta with extensive collateral network. Left ventricle gram was consistent with stress-induced cardiomyopathy. We noticed intermittent confusion and agitation. MRI of the brain showed areas of deep white matter ischaemia as well as microhaemorrhages, suggesting posterior reversible leucoencephalopathy syndrome. He unfortunately went into cardiac arrest and could not be revived.

BACKGROUND

This case is an example of an unusual presentation of rare congenital anomalies in an uncommon age group. Approximate incidence of interrupted aortic arch (IAA) is reported to be 3/1 000 000 live births.¹ It is usually associated with other congenital heart defects such as ventricular septal defect, patent ductus arteriosus (PDA) and truncus arteriosus.¹ Very rarely, it is an isolated anomaly as in this case. It is the first cardiac anomaly which demonstrates a genetic pattern which is homozygous deletion 1.5–3 Mb region of chromosome band 22q11.^{2,3}

In spite of being labelled as hypertensive in young age, unfortunately, our patient was not investigated for the cause of hypertension in his age group. Anticipating rare causes of common disease may have an impact on the outcome of the patient. IAA is a very rare cause of hypertension in adult population. It may be diagnosed early in young adults by anticipating on physical examination. Young patients presenting with uncontrolled hypertension should have a complete examination of the upper and lower limb pulses and blood pressure in all the four limbs. The diagnosis can be subsequently confirmed by relevant imaging.

CASE PRESENTATION

A 50-year-old man was brought in to the emergency department with a history of tearing interscapular pain, diaphoresis and being restless for 20 hours. He was initially taken to a local hospital where his blood pressure was found to be significantly elevated. He was treated as a case of hypertensive urgency. A plain CT scan brain was done to look for evidence of an acute cerebral insult, suspected due to his significantly elevated blood pressure and being restless but showed no abnormality. His medical history was significant for hypertension since the age of 16 years. However, he was not compliant with his antihypertensive medications.

On examination, he was restless and diaphoretic but oriented to time, place and person. He attributed his restlessness to chest pain. The pulse was 100 beats per min, regular and low volume in the right arm. Blood pressure in his left arm was 160/96 mm Hg and 120/80 mm Hg in the right arm. His lower extremities were cold and peripheral pulses were not palpable. Clinical examination of his chest and heart was unremarkable. He was electively intubated for further management.

INVESTIGATIONS

His first ECG showed ST segment elevation in leads V1–V6 but with no reciprocal changes (figure 1). With a presumed initial clinical diagnosis of aortic dissection due to the interarm blood pressure difference and sharp interscapular pain, an urgent bedside transthoracic echocardiography was performed which revealed severe left ventricular systolic dysfunction (visual estimation, basal segments were showing normal contraction, while all the mid and apical segments were akinetic) (see videos 1A and 1B in the online supplementary files 1 and 2, respectively); however, a dissection flap could not be appreciated. An urgent contrast CT of the chest was performed and did not show features of dissection. Lab investigation showed a white cell count of $25 \times 10^9/L$ and troponin-I value of 2.5 ng/mL. A repeat 12-lead ECG showed new T-wave inversions in leads V1–V6 (figure 2). On the basis of dynamic ECG changes, positive cardiac biomarkers and new left ventricular systolic dysfunction, we decided to proceed to coronary angiography to rule out coronary artery disease. Femoral access was used. The J-tipped guide wire could not be passed beyond the aortic arch. A radiocontrast injection showed complete occlusion of proximal segment of descending aorta (figure 3). The right femoral

CrossMark

To cite: Mari Baloch F, Tai JM, Hameed Khan A, et al. *BMJ Case Rep* Published Online First: [please include Day Month Year]. doi:10.1136/bcr-2017-219612

Figure 1 12-lead ECG showing ST segment elevation in leads V1–V6.

approach was switched to right radial approach. His aortogram with pigtail 6F catheter revealed complete occlusion of the aorta after origin of left subclavian artery associated with large collaterals arising from the arch of the aorta and subclavian artery and connecting to the descending aorta (figures 4 and 5). His coronary angiogram revealed mild disease in mid left anterior descending artery. The left ventricular gram findings were consistent with stress-induced cardiomyopathy (CMP). The repeat transthoracic echo after 72 hours showed normal left ventricular systolic function which further confirmed his diagnosis of takotsubo CMP (see video 2 in the online supplementary file 3). During his wean from sedation, while he was kept off sedation, we noticed intermittent confusion and agitation in his behaviour which was not explicable with his metabolic and medication profile. Neurology advised an MRI brain which showed focal area of diffusion restriction at the corpus callosum and deep white matter along the occipital horn of right lateral ventricle (figure 6a–d). The sequential weighted images showed evidence of signal dropout in the fourth ventricle and bilateral parietal sulci likely representing haemorrhage (figure 7).

DIFFERENTIAL DIAGNOSIS

Aortic dissection was the first differential diagnosis due to the clinical presentation. However, as the initial CT aortogram did not show evidence of dissection flap, the next differential to rule

out was acute myocardial infarction (MI) due to ECG changes and left ventricular dysfunction. The coronary angiogram did not show obstructive coronary lesion. The complete occlusion of the aorta distal to the origin of subclavian artery confirms the diagnosis of IAA.

TREATMENT

His coronary artery disease was not compatible with his severe left ventricular systolic dysfunction. He was treated medically for his acute left ventricular failure secondary to stress-induced CMP and was successfully extubated.

Cardiothoracic team was involved for possible surgical intervention for IAA and surgical options were discussed, but no immediate procedure was planned due to his fluctuating neurological status.

OUTCOME AND FOLLOW-UP

He was extubated after 48 hours and was managed conservatively for his brain injury. On day eight of his hospital stay, he developed apnoea followed by bradycardic cardiac arrest. The possible cause at that time was extension of his neurological injury up to the extent of acute cerebral compromise, but he was not stable enough to be moved for another brain imaging. He was attempted for resuscitation but could not revive.

Figure 2 12-lead ECG showing new T-wave inversion in leads V1–V6.

Figure 3 Radiocontrast injection across the descending aorta showing complete occlusion of the proximal part.

DISCUSSION

IAA is defined as the complete loss of luminal continuity between the ascending and the descending aorta. It is usually associated with other congenital cardiac defects like ventricular septal defects, bicuspid aortic valve and other aortopulmonary trunk anomalies.¹

IAA is classified into three types based on the site of interruption. Type A which is interruption distal to the origin of left subclavian artery (LSA), Type B which is interruption between left common carotid and LSA and Type C which is interruption between brachiocephalic trunk and left common carotid artery,⁴

Figure 4 Radiocontrast injection across the ascending aorta showing complete occlusion of distal part with luminal discontinuity.

Figure 5 Image of the CT aortogram showing extensive collateral circulation.

Majority of the cases are fatal in very early childhood due to the lack of extensive collaterals necessary to maintain the distal flow and closure of PDA.

Diagnosis of IAA can be made by non-invasive and invasive techniques. It can be anticipated by doing a careful examination of young adults with hypertension. Non-invasive techniques include echocardiography, CT and cardiac MRI. Invasive diagnosis is made in cardiac catheterisation laboratory by aortogram.

Figure 6 Apparent diffusion coefficient and diffusion weighted imaging images of MRI brain showing focal area of diffusion restriction along the occipital horn of lateral ventricle (a and b) and at the splenium of corpus callosum (c and d).

Figure 7 Sequential weighted imaging showing signal dropout in fourth ventricle and bilateral parietal sulci likely representing haemorrhage

Clinical presentation in IAA varies from failure to thrive in neonates to hypertension (most common), chronic fatigue, non-specific chest pain (due to collaterals) and heart failure (rarely) in adults. Presentation mimicking aortic dissection or acute MI has not been reported. On review of the literature, we found few case reports on IAA in adult patients. Isolated IAA without other congenital cardiac anomalies in adults is even rarer.⁵

Takotsubo CMP was initially recognised in Japan in 1990 and is an emerging diagnosis since 1998. Majority of the patients are women. Events are more prevalent in afternoon when stressful triggers are more likely to take place due to epinephrine surge which results in consequences such as acute coronary syndrome and left ventricular dysfunction. Common clinical presentation is with chest pain, dyspnoea and palpitation.⁶ To our knowledge, this CMP has not been reported with IAA.

Initial presentation of our patient raised a high suspicion of aortic dissection that turned out to be an IAA with associated stress-induced CMP. Literature shows that between 5% and 10% of dissections do not have an obvious intimal tear on the imaging and these are due to the rupture of aortic vasa vasorum with clinical sign and symptoms similar to acute intimal dissection.⁷

The posterior reversible encephalopathy syndrome (PRES) is a clinic-neuroradiological diagnosis with various aetiologies, severe hypertension being the one cause. It usually results in vasogenic oedema of posterior cerebral regions; however, the literature shows up to 15% incidence of intracranial haemorrhage (from minute haemorrhage to haematoma) in PRES as in our patient. The outcome varies, being reversible in majority of patients but can result in morbidity and mortality.⁸

We think that his clinical presentation was a combination of more than one disease; he may have had a tear in one of his collaterals due to the vasa vasorum rupture and therefore did not have a classical flap on CT scan. As he had high blood pressure since the age of 16 years, if he could have a comprehensive workup for secondary hypertension at young age, this diagnosis of IAA could have potentially been made and treated a long time before he presented in such an emergency situation.

This case emphasises the role of detailed history and physical examination in diagnosing diseases with high mortality which are rare, such as IAA in adults which join the two factors.

Learning points

- ▶ Stress-induced cardiomyopathy is a mimicker of myriad cardiac conditions.
- ▶ Detailed workup should be done for secondary hypertension before starting antihypertensive medications, especially in young patients.

Contributors FB is the corresponding author and writer of the manuscript. AH, JMT and AB contributed to the editing of the manuscript.

Competing interests None declared.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

© BMJ Publishing Group Ltd (unless otherwise stated in the text of the article) 2017. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

REFERENCES

- 1 Ponte M, Dias A, Dias Ferreira N, *et al*. Interrupted aortic arch: a misdiagnosed cause of hypertension. *Rev Port Cardiol* 2014;33:389–e1-5.
- 2 Rauch A, Hofbeck M, Leipold G, *et al*. Incidence and significance of 22q11.2 hemizygosity in patients with interrupted aortic arch. *Am J Med Genet* 1998;78:322–31.
- 3 Lewin MB, Lindsay EA, Jurecic V, *et al*. A genetic etiology for interruption of the aortic arch type B. *Am J Cardiol* 1997;80:493–7.
- 4 Javadzadegan H, Porhomayon J, Sadighi A, *et al*. Isolated interrupted aortic arch: unexpected diagnosis in a 63-year-old male. *Case Rep Crit Care* 2011;2011:1–3.
- 5 Lobato RF, Saliba LA, Ferreira CR, *et al*. Interrupted aortic arch with cardiac heart failure in young adult. *Arq Bras Cardiol* 2008;91:e4–6.
- 6 Sharkey SW, Lesser JR, Maron BJ, *et al*. Cardiology patient page. Takotsubo (stress) cardiomyopathy. *Circulation* 2011;124:e460–2.
- 7 Patel PD, Arora RR. Pathophysiology, diagnosis, and management of aortic dissection. *Ther Adv Cardiovasc Dis* 2008;2:439–68.
- 8 Hefzy HM, Barynski WS, Boardman JF, *et al*. Hemorrhage in posterior reversible encephalopathy syndrome: imaging and clinical features. *AJNR Am J Neuroradiol* 2009;30:1371–9.

Copyright 2017 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow