Published in final edited form as:

Food Chem Toxicol. 2017 November ; 109(Pt 1): 690–702. doi:10.1016/j.fct.2017.05.013.

Dose-response analysis of epigenetic, metabolic, and apical endpoints after short-term exposure to experimental hepatotoxics

Isabelle R. Miousse^a, Lynea A. Murphy^b, Haixia Lin^a, Melissa R. Schisler^b, Jinchun Sun^c, Marie-Cecile G. Chalbot^d, Radhakrishna Sura^b, Kamin Johnson^b, Matthew J. LeBaron^b, Ilias G. Kavouras^d, Laura K. Schnackenberg^c, Richard D. Beger^c, Reza J. Rasoulpour^{b,*}, and Igor Koturbash^{a,*}

^aDepartment of Environmental and Occupational Health, College of Public Health, University of Arkansas for Medical Sciences, Little Rock, AR, 72205, USA

^bToxicology and Environmental Research & Consulting, The Dow Chemical Company, Midland, MI, USA

^cDivision of Systems Biology, National Center for Toxicological Research, US Food and Drug Administration, Jefferson, AR 72079, USA

^dDepartment of Environmental Health Sciences, Ryals School of Public Health, University of Alabama at Birmingham, 1665 University Blvd, Birmingham, AL 35246, USA

Abstract

Identification of sensitive and novel biomarkers or endpoints associated with toxicity and carcinogenesis is of a high priority. There is increasing interest in the incorporation of epigenetic and metabolic biomarkers to complement apical data; however, a number of questions, including the tissue specificity, dose-response patterns, early detection of those endpoints, and the added value need to be addressed. In this study, we investigated the dose-response relationship between apical, epigenetic, and metabolomics endpoints following short-term exposure to experimental hepatotoxics, clofibrate (CF) and phenobarbital (PB). Male F344 rats were exposed to PB (0, 5, 25, and 100 mg/kg/day) or CF (0, 10, 50, and 250 mg/kg/day) for seven days. Exposure to PB or CF resulted in dose-dependent increases in relative liver weights, hepatocellular hypertrophy and proliferation, and increases in Cyp2b1 and Cyp4a1 transcripts. These changes were associated with altered histone modifications within the regulatory units of cytochrome genes, LINE-1 DNA hypomethylation, and altered microRNA profiles. Metabolomics data indicated alterations in the metabolism of bile acids. This study provides the first comprehensive analysis of the apical,

#The views expressed in this manuscript do not necessarily represent those of the US FDA.

*Correspondence to: Igor Koturbash: IKoturbash@uams.edu, 4301 W. Markham Str, Slot# 820-11, Department of Environmental and Occupational Health, College of Public Health, University of Arkansas for Medical Sciences, Little Rock, AR, 72205-7199, USA or Reza Rasoulpour: RRasoulpour@dow.com.

Disclaimer: The views presented in this article do not necessarily reflect those of the Food and Drug Administration.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

epigenetic and metabolic alterations, and suggests that the latter two occur within or near the dose response curve of apical endpoint alterations following exposure to experimental hepatotoxicants.

Keywords

Benchmark dose modeling; DNA methylation; LINE-1; miRNA; metabolomics; hepatotoxicants; dose-response

1. Introduction

The current product safety assessment paradigm identifies adverse apical effects to establish a dose-response relationship between exposure and health-related outcomes and has been used for many years in human risk assessment. The molecular mechanisms leading to adverse apical endpoints are not typically evaluated in the risk assessment process, although molecular changes underlying the apical effect may be considered. Accumulating evidence indicates that exposure to various toxicants and chemical carcinogens can affect the cellular epigenome (Hou et al., 2012; Koturbash et al., 2011c; Thomson et al., 2014); however, the lack of causal links between epigenetic and apical effects is an obstacle to the incorporation of epigenetic endpoints into the human risk assessment process (Alyea et al., 2014; Miousse et al., 2015b; Priestley et al., 2012). Furthermore, specific epigenetic endpoints, target tissue specificity, dose-response, and approaches for their evaluation need to be identified. This is particularly relevant in the case of hepatotoxicants and carcinogens with the non-genotoxic mode of action, where DNA damage cannot be utilized as an endpoint, and the reliance on apical endpoints is therefore greater. In these regards, experimental non-genotoxic rodent hepatocarcinogens, such as phenobarbital (PB) and clofibrate (CF), serve as suitable chemicals to investigate the relationship between the apical and epigenetic endpoints.

Epigenetics is defined as changes in gene expression and chromatin organization that occur without alterations in DNA sequence and are heritable through cell division. Disruption of the balanced cellular epigenetic status may result in the development of a variety of pathological states, including cancer (Baylin and Jones, 2011; Waldmann and Schneider, 2013). Exposure to numerous liver toxicants, including the rodent experimental hepatocarcinogen PB, has been associated with a variety of epigenetic alterations, such as changes in DNA methylation, histone modifications and non-coding RNAs (Koturbash et al., 2011b; Koufaris et al., 2013; Phillips et al., 2009; Phillips and Goodman, 2009; Thomson et al., 2012; Thomson et al., 2014; Watson and Goodman, 2002). Therefore, DNA methylation status of the Long Interspersed Nuclear Element-1 (LINE-1) and expression of microRNAs (miRNAs), were proposed to be utilized as early biomarkers of exposure to hepatotoxicants and carcinogens (Herceg et al., 2013; Koturbash et al., 2012; Koturbash et al., 2015; Lambert et al., 2015; Vliegenthart et al., 2015).

Metabolomics evaluates “the metabolite pool that exists within a cell under a particular set of conditions” (Fiehn, 2002). Metabolite profiling, typically using nuclear magnetic resonance (NMR) spectroscopy- or liquid chromatography/mass spectrometry (LC/MS)-based methods, has been utilized to evaluate biomarkers of hepatocarcinogenesis (Ohta et al., 2009; Tan et al., 2012; Unterberger et al., 2014). Metabolomics methods have been

applied previously to evaluate the toxicity of CF in rats (Ishihara et al., 2006; Strauss et al., 2012) and mice (Wheelock et al., 2007) and the effects of exposure to PB in rats (Rubtsov et al., 2010; Waterman et al., 2010). These approaches have the potential to identify metabolite biomarkers related to hepatotoxicity and hepatocarcinogenesis.

Currently, there are limited data available to establish the toxicological significance of epigenetic changes and their causal relationship to apical endpoints and alterations in metabolism in response to exposure to hepatotoxicants and non-genotoxic liver carcinogens, as assessed in regulatory guideline toxicology studies. Toxicological studies incorporating epigenetic endpoints are typically conducted at a single high-dose level of exposure and do not contain appropriate concurrent assessment of apical toxicity endpoints such as histopathology and clinical chemistry to correlate epigenetic changes to apical endpoints (Alyea et al., 2012; Miousse et al., 2015b; Rasoulpour et al., 2011). In addition, few studies have utilized multiple doses to investigate a dose-response relationship between apical effects, specific genes associated with alteration in epigenetic regulation, and metabolism and metabolite profiling. Currently, employing benchmark dose (BMD) response analysis has become a useful tool to estimate dose levels corresponding to specific response by selecting a suitable dose-response modeling in risk assessment. Therefore, this study was designed to explore the similarities and differences between the dose-response relationship and associated benchmark doses of apical and metabolomics endpoints, as well as transcriptional changes in the genes involved in epigenetic regulation following short-term exposure to experimental hepatotoxicants, CF and PB, and provide useful information on early toxicity at the multiple levels of biomarkers underpinning quantitative risk assessment.

2. Materials and Methods

2.1 Animals and treatment

Ten-week old male F344/DuCrI rats were purchased from Charles River Laboratories, Inc. (Kingston, New York). This strain was selected as it is commonly used in repeat dose toxicology studies and previous toxicity studies on CF and PB have used this strain. Rats were randomized according to body weight, identified via subcutaneously implanted transponders, housed one per cage in stainless steel cages, and provided LabDiet Certified Rodent Diet (PMI Nutrition International, St. Louis, MO) and water *ad libitum*. CF and PB were obtained from TOCRIS (BioScience, Bristol, United Kingdom) and Sigma-Aldrich (Saint Louis, MO), respectively. After acclimation, rats were administered PB or CF by oral gavage at dose levels of 0, 5, 25 or 100 mg/kg/day (PB) or 0, 10, 50, or 250 mg/kg/day (CF) in propylene glycol for seven days (n=5 per dose). Those doses are generally considered as NOAEL, LOAEL, and lowest carcinogenic doses, respectively. Animals were terminated 24 hrs after the last dosing. The animal studies were performed at The Dow Chemical Company, Toxicology and Environmental Research & Consulting (TERC), Midland, MI, which is accredited by the American Association for Accreditation of Laboratory Animal Care. All animal care and use activities were reviewed and approved by the Institutional Animal Care and Use Committee.

2.2. Osmotic pumps

Rats were implanted with mini-osmotic pumps (model 2ML1; Alzet Corporation, Palo Alto, CA) to allow for continuous administration of bromodeoxyuridine (BrdU; a structural analog of thymidine that incorporates into nuclear DNA and is used as a surrogate marker of cell proliferation) during the seven days of treatment (Eldridge et al., 1990). The pumps contained 20 mg/ml of BrdU in phosphate buffered saline (pH 7.6) and delivered at a rate of 10 µl/hour.

2.3. Clinical data and pathology

Rats submitted for necropsy were weighed (with the implanted osmotic pump *in situ*), anesthetized by inhalation of CO₂/O₂ or isoflurane/O₂, and blood samples were obtained from the orbital sinus. The animals were then euthanized by decapitation. The osmotic pumps were removed and weighed. The weights of the osmotic pumps were subtracted from the recorded body weights to determine the final body weight. The skin was removed from the carcass, the abdominal cavity opened, and the liver and kidneys excised and weighed. In addition, a 2–3 cm segment of the proximal duodenum was excised, flushed with fixative, and placed in the same fixative with the liver tissue, which served as a positive control for osmotic pump functionality and immunohistochemistry. The upper quarter of the left lateral liver lobe was processed in RNA_{later} for targeted gene expression analysis. The upper middle quarter of the left lateral lobe was trimmed and preserved in neutral phosphate-buffered 10% formalin and was used for histological examination and BrdU proliferation analysis. The lower quarter of the left lateral liver lobe and the lower portion of the median liver lobe were flash frozen and stored at –80 °C for further analyses. Livers from all rats were processed by standard histologic procedures. Paraffin embedded tissues were sectioned approximately 6 µm thick, stained with hematoxylin and eosin, and examined by a veterinary pathologist using a light microscope.

Serum was prepared as previously described (Hastings et al., 2012). Briefly, blood was collected into vacutainers containing sodium fluoride/potassium oxalate (BD Biosciences, Franklin Lakes, NJ) and centrifuged at 2000 xg for 15 min at 4 °C. The supernatant was transferred into a new tube and centrifuged twice at 800 xg for 15 min at 4 °C and kept at –80 °C for subsequent microRNA analysis.

2.4. Hepatocellular proliferation (BrdU)

At necropsy, sections of the liver were treated as described previously, processed by standard techniques, and mounted on glass slides. Tissue was stained for BrdU using the manufacturer's protocol (BD Biosciences, San Diego, California, USA) with modified heat-induced antigen retrieval. A small section of duodenum from each rat was also processed and stained to serve as a control for confirming systemic availability of BrdU and for immunohistochemical staining. Proliferation in rats was evaluated using light microscopy where positive nuclei were scored as percentages based on counting 1,000 hepatocytes in each of three hepatolobular zones: centrilobular, midzonal, and periportal regions. BrdU labelling has previously been used to analyze the DNA methylation within the repetitive elements, as well as histone modifications (Li et al., 2014), and, thus is not known to substantially affect epigenetic analysis. All BrdU slides were coded and read blind.

2.5. Gene expression analysis

Total RNA from liver and kidney was extracted using the Qiagen RNeasy kit following the manufacturer's protocol (Qiagen Inc., Valencia, CA). RNA quantity and quality were assessed by a Nanodrop 1000 or 2000 spectrophotometer (Thermo Scientific, Waltham, MA) and Agilent 2100 Bioanalyzer (Agilent Technologies, Palo Alto, CA), respectively. Only samples with an OD 260/280 ratio greater than 1.8, clearly defined 28S and 18S bands, and acceptable RNA Integrity Numbers (RIN; generally >7) from the BioAnalyzer were used for gene expression studies. Total RNA was treated with DNase enzyme to avoid DNA contamination. The RNA extraction samples were stored at -80 °C until used. cDNA was synthesized using random primers and a High Capacity cDNA Reverse Transcription Kit (Applied Biosystems, Foster City, CA) according to the manufacturer's protocol (Life Technologies, Carlsbad, CA). Gene expression analysis was performed on the following genes: *Cyp1a1*, *Cyp2b1*, *Cyp2b2*, *Cyp3a23/3a1*, *Cyp4a22*, *Dnmt1*, *Dnmt3a*, *Dnmt3b*, *MeCP2*, *Mbd1*, and *Mbd2*. *Actb*, *Gapdh*, and *Srp14* were used as control genes. Gene expression studies were conducted with an Applied Biosystems 7500 and ViiA 7™ fast Real-Time Polymerase Chain Reaction system (Applied Biosystems) using the TaqMan Gene Expression Assays (Life Technologies). Details on the gene expression analysis are provided in Supplementary Materials and Methods.

2.6. Chromatin immunoprecipitation (ChIP) analysis of gene promoters

Chromatin immunoprecipitation was performed using the Chromatin Immunoprecipitation Assay Kit (Millipore, Billerica, MA). Briefly, 30 mg of liver tissue was mechanically homogenized in PBS containing protease inhibitor, PMSF, and a final concentration of 1% formaldehyde and incubated at room temperature for 20 min. The reaction was stopped with a 1.25M glycine solution. The material was centrifuged 4 min at 5000 rpm at 4 °C, and the pellet was washed twice in PBS. The chromatin was then resuspended in SDS lysis buffer and sheared by sonication for 10 min in a water bath. The chromatin was resuspended in a total volume of 2 ml with ChIP dilution buffer and cleared with Protein A. Agarose/Salmon Sperm DNA. 200 µl of the cleared chromatin was used for incubation with each ChIP-grade antibodies against H3K18ac (Cat # ab1191), H3K4me3 (Cat #ab12209) (both Abcam), and H2A.Z (Cat # 17-10048, Millipore), following the manufacturer's protocol. Immunoprecipitated chromatin was quantified by qRT-PCR.

2.7. Western blot

Histones were acid-precipitated by homogenizing liver tissue in 1 ml lysis buffer (10 mM HEPES pH 7.9, 1.5 mM MgCl₂, 10 mM KCl) and centrifuging at 14,000 xg for 10 min at 4 °C. The acid-soluble fraction was used for immunodetection. For Dnmt1 immunodetection, the nuclear fraction was isolated from liver tissue using the EpiQuick Nuclear Extraction Kit (Epigentek, Farmingdale, NY) following the manufacturer's protocol. The histone-containing acid-soluble fraction was loaded on a 15% SDS-PAGE gel and the nuclear extract was loaded onto a 7.5% SDS-PAGE gel for western blotting. Histone antibodies, including total histone 3 for normalization, were purchased from Millipore and anti-Dnmt1 (Cat # AP1032b) antibody was purchased from Abgent (San Diego, CA). Gapdh mouse monoclonal antibody was used as a loading control (Santa Cruz, Cat # sc-32233).

2.8. Analysis of LINE-1 DNA methylation

Genomic DNA was extracted from frozen liver and kidney tissue using the AllPrep DNA/RNA extraction kit (Qiagen), according to the manufacturer's protocol. DNA concentrations and integrity were analyzed by the Nanodrop 2000 (Thermo Scientific). Only DNA samples with the 260/280 ratios between 1.8 and 1.9 and the 260/230 ratios above 1.5 were considered for further molecular analyses. Methylation-sensitive quantitative polymerase chain reaction (MS-PCR) was performed as described earlier (Koturbash et al., 2011c). For methylation analysis by pyrosequencing, genomic DNA was extracted as described above. Bisulfite conversion followed by pyrosequencing of 10 CpG sites in the rat LINE-1 transposable element was performed using the PyroMark96 (Qiagen) from 1 µg of gDNA. PyroMark96 software measured the percent methylation at each CpG within the LINE-1 ORF1. The results are presented as fold change from control.

2.9. miRNA analysis

MicroRNAs were extracted from flash-frozen livers and serum using the miRNeasy and miRNeasy Serum/Plasma kits, respectively (Qiagen). RNA concentration was assessed by Nanodrop 2000 (Thermo Scientific) and 25 ng of RNA containing miRNAs was used for reverse transcription with HiSpec buffer and miScript reverse transcriptase.

A custom miRNA array was designed with 21 miRNA targets (described in details in Supplementary Table 1) (Qiagen, custom cat#CMIRN02145). Arrays were prepared according to the manufacturer's indications, and run on a ViiA-7 instrument (Applied Biosystems). Results were analyzed using the web-based analysis tool (<http://pcrdataanalysis.sabiosciences.com/pcr/arrayanalysis.php>).

2.10. Metabolomics analysis by nuclear magnetic resonance (NMR) and UPLC/MS

Pre-weighed liver tissue (~150 mg) samples were homogenized in 3 mL ice cold methanol. The homogenates were then distributed into 1 mL aliquots. The aliquots were centrifuged for 12 min at 13,000 xg and 4 °C. The supernatant was transferred to a new Eppendorf tube and the solvent evaporated using a SpeedVac system. Detailed description of NMR and UPLC/MS analysis can be found in the Supplementary Materials and Methods.

2.11. Dose-response modeling of apical, transcriptional changes in genes involved in epigenetic regulation and metabolomic parameters

Benchmark dose (BMD)-response analysis was conducted to estimate the dose at which apical effects and specific genes associated with epigenetic regulation and metabolism, and metabolites associated with DNA methylation were altered above or below control. BMDs and BMDLs were calculated using benchmark dose software version 2.6 (BMDS 2.6.0.1) and BMDS Wizard 1.10 (<https://www.epa.gov/bmds/download-benchmark-dose-software-bmds#installing>) with the benchmark response set to one control standard deviation from the control mean (BMD1SD) or an extra risk of 10%, and included the lower 95% confidence limit (BMDL). Data sets were grouped according to dose-response type (e.g. continuous, quintal, and ordinal) which guided the choice of BMRs and the type of the models to calculate BMDs. The criterion for selecting a fitted model included global goodness-of-fit *p*-value (*p*>0.1), local scaled residuals (absolute value <2.0), the smallest Akaike's Information

Criterion (AIC) and visual inspection of model fitting (<https://www.epa.gov/risk/benchmark-dose-technical-guidance>). BMD and BMDL were estimated for apical, epigenetic and metabolomic endpoints or effects including: relative liver weight, hepatocellular hypertrophy and proliferation, liver cytochrome P450 gene expression, metabolite changes, and gene expression of DNA methyltransferases and methyl-binding proteins. If a chemical had more than one dose-response dataset, we selected the lowest BMD (without warnings, if available) and the BMDL or the lowest BMDL (if different from the previous BMDL). These were selected according to the endpoint or effect.

2.12. Statistical analysis

Five animals per group of treatment were used for each evaluated endpoint. The significance between experimental treatments was determined by one way ANOVA, followed by Dunnett's test to correct for multiple testing using the GraphPad Prism 6.02 software (GraphPad Software, La Jolla, CA). A p -value ≤ 0.05 was considered to be significant. Details on statistical analyses are provided in Supplementary Materials and Methods.

3. Results

3.1. Effects of CF and PB on relative liver weights and clinical biochemistry

Well-known associative events for PPAR α and CAR mediated liver effects include hepatomegaly and hepatocellular hypertrophy (Corton et al., 2014; Elcombe et al., 2014). Following a seven day exposure to CF, relative liver weight (liver weight to body weight ratio) was significantly increased by 9.7% and 54% at doses of 50 and 250 mg/kg/day, respectively (Fig. 1A). CF is a hypolipidemic drug and its effects on serum cholesterol were noted ranging from decreases in serum cholesterol of 13 to 33% at 50 and 250 mg/kg/day of CF, respectively (data not shown). Following seven days of PB exposure, a dose-dependent increase in relative liver weights at 5, 25, and 100 mg/kg/day of PB was detected (8%, 23%, and 41% higher than controls, respectively), which was statistically significant at 25 and 100 mg/kg/day (Fig. 1A).

3.2. Histopathological evaluation of liver

Administration of CF resulted in very slight centrilobular/midzonal hypertrophy at 50 mg/kg/day and panlobular hepatocellular hypertrophy at 250 mg/kg/day (Table 1). Very slight mitotic figures were observed in animals exposed to CF in the centrilobular and midzonal regions of rat livers at a dose of 50 mg/kg/day, and all animals had panlobular mitotic figures at 250 mg/kg/day (data not shown). Quantitative assessment of hepatocellular proliferation indicated a statistically significant increase in BrdU labeling indices (LI) compared to controls at the high dose of 250 mg/kg/day CF in the centrilobular (4.3-fold), midzonal (5.4-fold), and periportal (5.0-fold) regions (Fig. 1B).

For PB, increased BrdU LI were present in the centrilobular, midzonal, and periportal regions at doses of 25 and 100 mg/kg/day (Table 1). Except for the centrilobular regions, BrdU LI were identified at both 25 and 100 mg/kg/day PB. Despite the presence of hepatocellular hypertrophy and increased mitotic figures in all PB dose groups,

hepatocellular proliferation, as evidenced by an increase in LI, occurred only in the 25 and 100 mg/kg/day groups (Fig. 1B).

3.3. Liver cytochrome P450 gene expression

Gene expression responses for *Cyp1a1*, *Cyp2b1*, *Cyp3a1*, and *Cyp4a1* were assessed as biomarkers for activation of AhR (Whitlock, 1999), CAR (Honkakoski et al., 1998; Kawamoto et al., 1999; Wei et al., 2000), PXR (Kliewer et al., 1998; Xie et al., 2000), and PPAR- α (Aldridge et al., 1995; Palmer et al., 1994) signaling pathways, respectively. Treatment-related induction of *Cyp4a1* in the CF treated animals was observed at 50 and 250 mg/kg/day and consisted of a 5- to 23-fold induction in mRNA (Fig. 2A), which correlated with hepatomegaly and hepatocellular hypertrophy at both dose levels and increased hepatocellular proliferation at 250 mg/kg/day. In the PB treated animals, *Cyp2b1* mRNA was induced in a dose-dependent manner, consistent with CAR activation (Elcombe et al., 2014; Geter et al., 2014); in the low-, mid-, and high-dose groups, *Cyp2b1* mRNA was induced 11-, 48-, and 180-fold, respectively (Fig. 2A). The increase in *Cyp2b1* transcript levels correlated with the observation of hepatomegaly and hepatocellular proliferation at all dose levels and increased hepatocellular proliferation at 25 and 100 mg/kg/day of PB.

3.4. Chromatin immunoprecipitation (ChIP) analysis of gene promoters

In order to explore the potential epigenetic nature of the cytochrome gene expression induction, a ChIP analysis within the transcription start sites (TSS), and PB response element (PBRE) or peroxisome proliferative response element (PPRE) of the genes was performed on liver samples from the high dose PB and CF exposure groups. PB exposure was associated with the loss of histone H2A.Z, from the *Cyp2b1* TSS (Fig. 2B). Interestingly, increases in H2A.Z have previously been reported in transcriptional silencing of cytochromes (Chen et al., 2013). Similar effects were observed within the TSS of *Cyp4a1* after CF exposure (Fig. 2C). We also evaluated the PBRE of *Cyp2b1* and identified enrichment of transcriptional activation marks in this region – trimethylation of histone H3 lysine 4 (H3K4me3) and acetylation of histone H3 lysine 18 (H3K18ac). An insignificant increase in H3K4me3 was also observed at the TSS of *Cyp2b2* (data not shown).

3.5. Effects of exposure to CF and PB on DNA methylation machinery

Proper methylation status is facilitated by the DNA methylation machinery which includes DNA methyltransferases (*Dnmt1*, *Dnmt3a*, and *Dnmt3b*) and methyl-binding proteins (*Mecp2*, *Mbd1*). First, we aimed to investigate whether or not short-term administration of rodent non-genotoxic hepatotoxicants would target the DNA methylation machinery. Expression of methyltransferases *Dnmt1* and *Dnmt3a* was decreased after exposure to the highest doses of CF and PB, respectively, while the methyl-binding protein *Mbd1* mRNA was decreased in both treatments (Fig. 3A, B). Loss of *Dnmt1* mRNA expression was further confirmed at the protein level (Fig. 3C). Additionally, dose-dependent decreases of *Dnmt1* and *Mecp2* after treatment with CF and *Dnmt3b*, *Mecp2* and *Mbd1* after treatment with PB were observed (Fig. 3A, B). No significant differences in the expression of DNA methylation machinery were detected in the kidney, a non-target organ, of CF or PB-exposed rats (Supplementary Fig. 1).

3.6. Analysis of DNA methylation and expression of transposable element LINE-1

Given that CF or PB exposure altered the expression of DNA methyltransferases, we investigated whether these changes had consequences on DNA methylation levels. LINE-1 DNA methylation is considered to be a surrogate biomarker for global levels of DNA methylation, and acts as a “locking” mechanism that prevents LINE-1 from reactivation (Miousse et al., 2015a). DNA hypomethylation of LINE-1 and, associated with it, its increased expression that may lead to LINE-1 insertional mutagenesis, have been documented in hepatocarcinogenesis as well as in response to exposure to known hepatotoxicants and carcinogens (Hur et al., 2014; Miousse and Koturbash, 2015; Stribinskis and Ramos, 2006).

We analyzed the methylation status of the first open reading frame of LINE-1 (ORF1), which encodes the nucleic acid chaperone ORF1p. First, we utilized MS-PCR, as this methodology was widely used in previous studies (Koturbash et al., 2011c; Martens et al., 2005). We did not identify any significant changes in any of the experimental groups (Supplementary Fig. 2). We then employed a pyrosequencing approach, a more sensitive and accurate methodology for the assessment of the locus-specific DNA methylation. Following exposure to PB at a dose of 100 mg/kg/day, a slight but significant hypomethylation within LINE-1 ORF1 was detected in rat liver tissue (4%, $p<0.001$) (Fig. 4A). No significant changes were detected after exposure to lower doses of PB. Administration of CF did not change the methylation status of LINE-1 ORF1 (Fig. 4B). No significant changes in LINE-1 ORF1 methylation in kidneys were associated with exposure to either chemical (Supplementary Fig. 3). We did not identify any significant changes in LINE-1 ORF2 expression in the livers exposed to PB or CF rats (Supplementary Fig 3). Similarly, no effects were observed in the kidneys (Supplementary Fig 3).

3.7. miRNAs in liver tissue and serum

The expression of the panel of miRNAs previously reported altered in response to exposure to hepatotoxicants was evaluated in order to investigate whether or not miRNAs could serve as early biomarkers of the short-term exposure to known non-genotoxic hepatocarcinogens (Supplementary Table 1). Of 20 tested miRNAs, five were significantly down-regulated in the livers of rats following administration of CF or PB at their highest doses tested (250 mg/kg/day for CF and 100 mg/kg/day for PB): miR-122, miR-125a, miR-155, miR-199a-5p, and miR-199a-3p (Table 2). Furthermore, these five targets displayed a dose-dependent pattern of expression. Additionally, administration of PB was characterized by a decrease in miR-221 at the highest dose and an increase in miR-200b at the low and middle dose (Table 2).

Accumulating evidence indicates that circulating miRNAs may serve as non-invasive surrogate biomarkers of liver toxicity and hepatocarcinogenesis (Wang *et al.*, 2015, Koturbash *et al.*, 2012). In this study, we aimed to investigate whether the observed changes in liver miRNA expression could be also detected in the serum of rats exposed to CF or PB for seven days. Despite the detectable changes in miRNA expression in the liver, only miR-199a-3p was significantly changed in the serum of rats treated with 10 mg/kg/day of CF (1.9-fold, $p<0.05$) (Supplementary Table 2).

3.8. NMR Metabolomics Analyses

PLS-DA of the normalized, binned integrals from the NMR spectra of control and clofibrate liver samples showed the 250 mg/kg/day dose group separate along t[1] from the control, 10 mg/kg/day, and 50 mg/kg/day dose groups. Select metabolites detected by NMR spectroscopy were quantified and normalized to tissue weight, and trimethylamine-N-oxide was the only significantly changed metabolite in the 250 mg/kg/day dose group relative to control. The amino acids alanine, aspartate, betaine, glutamate, glutamine, leucine, lysine, and valine were also elevated after a high dose of CF, although not significantly. Choline was decreased in a dose-dependent manner relative to control. NMR analyses of the PB samples did not show significant changes compared to the control group.

3.9. LC/MS Metabolomics Analyses

Among the detected metabolites by LC/MS, distinct changes in bile acids (BAs) after administration of PB (Fig. 5A) and CF (Fig. 5B) were noted. At the 250 mg/kg/day CF, significant decreases were observed in the bile acid derivatives tauroursodeoxycholic acid (TUCA), taurodeoxycholic acid (TDCA), taurochenodeoxycholic acid (TCDCA), and tauroursodeoxycholic acid (TUDCA), while taurocholic acid (TCA) was significantly increased. For PB, TUDCA was significantly decreased and showed a dose-dependent trend at all three doses. TCA, TDCA, and TCDCA were decreased at the 100 mg/kg dose of PB, while only the TCDCA decrease was significant. At the highest dose of CF and PB, sulfolithcholyglycine and its isomer increased.

Fig. 5C and D shows changes in metabolites involved in oxidative stress and DNA methylation induced by PB or CF dosing. At the 250 mg/kg CF dose, sulfur-containing metabolites glutathione, methionine, taurine and S-adenosyl-homocysteine (SAH) decreased; among these, only taurine was significantly decreased. No changes were observed in any oxidative stress- or DNA methylation-related metabolites following dosing with PB.

3.10. Apical, metabolomics, and transcriptional changes in the expression of genes involved in epigenetic regulation benchmark dose-response analysis

Benchmark dose analysis was used to describe potency in the low-dose region, particularly to approximate the doses below which changes in endpoints/effects compared to background or controls. BMD estimates of the apical, transcriptional, and metabolomics endpoints are summarized in Tables 3, 4 and 5. As shown in those tables, BMD and BMDL estimates for relative liver weight were 23.4 and 18.2 mg/kg/day for CF and 18.7 and 14.1 mg/kg/day for PB, respectively. The hepatic biomarker (CYP 450 gene expression) BMD and BMDL for CF and PB were 17.2 and 13.5 mg/kg/day, and 7.6 and 6.0 mg/kg/day, respectively. The BMD and BMDL for hepatocyte cell proliferation was 90.2 and 63.4 mg/kg/day for CF and 31.8 and 22.8 mg/kg/day for PB, respectively. The BMD for transcriptional changes in genes involved in DNA methylation was 101.2 mg/kg/day for CF and 50.4 mg/kg/day for PB and the BMDL for CF and PB were 71.2 and 32.9 mg/kg/day. The BMD and BMDL for metabolites from CF and PB were 58.3 and 43.5 mg/kg/day and 28.7 and 20.9 mg/kg/day, respectively.

4. Discussion

Understanding the relationship between dose and apical endpoint responses is central to the human risk assessment process. The risk associated with chemical exposure is set by identifying a No-Observed-Adverse-Effect Level (NOAEL) for the most sensitive adverse apical effect, while adding safety factors to this NOAEL to account for interspecies and intraspecies differences, and comparing the derived value to human exposure levels. Before epigenetic or metabolomic parameters can be used to refine the risk assessment process, a number of criteria need to be satisfied. These include the identification of both a causal role of epigenetic or metabolomic changes in the apical endpoint mode of action and the NOAEL for this change. Furthermore, specific endpoints need to be selected from the variety of parameters (i.e., methylation of LINE-1, miRNA expression) and particular approaches for their determination (i.e., pyrosequencing, whole genome bisulfite sequencing) as well as the target organ specificity of alterations.

In the present study, we investigated the dose-response relationship among adverse apical effects, epigenetic alterations, and metabolite changes in a model of the rodent experimental hepatotoxicants and non-genotoxic hepatocarcinogens – PB and CF. To our knowledge, this is the first study that attempts to assess the dose-response relationship between those endpoints and further provides benchmark dose analysis.

In our study, exposure to either PB or CF resulted in a dose-dependent hepatocellular hypertrophy and alterations in the expression of cytochromes responsible for the metabolism of these compounds. It has been reported that the induction of CYP450 gene expression represents an adaptive response to protect against chemical exposure (Willson and Kliewer, 2002). Elevated levels of cytochrome P450 mRNAs were associated with alterations in histone modifications at the TSS, PBRE, and PPRE. Specifically, the PBRE element within *Cyp2b1* was characterized by an increase in trimethylation of histone H3 at lysine 4 and acetylation of histone H3 at lysine 18. Both of these histone marks are associated with a relaxed chromatin structure and transcriptional activation. We also observed a decrease in the levels of the histone variant H2A.Z at the transcriptional start site of both *Cyp2b1* and *Cyp4a1*. This variant, encoded by the gene *H2afz*, is generally thought to be associated with transcriptionally active genes in mammals; however, others have observed a decrease in its abundance at the TSS of stimulated genes, including at another cytochrome P450 – *Cyp26a1* (Chen et al., 2013). Our results show depletion of H2A.Z at this site in association with increased gene expression and in support of the hypothesis that for some genes, H2A.Z is present at the promoter during repressed and poised states and is dislodged upon activation. Interestingly, in both the present study and in the work by Chen and colleagues, these genes were cytochromes under the control of an enhancer containing a binding site for the retinoid X receptor.

In parallel, we evaluated the DNA methylation status of the LINE-1 retrotransposon, the most abundant repetitive element in mammalian genomes. While no significant differences were detected in DNA methylation of LINE-1 in any of the treatment groups using MS-PCR, pyrosequencing allowed the detection of a significant loss of LINE-1 ORF1 methylation in the livers of rats administered a potentially carcinogenic (100 mg/kg/day)

dose of PB (Carreira et al., 2014). Hypomethylation of LINE-1 is one of the major contributors to the global DNA hypomethylation observed in the cancerous tissue (Carreira et al., 2014; Estecio et al., 2007; Koturbash et al., 2011a; Miousse and Koturbash, 2015) and numerous studies have reported DNA hypomethylation of LINE-1 in response to exposure to various environmental toxicants and carcinogens [reviewed in (Herceg et al., 2013; Koturbash et al., 2011a)]. These results, however, need to be interpreted with caution, since one previous study failed to reproduce PB-induced hepatocarcinogenesis in F344 rats (Butler, 1978), while another observed dose-dependent increase in preneoplastic hepatocellular lesions (Hagiwara et al., 1999).

Importantly, in the current study, we observed loss of LINE-1 ORF1 DNA methylation only at a dose of PB (100 mg/kg/day) that was also associated with moderate centrilobular and midzonal hepatocellular hypertrophy and significant hepatocellular proliferation. Increased proliferation could potentially create the conditions where DNA methylation machinery is unable to cope with an increased metabolic demand to copy methylation patterns for newly synthesized DNA. Furthermore, mRNA levels for *Dnmt1* were significantly decreased after treatment with 250 mg/kg/day of CF. We also identified a decrease in the protein level of the maintenance DNA methyltransferase *Dnmt1* for both compounds. In addition, expression of two other genes involved in DNA methylation, *Dnmt3b* and *Mbd1*, showed a dose-dependent decreasing trend in expression after treatment with PB. Moreover, less pronounced hypertrophy and proliferation rates observed after exposure to CF did not result in the loss of LINE-1 ORF1 DNA methylation.

This loss of DNA methylation and alterations in the DNA methylation machinery observed in the current study are in good agreement with the previously reported data. Aberrant DNA methylation has been reported in mouse and rat livers after administration of PB (Phillips and Goodman, 2009; Thomson et al., 2014; Watson and Goodman, 2002). A previous study identified a decrease in DNA methylation with CF treatment, albeit at doses six times higher than those studied here (Nie et al., 2006). The data on effects of CF on DNA methylation is scarce, but it was hypothesized that CF may affect methyltransferase activity through the inducible nitric oxide synthase gene (Jiang et al., 2007). A decrease in *Dnmt3b* has been reported in mice after two weeks of treatment with PB, along with decreases in *Dnmt1* and *Dnmt3a* (Phillips et al., 2009). The increase in betaine and SAH, as evident from the metabolomics data (Fig 5B), further supports the detected disruption of DNA methylation in the liver. Based on our data and the existing literature, we can hypothesize that demethylation of repetitive elements will likely progress with extended treatment duration.

In addition to histone modifications and DNA methylation, we also turned our attention to miRNAs. After a thorough literature review, 20 miRNAs that were previously reported to be affected by various hepatotoxicants were selected (Supplementary Table S1). In this study, we identified five miRNAs that were significantly modulated in rat liver following treatment with the highest doses of both CF and PB. Importantly, the expression of all five miRNAs followed a dose-dependent pattern. Among these were a liver-specific miRNA miR-122, as well as several other miRNAs (miR-125a, miR-155, miR-199-5p and miR-199-3p) that have been reported to be modulated by a vast array of hepatotoxicants and carcinogens, including Wy-14,643, hexahydro-1,3,5-trinitro-1,3,5-triazine, acetaminophen, diethylnitrosamine,

carbon tetrachloride, and tamoxifen (Chen et al., 2012; Girard et al., 2008; Pogribny et al., 2010; Shah et al., 2007; Tryndyak et al., 2009; Wang et al., 2009). In addition, two miRNAs, miR-200b and miR-221, were affected only by PB. Dysregulation in the expression of miR-221 also followed a dose-dependent behavior and was previously associated with treatment with carbon tetrachloride and methyl-deficient diet. We also analyzed miRNA expression in the plasma but were not able to identify any significant alteration in the miRNAs studied. It is plausible to hypothesize that the lack of changes in serum miRNAs is related to the non-genotoxic mode of action of both chemicals and absence of systemic liver injury.

The flux of BAs, synthesized in the liver, is a well-known biomarker of liver function (Legido-Quigley et al., 2011), and blockade of bile flow from the liver to the intestine is one of the characteristics of liver disease or liver injury. The rate of bile flow has been reported as an indicator for the recovery process post-liver transplantation (Legido-Quigley et al., 2011). In the present study, both CF and PB at the highest dose induced decreases in TUCA and TUDCA, while inducing increases in sulfolithcholyglycine and its isomer. Decreases in tauro-conjugated bile acids indicated that either hepatocytes excrete these BAs rapidly, or that the liver cells might be injured resulting in reduced synthesis of these tauro-conjugated BAs. The analysis of BAs in liver tissue thus provided us with additional endpoints for the assessment of non-genotoxic hepatocarcinogens. Among the metabolites identified from both chemicals, changes in metabolic endpoints occurred at the lower doses of CF and PB (58 mg/kg/day for CF and 29 mg/kg/day for PB) compared to those of transcriptional changes.

Compared to NOAEL of 50 mg/kg/day for CF and 25 mg/kg/day for PB for epigenetic changes, we could not find a fitted model for calculating BMD and BMDL for LINE-1 and miRNAs as critical epigenetic endpoints due to the lack of a biologically significant dose-related trend. However, the dose-response relationship for the transcriptional changes in the genes involved in epigenetic regulation was evaluated. Specifically, the gene expression of examined DNA methyltransferases and methyl-binding proteins was altered at dose levels of 100 mg/kg/day for CF and 50 mg/kg/day for PB. It has been reported that following 7-day exposure to PB, the range of BMD for apical endpoints (e.g. liver weight, hepatic biomarkers, and hepatocyte cell proliferation) in male CD-1 mice was 2.4–30 mg/kg/day (Geter et al., 2014). For CF, the lowest tumorigenic dose is 250 mg/kg/day in rats (Reddy et al., 1979; Sargent et al., 2002). In the present study, estimated BMD for apical and hepatic biomarker endpoints was 7.6–46.4 mg/kg/day for PB and 17.2–142.8mg/kg/day for CF. Compared to NOAEL approach, BMD analysis provided a more refined dose-response assessment (Davis et al., 2011; Wignall et al., 2014). In addition, the current study identified the difference in BMD between CF and PB, and determined a lower point of departure (POD) for two kinds of hepatotoxicants at an early time-point which can be used to extrapolate human carcinogenic risk compared to a two-year bioassay and allow to identify hazard ranking of different chemicals in the quantitative risk assessment.

Altogether, the results from this seven day study suggest that liver epigenetic changes, such as LINE-1 DNA methylation and miRNA abundance, occur within the dose response curve of apical endpoint alterations following exposure to classical experimental hepatotoxicants.

However, it has to be taken into consideration that more sensitive approaches, such as pyrosequencing, need to be utilized in order to detect these changes, due to the lower magnitude of the latter at early time-points. Significant changes in liver epigenetic and metabolomics endpoints at the seven day time-point were detected only at higher, potentially carcinogenic, doses of PB or CF. Considering the BMD analysis, BMD values of both CF and PB for transcriptional changes in genes involved in epigenetic regulation were higher than for apical and metabolic endpoints. This finding may suggest the responsiveness of epigenetic endpoints at the level of exposure to potentially carcinogenic doses of chemicals. BMD for PB on transcriptional changes in genes involved in DNA methylation, apical, and metabolic endpoints is comparable and lower than a lowest tumorigenic dose (LTD>75 mg/kg/day) in rodents; while BMD for CF is lower than an LTD of CF (Fig. 6). Therefore, a short-term exposure to hepatotoxicants in rodents with a dose-response analysis of endpoints related to mode of action (e.g. apical, metabolic, and transcriptional changes in genes involved in epigenetic regulation) may predict liver carcinogenesis in experimental models. Analyses of additional time-points as well epigenetic and metabolomics endpoints will be required to further characterize these dose-response patterns.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Funding

This work was supported by the National Institutes of Health Center of Biological Research Excellence [grant number 1P20GM109005-01A1]; National Institutes of Health UAMS Clinical and Translational Science Award [grants number UL1TR000039, KL2TR000063]; and the Arkansas Biosciences Institute.

We are thankful to Drs. Kristy Kutanzi and Jessica LaRocca for critical reading and to Christopher Fettes for editing this manuscript.

Abbreviations

Ahr	Aryl Hydrocarbon receptor
BA	Bile Acid
BMD	Benchmark Dose
BrdU	Bromodeoxyuridine
CAR	Constitutive Androgen Receptor
CF	Clofibrate
ChIP	Chromatin Immunoprecipitation
DNA	Deoxyribonucleic Acid
Dnmt	DNA methyltransferase

LC/MS	Liquid Chromatography/Mass Spectrometry
LI	Labeling Indices
LINE-1	Long Interspersed Nuclear Element 1
MBD	Methyl-binding Domain
miRNA	microRNA
MS-PCR	Methylation-Sensitive quantitative Polymerase Chain Reaction
NMR	Nuclear Magnetic Resonance
NOAEL	No-Observed-Adverse-Effect Level
ORF	Open Reading Frame
PB	Phenobarbital
PBRE	Phenobarbital Response Element
PPARα	Peroxisome Proliferator-Activated Receptor alpha
PPRE	Peroxisome Proliferative Response Element
PXR	Pregnane X receptor
qRT-PCR	quantitative Real-Time Polymerase Chain Reaction
RNA	Ribonucleic Acid
SAH	S-adenosyl-homocysteine
TCA	Taurocholic Acid
TCDCA	Taurochenodeoxycholic Acid
TDCA	Taurodeoxycholic Acid
TSS	Transcription Start Site
TUCA	Tauroursocholic Acid
TUDCA	Tauroursodeoxycholic Acid

References

- Aldridge TC, Tugwood JD, Green S. Identification and Characterization of DNA Elements Implicated in the Regulation of Cyp4a1 Transcription. *Biochem J.* 1995; 306:473–479. [PubMed: 7887901]
- Alyea RA, Gollapudi BB, Rasoulpour RJ. Are we ready to consider transgenerational epigenetic effects in human health risk assessment? *Environ Mol Mutagen.* 2014; 55:292–298. [PubMed: 24259352]
- Alyea RA, Moore NP, LeBaron MJ, Gollapudi BB, Rasoulpour RJ. Is the current product safety assessment paradigm protective for epigenetic mechanisms? *J Pharmacol Tox Met.* 2012; 66:207–214.

- Baylin SB, Jones PA. A decade of exploring the cancer epigenome - biological and translational implications. *Nat Rev Cancer*. 2011; 11:726–734. [PubMed: 21941284]
- Butler WH. Long-Term Effects of Phenobarbitone-Na on Male Fischer Rats. *Brit J Cancer*. 1978; 37:418–423. [PubMed: 638020]
- Carreira PE, Richardson SR, Faulkner GJ. L1 retrotransposons, cancer stem cells and oncogenesis. *Febs J*. 2014; 281:63–73. [PubMed: 24286172]
- Chen P, Zhao JC, Wang Y, Wang M, Long HZ, Liang D, Huang L, Wen ZQ, Li W, Li X, Feng HL, Zhao HY, Zhu P, Li M, Wang QF, Li GH. H3.3 actively marks enhancers and primes gene transcription via opening higher-ordered chromatin. *Gene Dev*. 2013; 27:2109–2124. [PubMed: 24065740]
- Chen T, Li ZG, Yan J, Yang X, Salminen W. MicroRNA expression profiles distinguish the carcinogenic effects of riddelliine in rat liver. *Mutagenesis*. 2012; 27:59–66. [PubMed: 21976715]
- Corton JC, Cunningham ML, Hummer BT, Lau C, Meek B, Peters JM, Popp JA, Rhomberg L, Seed J, Klaunig JE. Mode of action framework analysis for receptor-mediated toxicity: The peroxisome proliferator-activated receptor alpha (PPAR alpha) as a case study. *Crit Rev Toxicol*. 2014; 44:1–49.
- Davis JA, Gift JS, Zhao QJ. Introduction to benchmark dose methods and US EPA's benchmark dose software (BMDS) version 2.1. 1. *Toxicol Appl Pharm*. 2011; 254:181–191.
- Sargent EV, Naumann BD, Dolan DG, Faria EC, Schulman L. The Importance of Human Data in the Establishment of Occupational Exposure Limits. *Human and Ecological Risk Assessment: An International Journal*. 2002; 8(4):805–822.
- Elcombe CR, Peffer RC, Wolf DC, Bailey J, Bars R, Bell D, Cattley RC, Ferguson SS, Geter D, Goetz A, Goodman JJ, Hester S, Jacobs A, Omiecinski CJ, Schoeny R, Xie W, Lake BG. Mode of action and human relevance analysis for nuclear receptor-mediated liver toxicity: A case study with phenobarbital as a model constitutive androstane receptor (CAR) activator. *Crit Rev Toxicol*. 2014; 44:64–82. [PubMed: 24180433]
- Eldridge SR, Tilbury LF, Goldsworthy TL, Butterworth BE. Measurement of Chemically-Induced Cell-Proliferation in Rodent Liver and Kidney - a Comparison of 5-Bromo-2'-Deoxyuridine and [H-3] Thymidine Administered by Injection or Osmotic Pump. *Carcinogenesis*. 1990; 11:2245–2251. [PubMed: 2265476]
- Estecio MRH, Gharibyan V, Shen LL, Ibrahim AEK, Doshi K, He R, Jelinek J, Yang AS, Yan PS, Huang THM, Tajara EH, Issa JJP. LINE-1 Hypomethylation in Cancer Is Highly Variable and Inversely Correlated with Microsatellite Instability. *Plos One*. 2007;2.
- Fiehn O. Metabolomics - the link between genotypes and phenotypes. *Plant Mol Biol*. 2002; 48:155–171. [PubMed: 11860207]
- Geter DR, Bhat VS, Gollapudi BB, Sura R, Hester SD. Dose-Response Modeling of Early Molecular and Cellular Key Events in the CAR-Mediated Hepatocarcinogenesis Pathway. *Toxicol Sci*. 2014; 138:425–445. [PubMed: 24449422]
- Girard M, Jacquemin E, Munnich A, Lyonnet S, Henrion-Caude A. miR-122, a paradigm for the role of microRNAs in the liver. *J Hepatol*. 2008; 48:648–656. [PubMed: 18291553]
- Hagiwara A, Miyata E, Tamano S, Sano M, Masuda C, Funae Y, Ito N, Fukushima S, Shirai T. Non-carcinogenicity, but dose-related increase in preneoplastic hepatocellular lesions, in a two-year feeding study of phenobarbital sodium in male F344 rats. *Food Chem Toxicol*. 1999; 37:869–879. [PubMed: 10506011]
- Hastings ML, Palma J, Duelli DM. Sensitive PCR-based quantitation of cell-free circulating microRNAs. *Methods*. 2012; 58:144–150. [PubMed: 22884953]
- Herceg Z, Lambert MP, van Veldhoven K, Demetriou C, Vineis P, Smith MT, Straif K, Wild CP. Towards incorporating epigenetic mechanisms into carcinogen identification and evaluation. *Carcinogenesis*. 2013; 34:1955–1967. [PubMed: 23749751]
- Honkakoski P, Zelko I, Sueyoshi T, Negishi M. The nuclear orphan receptor CAR-retinoid X receptor heterodimer activates the phenobarbital-responsive enhancer module of the CYP2B gene. *Mol Cell Biol*. 1998; 18:5652–5658. [PubMed: 9742082]
- Hou LF, Zhang X, Wang D, Baccarelli A. Environmental chemical exposures and human epigenetics. *Int J Epidemiol*. 2012; 41:79–105. [PubMed: 22253299]

- Hur K, Cejas P, Feliu J, Moreno-Rubio J, Burgos E, Boland CR, Goel A. Hypomethylation of long interspersed nuclear element-1 (LINE-1) leads to activation of proto-oncogenes in human colorectal cancer metastasis. *Gut*. 2014; 63:635–646. [PubMed: 23704319]
- Ishihara K, Katsutani N, Aoki T. A metabonomics study of the hepatotoxicants galactosamine, methylene dianiline and clofibrate in rats. *Basic Clin Pharmacol*. 2006; 99:251–260.
- Jiang YD, Zhang JZ, Xiong JT, Cao J, Li GZ, Wang SR. Ligands of peroxisome proliferator-activated receptor inhibit homocysteine induced DNA methylation of inducible nitric oxide synthase gene. *Acta Bioch Bioph Sin*. 2007; 39:366–376.
- Kawamoto T, Sueyoshi T, Zelko I, Moore R, Washburn K, Negishi M. Phenobarbital-responsive nuclear translocation of the receptor CAR in induction of the CYP2B gene. *Mol Cell Biol*. 1999; 19:6318–6322. [PubMed: 10454578]
- Kliwer SA, Moore JT, Wade L, Staudinger JL, Watson MA, Jones SA, McKee DD, Oliver BB, Willson TM, Zetterstrom RH, Perlmann T, Lehmann JM. An orphan nuclear receptor activated by pregnanes defines a novel steroid signaling pathway. *Cell*. 1998; 92:73–82. [PubMed: 9489701]
- Koturbash I, Beland FA, Pogribny IP. Role of epigenetic events in chemical carcinogenesis-a justification for incorporating epigenetic evaluations in cancer risk assessment. *Toxicol Mech Method*. 2011a; 21:289–297.
- Koturbash I, Beland FA, Pogribny IP. Role of microRNAs in the regulation of drug metabolizing and transporting genes and the response to environmental toxicants. *Expert Opin Drug Met*. 2012; 8:597–606.
- Koturbash I, Scherhag A, Sorrentino J, Sexton K, Bodnar W, Swenberg JA, Beland FA, de Villena FPM, Rusyn I, Pogribny IP. Epigenetic Mechanisms of Mouse Interstrain Variability in Genotoxicity of the Environmental Toxicant 1,3-Butadiene. *Toxicol Sci*. 2011b; 122:448–456. [PubMed: 21602187]
- Koturbash I, Scherhag A, Sorrentino J, Sexton K, Bodnar W, Tryndyak V, Latendresse JR, Swenberg JA, Beland FA, Pogribny IP, Rusyn I. Epigenetic Alterations in Liver of C57BL/6J Mice after Short-Term Inhalational Exposure to 1,3-Butadiene. *Environ Health Persp*. 2011c; 119:635–640.
- Koturbash I, Tolleson WH, Guo L, Yu DK, Chen S, Hong HX, Mattes W, Ning BT. microRNAs as pharmacogenomic biomarkers for drug efficacy and drug safety assessment. *Biomark Med*. 2015; 9:1153–1176. [PubMed: 26501795]
- Koufaris C, Wright J, Osborne M, Currie RA, Gooderham NJ. Time and dose-dependent effects of phenobarbital on the rat liver miRNAome. *Toxicology*. 2013; 314:247–253. [PubMed: 24157574]
- Lambert MP, Ancey PB, Esposti DD, Cros MP, Sklias A, Scoazec JY, Durantel D, Hernandez-Vargas H, Herceg Z. Aberrant DNA methylation of imprinted loci in hepatocellular carcinoma and after in vitro exposure to common risk factors. *Clin Epigenetics*. 2015; 7. [PubMed: 25628766]
- Legido-Quigley C, McDermott L, Vilca-Melendez H, Murphy GM, Heaton N, Lindon JC, Nicholson JK, Holmes E. Bile UPLC-MS fingerprinting and bile acid fluxes during human liver transplantation. *Electrophoresis*. 2011; 32:2063–2070. [PubMed: 21732555]
- Li B, Su T, Ferrari R, Li JY, Kurdiani SK. A unique epigenetic signature is associated with active DNA replication loci in human embryonic stem cells. *Epigenetics*. 2014; 9:257–267. [PubMed: 24172870]
- Martens JHA, O'Sullivan RJ, Braunschweig U, Opravil S, Radolf M, Steinlein P, Jenuwein T. The profile of repeat-associated histone lysine methylation states in the mouse epigenome. *Embo J*. 2005; 24:800–812. [PubMed: 15678104]
- Miousse IR, Chalbot MCG, Lumen A, Ferguson A, Kavouras IG, Koturbash I. Response of transposable elements to environmental stressors. *Mutat Res-Rev Mutat*. 2015a; 765:19–39.
- Miousse IR, Currie R, Datta K, Ellinger-Ziegelbauer H, French JE, Harrill AH, Koturbash I, Lawton M, Mann D, Meehan RR, Moggs JG, O'Lone R, Rasoulpour RJ, Pera RAR, Thompson K. Importance of investigating epigenetic alterations for industry, and regulators: An appraisal of current efforts by the Health and Environmental Sciences Institute. *Toxicology*. 2015b; 335:11–19. [PubMed: 26134581]
- Miousse IR, Koturbash I. The Fine LINE: Methylation Drawing the Cancer Landscape. *Biomed Res Int*. 2015

- Nie AY, McMillian M, Parker JB, Leone A, Bryant S, Yieh L, Bittner A, Nelson J, Carmen A, Wan J, Lord PG. Predictive toxicogenomics approaches reveal underlying molecular mechanisms of nongenotoxic carcinogenicity. *Mol Carcinogen*. 2006; 45:914–933.
- Ohta T, Masutomi N, Tsutsui N, Sakairi T, Mitchell M, Milburn MV, Ryals JA, Beebe KD, Guo L. Untargeted Metabolomic Profiling as an Evaluative Tool of Fenofibrate-Induced Toxicology in Fischer 344 Male Rats. *Toxicol Pathol*. 2009; 37:521–535. [PubMed: 19458390]
- Palmer CNA, Hsu MH, Muerhoff AS, Griffin KJ, Johnson EF. Interaction of the Peroxisome Proliferator-Activated Receptor-Alpha with the Retinoid-X Receptor-Alpha Unmasks a Cryptic Peroxisome Proliferator Response Element That Overlaps an Arp-1-Binding Site in the Cyp4a6 Promoter. *J Biol Chem*. 1994; 269:18083–18089. [PubMed: 8027069]
- Phillips JM, Burgoon LD, Goodman JJ. Phenobarbital Elicits Unique, Early Changes in the Expression of Hepatic Genes that Affect Critical Pathways in Tumor-Prone B6C3F1 Mice. *Toxicol Sci*. 2009; 109:193–205. [PubMed: 19270015]
- Phillips JM, Goodman JJ. Multiple Genes Exhibit Phenobarbital-Induced Constitutive Active/Androstane Receptor-Mediated DNA Methylation Changes during Liver Tumorigenesis and in Liver Tumors. *Toxicol Sci*. 2009; 108:273–289. [PubMed: 19233941]
- Pogribny IP, Starlard-Davenport A, Tryndyak VP, Han T, Ross SA, Rusyn I, Beland FA. Difference in expression of hepatic microRNAs miR-29c, miR-34a, miR-155, and miR-200b is associated with strain-specific susceptibility to dietary nonalcoholic steatohepatitis in mice. *Lab Invest*. 2010; 90:1437–1446. [PubMed: 20548288]
- Priestley CC, Anderton M, Doherty AT, Duffy P, Mellor HR, Powell H, Roberts R. Epigenetics - relevance to drug safety science. *Toxicol Res-Uk*. 2012; 1:23–31.
- Rasoulpour RJ, LeBaron MJ, Ellis-Hutchings RG, Klapacz J, Gollapudi BB. Epigenetic screening in product safety assessment: are we there yet? *Toxicol Mech Method*. 2011; 21:298–311.
- Reddy JK, Qureshi SA. Tumorigenicity of Hypolipemic peroxisome proliferator Ethyl-Alpha-P-Chlorophenosisobutyrate (Clobifate) in rats. *Br J Cancer*. 1979; 40:476–482. [PubMed: 508572]
- Rubtsov DV, Waterman C, Currie RA, Waterfield C, Salazar JD, Wright J, Griffin JL. Application of a Bayesian Deconvolution Approach for High-Resolution H-1 NMR Spectra to Assessing the Metabolic Effects of Acute Phenobarbital Exposure in Liver Tissue. *Anal Chem*. 2010; 82:4479–4485. [PubMed: 20446676]
- Shah YA, Morimura K, Yang Q, Tanabe T, Takagi M, Gonzalez FJ. Peroxisome proliferator-activated receptor alpha regulates a microRNA-mediated signaling cascade responsible for hepatocellular proliferation. *Mol Cell Biol*. 2007; 27:4238–4247. [PubMed: 17438130]
- Strauss V, Mellert W, Wiemer J, Leibold E, Kamp H, Walk T, Looser R, Prokoudine A, Fabian E, Krennrich G, Herold M, van Ravenzwaay B. Increased toxicity when fibrates and statins are administered in combination - A metabolomics approach with rats. *Toxicol Lett*. 2012; 211:187–200. [PubMed: 22484644]
- Stribinskis V, Ramos KS. Activation of human long interspersed nuclear element 1 retrotransposition by benzo(a)pyrene, an ubiquitous environmental carcinogen. *Cancer Res*. 2006; 66:2616–2620. [PubMed: 16510580]
- Tan YX, Yin PY, Tang L, Xing WB, Huang Q, Cao D, Zhao XJ, Wang WZ, Lu X, Xu ZL, Wang HY, Xu GW. Metabolomics Study of Stepwise Hepatocarcinogenesis From the Model Rats to Patients: Potential Biomarkers Effective for Small Hepatocellular Carcinoma Diagnosis. *Mol Cell Proteomics*. 2012;11.
- Thomson JP, Lempiainen H, Hackett JA, Nestor CE, Muller A, Bolognani F, Oakeley EJ, Schubeler D, Terranova R, Reinhardt D, Moggs JG, Meehan RR. Non-genotoxic carcinogen exposure induces defined changes in the 5-hydroxymethylome. *Genome Biol*. 2012;13.
- Thomson JP, Moggs JG, Wolf CR, Meehan RR. Epigenetic profiles as defined signatures of xenobiotic exposure. *Mutat Res-Gen Tox En*. 2014; 764:3–9.
- Tryndyak VP, Ross SA, Belland FA, Pogribny IP. Down-Regulation of the microRNAs miR-34a, miR-127, and miR-200b in Rat Liver During Hepatocarcinogenesis Induced by a Methyl-Deficient Diet. *Mol Carcinogen*. 2009; 48:479–487.
- Unterberger EB, Eichner J, Wrzodek C, Lempiainen H, Luisier R, Terranova R, Metzger U, Plummer S, Knorpp T, Braeuning A, Moggs J, Templin MF, Honndorf V, Piotto M, Zell A, Schwarz M. Ha-

- ras and beta-catenin oncoproteins orchestrate metabolic programs in mouse liver tumors. *Int J Cancer*. 2014; 135:1574–1585. [PubMed: 24535843]
- Vliegenthart ADB, Shaffer JM, Clarke JI, Peeters LEJ, Caporali A, Bateman DN, Wood DM, Dargan PI, Craig DG, Moore JK, Thompson AI, Henderson NC, Webb DJ, Sharkey J, Antoine DJ, Park BK, Bailey MA, Lader E, Simpson KJ, Dear JW. Comprehensive microRNA profiling in acetaminophen toxicity identifies novel circulating biomarkers for human liver and kidney injury. *Sci Rep-Uk*. 2015:5.
- Waldmann T, Schneider R. Targeting histone modifications - epigenetics in cancer. *Curr Opin Cell Biol*. 2013; 25:184–189. [PubMed: 23347561]
- Wang B, Majumder S, Nuovo G, Kutay H, Volinia S, Patel T, Schmittgen TD, Croce C, Ghoshal K, Jacob ST. Role of MicroRNA-155 at Early Stages of Hepatocarcinogenesis Induced by Choline-Deficient and Amino Acid-Defined Diet in C57BL/6 Mice. *Hepatology*. 2009; 50:1152–1161. [PubMed: 19711427]
- Waterman CL, Currie RA, Cottrell LA, Dow J, Wright J, Waterfield CJ, Griffin JL. An integrated functional genomic study of acute phenobarbital exposure in the rat. *Bmc Genomics*. 2010:11. [PubMed: 20053297]
- Watson RE, Goodman JI. Effects of phenobarbital on DNA methylation in GC-rich regions of hepatic DNA from mice that exhibit different levels of susceptibility to liver tumorigenesis. *Toxicol Sci*. 2002; 68:51–58. [PubMed: 12075110]
- Wei P, Zhang J, Egan-Hafley M, Liang SG, Moore DD. The nuclear receptor CAR mediates specific xenobiotic induction of drug metabolism. *Nature*. 2000; 407:920–923. [PubMed: 11057673]
- Wheelock CE, Goto S, Hammock BD, Newman JW. Clofibrate-induced changes in the liver, heart, brain and white adipose lipid metabolome of Swiss-Webster mice. *Metabolomics*. 2007; 3:137–145. [PubMed: 19079556]
- Whitlock JP. Induction of cytochrome P4501A1. *Annu Rev Pharmacol*. 1999; 39:103–125.
- Wignall JA, Shapiro AJ, Wright FA, Woodruff TJ, Chiu WA, Guyton KZ, Rusyn I. Standardizing Benchmark Dose Calculations to Improve Science-Based Decisions in Human Health Assessments. *Environ Health Persp*. 2014; 122:499–505.
- Willson TM, Kliewer SA. PXR, CAR and drug metabolism. *Nat Rev Drug Discov*. 2002; 1:259–266. [PubMed: 12120277]
- Xie W, Barwick JL, Simon CM, Pierce AM, Safe S, Blumberg B, Guzelian PS, Evans RM. Reciprocal activation of Xenobiotic response genes by nuclear receptors SXR/PXR and CAR. *Gene Dev*. 2000; 14:3014–3023. [PubMed: 11114890]

Highlights

- Exposure to phenobarbital or clofibrate resulted in dose-dependent increases in apical endpoints.
- Alterations in the expression of CYP450 genes were mediated by histone modifications.
- Hypomethylation of LINE-1 DNA and dose-dependent changes in the expression of microRNAs were detected.
- Metabolomics data indicated alterations in the metabolism of bile acids.

Fig. 1.

Relative liver weight and cellular proliferation after 7-day administration of clofibrate (CF) or phenobarbital (PB). A) At 50 and 250 mg/kg/day CF, relative liver weights were increased 9 and 20%. PB exposure resulted in dose-dependent increases in relative liver weights ranging from 8 to 41% at all dose levels and were statistically-identified at 25 and 100 mg/kg/day PB. Data are expressed as g of liver/100 g of body weight. Mean $\pm$ SD. *- Statistically different from control via one-way ANOVA and Dunnett's post-test ($p < 0.05$). B) Hepatocellular proliferation. Across all zones, CF increased hepatocellular proliferation at 250 mg/kg/day. PB exposure caused a dose-dependent increase in hepatocellular proliferation at all dose levels and was statistically-identified at 25 and 100 mg/kg/day. Data are mean percent labeling index per 1000 cells counted. CL = centrilobular, MZ = midzonal, PP = periportal, MKD – mg/kg/day. Mean $\pm$ SD. *- Statistically different from control via one-way ANOVA and Dunnett's post-test ($p < 0.05$).

Fig. 2.

Expression of key cytochrome P450 genes and histone modifications associated with their transcriptional activation. A) For CF, *Cyp4a1* mRNA was induced at 50 and 250 mg/kg/day, reaching 23-fold induction at 250 mg/kg/day. For PB, *Cyp2b1* mRNA was increased at all dose levels, attaining a 180-fold induction at the high dose. Data are fold induction compared to control. * - Statistically different from control via one-way ANOVA and Dunnett's post-test ($p < 0.05$). Chromatin Immunoprecipitation (ChIP) analysis of the (B) transcription start sites (TSS) and (C) phenobarbital-responsive element (PBRE) of *Cyp4a1* and *Cyp2b1* after 7-days exposure to CF or PB. * - Statistically different from control via one-way ANOVA and Dunnett's post-test ($p < 0.05$).

Fig. 3.

Expression of genes involved in methylation in the liver following 7-day exposure to CF (A) or PB (B). The methyltransferases *Dnmt1* and *Dnmt3b* were down-regulated in the highest doses of CF and PB, respectively, while the methyl-binding protein *Mbd1* was down-regulated in both treatments. C) Levels of *Dnmt1* in the livers of CF or PB treated rats as analyzed by Western blot. * - Statistically different from control via one-way ANOVA and Dunnett's post-test ($p < 0.05$).

Fig. 4. Pyrosequencing-based analysis of LINE-1 methylation in the liver and kidney following 7-day exposure to CF or PB. While CF did not alter LINE-1 element methylation, 100 mg/kg/day PB exposure caused a liver-specific and statistically significant reduction in LINE-1 element methylation. **** - Statistically different from control via one-way ANOVA and Dunnett's post-test ($p < 0.0001$).

Fig. 5.

Metabolic profiles of the rat livers following 7-day exposure to CF or PB. Effects of CF (A) and PB (B) on the bile acids and CF (C) and PB (D) on metabolites involved in oxidative stress and DNA methylation. * - Statistically different from control via one-way ANOVA and Dunnett's post-test ($p < 0.05$).

Fig. 6. Correlation of benchmark dose estimates for transcriptional changes in genes involved in DNA methylation, metabolic and apical endpoints in rat exposed to CF and PB. Horizontal dot line and hatched-line define the 75 mg/kg/day lowest tumorigenic dose (LTD) for PB and 250 mg/kg/day LTD for CF.

Table 1

Hepatocellular Hypertrophy Following Clofibrate or Phenobarbital Exposure

Hepatocellular Hypertrophy	Control	Clofibrate (mg/kg/day)			Phenobarbital (mg/kg/day)		
		10	50	250	5	25	100
Centrilobular/Midzonal							
Very slight	0/5	0/5	5/5	0/5	5/5	0/5	0/5
Slight	0/5	0/5	0/5	0/5	0/5	5/5	0/5
Moderate	0/5	0/5	0/5	0/5	0/5	0/5	5/5
Panlobular							
Very slight	0/5	0/5	0/5	5/5	0/5	0/5	0/5

Values displayed in bold were deemed to be due to clofibrate or phenobarbital exposure.

Table 2

miRNAs in rodent experimental hepatocarcinogenesis

miRNA	CF 10 MKD	CF 50 MKD	CF 250 MKD	PB 5 MKD	PB 25 MKD	PB 100 MKD
miR-122	-1.0	-1.2	-1.9**	-1.2	-1.2	-1.8*
miR-125a	-1.3	-1.4	-2.1**	-1.3	-1.5	-1.9*
miR-155	-1.7	-1.7	-2.2*	-1.3	-1.3	-2.0*
miR-199a-5p	1.0	-1.3	-1.9*	-1.1	-1.2	-1.7*
miR-199a-3p	-1.1	-1.3	-1.8*	-1.2	-1.3	-1.7**
miR-200b	1.1	-1.1	1.3	2.0*	2.3*	-1.6
miR-221	-1.2	-1.2	1.1	-1.2	-1.4	-2.1*

*
p 0.05;

**
p 0.01.

Measurements for which fold regulation than ± 1.5 and *p* value 0.05 are displayed in bold. CF – clofibrate, PB – phenobarbital; MKD – mg/kg/day.

Table 3

Benchmark dose estimates for apical and hepatic biomarker endpoints in rats exposed to CF and PB

	CF		PB	
	BMD (mg/kg/day)	BMDL (mg/kg/day)	BMD (mg/kg/day)	BMDL (mg/kg/day)
Relative liver weight	23.4155	18.2498	18.658	14.1473
<i>Cyp1a1</i> gene expression	142.798	90.2566	46.3974	30.8913
<i>Cyp2b1</i> gene expression	88.4215	62.3811	8.00736	6.25614
<i>Cyp3a1</i> gene expression	-	-	7.64078	5.97364
<i>Cyp4a1</i> gene expression	17.1943	13.4614	-	-
Hepatocyte cell proliferation				
Centrilobular	116.276	77.5801	19.4841	14.7285
Midzonal	77.6276	55.9088	27.4652	20.0801
Periportal	72.8438	52.9301	50.8573	33.0902
Total	90.2419	63.4394	31.7626	22.7607

Table 4

Benchmark dose estimates for transcriptional changes in genes involved in DNA methylation in rats exposed to CF and PB

	CF		PB	
	BMD (mg/kg/day)	BMDL (mg/kg/day)	BMD (mg/kg/day)	BMDL (mg/kg/day)
Dnmt1 gene expression	101.152	71.2198	77.4818	44.0625
Dnmt3a gene expression	163.575	99.1379	109.24	53.6435
Dnmt3b gene expression	-	-	50.4114	32.8757
<i>Mecp2</i> gene expression	148.732	92.8808	53.0571	34.1315
<i>Mbd1</i> gene expression	148.263	92.6761	51.0163	33.1664

Table 5

Metabolic benchmark dose estimates in rats exposed to CF and PB

	CF		PB	
	BMD (mg/kg/day)	BMDL (mg/kg/day)	BMD (mg/kg/day)	BMDL (mg/kg/day)
Cholic acid	140.081	89.0305	-	-
Deoxycholic acid	-	-	321.125	81.1221
Glycocholic acid	153.92	95.1166	95.7406	49.9375
Glycocholic acid isomer	-	-	93.8663	49.3835
Sulfolithcholyglycine	240.591	125.828	76.7072	43.7876
Sulfolithcholyglycine isomer	136.372	87.3312	43.6874	29.4951
Taurocholic acid (TCA)	177.45	104.618	91.1613	48.5655
Tauroursocholic acid (TUCA)	72.3732	52.6333	28.7302	20.8837
Taurodeoxycholic acid (TDCA)	87.4564	61.8163	-	-
Taurochenodeoxycholic acid (TCDCA)	140.209	89.0889	98.8181	50.8252
Tauroursodeoxycholic acid (TUDCA)	58.3373	43.475	45.3749	30.37
Methionine	230.008	122.645	-	-
Pyroglutamate	-	-	69.6838	41.1856
S-adenosyl-homocysteine (SAH)	-	-	75.7325	43.4383
Taurine	141.174	89.5258	-	-