Thoracic Spondylodiscitis Epidural Abscess in an Afebrile Navy Veteran: A Case Report

Zachary A. Cupler, DC, MS,^a Michael T. Anderson, DC, MS,^a and Thomas J. Stancik, DO^b

ABSTRACT

Objective: The purpose of this case study was to describe the differential diagnosis of a thoracic epidural abscess in a Navy veteran who presented to a chiropractic clinic for evaluation and management with acupuncture within a Veterans Affairs Medical Center.

Clinical Features: An afebrile 59-year-old man with acute thoracic spine pain and chronic low back pain presented to the chiropractic clinic at a Veterans Affairs Medical Center for consideration for acupuncture treatment.

Intervention and Outcome: The veteran elected to trial acupuncture once per week for 4 weeks. A routine thoracic magnetic resonance imaging scan without gadolinium detected a space-occupying lesion after the patient failed to attain 50% reduction of pain within 2 weeks with conservative care. The patient was diagnosed with a multilevel thoracic spondylodiscitis epidural abscess and was treated same day with emergency debridement and laminectomy of T7-8 with a T6-9 fusion. The patient had complete recovery without neurological compromise and completed an antibiotic regimen for 6 weeks.

Conclusion: A Navy veteran with acute thoracic spine and chronic low back pain appeared to respond initially but failed to achieve clinically meaningful outcomes. Follow-up advanced imaging detected a thoracic spondylodiscitis epidural abscess. Early diagnosis and immediate intervention are important to preserving neurological function and limiting morbidity in cases of spondylodiscitis epidural abscess. (J Chiropr Med 2017;16:246-251)

Key Indexing Terms: *Acupuncture Therapy; Chiropractic; Epidural Abscess; Infection; Thoracic; Veterans; Military Personnel*

INTRODUCTION

Back and neck pain are leading global causes of disability.¹ Back pain caused by serious underlying pathology accounts for less than 3% of all back pain.² A spinal epidural abscess (SEA) is a rare, life-threatening infection, with the incidence on the rise in recent decades. The rate of SEA in 1975 was 0.2 to 2.0 cases per 10000.³ In a study of tertiary referral centers in 1999, 12.5 cases per 10000 were reported.⁴ Early diagnosis is paramount to mitigate neurological compromise and limit morbidity. A history of diabetes mellitus (21%-38% of cases), intravenous (IV) drug use, end-stage renal disease, human immunodeficiency virus (HIV), or immunosuppressant use and iatrogenic causes, such as epidural spinal procedures, increase the risk of SEA.⁵⁻⁷

Cases of SEA have been documented as presenting to chiropractic offices often masquerading as non-specific neck or back pain.⁸⁻¹⁰ The classic clinical triad of back pain, fever, and neurological deficits is observed in only ~10% of cases.¹¹ Clinical progression of SEA has been observed to comprise four phases: (1) spinal ache, (2) root pain, (3) weakness of voluntary muscles and sphincters, followed by (4) paralysis.¹²

Because of the pervasive nature of neck and back pain, misdiagnosis of SEA on initial presentation is noteworthy, with estimates ranging from 11% to 75%.^{7,13} A SEA, on average, crosses 3.85 vertebral levels and is reported in the thoracic (15%-39.1%), lumbosacral (30%-54.7%), and cervical (9%-35.9%) regions.^{6,7,14,15} Magnetic resonance imaging (MRI) examination with and without gadolinium of the entire spine is the gold standard, as it differentiates SEA from other space-occupying lesions and malignancy. A computed tomography (CT) myelogram should be obtained when MRI is unavailable or is contraindicated because of an implanted electrical device (eg, pacemaker, spinal stimulator).^{5,11,15} The infectious agent causing SEA is most commonly of bacterial origin and overwhelmingly either methicillin-sensitive or methicillin-resistant *Staphylococcus aureus*.^{7,13} Once patients are diagnosed with SEA, best outcomes are achieved with rapid surgical intervention, abscess drainage, and concomitant antibiotic treatment. When monitored closely for clinical deterioration, there are select cases with limited neurological deficit that may be managed with antibiotic therapy only.^{11,13}

^a Physical Medicine & Rehabilitative Services, VA Butler Healthcare, Butler, Pennsylvania.

^b Primary Care, VA Butler Healthcare, Butler, Pennsylvania.

Corresponding Author: Zachary A Cupler, DC, MS, VA Butler Healthcare, 353 North Duffy Road, Butler, PA 16001. Tel.: +1 800-362-8262. (e-mail: Zachary.cupler@va.gov).

Paper submitted January 18, 2017; in revised form February 13, 2017; accepted March 28, 2017.

1556-3707

© 2017 National University of Health Sciences.

<http://dx.doi.org/10.1016/j.jcm.2017.03.002>

Multiple published SEA cases have suggested acupuncture to be the causative route of infection.¹⁶⁻³¹ In this report, we describe a case of SEA in a man who presented to a chiropractic office for acupuncture at a Veterans Affairs Medical Center. The purpose of this case study is to describe the differential diagnosis of a thoracic epidural abscess in a Navy veteran who presented to a chiropractic clinic for evaluation and management with acupuncture within a Veterans Affairs Medical Center.

CASE REPORT

A 59-year-old man with chronic low back pain and left leg pain below the knee was referred by his primary care physician (PCP) to a chiropractor for evaluation and consideration for acupuncture. The patient developed acute, non-traumatic left-sided thoracic pain 8 days prior to his initial appointment. He reported his thoracic spine pain was significantly relieved following a bowel movement the day prior, yet remained more prominent than low back and left leg pain.

The patient had Minor's sign and an antalgic gait. On a numeric pain scale from 0 to 10, he reported 7 to 8 for left-sided thoracic pain. He described the pain as sharp and "much better than yesterday" following his first bowel movement in 8 days. Valsalva maneuver exacerbated symptoms in the lumbar and thoracic regions, as did positions sustained longer than 30 minutes. Medical history was remarkable for left L5-S1 hemilaminotomy in 2011 and C2-6 cervical fusion in 1983. He smoked 1 pack per day and consumed alcohol occasionally. His medications included gabapentin, meloxicam, hydrocodone, and cyclobenzaprine. He denied recent fever, chills, nausea, vomiting, motor weakness, night pain, night sweats, and bowel or bladder dysfunction.

Vitals were collected by his PCP 3 days prior to evaluation. His temperature was 36.1°C, pulse rate was 88 beats per minute, and respirations were 20 per minute. Sensory and motor examinations were within normal limits. Straight-leg raise exacerbated chronic left radiculopathy. Dural slump test was positive for lumbar radiculopathy. Segmental palpation revealed tenderness throughout the lower thoracic spine, which was more pronounced on the left.

The veteran agreed to a course of acupuncture one time per week for 4 weeks. Short-term goals included increasing range of motion and reducing pain. Long-term goals included a sustained reduction of pain by at least 50% within 2 weeks and increased sustained positioning without aggravation of back pain from 30 to 60 minutes. On the initial visit, auricular acupuncture was applied bilaterally, and needles were retained for 10 to 15 minutes while the patient rested quietly. The National Acupuncture Detoxification Association (NADA) protocol is used primarily in mental health settings, as well as chronic

pain management. It is a standardized 5-point ear acupuncture protocol: shen men, sympathetic, liver, kidney and lung.³² Acupuncture was not applied to the thoracolumbar paraspinal regions. The patient exited with a verbalized pain score of 2 of 10.

On three subsequent appointments within 3 weeks after the initial encounter, acupuncture needles were inserted at an oblique angle and perpendicular to the thoracic paraspinals while the patient rested in a prone position. On the third visit, the thoracic pain radiated from the spine to the anterior rib cage with Valsalva-type maneuvers. In conjunction with acupuncture on the third visit, high-velocity, low-amplitude spinal manipulation was applied to the midthoracic region. The patient reported relief from thoracic and lumbar pain for the remainder of the day following each treatment. The pain returned to pretreatment intensity the evening after each appointment.

Vitals were collected again 3 weeks after initial consult with chiropractic services. Weight was stable at 67.4 kg. Temperature was 36.1°C, blood pressure was 118/62 mm Hg on the left arm, respirations were 20 breaths per minute, and heart rate was 68.5 beats per minute.

The patient's PCP ordered a thoracic MRI scan without contrast 4 weeks after symptom onset. The PCP chose to forgo plain film imaging because of the acute and non-traumatic nature of the thoracic pain, the long-term relationship with the patient, and the lack of sustainable response to conservative care after 3 sessions. Initial MRI findings without contrast appreciated abnormal T2 hyper-intense and T1 hypointense signal throughout T6 and T7 vertebral bodies with an epidural mass appreciated anterior and posterior to the cord with paraspinal extension and a dorsal mass effect of the cord (Figs 1-3).

Emergency thoracic MRI enhanced with gadolinium concluded there was intense enhancement of the T6 and T7 vertebral bodies and T6-7 disk with enhancing epidural soft tissue. This further confirmed a diagnosis of discitis and osteomyelitis with epidural abscess and phlegmon of the T6 and T7 regions. An abscess exhibits rim enhancement, whereas a phlegmon is granular tissue containing micro-abscesses without purulence, and MRI reveals a homogenous enhancement of inflammatory cells that causes high T2 and low T1 signals.³³ Complete blood count and comprehensive metabolic panel in the emergency room were negative for sepsis and non-specific inflammation. Temperature and neurological condition remained stable with hyporeflexia of the right patella.

He underwent same-day emergency debridement and laminectomy of T7-8 with a T6-9 fusion. Postsurgically, he was treated with an antibiotic regimen of cefepime, flagyl, and IV vancomycin. Tissue samples were positive for methicillin-sensitive *Staphylococcus aureus* (MSSA), and IV vancomycin was changed to oxacillin and rifampin. The patient was discharged in stable condition with a 6-week antibiotic home regimen.

Fig 1. Thoracic T2-weighted midsagittal image without gadolinium reveals an abnormal hyper-intense signal of the T6 and T7 vertebral bodies and T6-7 disk space. Epidural masses anterior and posterior to the cord with paraspinous extension are observed.

DISCUSSION

Acupuncture is commonly used to treat back, neck, and joint pain. Since 1982, there have been at least 16 reports in the English literature of SEA secondary to acupuncture

Fig 2. Thoracic T1-weighted midsagittal image without gadolinium reveals an abnormal hypointense signal throughout the T6 and T7 vertebral bodies.

treatment.¹⁶⁻³¹ It has been proposed that poorly controlled hygiene in acupuncture practice might lead to epidural infections.^{17-19,22,25} All invasive procedures that break the skin carry the risk of infection. Appropriate clean needle technique is required to minimize the risk of infection.³⁴ According to experimentation from the 1950s, injection of 7500000 abscess-derived *S. aureus* colonies is the minimum required to create a local skin infection in otherwise healthy volunteers. A liberal assumption suggests an acupuncture needle could push into the skin a maximum of 10000 organisms, less than reported to create an infection in otherwise healthy individuals.³⁵ Although acupuncture cannot be denied as a plausible explanation for the route of infection, it appears more likely than not in our case, considering the nature and timeline, that the SEA was the underlying pathology prior to presentation to the chiropractic clinic. Once the spine practitioner suspects an SEA, there can be no delay in care because of the high morbidity and mortality associated with this diagnosis.

This case of SEA was difficult to diagnose. The patient's history and physical exam findings limit clinical suspicion to consider an SEA. There was no recent history of infection, illness, or surgical or invasive procedure, nor did the individual have risk factors such as diabetes, IV drug use, or an immunocompromised state. The patient did not have an elevated temperature nor was sepsis detected, and there was no evidence of neurological deterioration. Twenty-two percent of cases have no discernable risk factors associated with SEA.⁷ Fever has been reported in 50% to 66% of patients diagnosed with SEA.^{6,7,15}

Neurologically intact SEA patients can be considered for IV antibiotic therapy and monitored closely for deterioration, which would necessitate surgical intervention.⁷ Antibiotics are administered for 4-6 weeks.⁵ Four risk factors have arisen to predict failure of medical management of SEA: diabetes mellitus, leukocytosis greater than $12.5 \times 10^3/\mu\text{L}$, positive blood cultures, and C-reactive protein $>115 \text{ mg/L}$.⁷

Delay in surgical intervention has been found to limit functional recovery when compared with immediate surgical intervention.³⁶ Operative management includes debridement and decompression and, if necessary, fixation of the compromised spinal column; laminectomy is the most common procedure.^{7,11} The mortality rate is 3% after surgical intervention, whereas post-surgical complications occur in 26% of cases.¹¹

Previous case reports have suggested acupuncture performed with poor hygiene may be the route of infection of SEA.^{17-19,22,25} SEA can be difficult to detect as in this case, no discernable infectious state was observed. The purpose of this case report was to highlight that SEA patients may present for acupuncture, and acupuncture may not be the cause of the SEA. Although it is not possible to clearly state acupuncture was not the route of infection, an otherwise healthy man without infection or immunocompromised state

Fig 3. A, T2-weighted axial image without gadolinium at the level of T6 vertebrae. The white arrow identifies a spinal cord signal. B, T2-weight axial image without gadolinium at the level of T7-8 disk reveals mass effect to the cord. The white arrow indicates the cord signal.

developed an SEA. There is a lack of clear imaging evidence to support route of infection to the superficial paraspinal tissue where acupuncture needles were inserted for treatment. There was no sudden onset of fever or erythema following the introduction of acupuncture, and neurologic status remained stable up until detection of the SEA. In addition, the complaint responded, though short-lived, to acupuncture and spinal manipulation, suggesting a combination of mechanical cause and chronic pain state.

It is pertinent for spine practitioners to assess for serious underlying pathology in spine-related disorders on initial evaluations as well as on follow-up visits. Spine practitioners, including those who perform acupuncture, are urged to consider SEA as a differential diagnosis when the patient presents with spinal pain and fever or infection.

Limitations

This study reports the findings from one patient. Other patients may have different findings and may respond differently to treatment. Each patient's case should be addressed in a unique manner. Thus, the findings in this case may not necessarily match those of other patients who have SEA.

CONCLUSION

A Navy veteran with acute thoracic spine and chronic low back pain appeared to initially respond to acupuncture, but failed to achieve a clinically meaningful outcome. Follow-up advanced imaging revealed a thoracic SEA, despite no elevation in temperature on multiple occasions or neurological deterioration. The spine practitioner is reminded that early diagnosis and immediate intervention are important to preserving neurological function and limiting morbidity in cases of SEA.

FUNDING SOURCES AND CONFLICTS OF INTEREST

This work was conducted at and supported by VA Butler Healthcare. No conflicts of interest were reported for this study.

CONTRIBUTORSHIP INFORMATION

Concept development (provided idea for the research): Z.A.C., M.T.A.
Design (planned the methods to generate the results): Z.A.C., M.T.A.
Supervision (provided oversight, responsible for organization and implementation, writing of the manuscript): Z.A.C.
Data collection/processing (responsible for experiments, patient management, organization, or reporting data): Z.A.C., T.J.S.
Analysis/interpretation (responsible for statistical analysis, evaluation, and presentation of the results): Z.A.C., M.T.A., T.J.S.
Literature search (performed the literature search): Z.A.C., M.T.A.
Writing (responsible for writing a substantive part of the manuscript): Z.A.C.
Critical review (revised manuscript for intellectual content, this does not relate to spelling and grammar checking): M.T.A., T.J.S.

Practical Applications

- Spine practitioners must be cognizant of risk factors in combination with history and exam findings of a spinal epidural abscess.
- This case demonstrates a spinal epidural abscess in a man in stable neurological condition without known risk factors, recent fever, or chills.
- Failure to respond to conservative treatment such as acupuncture resulted in advanced imaging that detected the spinal epidural abscess.

REFERENCES

- Vos T, Allen C, Arora M, et al. Global, regional, and national incidence, prevalence, and years lived with disability for 310 diseases and injuries, 1990–2015: a systematic analysis for the global burden of disease study 2015. *Lancet*. 2016;388(10053):1545-1602.
- Chou R, Qaseem A, Snow V, et al. Diagnosis and treatment of low back pain: a joint clinical practice guideline from the American college of physicians and the American pain society. *Ann Intern Med*. 2007;147(7):478-491.
- Baker AS, Ojemann RG, Swartz MN, Richardson Jr EP. Spinal epidural abscess. *N Engl J Med*. 1975;293(10):463-468.
- Rigamonti D, Liem L, Sampath P, et al. Spinal epidural abscess: contemporary trends in etiology, evaluation, and management. *Surg Neurol*. 1999;52(2):189-196 discussion 197.
- DeFroda SF, DePasse JM, Eltorai AE, Daniels AH, Palumbo MA. Evaluation and management of spinal epidural abscess. *J Hosp Med*. 2016;11(2):130-135.
- Adogwa O, Karikari IO, Carr KR, et al. Spontaneous spinal epidural abscess in patients 50 years of age and older: a 15-year institutional perspective and review of the literature. *J Neurosurg Spine*. 2014;20(3):344-349.
- Patel AR, Alton TB, Bransford RJ, Lee MJ, Bellabarba CB, Chapman JR. Spinal epidural abscess: risk factors, medical versus surgical management, a retrospective review of 128 cases. *Spine J*. 2014;24(2):326-330.
- Murphy DR, Carr BM. Spinal epidural abscess in a patient with chronic, recurrent low back pain. *J Neuromusc Sys*. 1999;7(2):70-73.
- Murphy DR, Morris NJ. Cervical epidural abscess in an afebrile patient: a case report. *J Manipulative Physiol Ther*. 2006;29(8):672-675.
- Wang CC, Kuo JR, Chio CC, Tsai TC. Acute paraplegia following chiropractic therapy. *J Clin Neurosci*. 2006;13(5):578-581.
- Singleton J, Edlow JA. Acute nontraumatic back pain: risk stratification, emergency department management and review of serious pathologies. *Emerg Med Clin North Am*. 2016;34(4):743-757.
- Heusner AP. Nontuberculous spinal epidural infections. *N Engl J Med*. 1948;239(23):845-854.
- Krishnamohan P, Berger JR. Spinal epidural abscess. *Curr Infect Dis Rep*. 2014;16(11):436.
- Schoenfeld AJ, Wahlquist TC. Mortality, complication risk, and total charges after the treatment of epidural abscess. *Spine J*. 2015;25(2):249-255.
- Reihnsaus E, Waldbaur H, Seeling W. Spinal epidural abscess: a meta-analysis of 915 patients. *Neurosurg Rev*. 2000;23(4):175-204 discussion 205.
- Godhania V. Lumbar spine osteomyelitis and epidural abscess formation secondary to acupuncture. *J Surg Case Rep*. 2016;2016(3).
- Chan JJ, Oh JJ. A rare case of multiple spinal epidural abscesses and cauda equina syndrome presenting to the emergency department following acupuncture. *Int J Emerg Med*. 2016;9(1):22.
- Yao Y, Hong W, Chen H, et al. Cervical spinal epidural abscess following acupuncture and wet-cupping therapy: a case report. *Complement Ther Med*. 2016;24:108-110.
- Singh LS, Alfshawy M, Singh N, Brennessel DJ. First reported case of methicillin-resistant staphylococcus aureus vertebral osteomyelitis with multiple spinal and paraspinal abscesses associated with acupuncture. *Case Rep Med*. 2015;2015:524241.
- Yin Z, He E, Ding H, Chen J. Brucella infection of the thoracic vertebral arch presenting with an epidural abscess: a case report. *J Med Case Rep*. 2015;9:237.
- Verma R, Chaudhari TS, Lachuriya G. Spontaneous extensive spinal epidural abscess presenting as acute quadriplegia. *BMJ Case Rep*. 2014;2014.
- Lee JH, Cho JH, Jo DJ. Cervical epidural abscess after cupping and acupuncture. *Complement Ther Med*. 2012;20(4):228-231.
- Yu HJ, Lee KE, Kang HS, Roh SY. Teaching NeuroImages: multiple epidural abscesses after acupuncture. *Neurology*. 2013;80(15):e169.
- Bang MS, Lim SH. Paraplegia caused by spinal infection after acupuncture. *Spinal Cord*. 2006;44(4):258-259.
- Daivajna S, Jones A, O'Malley M, Mehdian H. Unilateral septic arthritis of a lumbar facet joint secondary to acupuncture treatment – a case report. *Acupunct Med*. 2004;22(3):152-155.
- Chen MH, Chen MH, Huang JS. Cervical subdural empyema following acupuncture. *J Clin Neurosci*. 2004;11(8):909-911.
- Yang CW, Hsu SN, Liu JS, Hueng DY. Serratia marcescens spinal epidural abscess formation following acupuncture. *Intern Med*. 2014;53(15):1665-1668.
- Yazawa S, Ohi T, Sugimoto S, Satoh S, Matsukura S. Cervical spinal epidural abscess following acupuncture: successful treatment with antibiotics. *Intern Med*. 1998;37(2):161-165.
- Ha KY, Kim YH. Chronic inflammatory granuloma mimics clinical manifestations of lumbar spinal stenosis after acupuncture: a case report. *Spine (Phila Pa 1976)*. 2003;28(11):E217-E220.
- Ishibe M, Inoue M, Saitou K. Septic arthritis of a lumbar facet joint due to pyrexia. *Arch Orthop Trauma Surg*. 2001;121(1-2):90-92.
- Hadden WA, Swanson AJ. Spinal infection caused by acupuncture mimicking a prolapsed intervertebral disc. A case report. *J Bone Joint Surg Am*. 1982;64(4):624-626.
- Stuyt EB, Voyles CA. The national acupuncture detoxification association protocol, auricular acupuncture to support patients with substance abuse and behavioral health disorders: current perspectives. *Subst Abuse Rehabil*. 2016;7:169-180.
- Chao D, Nanda A. Spinal epidural abscess: a diagnostic challenge. *Am Fam Physician*. 2002;65(7):1341-1346.
- Brett J, editor. *Clean Needle Technique Manual: Best Practices for Acupuncture Needle Safety and Related Procedures* [Internet]. 7th ed. Baltimore, MD: Council of Colleges of Acupuncture and Oriental Medicine; 2015. Available at: <http://www.ccaom.org/>

- downloads/7thEditionManualEnglishPDFVersion.pdf. Accessed December 9, 2016.
35. Hoffman P. Skin disinfection and acupuncture. *Acupunct Med.* 2001;19(2):112-116.
36. Alton TB, Patel AR, Bransford RJ, Bellabarba C, Lee MJ, Chapman JR. Is there a difference in neurologic outcome in medical versus early operative management of cervical epidural abscesses? *Spine J.* 2015;15(1):10-17.