Published in final edited form as:

J Clin Psychiatry. 2017 July ; 78(7): e828–e836. doi:10.4088/JCP.17m11485.

A RISK ALGORITHM FOR THE PERSISTENCE OF SUICIDAL THOUGHTS AND BEHAVIORS DURING COLLEGE

Philippe Mortier, MD,

Research Group Psychiatry, Department of Neurosciences, KU Leuven University, Leuven, Belgium.

Glenn Kiekens, MSc,

Research Group Psychiatry, Department of Neurosciences, KU Leuven University, Leuven, Belgium.

Randy P. Auerbach, PhD, ABPP,

Department of Psychiatry, Harvard Medical School, Boston MA, USA; Center for Depression, Anxiety and Stress Research, McLean Hospital, Belmont, MA, USA.

Pim Cuijpers, PhD, MD,

Department of Clinical, Neuro- and Developmental Psychology, Vrije Universiteit Amsterdam, the Netherlands.

Koen Demyttenaere, PhD, MD,

Research Group Psychiatry, Department of Neurosciences, KU Leuven University, Leuven, Belgium.

Jennifer G. Green, PhD,

School of Education, Boston University, Boston, MA, USA.

Ronald C. Kessler, PhD,

Harvard Medical School, Department of Health Care Policy, Harvard University, Boston, MA, USA.

Matthew K. Nock, PhD,

Department of Psychology, Harvard University, Cambridge, MA, USA.

Alan M. Zaslavsky, PhD, and

Harvard Medical School, Department of Health Care Policy, Harvard University, Boston, MA, USA.

Ronny Bruffaerts, PhD

Research Group Psychiatry, Department of Neurosciences, KU Leuven University, Leuven, Belgium.

Abstract

Corresponding author: Philippe Mortier, M.D., Research Group Psychiatry, Department of Neurosciences, KU Leuven University, Herestraat 49, Leuven, Belgium. philippe.mortier@uzleuven.be. Tel: 0032/16348000. Fax: 0032/16348700.

Conflict of Interest Disclosures: The other authors report no biomedical financial interests or potential conflicts of interest.

OBJECTIVE—The primary aims are to: (a) identify patterns of suicidal thoughts and behaviors (STB) during college among students with lifetime pre-matriculation STB and (b) develop a risk-screening algorithm for persistence of pre-matriculation STB during college.

METHODS—Data come from the Leuven College Surveys, a series of prospective cohort studies of all incoming KU Leuven University freshmen. In the academic year 2012 and 2013, 4,889 incoming freshmen (73.2% response rate) provided baseline data on socio-demographic variables, childhood-adolescent traumatic experiences, 12-month stressful experiences, 12-month mental disorders, 12-month STB, and severity markers of pre-matriculation STB. A total of 2,566 students (69.3% conditional response rate) participated in 12- and 24-month follow-up surveys during the first two college years.

RESULTS—Thirteen percent (weighted $n=535$) of incoming freshmen reported lifetime pre-matriculation STB. Of those, 28.0% reported 12-month STB in one follow-up assessment and another 27.7% in both follow-up assessments. High persistence of STB (i.e., 12-month STB in two follow-up assessments) was most strongly associated with severity markers of pre-matriculation STB, with odds ratios in the 2.4–10.3 range and population attributable risk proportions between 9.2% and 50.8%. When aiming for less than 50% of false positive cases (PPV=54.4%), a multivariate predictive risk algorithm (cross-validated AUC=0.79) situated 59.9% of highly persistent cases among the 30% respondents with highest baseline predicted risk.

CONCLUSION—An individualized web-based screening approach is a promising strategy to identify students at the onset of university who may be at high risk for STB persistence during their academic career.

Keywords

suicide; suicidal thoughts and behaviors (STB); risk-screening algorithm; persistence; college students; cohort study

INTRODUCTION

Adolescence carries high risk for onset of suicidal thoughts and behaviors (STB).¹ Although STB decline when transitioning into young adulthood,^{2, 3} many adolescents (21–50%) continue to experience symptoms into their early twenties.^{4, 5} One potential prevention intervention for persistent STB includes web-based screening upon college entrance.^{6, 7} Given the high availability of internet access and geographic proximity to centralized student services, college campus may be ideally situated to access large groups of young adults for screening and referral to adequate care.⁷ Web-based screening provides a practical alternative for suicidal individuals who may be less likely to seek clinical services,⁸ and further, it may offer personalized feedback⁹ and access to online self-help interventions.¹⁰

Despite these potential advantages, it is currently unknown to what extent pre-matriculation onset STB persists into the college period. Of note, college student surveys suggest high lifetime STB prevalence (11–34%),^{11–13} and 12-month STB are comparable to same-aged peers (range 4.2–5.5%).¹⁴ In addition, screening has been criticized for overwhelming counselors with false-positive cases¹⁵ and imposing unrealistic demands on mental health centers.^{16, 17} Developing powerful risk screening algorithms for STB persistence may

remediate this.¹⁸ However, only one prior study developed such algorithm, restricted to depressed adults aged 18 years or older.¹⁹

The present study addresses these shortcomings by examining persistence of pre-matriculation onset STB during the college years in a large, longitudinal survey of college students (Leuven College Surveys [LCS],²⁰ part of the WHO World Mental Health Surveys International College Student project²¹). Consistent with recommendations to develop risk algorithms to target high-risk individuals for preventive interventions,²² we examine the strength of multivariate associations in our model of baseline predictors to determine whether a well-defined subset of students at highest risk of persistent STB can be detected. Baseline risk factors assessed at college entrance were selected based on available research on STB persistence, and include previous STB,²² childhood-adolescent trauma (e.g., domestic violence),²³ number of stressful life events,⁴ and mental disorders (e.g., internalizing and externalizing disorders).⁴

METHOD

Procedures

Full procedures of the LCS have been reported previously.²⁴ Briefly, the LCS consists of a series of web-based self-report surveys of KULeuven students. All 7,493 Dutch-speaking incoming freshmen aged 18 years or older were eligible for the baseline survey in the academic year 2012 and 2013. A total of 4,889 students completed the baseline survey (73.2% response rate after adjusting for non-participation due to college attrition). Students were contacted for the follow-up survey 12 and 24 months after the baseline assessment. A total of 2,566 of the original baseline respondents responded to at least one follow-up survey (69.3% conditional response rate after adjusting for non-participation due to college attrition). Informed consent was obtained from all subjects. Students with 12-month STB or non-suicidal self-injury (NSSI) received links to local mental health resources. The study's protocol was approved by the University Hospital Leuven Biomedical Ethical Board.

Measures

Socio-demographic variables—The university's students' administration office provided socio-demographic characteristics, including gender, age, nationality, parents' financial situation, parents' education, parental familial composition, university group membership, and secondary school educational type. Survey items assessed sexual orientation and college living situation.

Suicidal thoughts and behaviors (STB)—Items from the Self-Injurious Thoughts and Behaviors Interview (SITBI²⁵) assessed suicidal ideation (“*Did you ever in your life have thoughts of killing yourself?*”), suicide plans (“*Did you ever think about how you might kill yourself [e.g., taking pills, shooting yourself] or work out a plan of how to kill yourself?*”), and suicide attempts (“*Have you ever made a suicide attempt [i.e., purposefully hurt yourself with at least some intent to die?]*”). Baseline surveys assessed lifetime and 12-month STB (“*In the past 12 months, did you think/have you made...*”), whereas 12/24-m follow-up surveys each assessed 12-month STB to effectively cover the full follow-up period. At

baseline only, pre-matriculation lifetime STB severity markers included: (1) STB duration (> 5 years or more) and recency (remission at college entrance of 3 years or less) based on STB age-of-onset/offset measures, (2) proxy measures of intensity (e.g., “*How many days during that worst week did you have those thoughts?*”) and controllability of STB (“*How easy was it for you to control those thoughts or push them out of your mind when you wanted to?*”), referring to the week when STB was perceived as most severe,²⁶ and (3) STB-associated risk-taking behavior (“*People who think about killing themselves sometimes do dangerous things as a way to tempt fate [e.g., take a lot of drugs, drive too fast, volunteer for dangerous missions, or act recklessly]. How often in your life did you ever do dangerous things related to thinking about killing yourself?*”).²⁷

Traumatic experiences in childhood-adolescence—(i.e. prior to the age of 17) were assessed using 19 items adapted from the Composite International Diagnostic Interview (CIDI 3.0) childhood section,²⁸ the Adverse Childhood Experience Scale,²⁹ and the Bully Survey.³⁰ Items assessed parental psychopathology (i.e., any serious mental or emotional problems, substance use problems, STB or death by suicide, criminal activities, or interpersonal violence), physical abuse, emotional abuse, sexual abuse, neglect, bully victimization (i.e., either direct verbal or physical bullying, as well as indirect bullying [e.g., spreading rumors], or cyberbullying), and dating violence. Response options consisted of five-point Likert items (“*never*”, “*rarely*”, “*sometimes*”, “*often*”, and “*very often*”). Confirmatory factor analysis using our data showed excellent validity of the internal screener structure (RMSEA=0.019). To obtain dichotomously coded variables for PARP analysis of risk factors, cut-off values consisted of “*rarely*” for all items, except bully victimization which had a cut-off of “*sometimes*”, in line with previous recommendation.³¹

Risk for 12-month mental disorders—was assessed with the Global Appraisal of Individual Needs Short Screener (GAIN-SS³²) including: internalizing disorders, externalizing disorders, substance disorders, and crime/violence related disorders. The GAIN-SS sub-screens are very strongly correlated with the original corresponding subscales of the 60–120 minute DSM-IV-TR based GAIN structured interview (Pearson $r=0.84–0.93$ ³²). Confirmatory factor analysis using our data showed a very good validity of the internal GAIN-SS structure (RMSEA =0.032). For each screener, the recommended cut-off score for the highest probability of a 12-month diagnosis was used, i.e., three or more positive 12-month symptoms. We also assessed risk for other mental disorders or symptoms, including 12-month (hypo)mania, intermittent explosive disorder, psychosis (using CIDI-3.0 items),^{28, 33} eating disorder (using MINI items³⁴), and non-suicidal self-injury (using the SITBI,²⁵ cf. above).

Stressful events experienced in the past 12-months—were assessed using 12 items taken from well-validated screeners,^{35–37} and included relevant stressful experiences among young adults, including life-threatening illness or injury of a family member or close friend³⁸, accidents or death of a family member or close friend,³⁹ interpersonal events (e.g., break-up with a romantic partner, serious betrayal by someone other than one’s partner)⁴⁰, physical or sexual assault,^{41, 42} and legal problems (e.g., time spent in jail.)⁴³

Analyses

Nonresponse propensity weights⁴⁴ adjusted for potential non-response bias, and multiple imputation by chained equations⁴⁵ adjusted for survey attrition and within-survey item nonresponse. All analyses were restricted to the $n=535$ students who reported lifetime STB in the baseline survey. Adverse patterns of STB during the follow-up period were defined as *two-time STB* (i.e., reporting 12-month STB at both follow-up assessments), and *one-time STB* (i.e., reporting 12-month STB at one of the two follow-up assessments). Logistic regression analysis examined the strength of individual-level associations (i.e., odds ratios [OR]) between baseline risk factors with adverse patterns of STB. As this does not account for a population-level risk perspective⁴⁶ (i.e., high-prevalence risk factors carrying low individual risk for STB [i.e., low OR] may be equally or even more important to consider as low-prevalence risk factors carrying high risk for the affected individuals), population attributable risk proportions (PARP)⁴⁷ were calculated, thus allowing the risk factors that potentially attribute most to the persistence STB in student populations to be identified.

Finally, a series of multivariate models was estimated. Predictors were entered in blocks, beginning with socio-demographic variables, followed by childhood-adolescent traumatic experiences, 12-month stressful experiences, 12-month risk for mental disorders, and finally severity markers of pre-matriculation STB. Individual-level predicted probabilities based on the multivariate equation were created, and area under the curve (AUC) values calculated. Predicted probabilities were discretized into deciles and cross-classified with observed cases to visualize the concentration of risk associated with high composite predicted probabilities. Sensitivity was defined as the proportion of cases found among pre-defined proportions (e.g., 10%) of respondents with highest predicted probabilities. Positive Predictive Value (PPV) was defined as the probability of effectively developing the outcome when being among pre-defined proportions (e.g., 10%) of respondents with highest predicted probabilities. We used the method of leave-one-out cross-validation⁴⁸ to correct for the over-estimation of prediction accuracy when both estimating and evaluating model fit in a single sample.

RESULTS

Course of STB during college

The course of STB among the 535 students (13.0%; SE = 0.3) with lifetime STB is presented in Figure 1. Whereas about four out of ten students (44.3%) show a pattern of remission during the 2-year follow-up period, about three out of ten (28.0%) show *one-time STB* (i.e., one time 12-month STB during the two follow-up years), and three out of ten (27.7%) show *two-time STB* (i.e., two times 12-month STB during the two follow-up years). The importance of the distinction between *one-time* and *two-time STB* is seen in the fact that 12-month prevalence of suicide plans is significantly higher among those with *two-time STB* compared to those with *one-time STB* in each of the two follow-up surveys (51.1% vs. 14.1% in the first follow-up assessment; $\chi^2 = 16.1$; $p < 0.001$; and 78.9% vs. 33.2% in the second follow-up assessment; $\chi^2 = 29.2$; $p < 0.001$). Few 12-month suicide attempts were reported, i.e., eight at college entrance (two in the *one-time STB* group, six in the *two-time*

STB group), three during year 1 (one in the *one-time STB* group, two in the *two-time STB* group), and one during year 2 (in the *one-time STB* group).

Bivariate models for adverse course of STB during college

Inspecting results across risk factor domains (Tables 1–3) reveals that risk factors are almost exclusively associated with *two-time STB*, as opposed to *one-time STB*. Focusing on *two-time STB*, associations are strongest for severity markers of pre-matriculation STB (median OR=3.2; median PARP=17.9%; Table 3), including lifetime suicide attempts and 12-month STB at college entrance, but also STB duration (episodes spanning more than 5 years), recency (remission at college entrance for three years or less), perceived intensity (as determined by duration measures in worse week), and perceived controllability of STB. Considerable proportions of *two-time STB* are also associated with baseline stressful experiences (PARP=28.9%), as well as baseline risk for three or more mental disorders (PARP=20.1%; Table 2). Specific risk factors identified in these risk domains include interpersonal stressors (serious betrayal by someone else than one's partner, serious ongoing arguments or break-up with friends or family members, death of a friend or family member [median OR=2.1; median PARP=12.0%]), as well as symptoms of dysregulated mood (intermittent explosive disorder symptoms, [hypo]mania symptoms) and NSSI (median OR=4.2; median PARP=12.1%). In contrast, population-level risk of childhood-adolescent traumatic experiences is non-significant, and individual-level associations are only significant for physical abuse (OR=2.9) and sexual abuse (OR=5.1).

Multivariate models for adverse course of STB during college

Multivariate models (Supplementary eTables 1–3) also fit the data better for *two-time STB*, as opposed to *one-time STB* (drop in AIC of 41.0% versus 18.0%). Significant predictors for *two-time STB* in the final model adjusting for all risk domains (cross-validated AUC=0.79 [SE=0.04]) are lifetime suicide attempts (OR=11.1 [1.4–87.4]; PARP=7.2%) and 12-month STB (OR=4.7 [1.4–15.9]; PARP=12.3%). Of note, 12-month non-suicidal self-injury remains a strong independent risk factor (OR=5.7 [1.7–18.5]; PARP=8.7%), an effect which is only attenuated (OR=3.6 [0.9–14.9]; PARP=4.9%) in the final model adjusting for all risk domains. Table 4 shows cross-validated sensitivity and PPV estimates for different proportions of students at highest predicted risk based on the final model. One possible trade-off between these measures is considering the top 30% at highest risk, as this includes 59.9% of subsequent cases of *two-times STB*, with under 50% of false positive cases (PPV=54.4%).

DISCUSSION

In line with previous studies,^{4, 5, 23, 49, 50} we identified a small proportion of young adults that repeatedly report STB during the early college years (i.e., *two-times STB*; 27.7% of incoming freshmen with lifetime pre-matriculation STB, 3.6% of all incoming freshmen). As expected, robust predictors were previous STB,²² especially suicide attempts.⁵¹ We now find this adverse course also predicted by a wider range of severity markers related to previous STB, such as long-lasting symptoms, high intensity of ideation, and perceived loss of control over ideation. In contrast with one previous clinical study,⁵² these additional

severity markers were no longer significant in a full multivariate model adjusting for previous STB, pointing out the need for additional studies. Since we did not assess severity of STB during follow-up, future research should also deliver a more fine-grained picture of patterns of STB severity over time. This should clarify to what extent we identified a subgroup of young adults with an uninterrupted course of severe suicidal symptoms from childhood-adolescence into the college years, and to what extent these students are at risk for suicide. On the one hand, suicide attempts in our sample mainly occurred during childhood-adolescence, and were almost non-existing during follow-up in college. College attrition due to adverse mental health may explain this^{53, 54} but it is also compatible with studies that find risk for suicide attempt to be highest in the year after onset of suicidal ideation.⁵⁵ On the other hand, tendency towards suicidal planning remained high among those with more persistent STB (51–79%), and several prospective studies find a strong association between STB persistence and suicide attempts later in life.^{5, 56, 57}

Surprisingly, we found childhood-adolescent trauma generally not related to persistence of STB into young adulthood. Notable exceptions were physical and especially sexual abuse, which is in line with the detrimental effects of these experiences on early-life mental health.⁵⁸ However, due to low prevalence, the contribution of these risk factors to the total amount of persistent STB cases was limited. In contrast, many such cases were potentially attributable to interpersonal stressful experiences, and symptoms of dysregulated mood in the year before college entrance. This was especially the case for having arguments with family or friends, a risk factor ranking highest as precipitating event for suicide attempts.⁵⁹ When integrating these findings with existing research, one potential hypothesis is that earlier traumatic experiences initiate childhood-adolescent STB,⁶⁰ but also heightened levels of affective-behavioral dysregulation,⁶¹ and interpersonal stress sensitivity.⁶² These traits, in turn, may be predictive for persistent STB into young adulthood, relatively independent from the earlier trauma that initiated them. If confirmed, we may have identified a group at risk for developing borderline personality disorder, given the strong association of this disorder with persistent STB,⁶³ as well as symptoms of mood instability,⁶⁴ interpersonal stressful experiences,⁶⁵ and physical or sexual trauma.⁶⁶ For some students, persistent STB was also strongly related to NSSI, which has been conceptualized as a coping mechanism to deal with heightened interpersonal stress and unwanted internal emotional states,^{67, 68} and is also strongly associated with borderline personality characteristics.⁶⁹ In any case, these risk factors all emerged as important for persistence of childhood-adolescent STB into college, pointing out potential targets in college screening projects.

An important contribution of our study is the development of a prospective risk-screening algorithm that accurately predicts a persistent pattern of STB. Our algorithm performs equally-well as those developed for other multifactorial diseases, including coronary heart disease⁷⁰, common cancers^{71, 72}, diabetes⁵⁷, and all cause-mortality.⁷³ Together with a recently developed risk algorithm for onset of college STB,²⁴ these are very promising results with regard to individualized approaches to screening for STB in unrestricted populations of young people. Indeed, we demonstrate that a powerful prediction algorithm may optimize the link between high risk for STB and referral for intervention. For example, an intervention targeting the top 30% of freshmen at highest risk could effectively reach 60% of highly persistent—and possibly severe—STB cases during the following years. This

actually corresponds to an estimated 80 students each academic year in our university, and these are realistic proportions to refer to evidence-based care like psychiatric assessment and psychotherapeutic interventions.^{13, 74} While false-positive cases may continue to be an issue (an expected 36 cases each academic year in our university), it is worth taking into consideration that our estimations are based on a large number (> 40) of clinically significant predictors. High cumulative risk for STB based on our model may signal a need for referral *regardless* of a subsequent development of persistent STB, safeguarding against an inadequate inflow of students to specialized care.

Several limitations of this study deserve attention. First, response rates in the 69.3–73.2% range are good but do not rule out the possibility that we may have missed cases of STB in both baseline and follow-up assessments. However, our response rates (corrected for college attrition) are substantially higher than those of most cross-sectional web-based surveys (i.e. around 40%⁷⁵) or response rates in other recent large-scale college student surveys (39–44%^{13, 76}). Second, it is unknown to what extent baseline risk for mental disorders by using self-report measures effectively corresponds to mental disorders diagnosed by face-to-face clinical interviews. We addressed this limitation by selecting well-validated measures used in large surveys of the general population. Third, prediction accuracy measures were validated using the leave-one-out cross-validation technique, which delivers somewhat downward biased estimates of true prediction accuracy.⁷⁷ Therefore, cross-validated prediction accuracy measures in this study should be seen as underestimates, and future research should address this using larger sample sizes and better-equipped validation techniques.

CONCLUSION

We detected a small but significant proportion of students with a persistent pattern of STB, characterized by severe psychopathology and heightened sensitivity for interpersonal stressful experiences. Results are promising for accurately predicting highly persistent STB cases in a prospective way; however, results for relapsing-remitting STB were less favorable. Future research should address this by including additional risk and protective factors, by considering interactions between predictors, and by implementing machine learning techniques⁷⁸ to further improve prediction accuracy. Fortunately, the WMH-ICS project aims to collect prospective data on large samples of incoming freshmen worldwide, enabling these future endeavors.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Funding/Support: The Leuven College Survey was carried out in conjunction with the World Health Organization World Mental Health (WMH) survey initiative and is a part of the World Mental Health International College Student project. The WMH survey is supported by the National Institute of Mental Health (NIMH; R01MH070884), the John D. and Catherine T. MacArthur Foundation, the Pfizer Foundation, the US Public Health Service (R13-MH066849, R01-MH069864, and R01 DA016558), the Fogarty International Center (FIRCA R03-TW006481), the Pan American Health Organization, Eli Lilly and Company, Ortho-McNeil Pharmaceutical, GlaxoSmithKline, and Bristol-Myers Squibb. A complete list of all within-country and cross-national WMH

publications can be found at <http://www.hcp.med.harvard.edu/wmh/>. In Belgium specifically, these activities were supported by the Belgian Fund for Scientific Research (11N0514N/11N0516N/ 1114717N), the King Baudouin Foundation (2014-J2140150-102905), and Eli Lilly (IIT-H6U-BX-I002). Auerbach's work was supported by the National Institute of Mental Health (R56 MH109566).

Role of the Sponsor: The funding sources had no role in the design and conduct of the study; collection, management, analysis, interpretation of the data; preparation, review, or approval of the manuscript; and decision to submit the manuscript for publication.

In the past 3 years, Dr. Kessler received support for his epidemiological studies from Sanofi Aventis, was a consultant for Johnson & Johnson Wellness and Prevention, and served on an advisory board for the Johnson & Johnson Services Inc. Lake Nona Life Project. Kessler is a co-owner of DataStat, Inc., a market research firm that carries out healthcare research.

References

1. Nock MK, Green JG, Hwang I, et al. Prevalence, correlates, and treatment of lifetime suicidal behavior among adolescents: results from the National Comorbidity Survey Replication Adolescent Supplement. *JAMA Psychiatry*. 2013 Mar; 70(3):300–310. [PubMed: 23303463]
2. Lee CG, Seo D-C. Trajectory of suicidal ideation in relation to perceived overweight from adolescence to young adulthood in a representative United States sample. *J Adolesc Health*. 2013; 53(6):712–716. [PubMed: 23910569]
3. Musci RJ, Hart SR, Ballard ED, et al. Trajectories of Suicidal Ideation from Sixth through Tenth Grades in Predicting Suicide Attempts in Young Adulthood in an Urban African American Cohort. *Suicide Life Threat Behav*. 2016 Jun; 46(3):255–265. [PubMed: 26395337]
4. Steinhausen HC, Bosiger R, Metzke CW. Stability, correlates, and outcome of adolescent suicidal risk. *J Child Psychol Psychiatry*. 2006 Jul; 47(7):713–722. [PubMed: 16790006]
5. Thompson M, Kuruwita C, Foster EM. Transitions in suicide risk in a nationally representative sample of adolescents. *J Adolesc Health*. 2009 May; 44(5):458–463. [PubMed: 19380093]
6. Zalsman G, Hawton K, Wasserman D, et al. Suicide prevention strategies revisited: 10-year systematic review. *The Lancet Psychiatry*. 2016 Jul; 3(7):646–659. [PubMed: 27289303]
7. Harrod CS, Goss CW, Stallones L, DiGuseppi C. Interventions for primary prevention of suicide in university and other post-secondary educational settings. *Cochrane Database Syst Rev*. 2014 Oct 29. (10):CD009439. [PubMed: 25353703]
8. Hom MA, Stanley IH, Joiner TE Jr. Evaluating factors and interventions that influence help-seeking and mental health service utilization among suicidal individuals: A review of the literature. *Clin Psychol Rev*. 2015 Aug; 40:28–39. [PubMed: 26048165]
9. King CA, Eisenberg D, Zheng K, et al. Online suicide risk screening and intervention with college students: a pilot randomized controlled trial. *J Consult Clin Psychol*. 2015 Jun; 83(3):630–636. [PubMed: 25688811]
10. van Spijker BA, van Straten A, Kerkhof AJ. Effectiveness of online self-help for suicidal thoughts: results of a randomised controlled trial. *PloS One*. 2014; 9(2):e90118. [PubMed: 24587233]
11. Klonsky ED, May A. Rethinking Impulsivity in Suicide. *Suicide Life Threat Behav*. 2010 Dec; 40(6):612–619. [PubMed: 21198330]
12. Drum DJ, Brownson C, Burton Denmark A, Smith SE. New data on the nature of suicidal crises in college students: Shifting the paradigm. *Prof Psychol-Res Pr*. 2009; 40(3):213–222.
13. Paul E, Tsypes A, Eidlitz L, Ernhout C, Whitlock J. Frequency and functions of non-suicidal self-injury: Associations with suicidal thoughts and behaviors. *Psychiatry Res*. 2014 Dec 24.
14. Han B, Compton WM, Eisenberg D, Milazzo-Sayre L, McKeon R, Hughes A. Prevalence and mental health treatment of suicidal ideation and behavior among college students aged 18–25 years and their non-college-attending peers in the United States. *J Clin Psychiatry*. 2016 May 10.
15. Hallfors D, Brodish PH, Khatapoush S, Sanchez V, Cho H, Steckler A. Feasibility of screening adolescents for suicide risk in "real-world" high school settings. *Am J Public Health*. 2006 Feb; 96(2):282–287. [PubMed: 16380568]
16. Schwartz AJ. Four eras of study of college student suicide in the United States: 1920–2004. *J Am Coll Health*. 2006 May-Jun; 54(6):353–366. [PubMed: 16789651]

17. Ream GL. The Interpersonal-Psychological Theory of Suicide in College Student Suicide Screening. *Suicide Life Threat Behav.* 2016 Apr; 46(2):239–47. [PubMed: 26317690]
18. Kessler RC, Warner CH, Ivany C, et al. Predicting suicides after psychiatric hospitalization in US Army soldiers: the Army Study To Assess Risk and rEsilience in Servicemembers (Army STARRS). *JAMA Psychiatry.* 2015 Jan; 72(1):49–57. [PubMed: 25390793]
19. Liu Y, Sareen J, Bolton JM, Wang JL. Development and validation of a risk prediction algorithm for the recurrence of suicidal ideation among general population with low mood. *J Affect Disord.* 2016 Mar 15; 193:11–17. [PubMed: 26766029]
20. The Leuven College Surveys (LCS). [Accessed April 17, 2017] 2015. Available from: <http://www.mindmates.be/page.phpid28>
21. The WHO World Mental Health Surveys International College Student Project (WMH-ICS). [Accessed April 17, 2017] 2015. Available from: http://www.hcp.med.harvard.edu/wmh/college_student_survey.php
22. Ribeiro JD, Franklin JC, Fox KR, et al. Self-injurious thoughts and behaviors as risk factors for future suicide ideation, attempts, and death: a meta-analysis of longitudinal studies. *Psychol Med.* 2015 Sep; 15:1–12.
23. Wilcox HC, Arria AM, Caldeira KM, Vincent KB, Pinchevsky GM, O'Grady KE. Prevalence and predictors of persistent suicide ideation, plans, and attempts during college. *J Affect Disord.* 2010; 127:287–294. [PubMed: 20471691]
24. Mortier P, Demyttenaere K, Auerbach RP, et al. First onset of suicidal thoughts and behaviours in college. *J Affect Disord.* 2016 Sep 28; 207:291–299. [PubMed: 27741465]
25. Nock MK, Holmberg EB, Photos VI, Michel BD. Self-Injurious Thoughts and Behaviors Interview: development, reliability, and validity in an adolescent sample. *Psychol Assess.* 2007 Sep; 19(3):309–317. [PubMed: 17845122]
26. Posner K, Brown GK, Stanley B, et al. The Columbia-Suicide Severity Rating Scale: initial validity and internal consistency findings from three multisite studies with adolescents and adults. *Am J Psychiatry.* 2011 Dec; 168(12):1266–1277. [PubMed: 22193671]
27. Heeringa SG, Gebler N, Colpe LJ, et al. Field procedures in the Army Study to Assess Risk and Resilience in Servicemembers (Army STARRS). *Int J Methods Psychiatr Res.* 2013 Dec; 22(4): 276–287. [PubMed: 24038395]
28. Kessler RC, Ustun TB. The World Mental Health (WMH) Survey Initiative Version of the World Health Organization (WHO) Composite International Diagnostic Interview (CIDI). *Int J Methods Psychiatr Res.* 2004; 13(2):93–121. [PubMed: 15297906]
29. Felitti VJ, Anda RF, Nordenberg D, et al. Relationship of childhood abuse and household dysfunction to many of the leading causes of death in adults. The Adverse Childhood Experiences (ACE) Study. *Am J Prev Med.* 1998 May; 14(4):245–258. [PubMed: 9635069]
30. Swearer S, Cary P. Perceptions and Attitudes Toward Bullying in Middle School Youth. *Journal of Applied School Psychology.* 2003; 19(2):63–79.
31. Nansel TR, Overpeck M, Pilla RS, Ruan WJ, Simons-Morton B, Scheidt P. Bullying behaviors among US youth: prevalence and association with psychosocial adjustment. *JAMA.* 2001 Apr 25; 285(16):2094–2100. [PubMed: 11311098]
32. Dennis ML, Chan YF, Funk RR. Development and validation of the GAIN Short Screener (GSS) for internalizing, externalizing and substance use disorders and crime/violence problems among adolescents and adults. *Am J Addict.* 2006; 15(Suppl 1):80–91. [PubMed: 17182423]
33. Haro JM, Arbabzadeh-Bouchez S, Brugha TS, et al. Concordance of the Composite International Diagnostic Interview Version 3.0 (CIDI 3.0) with standardized clinical assessments in the WHO World Mental Health surveys. *Int J Methods Psychiatr Res.* 2006; 15(4):167–180. [PubMed: 17266013]
34. Sheehan DV, Lecrubier Y, Sheehan KH, et al. The Mini-International Neuropsychiatric Interview (M.I.N.I.): the development and validation of a structured diagnostic psychiatric interview for DSM-IV and ICD-10. *J Clin Psychiatry.* 1998; 59(Suppl 20):22–33. quiz 34–57.
35. Brugha TS, Cragg D. The List of Threatening Experiences: the reliability and validity of a brief life events questionnaire. *Acta Psychiatr Scand.* 1990 Jul; 82(1):77–81. [PubMed: 2399824]

36. Bray RM, Hourani LL. Substance use trends among active duty military personnel: findings from the United States Department of Defense Health Related Behavior Surveys, 1980–2005. *Addiction*. 2007 Jul; 102(7):1092–1101. [PubMed: 17567397]
37. Vogt DS, Proctor SP, King DW, King LA, Vasterling JJ. Validation of scales from the Deployment Risk and Resilience Inventory in a sample of Operation Iraqi Freedom veterans. *Assessment*. 2008 Dec; 15(4):391–403. [PubMed: 18436857]
38. Stoeckel M, Weissbrod C. Growing up with an ill parent: An examination of family characteristics and parental illness features. *Fam Syst Health*. 2015 Dec; 33(4):356–362. [PubMed: 26120764]
39. Rostila M, Berg L, Arat A, Vinnerljung B, Hjern A. Parental death in childhood and self-inflicted injuries in young adults—a national cohort study from Sweden. *Eur Child Adolesc Psychiatry*. 2016 Oct; 25(10):1103–1111. [PubMed: 26932156]
40. Buitron V, Hill RM, Pettit JW, Green KL, Hatkevich C, Sharp C. Interpersonal stress and suicidal ideation in adolescence: An indirect association through perceived burdensomeness toward others. *J Affect Disord*. 2016 Jan 15; 190:143–149. [PubMed: 26519633]
41. Trotman GE, Young-Anderson C, Deye KP. Acute Sexual Assault in the Pediatric and Adolescent Population. *J Pediatr Adolesc Gynecol*. 2015 May 16.
42. Viswanathan S, Datta S, Sheridan P, Lax-Pericall T. "Too Young to be Worried!" Psychiatric Assessment and Follow-up of Young People After Severe Physical Assault in an Inner City Hospital of South London. *Ann Med Health Sci Res*. 2014 Jan; 4(1):85–89. [PubMed: 24669337]
43. Salekin RT. Psychopathy and recidivism from mid-adolescence to young adulthood: cumulating legal problems and limiting life opportunities. *J Abnorm Psychol*. 2008 May; 117(2):386–395. [PubMed: 18489214]
44. Rosenbaum PR, Rubin DB. The Central Role of the Propensity Score in Observational Studies of Causal Effects. *Biometrika*. 1983; 70:41–55.
45. van Buuren S. Multiple imputation of discrete and continuous data by fully conditional specification. *Stat Methods Med Res*. 2007 Jun; 16(3):219–242. [PubMed: 17621469]
46. Christensen H, Cuijpers P, Reynolds CF 3rd. Changing the Direction of Suicide Prevention Research: A Necessity for True Population Impact. *JAMA Psychiatry*. 2016 May 1; 73(5):435–436. [PubMed: 26982348]
47. Kryszinska K, Martin G. The struggle to prevent and evaluate: application of population attributable risk and preventive fraction to suicide prevention research. *Suicide Life Threat Behav*. 2009 Oct; 39(5):548–557. [PubMed: 19929154]
48. Efron B, Gong G. A leisurely look at the bootstrap, the jackknife and cross-validation. *The American Statistician*. 1983; 37:36–48.
49. Nkansah-Amankra S. Adolescent suicidal trajectories through young adulthood: prospective assessment of religiosity and psychosocial factors among a population-based sample in the United States. *Suicide Life Threat Behav*. 2013 Aug; 43(4):439–459. [PubMed: 23601148]
50. Giletta M, Prinstein MJ, Abela JR, Gibb BE, Barrocas AL, Hankin BL. Trajectories of suicide ideation and nonsuicidal self-injury among adolescents in mainland China: Peer predictors, joint development, and risk for suicide attempts. *J Consult Clin Psychol*. 2015 Apr; 83(2):265–279. [PubMed: 25688812]
51. Borges G, Angst J, Nock MK, Ruscio AM, Kessler RC. Risk factors for the incidence and persistence of suicide-related outcomes: a 10-year follow-up study using the National Comorbidity Surveys. *J Affect Disord*. 2008 Jan; 105(1–3):25–33. [PubMed: 17507099]
52. Yen S, Weinstock LM, Andover MS, Sheets ES, Selby EA, Spirito A. Prospective predictors of adolescent suicidality: 6-month post-hospitalization follow-up. *Psychol Med*. 2013 May; 43(5):983–993. [PubMed: 22932393]
53. Auerbach RP, Alonso J, Axinn WG, et al. Mental disorders among college students in the World Health Organization World Mental Health Surveys. *Psychol Med*. 2016 Aug; 3:1–16.
54. Mortier P, Demyttenaere K, Auerbach RP, et al. The impact of lifetime suicidality on academic performance in college freshmen. *J Affect Disord*. 2015 Jul 31; 186:254–260. [PubMed: 26254617]
55. Nock MK, Borges G, Bromet EJ, et al. Cross-national prevalence and risk factors for suicidal ideation, plans and attempts. *Br J Psychiatry*. 2008 Feb; 192(2):98–105. [PubMed: 18245022]

56. Rueter MA, Holm KE, McGeorge CR, Conger RD. Adolescent suicidal ideation subgroups and their association with suicidal plans and attempts in young adulthood. *Suicide Life Threat Behav.* 2008 Oct; 38(5):564–575. [PubMed: 19014308]
57. Brezo J, Paris J, Tremblay R, Vitaro F, Hebert M, Turecki G. Identifying correlates of suicide attempts in suicidal ideators: a population-based study. *Psychol Med.* 2007 Nov; 37(11):1551–1562. [PubMed: 17537281]
58. O'Brien BS, Sher L. Child sexual abuse and the pathophysiology of suicide in adolescents and adults. *Int J Adolesc Med Health.* 2013; 25(3):201–205. [PubMed: 23843572]
59. Blasco-Fontecilla H, Delgado-Gomez D, Legido-Gil T, de Leon J, Perez-Rodriguez MM, Baca-Garcia E. Can the Holmes-Rahe Social Readjustment Rating Scale (SRRS) be used as a suicide risk scale? An exploratory study. *Arch Suicide Res.* 2012; 16(1):13–28. [PubMed: 22289025]
60. Miller AB, Esposito-Smythers C, Weismore JT, Renshaw KD. The relation between child maltreatment and adolescent suicidal behavior: a systematic review and critical examination of the literature. *Clin Child Fam Psychol Rev.* 2013 Jun; 16(2):146–172. [PubMed: 23568617]
61. Carvalho Fernando S, Beblo T, Schlosser N, et al. The impact of self-reported childhood trauma on emotion regulation in borderline personality disorder and major depression. *J Trauma Dissociation.* 2014; 15(4):384–401. [PubMed: 24283697]
62. Huh HJ, Kim SY, Yu JJ, Chae JH. Childhood trauma and adult interpersonal relationship problems in patients with depression and anxiety disorders. *Ann Gen Psychiatry.* 2014; 13:26. [PubMed: 25648979]
63. Nakar O, Brunner R, Schilling O, et al. Developmental trajectories of self-injurious behavior, suicidal behavior and substance misuse and their association with adolescent borderline personality pathology. *J Affect Disord.* 2016 Jun; 197:231–238. [PubMed: 26995466]
64. Koenigsberg HW, Harvey PD, Mitropoulou V, et al. Characterizing affective instability in borderline personality disorder. *Am J Psychiatry.* 2002 May; 159(5):784–788. [PubMed: 11986132]
65. Pagano ME, Skodol AE, Stout RL, et al. Stressful life events as predictors of functioning: findings from the collaborative longitudinal personality disorders study. *Acta Psychiatr Scand.* 2004 Dec; 110(6):421–429. [PubMed: 15521826]
66. Ogata SN, Silk KR, Goodrich S, Lohr NE, Westen D, Hill EM. Childhood sexual and physical abuse in adult patients with borderline personality disorder. *Am J Psychiatry.* 1990 Aug; 147(8):1008–1013. [PubMed: 2375434]
67. Kiekens G, Bruffaerts R, Nock MK, et al. Non-suicidal self-injury among Dutch and Belgian adolescents: Personality, stress and coping. *Eur Psychiatry.* 2015 Sep; 30(6):743–749. [PubMed: 26260264]
68. Prinstein MJ, Guerry JD, Browne CB, Rancourt D. Interpersonal models of nonsuicidal self-injury. In: Nock MK, editor. *Understanding nonsuicidal self-injury: Origins, assessment, and treatment.* Washington, DC: US: American Psychological Association; 2009. p. 79-98.
69. Leichsenring F, Leibing E, Kruse J, New AS, Leweke F. Borderline personality disorder. *Lancet.* 2011 Jan 01; 377(9759):74–84. [PubMed: 21195251]
70. Backholer K, Hirakawa Y, Tonkin A, et al. Development of an Australian cardiovascular disease mortality risk score using multiple imputation and recalibration from national statistics. *BMC Cardiovasc Disord.* 2017 Jan 06; 17(1):17. [PubMed: 28061760]
71. Hippisley-Cox J, Coupland C. Development and validation of risk prediction algorithms to estimate future risk of common cancers in men and women: prospective cohort study. *BMJ open.* 2015 Mar 17; 5(3):e007825.
72. Liu X, Chen Z, Fine JP, et al. A competing-risk-based score for predicting twenty-year risk of incident diabetes: the Beijing Longitudinal Study of Ageing study. *Sci Rep.* 2016 Nov 16; 6:37248. [PubMed: 27849048]
73. Berard E, Bongard V, Arveiler D, et al. Ten-year risk of all-cause mortality: assessment of a risk prediction algorithm in a French general population. *Eur J Epidemiol.* 2011 May; 26(5):359–368. [PubMed: 21188478]
74. Jobes DA, Au JS, Siegelman A. Psychological approaches to suicide treatment and prevention. *Curr Treat Options Psychiatry.* 2015; 2(4):363–370.

75. Cook C, Heath F, Thompson RL. A meta-analysis of response rates in Web- or internet-based surveys. *Educ Psychol Meas.* 2000 Dec; 60(6):821–836.
76. Eisenberg D, Hunt J, Speer N. Mental health in American colleges and universities: variation across student subgroups and across campuses. *J Nerv Ment Dis.* 2013 Jan; 201(1):60–67. [PubMed: 23274298]
77. Smith GC, Seaman SR, Wood AM, Royston P, White IR. Correcting for optimistic prediction in small data sets. *Am J Epidemiol.* 2014 Aug 01; 180(3):318–324. [PubMed: 24966219]
78. Franklin JC, Ribeiro JD, Fox KR, et al. Risk Factors for Suicidal Thoughts and Behaviors: A Meta-Analysis of 50 Years of Research. *Psychol Bull.* 2016 Nov 14.

CLINICAL POINTS

- There is a need for validated risk-screening algorithms to identify young adults with high risk for STB persistence against low amounts of false positive cases.
- Web-based risk-screening for a wide range of clinically relevant risk factors at college entrance is a promising approach to optimize the link between high risk for STB persistence and subsequent evidence-based care.

Figure 1.

Course of STB during college among students with a prematriculation lifetime history of STB (weighted n=535).

Note: proportions indicate the number of participants that report (no) 12-month STB within the group where the arrow originates from. Groups with (no) 12-month STB in follow-up 1 were grouped together to calculate proportions in follow-up 2 (as indicated by the vertical bars).

^a Reporting one time 12-month STB during two-year follow-up

^b Reporting two times 12-month STB during two year follow-up

Abbreviations: 12-m = 12-month.

Table 1

Socio-demographic variables as baseline predictors for adverse clinical patterns of STB during follow-up.

	Prevalence		Bivariate model ^a					
			One-time STB ^b			Two-time STB ^b		
	n(w)	%(w)	OR	95% CI	PARP(%)	OR	95% CI	PARP(%)
I. Socio-demographic variables								
Being male	245	45.8	2.2	1.5 (0.8–2.6)	9.5	1.3	(0.7–2.2)	2.3
Age > 18 years	189	35.3	2.1	1.4 (0.8–2.6)	0.4	2.3*	(1.2–4.4)	16.5*
Non-Belgian nationality	58	10.8	1.4	1.2 (0.5–3.0)	0.9	1.1	(0.4–3.4)	0.5
Parents' financial situation difficult	138	25.7	2.1	1.1 (0.6–2.1)	1.4	1.0	(0.5–2.0)	–0.2
Parental education ^c								
- both parents high	295	55.2	2.5	(ref)	-	(ref)	-	-
- only one parent high	135	25.3	2.1	1.2 (0.6–2.4)	2.6	1.2	(0.6–2.4)	2.3
- none of parents high	104	19.5	2.1	1.5 (0.7–3.4)	2.5	1.9	(0.9–4.2)	6.8
Non-intact familial composition ^d	180	33.6	2.2	1.2 (0.6–2.1)	1.6	1.2	(0.6–2.3)	3.3
Non-heterosexual orientation	78	14.7	1.9	1.2 (0.5–2.7)	–0.3	1.6	(0.7–3.8)	4.7
College-related socio-demographics								
University Group membership								
- Human Sciences	318	59.4	2.1	(ref)	-	(ref)	-	-
- Science & Technology	139	25.9	1.9	1.1 (0.6–2.0)	2.0	0.9	(0.5–1.7)	–2.4
- Biomedical Sciences	78	14.7	1.5	1.2 (0.5–2.6)	1.1	1.1	(0.5–2.6)	0.8
Non-GSE pre-educational level	65	12.2	1.6	1.2 (0.4–3.4)	2.0	0.7	(0.2–2.8)	–2.2
Living with parents	174	32.5	2.4	1.2 (0.7–2.2)	4.7	1.0	(0.5–2.0)	–1.6

Note: significant OR/PARP are shown in bold and are marked with

* ($\alpha=0.05$).

^aThe bivariate associations are based on a separate model for each row, with the variable in the row as the only predictor in the model.

^bno STB during follow-up is the reference category

^cHigh degree of parental education defined as holding a college bachelor degree or more.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

^pNon-intact familial composition defined as parents being divorced or separated.

Abbreviations: GSE = general secondary education; OR = Odds ratio; PARP = population attributable risk proportion; STB = Suicidal Thoughts and Behaviors.

Childhood-adolescent traumatic experiences and 12-month stressful experiences as baseline predictors for adverse clinical patterns of STB during follow-up.

Table 2

		Bivariate model ^a								
		Prevalence			One-time STB ^b			Two-time STB ^b		
		n(w)	%(w)	(SE)	OR	95% CI	PARP(%)	OR	95% CI	PARP(%)
II. Traumatic experiences (< age 17)										
	parental psychopathology	286	53.5	2.6	1.4	(0.8–2.4)	4.3	1.7	(0.9–3.1)	15.4
	physical abuse	86	16.0	1.9	1.6	(0.7–3.9)	–0.1	2.9*	(1.2–7.0)	10.8*
	emotional abuse	209	39.1	2.5	1.1	(0.6–2.1)	–1.4	1.6	(0.9–3.0)	12.4
	sexual abuse	22	4.1	1.2	1.8	(0.3–10.8)	–0.8	5.1*	(1.1–22.9)	4.9*
	neglect	84	15.8	2.0	1.9	(0.8–4.5)	4.0	2.0	(0.8–5.0)	4.6
	bully victimization	332	62.2	2.6	1.3	(0.7–2.2)	6.9	1.2	(0.7–2.2)	3.9
	dating violence	74	13.9	1.9	1.1	(0.5–2.5)	0.4	1.1	(0.5–2.7)	1.0
	Any traumatic experience	458	85.6	1.8	1.3	(0.6–3.0)	8.8	1.3	(0.6–2.9)	10.6
	Number of traumatic experiences									
0	77	14.4	1.8	(ref)	-	-	(ref)	-	-	
1	158	29.5	2.3	1.0	(0.4–2.5)	0.7	0.8	(0.3–2.0)	–3.7	
2	116	21.7	2.2	1.5	(0.6–3.9)	4.7	1.3	(0.5–3.3)	1.4	
3+	184	34.4	2.4	1.6	(0.6–3.9)	3.5	2.0	(0.8–4.9)	13.2	
F-test (p-value) ^c					F=0.32 (0.81)			F=1.16 (0.32)		
III. Twelve-month stressful experiences										
	life-threatening illness or injury of a friend or family member	131	24.5	1.9	1.0	(0.5–1.9)	–0.4	1.0	(0.5–2.0)	0.9
	death of a friend or family member	117	21.9	1.9	1.3	(0.7–2.7)	–1.1	2.3*	(1.1–4.8)	12.0*
	break-up with a romantic partner	147	27.5	2.0	1.0	(0.5–1.9)	0.2	1.1	(0.5–2.1)	1.2
	romantic partner cheated	40	7.4	1.2	0.9	(0.3–2.7)	0.1	0.8	(0.3–2.3)	–1.1
	serious betrayal someone else than partner	157	29.4	2.1	1.8	(1.0–3.4)	6.1	2.1*	(1.1–4.1)	11.1*
	serious ongoing arguments or break-up with friend or family member	187	34.9	2.2	1.4	(0.8–2.5)	0.8	2.0*	(1.1–3.8)	15.1*

	Bivariate model ^a									
	Prevalence					One-time STB ^b				
	n(w)	%(w)	(SE)	OR	95% CI	PARP(%)	OR	95% CI	PARP(%)	Two-time STB ^b
life-threatening accident	18	3.4	0.8	8.5	(0.4–166.1)	1.8	10.0	(0.5–198.6)	2.6	
seriously physically assaulted	28	5.3	1.0	5.1	(0.5–57.1)	1.5	7.2	(0.4–123.9)	4.3	
sexually assaulted or raped	18	3.3	0.8	3.6	(0.6–21.9)	1.5	4.0	(0.7–22.8)	2.0	
any serious legal problem	38	7.0	1.2	1.6	(0.4–7.1)	–0.2	3.0	(0.6–13.7)	4.5	
Any stressful event	389	72.7	2.0	1.6	(0.9–3.0)	9.9	2.2*	(1.1–4.2)	28.9*	
Number of stressful experiences										
0	146	27.3	2.0	(ref)	-	-	(ref)	-	-	
1	141	26.4	2.0	1.4	(0.7–2.9)	3.6	1.6	(0.7–3.2)	5.2	
2	106	19.8	1.8	1.6	(0.7–3.7)	4.1	1.8	(0.8–4.2)	5.5	
3+	141	26.5	2.0	1.9	(0.9–4.0)	2.6	3.4*	(1.5–7.6)	18.3*	
F-test (p-value) ^c				F=0.53 (0.66)			F=2.52 (0.057)			

Note: significant OR/PARP are shown in bold and are marked with

* (α=0.05).

^a The bivariate associations are based on a separate model for each row, with the variable in the row as the only predictor in the model.

^b no STB during follow-up is the reference category

^c Cochran-Armitage trend test. The F-test evaluates significance (α=0.05) of 200 pooled Cochran-Armitage $\chi^2(3)$ linear trend tests.

Abbreviations: GSE = general secondary education; OR = Odds ratio; PARP = population attributable risk proportion; STB = Suicidal Thoughts and Behaviors.

Table 3

Twelve-month mental disorders and severity markers of pre-matriculation STB as baseline predictors for adverse clinical patterns of STB during follow-up.

	Bivariate model ^a									
	Prevalence			One-time STB ^b			Two-time STB ^b			
	n(w)	%(w)	(SE)	OR	95% CI	PARP(%)	OR	95% CI	PARP(%)	
IV. Twelve-month mental disorders										
risk for internalizing disorder	313	58.6	2.2	1.2	(0.7–2.1)	0.9	1.5	(0.8–2.7)	14.4	
risk for externalizing disorder	181	33.8	2.2	1.3	(0.7–2.4)	3.2	1.5	(0.8–2.9)	6.7	
risk for substance use disorder	57	10.6	1.4	1.7	(0.6–4.7)	1.6	1.9	(0.6–5.8)	3.4	
risk for crime/violence disorder	1	0.2	0.2	/	/	/	/	/	/	
IED item positive	78	14.6	1.6	1.8	(0.8–4.4)	0.7	3.1*	(1.1–8.3)	9.6*	
(hypo)mania item positive	101	18.9	1.8	3.4*	(1.5–8.1)	7.0*	4.2*	(1.5–12.1)	12.1*	
any eating disorder item positive	104	19.4	1.8	1.3	(0.7–2.7)	0.6	1.8	(0.9–3.8)	7.1	
any psychotic item positive	56	10.4	1.4	1.9	(0.7–5.1)	2.9	1.9	(0.7–5.4)	2.7	
non-suicidal self-injury	76	14.2	1.5	2.0	(0.8–4.6)	–0.5	4.2*	(1.9–9.6)	13.6*	
Any positive screen	403	75.4	1.9	1.5	(0.8–2.8)	8.2	1.8	(0.9–3.6)	24.0	
Number of positive screens										
0	132	24.6	1.9	(ref)	-	-	(ref)	-	-	
1	142	26.5	2.0	1.0	(0.5–2.2)	–0.9	1.2	(0.6–2.5)	3.0	
2	109	20.3	1.8	1.4	(0.6–3.2)	4.2	1.2	(0.5–2.7)	0.8	
3+	153	28.6	2.0	2.3*	(1.0–5.1)	5.2*	3.7*	(1.6–9.0)	20.1*	
F-test (p-value) ^c				F=1.31 (0.27)			F=2.78 (0.040)			
V. Severity markers of pre-matriculation STB										
Twelve-month STB at baseline	225	42.1	2.2	1.7	(0.9–3.1)	–1.8	4.4*	(2.3–8.4)	38.1*	
Pre-matriculation lifetime STB										
- no pre-matriculation suicide plans or attempts (i.e., ideation only) ^d	236	44.1	2.2	(ref)	-	-	(ref)	-	-	
- pre-matriculation suicide plans; no attempts	229	42.9	2.2	1.6	(0.9–2.9)	5.1	2.4*	(1.3–4.7)	19.7*	

		Bivariate model ^a									
		Prevalence			One-time STB ^b				Two-time STB ^b		
		n(w)	%(w)	(SE)	OR	95% CI	PARP(%)	OR	95% CI	PARP(%)	
- pre-matriculation suicide attempts		69	13.0	1.5	3.5 *	(1.1–11.2)	0.8 *	10.3 *	(2.9–37.1)	16.9 *	
STB characteristics in worse week of pre-matriculation lifetime STB											
duration STB 4 days or more/week		248	46.3	2.2	1.4	(0.8–2.5)	3.3	1.9 *	(1.1–3.4)	17.3 *	
duration STB 5 hours or more/day		47	8.7	1.2	2.1	(0.7–6.3)	–0.7	4.9 *	(1.9–12.6)	9.2 *	
control STB very difficult/impossible		89	16.7	1.6	1.9	(0.9–4.0)	1.4	3.1 *	(1.5–6.4)	11.0 *	
any STB-associated risky behavior		167	31.2	2.0	1.2	(0.7–2.3)	0.4	1.7	(0.9–3.1)	9.8	
Course of pre-matriculation lifetime STB											
duration 5 years or more		141	26.4	2.0	1.4	(0.7–2.8)	–1.7	2.9 *	(1.5–5.7)	18.4 *	
remission at college entrance 3 years or less		423	79.0	1.8	1.4	(0.7–3.1)	–1.5	3.4 *	(1.3–9.0)	50.8 *	

Note: significant OR/PARP are shown in bold and are marked with

* ($\alpha=0.05$).

^aThe bivariate associations are based on a separate model for each row, with the variable in the row as the only predictor in the model.

^bno STB during follow-up is the reference category

^cCochran-Armitage trend test. The F-test evaluates significance ($\alpha=0.05$) of 200 pooled Cochran-Armitage $\chi^2(3)$ linear trend tests.

^dStudents with a pre-matriculation history of ideation but not plans or attempts are the reference category for the 3-category predictor

Abbreviations: GSE = general secondary education; IED = intermittent explosive disorder; OR = Odds ratio; PARP = population attributable risk proportion; STB = Suicidal Thoughts and Behaviors; / = could not be estimated.

Table 4

Concentration of risk of persistent STB cases in different proportions of incoming freshmen at highest predicted risk based on a multivariate model^a including all risk factors.

% at highest predicted risk	One-time STB		Two-time STB	
	Sensitivity(%[SE]) ^b	PPV(%[SE]) ^c	Sensitivity(%[SE]) ^b	PPV(%[SE]) ^c
100	100.0 (0.0)	28.0 (3.6)	100.0 (0.0)	27.7 (3.2)
90	90.1 (3.5)	27.9 (3.7)	98.7 (1.4)	30.3 (3.5)
80	80.8 (4.9)	28.1 (3.9)	96.2 (2.5)	33.2 (3.9)
70	71.9 (5.9)	28.6 (4.4)	92.8 (3.3)	36.5 (4.3)
60	62.4 (6.5)	28.8 (4.8)	87.6 (4.0)	40.0 (4.8)
50	53.3 (6.5)	29.4 (5.2)	80.6 (4.9)	44.1 (5.3)
40	43.4 (6.4)	29.8 (5.8)	71.6 (5.4)	48.8 (6.1)
30	33.5 (6.0)	30.4 (6.4)	59.9 (5.8)	54.4 (7.1)
20	23.0 (5.5)	31.1 (8.1)	45.4 (5.6)	61.6 (8.5)
10	12.0 (4.3)	31.9 (11.5)	27.1 (4.6)	73.1 (10.5)

^aSee Model 5 in the Supplementary materials covering multivariate model construction (Supplementary Tables 1–3). Model-based AUC values were 0.71 [SE=0.03] for *one-time* STB, and 0.87 [SE=0.03] for *two-time* STB. Cross-validated AUC values were 0.52 [SE=0.05] for *one-time* STB, and 0.79 [SE=0.04] for *two-time* STB.

^bSensitivity = Proportion of persistent STB outcome cases found among the row % of respondents at highest predicted risk, based on cross-validated predicted probabilities.

^cPositive Predictive Value (PPV) = Probability of effectively developing persistent STB when being among the row % at highest predicted risk, based on cross-validated predicted probabilities.