CASE REPORT

Secondary aortoduodenal fistula and the unrecognised herald bleed

Francesc Simó Alari,¹ Esther Molina González,² Israel Gutierrez,¹ Aïcha Ahamdanech-Idrissi³

¹General surgery, Centre Hospitalier de la val d'Ariege, Foix, France

²Intensive Care Medicine, Centre hospitalier du val d'Ariege, Foix, France

³Anesthesiology, Centre hospitalier du val d'Ariege, Foix, Midi Pyrénées, France

Correspondence to
Dr Francesc Simó Alari,
f.simo.alari@gmail.com

Accepted 9 October 2017

SUMMARY

A 65-year-old man with history of aortobifemoral surgery 4 years ago was admitted to hospital after sudden abdominal pain. Initially misdiagnosed as renal colic, he was treated with analgesics, and while on observation he started with haematemesis, rapidly responding to volume infusion. Upper gastrointestinal endoscopy showed fresh blood in the stomach with no visible active bleeding. CT scan showed an important contrast extravasation from the aorta to the third duodenal portion, restarting haemodynamic instability and a cardiac arrest. Cardiopulmonary resuscitation (CPR) manoeuvres followed by intense fluid resuscitation and urgent laparotomy with a fast transabdominal supraceliac aortic clamping was performed. After multiple blood and plasma unit transfusion and intravenous norepinephrine, two more cardiac arrests were recovered. Unfortunately, after aggressive management, the patient rapidly deteriorated and deceased on the table. Aortoduodenal fistula is a rare entity causing life-threatening bleeding. Its diagnosis requires high clinical suspicion and surgery offers the only hope for survival.

BACKGROUND

Aortoduodenal fistula (ADF) is a very rare and dreaded complication after aortic graft surgery. It is usually diagnosed postmortem due to the massive upper gastrointestinal (GI) bleeding that presents after a sudden and intense abdominal pain.

Our case had a clear 'herald bleed' that was not identified until it was out of control, allowing a CT scan to be performed the exact moment the haemorrhage restarted. The spectacularity of the images shows a clear contrast extravasation into the duodenal frame.

After a bibliographical research, we noticed that there are images of ADF published in medical papers, but none of them showing an active bleeding that could be visualised in a three-dimensional reconstruction.

We think this case illustrates in a very graphic way the massive GI haemorrhage that occurs when this type of fistula is formed, which correlates with its high mortality rate.

This case also illustrates that any patient with a history of aortic surgery or a stent graft who develops an upper GI bleed has an aortoenteric fistula (AEF) until proven otherwise. A high suspicion index is crucial to recognise and avoid treatment delays.

An immediate CT scan should be performed, and in case of being inconclusive, a duodenoscopy should be performed ideally in the operating theatre.

CASE PRESENTATION

A 65-year-old man with a history of open aorto-bifemoral Dacron graft surgery due to abdominal aortic aneurysm 4 years ago consulted the emergency department after a sudden intense diffuse abdominal pain 3 hours earlier with lumbar discomfort. On arrival, he was haemodynamically stable, with a diffusely tender abdomen but with no defence. Blood test showed an haemoglobin (Hb) of 13.8 g/dL and creatinine of 199 mmol/L and coagulation parameters were normal.

Initially oriented as a renal colic, he received intravenous painkillers and was taken to the observation area. Three hours later he presented hypotension and moderate haematemesis. A new Hb control had dropped to 9.5 g/dL, and fluid resuscitation and red blood transfusion were initiated. An upper GI gastroscopy was performed until the first duodenal portion due to the lack of duodenoscopy after haemodynamic stabilisation, showing fresh blood in the stomach with no clots. After aspiration and saline wash, no sign of oesophageal varices, peptic ulcer or any other active bleeding source was evidenced.

The patient was hospitalised in the intensive care unit for clinical control, at which moment the case was presented to our department. Being haemodynamically stable, a CT with intravenous contrast scan was performed 20 min after our visit, during which haemodynamic instability was restarted followed by cardiac arrest. CT images showed an important contrast leakage from the aorta directly to the third duodenal portion (figures 1–3). While performing cardiopulmonary resuscitation (CPR) manoeuvres and intense fluid perfusion, the patient was rapidly transferred to the operating room despite having no vascular surgeon in our hospital and only disposing of a single aortofemoral Dacron graft on stock, in a desperate attempt to control the massive GI bleeding.

A rapid induction and intubation protocol followed by an urgent xifo-pubix laparotomy with a fast liberation of adhesions (due to aortic surgery) allowed access to the oesophagogastric junction to perform a transabdominal supraceliac aortic clamping. A Cattel-Braasch manoeuvre

CrossMark

To cite: Simó Alari F, Molina González E, Gutierrez I, et al. *BMJ Case Rep* Published Online First: [please include Day Month Year]. doi:10.1136/bcr-2017-220186

Figure 1 Sagittal CT scan showing a contrast leakage from the aorta to the third duodenal portion.

permitted a complete duodenal exposure, evidencing a 1.5 cm diameter defect in the anterior aortic wall fistulising to the posterior wall of the third duodenal portion. Primary suture of the defect was performed on a nearly empty aorta. Simultaneously, CPR manoeuvres were continued with multiple red blood, fresh frozen plasma unit transfusions and autotransfusion with a Cell Saver unit and fibrinogen, high-dose norepinephrine, and bicarbonate infusion. Unfortunately and besides all efforts and aggressive management, two more cardiac arrests led to a rapidly deteriorating state, and the patient was deceased on the table.

DIFFERENTIAL DIAGNOSIS

Mallory-Weiss syndrome, gastric/duodenal ulcer, perforation or neoplasm, oesophageal varices and abdominal aortic dissection.

TREATMENT

AEF treatment includes surgical aorta repair, via in situ graft placement or extra-anatomic bypass. Primary repair of the

Figure 2 Coronal CT scan showing a contrast leakage from the aorta to the third duodenal portion.

Figure 3 Three-dimensional CT scan reconstruction. Aortoduodenal fistula connection with active extravasation of contrast to the duodenum.

small bowel is usually performed, but may require resection and anastomosis.

OUTCOME AND FOLLOW-UP

Unfortunately, after aggressive management and all efforts, the patient rapidly deteriorated and was deceased intraoperatively during surgery.

No prodromal symptoms were registered due to the rapid development of events.

DISCUSSION

AEF is a rare entity first described in 1829 by Sir Astley Cooper as an open link between the aorta and the GI tract.¹ Secondary AEF (SAEF) occurs in patients with previous aortic graft surgery. The main cause is thought to be by mechanical pressure of the graft against the bowel or a pseudoaneurysm due to perigraft infection or haematoma, each pulsation producing shear stress in the GI mucosa provoking erosion.² Despite its low incidence (1% approximately) historic mortality rates reached 57% depending on the surgical technique.³

In our case no symptoms were registered due to the rapid development of events. Some patients present prodromal symptoms prior to presentation such as fever, fatigue, malaise or even weight loss, describing a vague constitutional syndrome. Due to its life-threatening potential, any patient with a history of aortic surgery or a stent graft who develops an upper GI bleed has an AEF until proven otherwise.

Fistula can occur at any point of the GI tract, the third duodenal portion being the most frequent (two-thirds) due to its retroperitoneal location. Because of it, an enteroscope or a paediatric colonoscope could be used instead of a regular gastroscope so as to gain visualisation to distal duodenum and proximal jejunum.⁴ Signs such as visible graft, adherent clot, active bleeding, ulcer or pulsating mass suggest SAEF.⁵ Clinical presentation varies through patients, but the most common reported presentation is GI bleeding, which can start with a mild to moderate upper or lower bleeding; the classic triad of palpable abdominal mass,

abdominal pain and GI bleeding is infrequent.⁶ Two-thirds can achieve haemostasis themselves; the so called 'herald bleed' temporarily seals off through a thrombus formation, lasting no more than 24 hours, which can precede lethal exsanguination.⁷ In case no surgery is performed, mortality is near 100%;⁸ when surgery is performed on time, it often requires complex surgical repair associated with highly considerable morbidity and mortality.²

SAEF diagnosis is based on clinical grounds aided by readily available radiographical modalities. CT scan has an approximate 90% sensitivity and specificity; signs of gas shadow around the graft, focal wall thickening and inflammation, periaortic haematoma or soft tissue collection, or intravenous contrast within the GI lumen orientates its diagnosis.^{9–10} Clinical suspicion is key to prompt diagnosis and treatment, the treatment of choice being an immediate surgical repair.¹¹

In our case, the patient was not reconstructed. Nowadays, there are no established guidelines for repair of SAEF, and there is no gold standard technique for reconstruction, the most common being in-line reconstruction or extra-anatomic bypass. But the three major factors are most important when performing a surgical approach: bleeding control, infection prevention, and bowel and vascular reconstruction. Massive bleeding is a major cause of intraoperative and postoperative death. The surgeon's duty is by performing damage control surgery to obtain rapid control of the haemorrhage before the lethal triad starts (hypothermia, metabolic acidosis and coagulopathy), which has been demonstrated to drastically worsen the results and increase mortality.^{12–13} In our case, the Cattel-Braasch manoeuvre (or extended Kocher manoeuvre), usually performed in patients with trauma, offered a rapid access to the retroperitoneum. It consists of an en bloc medial rotation of all the right intraperitoneal and retroperitoneal organs to the middle line. It gives an excellent exposition of the aortic bifurcation, vena cava, right ureter and kidney.^{14–15} Once the haemorrhage point has been located, direct compression on the smaller area using gauzes or fingers can gain precious time for vascular clamping.

Clinical outcomes may be affected by the different grafts used. Femoral vein and cryopreserved homografts have a lower risk of leaking when compared with fresh allografts such as femoral vein. Rifampicin-soaked endografts have been reported to have successful outcomes.¹⁶ Postimplantation syndrome may be present after SAEF treatment, characterised as persistent fever, leucocytosis with no confirmed source of infection and negative cultures.¹⁷

An open approach implies graft excision and extra-anatomic bypass or in-line reconstruction. Extra-anatomic bypass avoids placing a prosthetic graft in a contaminated field and allows simultaneous bowel repair. However, they have a reduced patency and the patient is at risk of aortic stump blow-out, infection and thrombosis. Autologous vein repair remains, for some authors, the treatment of choice in infected aortic graft surgery, using a femoropopliteal vein, which is more time-consuming, or saphenous vein spiral graft.^{18–19}

Endovascular aortic repair consists of endovascular in situ graft replacement and partial or complete graft excision. It offers a shorter duration and requires minimal anaesthesia. It also allows fistula bleeding control while planning a definitive therapy. A multicentre study comparing open vs endovascular repair showed lower overall morbidity in endovascular patients (25% vs 77%), but secondary infection relapse persisted due to the incomplete removal of the infected graft.²⁰

Current advances in interventional radiology enable a two-staged approach, delaying definitive surgery until after

bleeding control through a stent graft.^{21–22} When available, aortic occluders such as intra-aortic balloon occlusion or resuscitative endovascular balloon occlusion of the aorta are a recognised feasible strategy for fast temporary control of intra-abdominal bleeding or shock in patients with trauma. It is relatively easy to place and allows aortic bleeding control faster and easier than other techniques; its complications are organ ischaemia and reperfusion damage, recommending an intermittent deflation of the balloon and direct hand compression to allow limb blood flow.²³ Once haemodynamic stability is achieved, vascular repair through abdominal or extra-anatomic bypass can be attempted by an experienced vascular surgeon. Vascular reconstruction is usually decided according to vascular surgeons' preferences. Usually the chosen technique is an extra-anatomic bypass (axillobifemoral bypass) followed by infected graft extraction and creation of an aortic stump. In situ reconstruction is more frequently performed in cases with negligible infrarenal aortic neck or young healthy patients.³

Bowel reconstruction strategies also play an important role on mortality. Endovascular management without later bowel reconstruction, isolation of the affected duodenal area and antibiotic therapy are associated with a high incidence of persistent or recurrent infection.¹⁸ Duodenal reconstructive techniques include primary repair, resection with primary anastomosis and exclusion. A retrospective review by Howard *et al*²⁴ compared different approaches; duodenal leaks remained the major complication, delaying recovery and increasing morbidity and mortality. Duodenal exclusion with gastrojejunostomy had the lowest rate of duodenal leaking.

Broad-spectrum antibiotics should be administered early, covering Gram-positive, Gram-negative and anaerobes, after the excision of the infected graft. Piperacillin/tazobactam, vancomycin and gentamicin provide an empirical coverage for the majority of the germs usually identified. Antibiotic therapy should be continued intravenously at least for 4–6 weeks, followed by long-term oral antibiotics.^{25–26} Failure to treat sepsis may result in 60% mortality.

AEF is a rare and dreaded complication of aortic graft replacement in which clinical presentation may vary, making its diagnosis difficult. A multidisciplinary team approach is usually requested, including vascular surgeon, general surgeon and interventional radiology. Any patient with a history of aortic graft repair presenting with upper GI bleed has an ADF until proven otherwise. High index suspicion and clinical judgement is crucial to establish prompt diagnosis that directly affects prognosis, which will depend on the haemodynamic status on presentation, surgical time and technique, and postoperative complications.

Learning points

- ▶ Any patient with a history of aortic graft repair presenting with upper gastrointestinal bleed has an aortoduodenal fistula until proven otherwise.
- ▶ Secondary aortoenteric fistula remains a highly lethal condition with mortality rates over 50%.
- ▶ Recognise the 'herald bleed' as an alarm sign.
- ▶ High index suspicion is crucial for an early diagnosis.
- ▶ Endovascular approach can be used as a temporary or definitive method to stop bleeding.

Contributors FSA: dealt directly with case, writing the report, design and data collection. EMG, IG and AA-I: dealt directly with case.

Funding The authors declare no contributorship nor external funding sources for the development of this document.

Competing interests None declared.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

© BMJ Publishing Group Ltd (unless otherwise stated in the text of the article) 2017. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

REFERENCES

- Davis JH. Complications of surgery of the abdominal aorta. *Am J Surg* 1975;130:523–7.
- Smeds MR, Duncan AA, Harlander-Locke MP, et al. Treatment and outcomes of aortic endograft infection. *J Vasc Surg* 2016;63:332–40.
- McCarthy MJ. Open or endovascular repair of secondary aortoenteric fistulae? *Eur J Vasc Endovasc Surg* 2011;41:635–6.
- Gerson LB, Fidler JL, Cave DR, et al. ACG clinical guideline: diagnosis and management of small bowel bleeding. *Am J Gastroenterol* 2015;110:1265–87.
- Batt M, Jean-Baptiste E, O'Connor S, et al. Early and late results of contemporary management of 37 secondary aortoenteric fistulae. *Eur J Vasc Endovasc Surg* 2011;41:748–57.
- Saers SJ, Scheltinga MR. Primary aortoenteric fistula. *Br J Surg* 2005;92:143–52.
- Busuttill SJ, Goldstone J. Diagnosis and management of aortoenteric fistulas. *Semin Vasc Surg* 2001;14:302–11.
- Deijen CL, Smulders YM, Coveliers HM, et al. The importance of early diagnosis and treatment of patients with aortoenteric fistulas presenting with herald bleeds. *Ann Vasc Surg* 2016;36:28–34.
- Hughes FM, Kavanagh D, Barry M, et al. Aortoenteric fistula: a diagnostic dilemma. *Abdom Imaging* 2007;32:398–402.
- Hagspiel KD, Turba UC, Bozlar U, et al. Diagnosis of aortoenteric fistulas with CT angiography. *J Vasc Interv Radiol* 2007;18:497–504.
- Gesumaria RC, Brown JM, Espinosa J, et al. When to expect aortoduodenal fistula as a cause of abdominal pain in ED patients. *Am J Emerg Med* 2016;34:e1–2468.e4:2468.e1–2468.e4.
- Varghese M, Jorgensen GT, Aune C, et al. Primary aortoduodenal fistula—a case report and a review of the literature. *Ann Vasc Surg* 2016;34:e1–4:271.e1–271.e4.
- Shapiro MB, Jenkins DH, Schwab CW, et al. Damage control: collective review. *J Trauma* 2000;49:969–78.
- Hirshberg A, Mattox KL. *Top Knife: The Art and Craft of Trauma Surgery. Chap 2.* Shrewsbury: TFM publishing, 2005.
- Jacobs LM, Gross R, Luk S. *Advanced trauma operative management. Trauma laparotomy.* Woodbury, CT: Cine Med, Inc, 2004:14. 2–24.
- Kieffer E, Gomes D, Chiche L, et al. Allograft replacement for infrarenal aortic graft infection: early and late results in 179 patients. *J Vasc Surg* 2004;39:1009–17.
- Arnaoutoglou E, Kouvelos G, Papa N, et al. Prospective evaluation of post-implantation inflammatory response after EVAR for AAA: influence on patients' 30 day outcome. *Eur J Vasc Endovasc Surg* 2015;49:175–83.
- Daenens K, Fourneau I, Nevelsteen A. Ten-year experience in autogenous reconstruction with the femoral vein in the treatment of aortofemoral prosthetic infection. *Eur J Vasc Endovasc Surg* 2003;25:240–5.
- Heikens JT, Coveliers HM, Burger DH, et al. Saphenous vein spiral graft: successful emergency repair of a mycotic aneurysm with aortoduodenal fistula. *Eur J Vasc Endovasc Surg* 2006;32:408–10.
- Kakkos SK, Antoniadis PN, Klonaris CN, et al. Open or endovascular repair of aortoenteric fistulas? A multicentre comparative study. *Eur J Vasc Endovasc Surg* 2011;41:625–34.
- Silingardi R, Coppi G, Saitta G, et al. Use of aortic occluders and endovascular plugs in a 2-step hybrid treatment of secondary aortoenteric fistulas. *Ann Vasc Surg* 2016;30:e9–15.:310.e9–310.e15.
- Marone EM, Mascia D, Kahlberg A, et al. Emergent endovascular treatment of a bleeding recurrent aortoenteric fistula as a "bridge" to definitive surgical repair. *J Vasc Surg* 2012;55:1160–3.
- Miyamoto K, Inaba M, Kojima T, et al. Intra-Aortic Balloon Occlusion (IABO) may be useful for the management of secondary aortoduodenal fistula (SADF): A case report. *Int J Surg Case Rep* 2016;25:234–7.
- Howard R, Kurz S, Sherman MA, et al. Contemporary management of secondary aortoduodenal fistula. *Ann Vasc Surg* 2015;29:1614–8.
- Herdlich BJ, Fairman RM. How to manage infected aortic endografts. *J Cardiovasc Surg* 2013;54:595–604.
- Murphy EH, Szeto WY, Herdlich BJ, et al. The management of endograft infections following endovascular thoracic and abdominal aneurysm repair. *J Vasc Surg* 2013;58:1179–85.

Copyright 2017 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- Submit as many cases as you like
- Enjoy fast sympathetic peer review and rapid publication of accepted articles
- Access all the published articles
- Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow