Published in final edited form as:

Mol Neurobiol. 2018 April ; 55(4): 3196–3210. doi:10.1007/s12035-017-0584-5.

Cocaine-mediated downregulation of miR-124 activates microglia by targeting KLF4 and TLR4 signaling

Palsamy Periyasamy, Ke Liao, Yeon Hee Kook, Fang Niu, Shannon E. Callen, Ming-Lei Guo*, and Shilpa Buch*

Department of Pharmacology and Experimental Neuroscience, 985880 Nebraska Medical Center, University of Nebraska Medical Center, Omaha, NE 68198, USA

Abstract

Cocaine is known to activate microglia both *in vitro* and *in vivo*. High expression of microglial Toll-like receptors (TLRs) and their downstream signal transducers play critical roles in determining microglial activation status. Emerging reports have also demonstrated that cocaine can enhance the strength of TLR signaling. Detailed molecular mechanisms underlying this phenomenon, however, remain elusive. In this study, we investigated the role(s) of miR-124 in regulating microglial TLR4 signaling in the context of cocaine. Herein, we found a dose- and time-dependent upregulation of KLF4 in cocaine-exposed BV-2 cells and rat primary microglial cells (rPMs). KLF4 also identified as a novel 3'-UTR target directly regulated by miR-124. In parallel, miR-124 regulated multiple TLR4 signaling molecules including TLR4, MyD88, TRAF6, and IRAK1. Repeated doses of cocaine (20 mg/kg; *i.p.*) administration in mice for 7 days further validated the *in vitro* key findings. Also, miR-124 overexpression significantly blocked the cocaine-mediated upregulation of pro-inflammatory cytokines. In contrast, miR-124 overexpression notably increased the expression of anti-inflammatory mediators in cocaine-exposed microglial cells. Intriguingly, stereotactic administration of lentivirus-miR-124 in the striatum significantly inhibited cocaine-mediated microglial activation and locomotor hyperactivity *in vivo*. In summary, these findings implicate the role of miR-124 in regulating TLR4 signaling, thereby indicating a new pathway responsible for cocaine-mediated microglial activation.

Keywords

neuroinflammation; cocaine; TLR4; microglial cells; MyD88; TRAF6; miR-124

Introduction

Microglia, the brain resident macrophages, are the primary source of innate immune responses in the central nervous system (CNS). The activation status of microglia determines the intensity of inflammation in the CNS [1,2]. Microglia expresses a variety of pattern-

*Corresponding authors at: Department of Pharmacology and Experimental Neuroscience, 985880 Nebraska Medical Center, University of Nebraska Medical Center, Omaha, NE 68198, USA. sbuch@unmc.edu (Shilpa Buch), minglei.guo@unmc.edu (Ming-Lei Guo).

Conflict of interest: The authors declare no competing financial interests.

recognition receptors (PRR). Of these, the prototypic PRR, toll-like receptor 4 (TLR4) and its downstream signal transducers including myeloid differentiation primary-response protein 88 (MyD88), IL-1R-associated kinase 1 (IRAK1), TNF receptor-associated factor 6 (TRAF6), and NF- κ B, are well-established players in modulating microglial activation [3-6]. TLR4-induced NF- κ B translocation causes transcriptional upregulation of a plethora of pro-inflammatory cytokines [7,8]. Krüppel like factor 4 (KLF4) is also a downstream transcriptional factor of the TLR4 pathway, and its expression is upregulated in microglial cells following lipopolysaccharide (LPS) stimulation [9]. Moderate activation of microglia is a necessary initial response to brain insult. Persistent activation of microglia, on the other hand, leading to a state of exacerbated neuroinflammation results in neuronal death [10-12]. Increased microglial activation has been shown in multiple neurodegenerative diseases such as Parkinson's disease (PD), Alzheimer's diseases, and amyotrophic lateral sclerosis [13-15] and drug addiction [16,17].

Emerging evidence suggests a close link between drug addiction and the dysregulation of central immune pathways [18,19]. Addictive drugs including cocaine have been consistently shown to activate microglia both *in vitro* and *in vivo* [19-22,17]. Interestingly, inhibition of glial cell activation was shown to block cocaine- and methamphetamine-mediated behavioral changes in rodents, implying thereby that elevated neuroinflammation plays causative roles in the development of drug addiction [23,24]. Previous reports have demonstrated TLR4 signaling [25] as well as reactive oxygen species (ROS), ER stress/autophagy axis to be involved in cocaine-mediated microglial activation [17].

MicroRNAs (miRs) are an important class of biological molecules with critical roles in many fundamental cellular processes such as proliferation, differentiation, and apoptosis [26-28]. The brain is one of the miR-enriched organs comprising of several miRs that are vital for its functioning including development, neurogenesis, and neuroinflammation [29]. Several brain-enriched miRs including miR-124 have been demonstrated to modulate microglial functions. Under normal conditions, miR-124 is highly expressed constitutively in microglia and contributes to its quiescence [30,31]. Downregulation of miR-124 has been associated with multiple neuroinflammatory disorders such as PD, dementia and Multiple Sclerosis [32,33,31,34]. Interestingly, the expression levels of miR-124 can be decreased by the classic TLR4 activator, LPS and correspondingly, miR-124 overexpression has been shown to block LPS-induced microglial activation [35].

The current study was aimed at examining the modulation of miR-124 in microglia and to identify potential targets of miR-124 in the context of cocaine exposure. Our findings demonstrated that KLF4 and TLR4 signaling mediators are novel targets directly regulated by miR-124. Overexpression of miR-124, resulted in amelioration of cocaine-mediated microglial activation *in vitro* as evidenced by the downregulation of proinflammatory factors and upregulation of anti-inflammatory factors in cocaine-exposed microglia transfected with miR-124. Furthermore, intrastriatal administration of lentivirus-miR-124 injection in mice resulted in abrogation of cocaine-mediated microglial activation. Taken together, our findings implicate the role of miR-124 in regulating TLR4 signaling, thereby indicating a new pathway responsible for cocaine-mediated microglial activation.

Materials and methods

Reagents

Cocaine hydrochloride (C5776), and 5-Azacytidine (5-AZA; A3656) were purchased from Sigma-Aldrich. TaqMan® microRNA assays for miR-124 (001182) quantification was purchased from Applied Biosystems; miR-124 mimic and inhibitor and negative control were purchased from Dharmacon; Lentivirus-miR-124 and control lentivirus were purchased from BioSetia (miR-LV084 and LV000). Antibodies to TLR4 (sc-10741), IRAK1 (sc-7883), TRAF6 (sc-7221), MyD88 (sc-11356) and KLF4 (sc-20691) were purchased from Santa Cruz Biotechnology. Goat anti-rabbit (sc-2004) and goat anti-mouse (sc-2005) secondary antibodies were also from Santa Cruz Biotechnology. Realtime qPCR primers for TNF α , IL6, IL1 β , CCL2, TGF β , IL4, IL10 and Arg1 were commercially available from Invitrogen.

BV-2 cells and HEK293 cell culture

BV-2 immortalized cell line was obtained from Dr. Sanjay Maggirwar (University of Rochester Medical Center, Rochester, NY, USA) and was grown and routinely maintained in DMEM (Invitrogen, 11995-065) with 10% heat-inactivated fetal bovine serum (FBS, Invitrogen, 16000-044) at 37°C and 5% CO₂ and used up to 20 passages. HEK293 cells were purchased from ATCC (CRL-1573) and grown and routinely maintained in DMEM with 10% FBS at 37°C and 5% CO₂.

Rat primary microglial cell culture

Rat primary microglial cells were obtained from Sprague-Dawley newborn (1-3 days) pups and isolated according to the protocol described by Ni and Aschner [36]. Briefly, after digestion and dissociation of the dissected brain cortices in Hanks buffered salt solution (Invitrogen, 14025076) supplemented with 0.25% trypsin (Invitrogen, 25300-054), mixed glial cultures were prepared by resuspending the cell suspension in DMEM supplemented with 10% FBS with 100 U/ml penicillin, and 0.1 mg/ml streptomycin. Cells were plated at 10×10^6 cells/flask density onto 75 cm² cell culture flasks. The cell medium was replaced every 5 days, and after the first medium change, macrophage colony-stimulating factor (0.25 ng/ml; Invitrogen, PHC9504) was added to the flasks to promote microglial proliferation. When confluent (7 to 10 days), mixed glial cultures were subjected to shaking at 37°C at 220 g for 2 h, to promote microglia detachment from the flasks. The cell medium, containing the released microglia cells was collected from each flask and centrifuged at 1000 g for 5 min to collect cells and plated onto cell culture plates for all subsequent experiments. The purity of microglial cultures was evaluated by immunocytochemistry using the antibody specific for Iba-1 (Wako Pure Chemical Industries, 019-19741) and used if >95% pure.

Western blotting

Treated cells were lysed using the Mammalian Cell Lysis kit (Sigma, MCL1-1KT). 10 μ g of the proteins were electrophoresed on a 10 % sodium dodecyl sulfate-polyacrylamide gel under reducing conditions followed by transfer to PVDF membranes (Millipore, IPVH00010). The blots were blocked with 5 % nonfat dry milk (in 1 $\times$ TTPS buffer).

Western blots were then probed with antibodies specific for the indicated proteins. The protein amounts loaded were normalized according to the β -actin signal, using an anti- β -actin antibody (Sigma, A5441). The secondary antibodies were HRP conjugated to goat anti-mouse/rabbit IgG.

Quantitative polymerase chain reaction (qPCR)

Total RNA was extracted using Quick-RNA™ MiniPrep Plus (Zymo Research, R1058) as per manufacturer's protocol and quantified using Nanodrop. Reverse transcription reactions were performed using a Verso cDNA kit (Invitrogen, AB-1453/B), as per manufacturer's instructions. qPCRs were performed by using SYBR Green ROX qPCR Mastermix (Qiagen, 330510). Reaction systems were set up as follows: 10 μ l SYBR Green Mastermix, 0.5 μ l forward primers, 0.5 μ l reverse primers and 9 μ l distilled water. 96-well plates were placed into a 7500 Fast Real-Time PCR system (Applied Biosystems, Grand Island, NY) for a program running. Actin levels were set as an internal control for calculation.

TaqMan® microRNA assays for miR-124

TaqMan® microRNA assays for miR-124 were commercially available from Applied Biosystems. TaqMan® MicroRNA Reverse Transcription Kit (PN 4366596) was utilized according to indicated protocol. The reaction system (15 μ l) was set as followed: 10 mM dNTPs (with dTTP) 1.5 μ l, MultiScribe™ Reverse Transcriptase 50 U/ μ l 1.0 μ l, 10 $\times$ Reverse Transcription Buffer 1.5 μ l, RNase Inhibitor, 20U/ μ l 0.2 μ l, Nuclease-free water to 7 μ l, total RNA 5 μ l (~50 ng), 3 μ l of RT (5 $\times$) primer. The tube containing the reaction mixture was centrifuged and loaded onto the thermal cycler: 16°C for 30 min; 42 °C for 30 min; 85 °C for 5 min. The reverse transcription product was then diluted 1:10 for the following PCR reaction: TaqMan® PCR primer (20 $\times$) 1 μ l, RT reaction product 1.5 μ l, TaqMan® 2 $\times$ Universal PCR Master Mix, No AmpErase UNGa 10 μ l (PN 4324018) and distilled water up to 20 μ l. The PCR conditions were as follows: hold 95°C for 10 min, then 40 cycles for 95°C for 15 second and 60°C for 1 min. All reactions were run in triplicate. The expression level of miR-124 was calculated by normalizing to U6 snRNA.

Luciferase reporter assays

KLF4 plasmids containing different 3'-UTR regions were provided by Dr. Michael Ruppert (pMIR-Report-Luc-KLF4, addgene). Briefly, the HEK293 cells were seeded into 96-well plates and co- transfected with KLF4 3'-UTR plasmids and miR-124 mimics/miR-scramble in a molar ratio 10:1. The luciferase activity was determined 24 h posttransfection, and the reporter assay was performed according to the manufacturer's protocol (Promega). Renilla luciferase activity was normalized to firefly luciferase and expressed as a percentage of the control.

Ago2 Immunoprecipitation

BV-2 cells were plated onto 6-well plates and transfected with either miR-control or miR-124 mimics for 24 h. After treatment, cells were washed in the cold 1 $\times$ PBS, scraped, and then lysed with a buffer containing 0.5% NP40, 150 mM KCL, 25 mM Tris-glycine pH 7.5, 2 mM EDTA, 0.5 mM DTT, and inhibitors of RNases, proteases, and phosphatases.

10% of total lysate was removed and kept as the input samples and the remainder used for immunoprecipitation. 10 µg of anti-Ago2 (11A9, Sigma-Aldrich, SAB4200085) or anti-FLAG (M2, F1804, Sigma-Aldrich) antibodies were incubated overnight with protein A/G agarose beads (Thermo Scientific, 20423) at 4° C. Precleared lysates were then incubated with the appropriate antibody-bound beads, and the immunoprecipitated proteins were washed and incubated with DNase I (Invitrogen, 18068015) followed by digestion with proteinase K (Zymo Research, D3001-2) for 15 mins. RNA extraction was then performed using Quick-RNA™ MiniPrep Plus (Zymo Research, R1058) and quantified using Nanodrop. Total, as well as Ago2 immunoprecipitated RNA samples, were then used for determining the binding targets of miR-124, such as KLF4, MyD88, IRAK1, TRAF6 and TLR4 by qPCR.

miR-124 oligo transfection

MiR-124 mimics, inhibitors, and control were purchased from GE Health Corporation. BV-2 cells were seeded into 24-well plates (1×10^5 /well) and transfected with 30 pmol of various oligos by Lipofectamine 2000 according to manufacturer's protocol (Invitrogen). 24 h later cellular homogenates were extracted to detect various proteins levels as indicated.

Chronic cocaine administration

Sprague-Dawley rats and C57BL/6N mice were purchased from Charles River Laboratories (Wilmington, MA, USA). All the animals were housed under conditions of constant temperature and humidity on a 12-h light, 12-h dark cycle, with lights on at 0700 h. Food and water were available ad libitum. All animal procedures were performed according to the protocols approved by the Institutional Animal Care and Use Committee of the University of Nebraska Medical Center and the National Institutes of Health. For repeated cocaine administration, C57BL/6N mice were injected with cocaine (20mg/kg, *i.p.*) or saline for 7 days. 1 h after the last injection, mice were sacrificed for brain removal. The striatal protein homogenates were extracted and assessed for the levels of various proteins as indicated.

Stereotactic injection of Lentivirus-miR-124

Male 8-week-old C57BL/6N mice were divided into four groups ($n = 8$) receiving the following treatments: (1) saline with miR-control lentivirus; (2) saline with miR-124 lentivirus; (3) cocaine with miR-control lentivirus; (4) cocaine with miR-124 lentivirus. The lentiviruses (1.5 µl) were microinjected into the mouse brain striatum (at least 10^7 infectious units per ml) using the microinjection parameters (coordinates 2.8 mm behind the bregma, 1.7 mm lateral from the sagittal midline at a depth of 4.5 mm to skull surface). To increase the precision, a 10-ml Hamilton syringe with a 33G needle was utilized for intrastriatal injection as described.

Open-field test

Open-field consisted of the square arena (40 cm × 40 cm) and wall (35 cm high). Mice were carried to the test room in their home cages and were handled by the base of their tails at all times. Mice were placed at one of the four corners of the open field and allowed to explore the apparatus for 5 minutes, following which mice were returned to their home cages and

received the final injection of cocaine/saline. The mice were then placed at the same corner of the arena, and a video camera was suspended directly over the arena for recording behavioral activities for 20 mins using an automated video tracking system (EthoVision 3.1, Noldus Information Technology, Inc, Leesburg, VA). The parameters such as the total distance of running, latency time and the average speed were included and calculated using the Ethovision 3.1 software. After each test, the apparatus was thoroughly cleaned with cotton pad wetted with 70% ethanol. A single primary observer blinded to the experimental condition conducted the behavioral observations.

Statistical analysis

All the data were expressed as mean $\pm$ SEM, and appropriate statistical significance was chosen based on the experimental design using GraphPad Prism version 5 (San Diego, CA, USA). The specific statistical analysis used is indicated in the text and each figure caption for all studies. Bonferroni post hoc test was used for one-way analysis of variance (ANOVA) and Tukey post hoc test was used for two-way ANOVA. Values were considered statistically significant when $P < 0.05$.

Results

Cocaine upregulated KLF4 levels *in vitro* and *in vivo*

KLF4, characterized as a zinc finger-containing transcriptional factor, belongs to the KLF protein family which includes 17 different members [37]. KLF4 is essential for embryonic development and regulates a variety of processes including cell proliferation, differentiation, apoptosis, migration, and invasion [34]. KLF4 is expressed in a variety of cell types including monocytes/macrophages. Previous reports have demonstrated that KLF4 is expressed in microglia and functions as a downstream factor of TLR signaling, plays critical roles in microglial activation and ensuing neuroinflammation [38-41]. Multiple studies have shown that psychostimulants including cocaine can activate microglia both *in vitro* and *in vivo* [19-22]. To investigate whether cocaine-mediated microglial activation involved KLF4, BV-2 cells were exposed to varying doses of cocaine (1, 10, and 100 μ M) for 24 h. Following cocaine exposure, total cellular proteins were isolated to determine levels of KLF4 expression by western blotting. Our findings (Fig. 1A) showed that cocaine dose-dependently upregulated KLF4 protein levels ($P < 0.05$). Next, we isolated rPMs from the brain tissues of newborn rat pups and exposed them to cocaine, as was done for BV-2 cells. Consistent with the BV-2 cells, cocaine also upregulated KLF4 levels in rPMs ($P < 0.05$; Fig. 1B). Based on these experiments, we have chosen 10 μ M cocaine as a dose for ensuing experiments. The rationale for choosing 10 μ M cocaine is based on the fact that following intranasal cocaine administration plasma levels of cocaine range between 0.4 to 1.6 μ M [42], while the plasma cocaine concentrations in tolerant abusers reach levels up to 13 μ M [43]. Additionally, cocaine levels in postmortem brains of chronic cocaine users following acute intoxications have been reported higher than 100 μ M [44]. Next, we performed time-course experiments to determine KLF4 protein levels in both BV-2 cells and rPMs exposed to 10 μ M cocaine for varying time points. Our findings clearly showed that cocaine significantly ($P < 0.05$) increased KLF4 expression in a time-dependent manner. This upregulation was evident as early as 6 h post-treatment and persisted for 24 h in both BV-2 cells (Fig. 1C) and

rPMs (Fig. 1D). We next sought to investigate whether cocaine also upregulated KLF4 levels *in vivo*. For this wild-type, C57BL/6N mice were injected with cocaine (20 mg/kg, *i.p.*) for 7 consecutive days and the striatum was removed for assessment of KLF4 protein levels. Our findings demonstrated that KLF4 levels were significantly ($P < 0.05$) elevated in cocaine-treated mouse brains compared with saline-treated controls (Fig. 1E).

KLF4 is a novel 3'-UTR target of miR-124

We next explored the mechanisms underlying cocaine-mediated upregulation of KLF4. miRs have been demonstrated to play critical roles in regulating microglial inflammatory responses [45-47,29]. Our previous work showed that cocaine downregulated the expression of miR-124 levels via promoter DNA methylation in microglia [48]. Herein we replicated these findings and again demonstrated that exposure of both BV-2 cells and rPMs to cocaine downregulated miR-124 levels in a time-dependent manner (Supplementary Fig. S1). We next sought to examine a possible functional linkage between miR-124 and KLF4 in the context of cocaine. Bioinformatics analyses (miRanda, and TargetScan) suggested there are three binding sites located in the 3'-UTR of KLF4 mRNA (nucleotide position: 456-475, 772-791, and 814-833) indicating KLF4 as a direct target regulated by miR-124 (Fig. 2A). To investigate the effects of miR-124 on KLF4 levels, BV-2 cells were transfected with miR-124 mimics/scrambles followed by exposing the transfected cells to cocaine (10 μ M) for 24 h. As shown in Fig. 2B, miR-124 overexpression significantly ($P < 0.05$) inhibited endogenous KLF4 levels while also attenuating cocaine-mediated upregulation of KLF4 in BV-2 cells. Cocaine-mediated upregulation of KLF4 levels and the inhibitory effects of miR-124 on KLF4 expression was also validated in rPMs (Fig. 2C). Reciprocally, BV-2 cells and rPMs transfected with miR-124 inhibitors demonstrated a significant ($P < 0.05$) upregulation of KLF4 expression levels (Fig. 2D and 2E). To confirm that miR-124 could directly bind to the KLF4 3'-UTR, we next cotransfected the recombinant KLF4 3'-UTR (fragment -1 or -2) luciferase reporter plasmids (schematic shown in Fig. 2F) with miR-124 mimics and miR-124 inhibitor, respectively, into HEK-293 cells, followed by an assessment of luciferase activity. Our findings demonstrated a decrease in luciferase activity in cells cotransfected with 3'-UTR-2 (containing the binding sites) and miR-124 mimics compared with cells cotransfected with 3'-UTR-1 and miR-124 mimics (Fig. 2F). In contrast, cells cotransfected with miR-124 mimics and KLF4 3'-UTR-1 had a negligible effect on luciferase activity. However, cells cotransfected with KLF4 3'-UTR plasmids and miR-124 inhibitors had no effects on luciferase activity. The binding of miR-124 with its target, KLF4 was further confirmed using Argonaute 2 immunoprecipitation assay (Ago2 IP). In principle, Ago2 is an indispensable component of the RNA-induced silencing complex (RISC) that is directed to target mRNAs by the miRNA guide strand. This complex is then immunoprecipitated and the mRNA levels quantified by qPCR. Based on this, we performed Ago2 IP in BV-2 cells that were transfected with either miR-124 mimics or miR-control for 24 h. As shown in Fig. 2H, Ago2 IP confirmed KLF4 as a direct target of miR-124 because KLF4 mRNA showed increased Ago2 binding in a miR-124-dependent manner. Taken together, these findings demonstrated KLF4 as a direct 3'-UTR target regulated by miR-124.

Overexpression of miR-124 inhibited cocaine-mediated TLR4 upregulation

A striking feature of miR regulation is that a single miR can modulate several hundred mRNAs simultaneously to modify cellular activity. In addition to KLF4, we also assessed other molecules which are potential direct targets of miR-124 and also known to play a role in cocaine-mediated microglial activation. Recently, the interaction and activation of TLR4 following cocaine exposure was shown to be critical for cocaine-mediated neuroinflammation *in vivo* [49,25]. In the current study, we sought to examine whether cocaine also regulated expression levels of TLR4 protein in microglial cells. BV-2 cells were exposed to cocaine at different doses (1, 10, and 100 μ M for 24 h) for the assessment of TLR4 protein levels. Our findings demonstrated that cocaine exposure to BV-2 cells resulted in increased expression of TLR4 protein levels in a dose-dependent manner with highest expression at cocaine concentrations of 10 and 100 μ M ($P < 0.05$; Fig. 3A). Next, we determined the time-dependent expression of TLR4 levels, in which BV-2 cells were exposed to 10 μ M cocaine for different time periods (0-24 h). As shown in Fig. 3B, exposure of BV-2 cells to 10 μ M cocaine significantly ($P < 0.05$) increased TLR4 protein levels on 6 h onwards while negligible effects at the earlier time point (3 h). Furthermore, upregulation of TLR4 expression was also observed in cocaine-administered mouse brains compared to saline-administered controls. Similar to the cocaine-mediated upregulation of KLF4 expression, we also observed upregulation of TLR4 in the presence of cocaine (Fig. 3C). Based on the bioinformatics analyses (miRanda and TargetScan) that TLR4 has two binding sites for miR-124 in its 3'-UTR region (Fig. 3D), we next sought to confirm that TLR4 is indeed a target of miR-124. BV-2 cells were transfected with miR-124 mimics followed with cocaine exposure for 24 h. Protein homogenates were assessed for detection of TLR4 levels and as shown in Fig. 3E, overexpression of miR-124 significantly ($P < 0.05$) inhibited endogenous TLR4 levels while also abrogating cocaine-mediated upregulation of TLR4 levels. In contrast and, as expected, transfection of BV-2 cells with miR-124 inhibitors resulted in increased expression of TLR4 (Fig. 3F). The regulation of TLR4 levels by miR-124 was further validated in rPMs wherein overexpression of miR-124 followed by cocaine exposure resulted in decreased levels of endogenous TLR4 (Fig. 3G). On the other hand, transfection of rPMs with miR-124 inhibitors followed by cocaine exposure resulted in statistically increased ($P < 0.05$) expression of TLR4 (Fig. 3G). Ago2 IP assay further confirmed the binding of miR-124 with TLR4 in BV-2 cells transfected with miR-124 mimics compared with miR-control (Fig. 3H). Taken together, these results provide evidence that cocaine exposure enhanced TLR4 levels both *in vitro* and *in vivo* and that overexpression of miR-124 blocked these effects.

Overexpression of miR-124 also inhibited the levels of MyD88, IRAK1, and TRAF6

Previous reports demonstrated that miR-124 negatively regulated TLR4 signaling by modulating the levels of MyD88 and TRAF6 in both alveolar macrophage and epithelial cells [50,51]. In addition to these two TLR4 signaling transducers, we found that IRAK1 has one miR-124 binding site in its 3'-UTR indicating IRAK1 could be yet another target regulated by miR-124 (Fig. 4A). The next step was to explore whether cocaine exposure could also lead to increased expression levels of these three signal transducers (MyD88, IRAK1, and TRAF6) and if so, whether miR-124 overexpression could block these effects. BV-2 cells were transfected with either miR-124 mimics or scrambled siRNA followed by

cocaine exposure for 24 h and protein homogenates prepared for targeted proteins detection. Our results clearly showed that cocaine significantly ($P < 0.05$) increased the levels of MyD88, IRAK1, and TRAF6 and that miR-124 overexpression abrogated cocaine-mediated upregulation of these mediators (Figs. 4B, 4D, and 4F). In contrast and, as expected, transfection of BV-2 cells with miR-124 inhibitor resulted in increased expression of MyD88 (Fig. 4C), IRAK1 (Fig. 4E) and TRAF6 (Fig. 4G) compared with cells transfected with miR-control. These findings were also validated in rPMs transfected with miR-124 mimics or miR-124 inhibitors (Fig. 4H and 4I). Ago2 IP assay further confirmed the binding of miR-124 with MyD88, IRAK1, and TRAF6 in BV-2 cells transfected with miR-124 mimics compared with cells transfected with miR-control (Fig. 4J). Taken together, our results suggest that in addition to KLF4 and TLR4, miR-124 also modulates the levels of other signaling transducers such as MyD88, IRAK1, and TRAF6 in microglial cells.

Overexpression of miR-124 inhibits cocaine-mediated microglial activation *in vitro*

The findings presented above demonstrate that miR-124 modulates the expression levels of multiple mediators in TLR4 signaling. Since miR-124 is known to maintain microglia in a quiescent state and also promote the transition of microglia from a proinflammatory to an anti-inflammatory phenotype [35,52], we next sought to investigate the effect of miR-124 overexpression on cocaine-mediated microglial activation. BV-2 cells were transfected with miR-124 mimics or scrambled siRNA followed by exposure to cocaine and assessed for expression levels of M1 (pro-inflammatory) and M2 (anti-inflammatory) markers by qPCR. The representative M1 markers included TNF α , CCL2, NOS2 while the M2 markers included TGF β , IL4, and IL10. As shown in Figs. 5A, 5B, and 5C, cocaine exposure of BV-2 cells resulted in a significant ($P < 0.05$) increase in mRNA levels of the M1 markers (TNF α : 2.68 ± 0.17 ; CCL2: 2.10 ± 0.09 ; NOS2: 3.63 ± 0.20 folds; respectively, $P < 0.05$) and overexpression of miR-124 abrogated cocaine-mediated effects on these M1 markers. Concomitantly, there was a substantial increase in expression of M2 anti-inflammatory markers in miR-124 overexpressing microglia (TGF β : 1.56 ± 0.03 ; IL4: 3.82 ± 0.14 ; IL10: 3.32 ± 0.07 folds, respectively, $P < 0.05$) as shown in Figs. 5D, 5E, and 5F. Our results thus imply that miR-124 overexpression blocked cocaine-mediated microglial activation by promoting microglia from a proinflammatory to an anti-inflammatory phenotype. In addition to M1 and M2 markers, we also demonstrated that the levels of two other miR-124 targets c/EBP (Fig. 5G) and ARG1 (Fig. 5H) were also decreased in cells over-expressing miR-124, a finding that is in agreement with previous reports [31].

Overexpression of Lentivirus miR-124 inhibits cocaine-mediated microglial activation *in vivo*

Based on our findings that miR-124 overexpression inhibited cocaine-mediated microglial inflammatory responses *in vitro*, we next sought to validate these observations *in vivo*. For this wild-type C57BL/6N mice (n=8) were stereotactically injected with RFP-lentivirus (LV) containing miR-124 or miR-control into the striatum. Two weeks later mice were administered with cocaine (20 mg/kg; *i.p.*) for 7 consecutive days (schematic shown in Fig. 6A). 1 h after the last injection, mice were sacrificed and striatal tissues processed for assessment of RFP expression to examine the efficacy of miR-124 expression. Interestingly, mice that received LV injection revealed strong RFP expression co-localized with microglia

in the striatum indicating successful virus transfection and expression *in vivo* (Fig. 6B). As expected, mice injected with LV-miR-124 exhibited increased miR-124 levels in the striatum compared to mice receiving control virus injection (Fig. 6C). Cocaine administration significantly ($P < 0.05$) downregulated miR-124 levels in the mice brains compared with mice receiving LV-miR-control ($P < 0.05$, Fig. 6C). We next wanted to assess the effect of miR-124 overexpression on KLF4 and TLR4 levels in the striatum in the presence of cocaine. Cocaine administration significantly increased KLF4 and TLR4 levels in the LV-miR-control group as expected. Interestingly, there was no significant changes in KLF4 and TLR4 levels in LV-miR-124 injected mice indicating thereby that miR-124 overexpression blocked cocaine-mediated upregulation of KLF4 and TLR4 (Figs. 6D, 6E and 6F). We also observed that there is no difference in KLF4 levels between the two saline groups following miR-124 overexpression. One possible explanation for this discrepancy could be that the constitutive expression of miR-124 is already high in the saline group (maintaining a basal level of KLF4) and that further upregulation of miR-124 has little effect in modulating the KLF4 levels. Our findings also demonstrated that cocaine-mediated upregulation of microglial activation (increased Iba-1 levels) was abrogated in mice with LV-miR-124 injection (Figs. 6D and 6G). Additionally, to assess the role of miR-124 on cocaine-mediated locomotor activity, mice overexpressing LV-miR-124 or miR-control were administered cocaine for 7 days following which the mice were monitored for locomotor activity using the open-field test. Fig. 6H shows the total distance traveled for 20 min following the last saline or cocaine injection. Regarding the distance traveled there was no statistical difference observed between the saline administered LV-miR-control and LV-RFP-miR-124 groups. Interestingly, however, cocaine injection had a notable stimulating effect on locomotor activity in both LV-miR-control as well as LV-RFP-124 injected mice, with a significant downregulation of locomotor activity in LV-miR-124 mice compared with the LV-miR-control group ($P > 0.05$). Taken together, these findings demonstrated that miR-124 overexpression blocked cocaine-mediated microglial activation *in vivo*, thereby lending support to the idea that miR-124 maintains microglia in a quiescent state by inhibiting neuroimmune signaling induced by cocaine.

Discussion

Although cocaine is well-known to activate microglia leading to increased neuroinflammation, the detailed molecular mechanism(s) underlying this phenomenon remain poorly understood. It is now being recognized that development of addiction is associated with a neuroinflammatory phenotype, specifically with the implication of activation of microglia. MiR-124 is a brain-enriched miR that is highly expressed in microglia. This specific miR has been well established as a master regulator of microglial quiescence. Downregulation of miR-124 has been suggested in multiple neuroinflammatory diseases including PD, Amyotrophic Lateral Sclerosis, with miR-124 overexpression shown to mitigate the pathogenesis associated with these brain diseases. Since cocaine addiction is linked to microglial activation and the fact that miR-124 is downregulated in response to cocaine exposure both *in vitro* and *in vivo*, we infer that miR-124 downregulation could be critical in the development of drug addiction. Following our previous studies showing that cocaine downregulated microglial miR-124 [48], herein we demonstrate that miR-124 can

modulate the strength of TLR4 signaling by regulating multiple molecules including TLR4, MyD88, IRAK1, TRAF6, and KLF4 and that microglial over-expression of miR-124 inhibited cocaine-mediated activation both *in vitro* and *in vivo*. Since elevated neuroinflammatory levels are believed to contribute to the development of drug addiction, our results suggest that miR-124 can be envisioned as a potential therapeutic agent for the treatment of drug addiction via its inhibition of microglial activation.

Over the last several decades, much attention has been focused on the effects of abused drugs, such as cocaine, on signal transmission between neurons and the ensuing neuronal excitability in the context of addiction. Recent emerging evidence has also shed light on the cross-talk between neurons and glia (neuroimmune signaling), as a contributor to the development of drug addiction. Abused drugs including cocaine, amphetamine, and alcohol have long been recognized to activate microglia/astrocytes in multiple *in vitro* and *in vivo* model systems. Findings from our lab have demonstrated the ability of cocaine to activate microglia and rPM, leading in turn, to the production and release of multiple pro-inflammatory factors including TNF α , IL6, and CCL2 [17,53,19]. Various other reports have also demonstrated increased neuroinflammation following exposure of rodents to cocaine and other abused drugs. For example, amphetamine administration (4 mg/kg at 2-h intervals, four times) was shown to activate microglia in mice [54]. In another study acute, high dose (30 mg/kg) of methamphetamine exposure increased the mRNA levels of pro-inflammatory cytokines IL-6 and TNF α in the striatum and hippocampus of mice [55]. Another study demonstrated that even lower doses of methamphetamine could activate microglia [56]. Furthermore, the relationship between psychostimulant exposure and ensuing neuroinflammation has also been confirmed by analyzing annotated datasets which are available in the GEO public database [57-59]. Clinical evidence has also revealed a close linkage between neuroinflammation and drug addiction. For example, it has also been shown that there is a statistically significant increase in activated macrophages/microglia in chronic cocaine abusers [60]. In another study comparing cognitive function with peripheral markers of immune activation, significant correlations were observed in methamphetamine users but not in matched controls [61]. Another study using positron emission tomography suggested a diffuse persistent increase in activated microglia in all brain regions of human methamphetamine abusers compared to control subjects [20]. Studies by Watkin's group has elegantly demonstrated that cocaine can directly interact with TLR4 to enhance microglial innate immune responses including increased secretion of IL1 β , leading in turn, to upregulation of synaptic dopamine concentrations in the striatum. In this study using the TLR4 KO mice, the authors demonstrated that there was a decrease in cocaine-induced behavioral changes in both conditioned place preference (CPP) and self-administration assays, implying thereby that microglial activation via TLR4 was critical for cocaine addiction [25]. Intriguingly, inhibitors of glial activation have recently been suggested as potential treatment strategies for psychostimulant abuse [62]. Taken together, glial-neuronal cross-talk induced by drug abuse further facilitates the inflammatory factors that are critical targets for the development of psychostimulant abuse medications.

Herein we have provided a novel molecular mechanism responsible for cocaine-mediated microglia activation (neuroinflammation). Previously we demonstrated that cocaine exposure in microglia leads to downregulation of microglial miR-124 levels via an increase

in the methylation status of the miR-124 promoter. Downregulation of miR-124, in turn, resulted in activation of the TLR4-mediated signaling via elevation of multiple miR-124 regulated mediators such as KLF4, TLR4, MyD88, TRAK1, and TRAF6. Since both microglial activation and elevated neuroimmune signaling have been suggested to activate the dopaminergic neuronal activity in the brain striatum, miR-124 could be envisioned to play critical roles in cocaine-associated reward circuitry and possibly, also in addiction to cocaine.

It is widely recognized that small noncoding RNAs, such as microRNAs play critical roles in modulating glial activation and other cellular processes. As an example, miR-124, a brain-enriched miRNA, plays key roles in neurogenesis, synapse plasticity, and glia-neuronal interactions to maintain brain homeostasis. MiR-124 was first identified as a neuroprotective factor promoting neurogenesis during brain development and maintenance of synaptic homeostasis in the adult brain [63,64]. There have been reports indicating miR-124 as one of the miRs that are highly expressed in resting microglia and, its expression levels were found to correlate with microglia quiescence [31,30] closely. In another study, it was shown that overexpression of miR-124 blocked LPS-induced microglial activation *in vitro* [35]. Decreased miR-124 levels have been reported in various neuroinflammatory disorders including PD, dementia and Multiple Sclerosis [32,33,31,34]. Drug addiction is recognized as a neuroinflammation-related disease with several abused drugs having demonstrated the ability to modulate miR-124 levels. It has been shown that chronic administration of cocaine decreased miR-124 levels in the nucleus accumbens (NAc) and that down-regulation of miR-124 was critical for cocaine-induced synaptic plasticity and behavioral changes [65]. These findings are in agreement with our results that cocaine exposure leads to downregulation of miR-124, culminating ultimately into microglial activation. The role of miR-124 in drug addiction has also been reported by overexpression studies wherein lentivirus-miR-124 transduction in the NAc resulted in attenuation of cocaine-induced CPP [66]. Interestingly, reduced levels of miR-124 have also been found in the ventral tegmental area of methamphetamine self-administering rats by microarray analysis [67]. There are also reports demonstrating decreased miR-124 levels in ethanol-withdrawn rats, which was likely attributed to histone acetylation [68]. All these studies allude to the fact that miR-124 downregulation is critical for reward-related behavior changes induced by abused drugs, but detailed mechanisms remain unexplored. Our behavioral findings are also consistent with the published reports and lend further credence to the fact that cocaine-mediated downregulation of microglial miR-124 was involved in increased neuroimmune signaling mediated by TLR4 signaling, leading in turn, to addictive behavior [25,69]. We also acknowledge that BV-2 cells and rat primary microglia may not recapitulate the function of microglia *in vivo*. However, the novelty of our findings are: (1) KLF4, an important pro-inflammatory transcription factor, that was identified in this study as a novel miR-124 target; (2) miR-124 can modulate multiple molecules in TLR4 signaling including TLR4, MyD88, IRAK1, and TRAF6; (3) overexpression of miR-124 can inhibit cocaine-mediated microglial activation *in vitro* and *in vivo*.

In summary, we have demonstrated that miR-124 regulated microglial TLR4 signaling and overexpression of miR-124 inhibited cocaine-mediated microglial activation, and this could have an impact on the ensuing cross-talk between glia and neurons (Fig. 7). Overexpression

of miR-124 has already been shown to prevent microglial activation through inhibition of reactive oxygen species production [70]. Taken together, miR-124 can be an envisioned as a potential target for modulating drug abuse mediated neuroinflammation and for ameliorating drug addiction.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This work was supported by NIH grants: DA043138, DA033150, DA035203.

References

1. Salter MW, Beggs S. Sublime microglia: expanding roles for the guardians of the CNS. *Cell*. 2014; 158(1):15–24. DOI: 10.1016/j.cell.2014.06.008 [PubMed: 24995975]
2. Ransohoff RM, Cardona AE. The myeloid cells of the central nervous system parenchyma. *Nature*. 2010; 468(7321):253–262. DOI: 10.1038/nature09615 [PubMed: 21068834]
3. Heiman A, Pallott A, Heary RF, Elkabes S. Toll-like receptors in central nervous system injury and disease: a focus on the spinal cord. *Brain Behav Immun*. 2014; 42:232–245. DOI: 10.1016/j.bbi.2014.06.203 [PubMed: 25063708]
4. Pedras-Vasconcelos J, Puig M, Verthelyi D. TLRs as therapeutic targets in CNS inflammation and infection. *Front Biosci (Elite Ed)*. 2009; 1:476–487. [PubMed: 19482661]
5. O'Neill LA, Golenbock D, Bowie AG. The history of Toll-like receptors - redefining innate immunity. *Nat Rev Immunol*. 2013; 13(6):453–460. DOI: 10.1038/nri3446 [PubMed: 23681101]
6. Fitzgerald KA, Rowe DC, Barnes BJ, Caffrey DR, Visintin A, Latz E, Monks B, Pitha PM, Golenbock DT. LPS-TLR4 signaling to IRF-3/7 and NF-kappaB involves the toll adapters TRAM and TRIF. *J Exp Med*. 2003; 198(7):1043–1055. DOI: 10.1084/jem.20031023 [PubMed: 14517278]
7. Papageorgiou IE, Lewen A, Galow LV, Cesetti T, Scheffel J, Regen T, Hanisch UK, Kann O. TLR4-activated microglia require IFN-gamma to induce severe neuronal dysfunction and death in situ. *Proc Natl Acad Sci U S A*. 2016; 113(1):212–217. DOI: 10.1073/pnas.1513853113 [PubMed: 26699475]
8. Rivest S. Molecular insights on the cerebral innate immune system. *Brain Behav Immun*. 2003; 17(1):13–19. [PubMed: 12615045]
9. Feinberg MW, Cao Z, Wara AK, Lebedeva MA, Senbanerjee S, Jain MK. Kruppel-like factor 4 is a mediator of proinflammatory signaling in macrophages. *J Biol Chem*. 2005; 280(46):38247–38258. DOI: 10.1074/jbc.M509378200 [PubMed: 16169848]
10. Pena-Altamira E, Prati F, Massenzio F, Virgili M, Contestabile A, Bolognesi ML, Monti B. Changing paradigm to target microglia in neurodegenerative diseases: from anti-inflammatory strategy to active immunomodulation. *Expert Opin Ther Targets*. 2016; 20(5):627–640. DOI: 10.1517/14728222.2016.1121237 [PubMed: 26568363]
11. Walker DG, Lue LF. Immune phenotypes of microglia in human neurodegenerative disease: challenges to detecting microglial polarization in human brains. *Alzheimers Res Ther*. 2015; 7(1): 56.doi: 10.1186/s13195-015-0139-9 [PubMed: 26286145]
12. Houdek HM, Larson J, Watt JA, Rosenberger TA. Bacterial lipopolysaccharide induces a dose-dependent activation of neuroglia and loss of basal forebrain cholinergic cells in the rat brain. *Inflamm Cell Signal*. 2014; 1(1)doi: 10.14800/ics.47
13. Moehle MS, West AB. M1 and M2 immune activation in Parkinson's Disease: Foe and ally? *Neuroscience*. 2015; 302:59–73. DOI: 10.1016/j.neuroscience.2014.11.018 [PubMed: 25463515]
14. Calsolaro V, Edison P. Neuroinflammation in Alzheimer's disease: Current evidence and future directions. *Alzheimers Dement*. 2016; 12(6):719–732. DOI: 10.1016/j.jalz.2016.02.010 [PubMed: 27179961]

15. Brites D, Vaz AR. Microglia centered pathogenesis in ALS: insights in cell interconnectivity. *Front Cell Neurosci.* 2014; 8:117.doi: 10.3389/fncel.2014.00117 [PubMed: 24904276]
16. Lacagnina MJ, Rivera PD, Bilbo SD. Glial and Neuroimmune Mechanisms as Critical Modulators of Drug Use and Abuse. *Neuropsychopharmacology.* 2017; 42(1):156–177. DOI: 10.1038/npp.2016.121 [PubMed: 27402494]
17. Guo ML, Liao K, Periyasamy P, Yang L, Cai Y, Callen SE, Buch S. Cocaine-mediated microglial activation involves the ER stress-autophagy axis. *Autophagy.* 2015; 11(7):995–1009. DOI: 10.1080/15548627.2015.1052205 [PubMed: 26043790]
18. Cui C, Shurtleff D, Harris RA. Neuroimmune mechanisms of alcohol and drug addiction. *Int Rev Neurobiol.* 2014; 118:1–12. DOI: 10.1016/B978-0-12-801284-0.00001-4 [PubMed: 25175859]
19. Liao K, Guo M, Niu F, Yang L, Callen SE, Buch S. Cocaine-mediated induction of microglial activation involves the ER stress-TLR2 axis. *J Neuroinflammation.* 2016; 13(1):33.doi: 10.1186/s12974-016-0501-2 [PubMed: 26860188]
20. Sekine Y, Ouchi Y, Sugihara G, Takei N, Yoshikawa E, Nakamura K, Iwata Y, Tsuchiya KJ, Suda S, Suzuki K, Kawai M, Takebayashi K, Yamamoto S, Matsuzaki H, Ueki T, Mori N, Gold MS, Cadet JL. Methamphetamine causes microglial activation in the brains of human abusers. *J Neurosci.* 2008; 28(22):5756–5761. DOI: 10.1523/JNEUROSCI.1179-08.2008 [PubMed: 18509037]
21. Grace PM, Strand KA, Galer EL, Urban DJ, Wang X, Baratta MV, Fabisiak TJ, Anderson ND, Cheng K, Greene LI, Berkelhammer D, Zhang Y, Ellis AL, Yin HH, Campeau S, Rice KC, Roth BL, Maier SF, Watkins LR. Morphine paradoxically prolongs neuropathic pain in rats by amplifying spinal NLRP3 inflammasome activation. *Proc Natl Acad Sci U S A.* 2016; 113(24):E3441–3450. DOI: 10.1073/pnas.1602070113 [PubMed: 27247388]
22. Crews FT, Sarkar DK, Qin L, Zou J, Boyadjieva N, Vetreno RP. Neuroimmune Function and the Consequences of Alcohol Exposure. *Alcohol Res.* 2015; 37(2):331–341. 344–351. [PubMed: 26695754]
23. Snider SE, Hendrick ES, Beardsley PM. Glial cell modulators attenuate methamphetamine self-administration in the rat. *Eur J Pharmacol.* 2013; 701(1-3):124–130. DOI: 10.1016/j.ejphar.2013.01.016 [PubMed: 23375937]
24. Chen H, Uz T, Manev H. Minocycline affects cocaine sensitization in mice. *Neurosci Lett.* 2009; 452(3):258–261. DOI: 10.1016/j.neulet.2009.01.078 [PubMed: 19348734]
25. Northcutt AL, Hutchinson MR, Wang X, Baratta MV, Hiranita T, Cochran TA, Pomrenze MB, Galer EL, Kopajtic TA, Li CM, Amat J, Larson G, Cooper DC, Huang Y, O'Neill CE, Yin H, Zahniser NR, Katz JL, Rice KC, Maier SF, Bachtell RK, Watkins LR. DAT isn't all that: cocaine reward and reinforcement require Toll-like receptor 4 signaling. *Mol Psychiatry.* 2015; 20(12):1525–1537. DOI: 10.1038/mp.2014.177 [PubMed: 25644383]
26. Carthew RW, Sontheimer EJ. Origins and Mechanisms of miRNAs and siRNAs. *Cell.* 2009; 136(4):642–655. DOI: 10.1016/j.cell.2009.01.035 [PubMed: 19239886]
27. Bartel DP. MicroRNAs: target recognition and regulatory functions. *Cell.* 2009; 136(2):215–233. DOI: 10.1016/j.cell.2009.01.002 [PubMed: 19167326]
28. Hobert O. Gene regulation by transcription factors and microRNAs. *Science.* 2008; 319(5871):1785–1786. DOI: 10.1126/science.1151651 [PubMed: 18369135]
29. Cardoso AL, Guedes JR, de Lima MC. Role of microRNAs in the regulation of innate immune cells under neuroinflammatory conditions. *Curr Opin Pharmacol.* 2016; 26:1–9. DOI: 10.1016/j.coph.2015.09.001 [PubMed: 26410391]
30. Svahn AJ, Giacomotto J, Graeber MB, Rinkwitz S, Becker TS. miR-124 Contributes to the functional maturity of microglia. *Dev Neurobiol.* 2016; 76(5):507–518. DOI: 10.1002/dneu.22328 [PubMed: 26184457]
31. Ponomarev ED, Veremeyko T, Barteneva N, Krichevsky AM, Weiner HL. MicroRNA-124 promotes microglia quiescence and suppresses EAE by deactivating macrophages via the C/EBP- α -PU.1 pathway. *Nat Med.* 2011; 17(1):64–70. DOI: 10.1038/nm.2266 [PubMed: 21131957]
32. Gascon E, Lynch K, Ruan H, Almeida S, Verheyden JM, Seeley WW, Dickson DW, Petrucelli L, Sun D, Jiao J, Zhou H, Jakovcevski M, Akbarian S, Yao WD, Gao FB. Alterations in

- microRNA-124 and AMPA receptors contribute to social behavioral deficits in frontotemporal dementia. *Nat Med*. 2014; 20(12):1444–1451. DOI: 10.1038/nm.3717 [PubMed: 25401692]
33. Kanagaraj N, Beiping H, Dheen ST, Tay SS. Downregulation of miR-124 in MPTP-treated mouse model of Parkinson's disease and MPP iodide-treated MN9D cells modulates the expression of the calpain/cdk5 pathway proteins. *Neuroscience*. 2014; 272:167–179. DOI: 10.1016/j.neuroscience.2014.04.039 [PubMed: 24792712]
 34. Wang H, Ye Y, Zhu Z, Mo L, Lin C, Wang Q, Wang H, Gong X, He X, Lu G, Lu F, Zhang S. MiR-124 Regulates Apoptosis and Autophagy Process in MPTP Model of Parkinson's Disease by Targeting to Bim. *Brain Pathol*. 2016; 26(2):167–176. DOI: 10.1111/bpa.12267 [PubMed: 25976060]
 35. Veremeyko T, Siddiqui S, Sotnikov I, Yung A, Ponomarev ED. IL-4/IL-13-dependent and independent expression of miR-124 and its contribution to M2 phenotype of monocytic cells in normal conditions and during allergic inflammation. *PLoS One*. 2013; 8(12):e81774.doi: 10.1371/journal.pone.0081774 [PubMed: 24358127]
 36. Ni M, Aschner M. Neonatal rat primary microglia: isolation, culturing, and selected applications. *Curr Protoc Toxicol*. 2010; Chapter 12:Unit 12 17. doi: 10.1002/0471140856.tx1217s43
 37. Black AR, Black JD, Azizkhan-Clifford J. Sp1 and kruppel-like factor family of transcription factors in cell growth regulation and cancer. *J Cell Physiol*. 2001; 188(2):143–160. DOI: 10.1002/jcp.1111 [PubMed: 11424081]
 38. Kaushik DK, Thounaojam MC, Kumawat KL, Gupta M, Basu A. Interleukin-1beta orchestrates underlying inflammatory responses in microglia via Kruppel-like factor 4. *J Neurochem*. 2013; 127(2):233–244. DOI: 10.1111/jnc.12382 [PubMed: 23895397]
 39. Kaushik DK, Gupta M, Das S, Basu A. Kruppel-like factor 4, a novel transcription factor regulates microglial activation and subsequent neuroinflammation. *J Neuroinflammation*. 2010; 7:68.doi: 10.1186/1742-2094-7-68 [PubMed: 20946687]
 40. Kaushik DK, Mukhopadhyay R, Kumawat KL, Gupta M, Basu A. Therapeutic targeting of Kruppel-like factor 4 abrogates microglial activation. *J Neuroinflammation*. 2012; 9:57.doi: 10.1186/1742-2094-9-57 [PubMed: 22429472]
 41. Rosenzweig JM, Glenn JD, Calabresi PA, Whartenby KA. KLF4 modulates expression of IL-6 in dendritic cells via both promoter activation and epigenetic modification. *J Biol Chem*. 2013; 288(33):23868–23874. DOI: 10.1074/jbc.M113.479576 [PubMed: 23846700]
 42. Van Dyke C, Barash PG, Jatlow P, Byck R. Cocaine: plasma concentrations after intranasal application in man. *Science*. 1976; 191(4229):859–861. [PubMed: 56036]
 43. Stephens BG, Jentzen JM, Karch S, Mash DC, Wetli CV. Criteria for the interpretation of cocaine levels in human biological samples and their relation to the cause of death. *Am J Forensic Med Pathol*. 2004; 25(1):1–10. [PubMed: 15075680]
 44. Kalasinsky KS, Bosy TZ, Schmunk GA, Ang L, Adams V, Gore SB, Smialek J, Furukawa Y, Guttman M, Kish SJ. Regional distribution of cocaine in postmortem brain of chronic human cocaine users. *J Forensic Sci*. 2000; 45(5):1041–1048. [PubMed: 11005179]
 45. Karthikeyan A, Patnala R, Jadhav SP, Eng-Ang L, Dheen ST. MicroRNAs: Key Players in Microglia and Astrocyte Mediated Inflammation in CNS Pathologies. *Curr Med Chem*. 2016; 23(30):3528–3546. [PubMed: 27528056]
 46. Thome AD, Harms AS, Volpicelli-Daley LA, Standaert DG. microRNA-155 Regulates Alpha-Synuclein-Induced Inflammatory Responses in Models of Parkinson Disease. *J Neurosci*. 2016; 36(8):2383–2390. DOI: 10.1523/JNEUROSCI.3900-15.2016 [PubMed: 26911687]
 47. Brites D, Fernandes A. Neuroinflammation and Depression: Microglia Activation, Extracellular Microvesicles and microRNA Dysregulation. *Front Cell Neurosci*. 2015; 9:476.doi: 10.3389/fncel.2015.00476 [PubMed: 26733805]
 48. Guo ML, Periyasamy P, Liao K, Kook YH, Niu F, Callen SE, Buch S. Cocaine-mediated downregulation of microglial miR-124 expression involves promoter DNA methylation. *Epigenetics*. 2016; 11(11):819–830. DOI: 10.1080/15592294.2016.1232233 [PubMed: 27786595]
 49. Lewitus GM, Konefal SC, Greenhalgh AD, Pribrag H, Augereau K, Stellwagen D. Microglial TNF-alpha Suppresses Cocaine-Induced Plasticity and Behavioral Sensitization. *Neuron*. 2016; 90(3):483–491. DOI: 10.1016/j.neuron.2016.03.030 [PubMed: 27112496]

50. Ma C, Li Y, Li M, Deng G, Wu X, Zeng J, Hao X, Wang X, Liu J, Cho WC, Liu X, Wang Y. microRNA-124 negatively regulates TLR signaling in alveolar macrophages in response to mycobacterial infection. *Mol Immunol*. 2014; 62(1):150–158. DOI: 10.1016/j.molimm.2014.06.014 [PubMed: 24995397]
51. Ma C, Li Y, Zeng J, Wu X, Liu X, Wang Y. Mycobacterium bovis BCG triggered MyD88 induces miR-124 feedback negatively regulates immune response in alveolar epithelial cells. *PLoS One*. 2014; 9(4):e92419.doi: 10.1371/journal.pone.0092419 [PubMed: 24705038]
52. Sun Y, Li Q, Gui H, Xu DP, Yang YL, Su DF, Liu X. MicroRNA-124 mediates the cholinergic anti-inflammatory action through inhibiting the production of pro-inflammatory cytokines. *Cell Res*. 2013; 23(11):1270–1283. DOI: 10.1038/cr.2013.116 [PubMed: 23979021]
53. Yao H, Yang Y, Kim KJ, Bethel-Brown C, Gong N, Funa K, Gendelman HE, Su TP, Wang JQ, Buch S. Molecular mechanisms involving sigma receptor-mediated induction of MCP-1: implication for increased monocyte transmigration. *Blood*. 2010; 115(23):4951–4962. DOI: 10.1182/blood-2010-01-266221 [PubMed: 20354174]
54. Asanuma M, Miyazaki I, Higashi Y, Tsuji T, Ogawa N. Specific gene expression and possible involvement of inflammation in methamphetamine-induced neurotoxicity. *Ann N Y Acad Sci*. 2004; 1025:69–75. DOI: 10.1196/annals.1316.009 [PubMed: 15542702]
55. Goncalves J, Martins T, Ferreira R, Milhazes N, Borges F, Ribeiro CF, Malva JO, Macedo TR, Silva AP. Methamphetamine-induced early increase of IL-6 and TNF-alpha mRNA expression in the mouse brain. *Ann N Y Acad Sci*. 2008; 1139:103–111. DOI: 10.1196/annals.1432.043 [PubMed: 18991854]
56. Wisor JP, Schmidt MA, Clegern WC. Cerebral microglia mediate sleep/wake and neuroinflammatory effects of methamphetamine. *Brain Behav Immun*. 2011; 25(4):767–776. DOI: 10.1016/j.bbi.2011.02.002 [PubMed: 21333736]
57. Piechota M, Korostynski M, Solecki W, Gieryk A, Slezak M, Bilecki W, Ziolkowska B, Kostrzewa E, Cymerman I, Swiech L, Jaworski J, Przewlocki R. The dissection of transcriptional modules regulated by various drugs of abuse in the mouse striatum. *Genome Biol*. 2010; 11(5):R48.doi: 10.1186/gb-2010-11-5-r48 [PubMed: 20459597]
58. Renthal W, Maze I, Krishnan V, Covington HE 3rd, Xiao G, Kumar A, Russo SJ, Graham A, Tsankova N, Kippin TE, Kerstetter KA, Neve RL, Haggarty SJ, McKinsey TA, Bassel-Duby R, Olson EN, Nestler EJ. Histone deacetylase 5 epigenetically controls behavioral adaptations to chronic emotional stimuli. *Neuron*. 2007; 56(3):517–529. DOI: 10.1016/j.neuron.2007.09.032 [PubMed: 17988634]
59. Ahmed SH, Lutjens R, van der Stap LD, Lekic D, Romano-Spica V, Morales M, Koob GF, Repunte-Canonigo V, Sanna PP. Gene expression evidence for remodeling of lateral hypothalamic circuitry in cocaine addiction. *Proc Natl Acad Sci U S A*. 2005; 102(32):11533–11538. DOI: 10.1073/pnas.0504438102 [PubMed: 16076954]
60. Little KY, Ramssen E, Welchko R, Volberg V, Roland CJ, Cassin B. Decreased brain dopamine cell numbers in human cocaine users. *Psychiatry Res*. 2009; 168(3):173–180. DOI: 10.1016/j.psychres.2008.10.034 [PubMed: 19233481]
61. Loftis JM, Choi D, Hoffman W, Huckans MS. Methamphetamine causes persistent immune dysregulation: a cross-species, translational report. *Neurotox Res*. 2011; 20(1):59–68. DOI: 10.1007/s12640-010-9223-x [PubMed: 20953917]
62. Beardsley PM, Hauser KF. Glial modulators as potential treatments of psychostimulant abuse. *Adv Pharmacol*. 2014; 69:1–69. DOI: 10.1016/B978-0-12-420118-7.00001-9 [PubMed: 24484974]
63. Deo M, Yu JY, Chung KH, Tippens M, Turner DL. Detection of mammalian microRNA expression by in situ hybridization with RNA oligonucleotides. *Dev Dyn*. 2006; 235(9):2538–2548. DOI: 10.1002/dvdy.20847 [PubMed: 16736490]
64. Krichevsky AM, Sonntag KC, Isacson O, Kosik KS. Specific microRNAs modulate embryonic stem cell-derived neurogenesis. *Stem Cells*. 2006; 24(4):857–864. DOI: 10.1634/stemcells.2005-0441 [PubMed: 16357340]
65. Chandrasekar V, Dreyer JL. microRNAs miR-124, let-7d and miR-181a regulate cocaine-induced plasticity. *Mol Cell Neurosci*. 2009; 42(4):350–362. DOI: 10.1016/j.mcn.2009.08.009 [PubMed: 19703567]

66. Chandrasekar V, Dreyer JL. Regulation of MiR-124, Let-7d, and MiR-181a in the accumbens affects the expression, extinction, and reinstatement of cocaine-induced conditioned place preference. *Neuropsychopharmacology*. 2011; 36(6):1149–1164. DOI: 10.1038/npp.2010.250 [PubMed: 21307844]
67. Bosch PJ, Benton MC, Macartney-Coxson D, Kivell BM. mRNA and microRNA analysis reveals modulation of biochemical pathways related to addiction in the ventral tegmental area of methamphetamine self-administering rats. *BMC Neurosci*. 2015; 16:43.doi: 10.1186/s12868-015-0186-y [PubMed: 26188473]
68. Mizuo K, Katada R, Okazaki S, Tateda K, Watanabe S, Matsumoto H. Epigenetic regulation of MIR-124 under ethanol dependence and withdrawal. *Nihon Arukoru Yakubutsu Igakkai Zasshi*. 2012; 47(3):155–163. [PubMed: 22894056]
69. Bachtell R, Hutchinson MR, Wang X, Rice KC, Maier SF, Watkins LR. Targeting the Toll of Drug Abuse: The Translational Potential of Toll-Like Receptor 4. *CNS Neurol Disord Drug Targets*. 2015; 14(6):692–699. [PubMed: 26022268]
70. Louw AM, Kolar MK, Novikova LN, Kingham PJ, Wiberg M, Kjems J, Novikov LN. Chitosan polyplex mediated delivery of miRNA-124 reduces activation of microglial cells in vitro and in rat models of spinal cord injury. *Nanomedicine*. 2016; 12(3):643–653. DOI: 10.1016/j.nano.2015.10.011 [PubMed: 26582736]

Figure 1.

Cocaine increased KLF4 levels *in vitro* and *in vivo*. Representative western blots showing the cocaine-mediated dose-dependent upregulation of KLF4 levels in both the BV-2 cells (A) and the rPMs (B). Representative western blots showing the cocaine-mediated time-dependent upregulation of KLF4 levels in both the BV-2 cells (C) and the rPMs (D). (E) Representative western blots showing the increased KLF4 expression levels in the striatal brain region of mice chronically administered with cocaine (20 mg/kg, *i.p.*) for 7 days. β -actin was used as a protein loading control for all the experiments. The data are presented as

mean $\pm$ SEM from 6 independent experiments. One-way ANOVA followed by Bonferroni *post hoc* test was used to determine the statistical significance. *, $P < 0.05$ vs control.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Figure 2.

KLF4 is a novel target regulated by miR-124. (A) Putative miR-124 binding sites in *KLF4* gene. The potential complementary residues are shown in red color. (B) Representative western blots showing the overexpression of miR-124 significantly blocked the upregulation of KLF4 in BV-2 cells transfected with miR-124 mimics/control followed with cocaine (10 μ M) exposure for 24 h. (C) Representative western blots showing the overexpression of miR-124 significantly blocked the upregulation of KLF4 in rPMs transfected with miR-124 mimics/control followed with cocaine (10 μ M) exposure for 24 h. (D) Representative western blots showing the miR-124 inhibitor significantly increased the KLF4 in BV-2 cells transfected with miR-124 inhibitor/control followed with cocaine (10 μ M) exposure for 24 h. (E) Representative western blots showing the miR-124 inhibitor significantly increased the KLF4 in rPMs transfected with miR-124 inhibitor/control followed with cocaine (10 μ M) exposure for 24 h. (F) Schematic diagram of KLF4-3'-UTR-1 and KLF4-3'-UTR-2 plasmids. KLF4-3'-UTR-1 plasmid has no miR-124 binding sites while KLF4-3'-UTR-2 plasmid has three miR-124 binding sites. (G) Relative luciferase activity of 3'-UTR-1 and 3'-UTR-2 constructs of KLF4 co-transfected with miR-control, miR-124 mimics, and miR-124 inhibitor in HEK293 cells. Overexpression of miR-124 mimics/inhibitor had no effect on the luciferase activity of KLF4-3'-UTR-1 plasmids while miR-124 mimics significantly inhibited KLF4-3'-UTR-2 luciferase activity. (H) Ago2 IP confirmed the enrichment of the miR-124 target mRNA, KLF4 in Ago2 IP as compared with total RNA isolated from BV-2 cells transfected with miR-124 mimics/control. β -actin was used as a loading control for all the experiments. The data are presented as mean $\pm$ SEM from 6

independent experiments. One-way ANOVA followed by Bonferroni *post hoc* test was used to determine the statistical significance for multiple groups, and a Student's t test was used to determine the statistical significance for between 2 groups. *, $P < 0.05$ vs. control; #, $P < 0.05$ vs. cocaine.

Figure 3.

Overexpression of miR-124 inhibited the cocaine-mediated TLR4 upregulation. (A) Representative western blots showing the cocaine-mediated dose-dependent upregulation of TLR4 levels in BV-2 cells. (B) Representative western blots showing the cocaine-mediated time-dependent upregulation of TLR4 levels in BV-2 cells (C) Representative western blots showing the increased TLR4 expression levels in the striatal brain region of mice chronically administered with cocaine (20 mg/kg, i.p.) for 7 days. (D) Putative miR-124 binding sites in *TLR4* gene. The potential complementary residues are shown in red color. (E)

Representative western blots showing the overexpression of miR-124 significantly blocked the upregulation of TLR4 in BV-2 cells transfected with miR-124 mimics/control followed with cocaine (10 μ M) exposure for 24 h. (F) Representative western blots showing the miR-124 inhibitors significantly increased the TLR4 in rPMs transfected with miR-124 inhibitor/control for 24 h. (G) Representative western blots showing the overexpression of miR-124 significantly blocked the upregulation of TLR4 in rPMs transfected with miR-124 mimics while inhibition of miR-124 increased the TLR4 upregulation in rPMs transfected with the miR-124 inhibitor for 24 h. (H) Ago2 IP confirmed the enrichment of the miR-124 target mRNA, TLR4 in Ago2 IP as compared with total RNA isolated from BV-2 cells transfected with miR-124 mimics/control. β -actin was used as a loading control for all the experiments. The data are presented as mean $\pm$ SEM from 6 independent experiments. One-way ANOVA followed by Bonferroni *post hoc* test was used to determine the statistical significance for multiple groups and a Student's t test was used to determine the statistical significance for between 2 groups. *, $P < 0.05$ vs. control; #, $P < 0.05$ vs. cocaine.

Figure 4.

Overexpression of miR-124 blocked the cocaine-mediated upregulation of MyD88, IRAK1, and TRAF6. (A) Putative miR-124 binding sites in *MyD88*, *IRAK1*, and *TRAF6* genes. The potential complementary residues are shown in red color. Representative western blots showing the overexpression of miR-124 significantly blocked the upregulation of MyD88 (B), IRAK1 (D), and TRAF6 (F) in BV-2 cells transfected with miR-124 mimics/control followed with cocaine (10 μ M) exposure for 24 h. Representative western blots showing the inhibition of miR-124 significantly increased the upregulation of MyD88 (C), IRAK1 (E),

and TRAF6 (G) in BV-2 cells transfected with miR-124 inhibitor/control followed with cocaine (10 μ M) exposure for 24 h. (H, I) Representative western blots showing the overexpression of miR-124 significantly blocked the upregulation of MyD88, IRAK1, and TRAF6 in rPMs transfected with miR-124 mimics while inhibition of miR-124 increased the upregulation of MyD88, IRAK1, and TRAF6 in rPMs transfected with the miR-124 inhibitor for 24 h. (J) Ago2 IP confirmed the enrichment of the miR-124 target mRNAs, MyD88, IRAK1, and TRAF6 in Ago2 IP as compared with total RNA isolated from BV-2 cells transfected with miR-124 mimics/control. β -actin was used as a loading control for all the experiments. The data are presented as mean $\pm$ SEM from 6 independent experiments. One-way ANOVA followed by Bonferroni post hoc test was used to determine the statistical significance for multiple groups, and a Student's t test was used to determine the statistical significance for between 2 groups. *, $P < 0.05$ vs. control; #, $P < 0.05$ vs. cocaine.

Figure 5.

Overexpression of miR-124 inhibited cocaine-mediated microglial activation *in vitro*. qPCR analysis showing the overexpression of miR-124 significantly inhibited the increased mRNA expressions of various proinflammatory mediators such as TNF α (A), CCL2 (B), and NOS2 (C), in BV-2 cells transfected with miR-124 mimics/control followed with cocaine (10 μ M) exposure for 24 h. In contrast, qPCR analysis showing the overexpression of miR-124 significantly increased mRNA expressions of various anti-inflammatory mediators such as TGF β (D), IL4 (E), and IL10 (F), in BV-2 cells transfected with miR-124 mimics/control followed with cocaine (10 μ M) exposure for 24 h. (G and H) qPCR analysis showing the overexpression of miR-124 significantly inhibited the mRNA expressions of c/EBP and ARG1 in BV-2 cells transfected with miR-124 mimics/control followed with cocaine (10 μ M) exposure for 24 h. β -actin was used as an internal housekeeping gene for all the experiments. The data are presented as mean $\pm$ SEM from 6 independent experiments. One-way ANOVA followed by Bonferroni post hoc test was used to determine the statistical significance for multiple groups, and a Student's t test was used to determine the statistical significance for between 2 groups. *, $P < 0.05$ vs. control; #, $P < 0.05$ vs. cocaine.

Figure 6.

Overexpression of miR-124 inhibited cocaine-mediated microglial activation *in vivo*. (A) Schematic showing the stereotactic injection of lentivirus (LV)-miR-124 followed by chronic cocaine administration (20 mg/kg; *i.p.* for 7 days) in mice. (B) Representative fluorescent-microscopic image showing the overexpression of miR-124 colocalized with microglia in the striatum region of mice stereotactically injected with LV-miR-124. (C) qPCR analysis showing the expression levels of miR-124 in the striatal region of mice stereotactically injected with LV-miR-124 followed by cocaine administration. (D-G) Representative western blots and quantifications showing the expression levels of KLF4, TLR4, and Iba-1 in the striatal region of mice stereotactically injected with LV-miR-124 followed by cocaine administration. β -actin was used as a protein loading control for all the experiments. The data are presented as mean $\pm$ SEM from 8 independent experiments. Two-way ANOVA followed by Tukey post hoc test was used to determine the statistical significance for multiple groups, and a Student's *t* test was used to determine the statistical significance for between 2 groups. *, $P < 0.05$ vs. saline; #, $P < 0.05$ vs. cocaine.

Figure 7.
Schematic diagram representing the miR-124-mediated regulation of TLR4 signaling in the context of cocaine.