Published in final edited form as:

J Acquir Immune Defic Syndr. 2017 December 15; 76(5): 493–500. doi:10.1097/QAI.0000000000001549.

Acute Alcohol Consumption Directly Increases HIV Transmission Risk: A Randomized Controlled Experiment

Paul A. Shuper, Ph.D.^{1,2}, Narges Joharchi, M.Sc.¹, Peter M. Monti, Ph.D.³, Mona Loutfy, MD, FRCPC, MPH^{4,5,6}, and Jürgen Rehm, Ph.D.^{1,2,7,8,9,10,11,12}

¹Institute for Mental Health Policy Research, Centre for Addiction and Mental Health, Toronto, Canada

²Dalla Lana School of Public Health, University of Toronto, Toronto, Canada

³Brown University Center for Alcohol and Addiction Studies and Department of Behavioral and Social Sciences; Brown University, Providence, RI, USA

⁴Women's College Hospital, Toronto, Canada

⁵Department of Medicine, University of Toronto, Toronto, Canada

⁶Maple Leaf Medical Clinic, Toronto, Canada

⁷Campbell Family Mental Health Research Institute, Centre for Addiction and Mental Health, Toronto, Canada

⁸Department of Psychiatry, Faculty of Medicine, University of Toronto, Toronto, Canada

⁹PAHO/WHO Collaborating Centre for Mental Health & Addiction, Centre for Addiction and Mental Health, Toronto, Canada

¹⁰Epidemiological Research Unit, Technische Universität Dresden, Dresden, Germany

¹¹Klinische Psychologie & Psychotherapie, Dresden, Germany

¹²Graduate Department of Community Health and Institute of Medical Science, University of Toronto, Toronto, Canada

Abstract

Background—Alcohol consumption has frequently been purported as a driver of condomless sex and HIV transmission, but to date, experimental evidence for the causal risk-taking impact of alcohol among HIV-positive populations is lacking. The present experiment sought to determine whether acute alcohol consumption has a direct causal impact on condomless sex intentions among HIV-positive men-who-have-sex-with-men (MSM), and to assess whether alcohol's impact differs between MSM who are HIV-positive versus HIV-negative.

Methods—In a randomized controlled alcohol administration experiment, HIV-positive and HIV-negative MSM were brought into a specialized barroom laboratory and randomly assigned to

Corresponding Author: Paul A. Shuper, Ph.D., Centre for Addiction and Mental Health, 33 Russell Street, Toronto, ON, M5S 2S1, Canada, Phone: 416-535-8501, x34097; paul.shuper@camh.ca.

Conflict of Interest: The authors declare no conflicts of interest.

beverage consumption condition: alcohol (target BAC=.080%), placebo alcohol (target BAC=.000%), or water (control). Participants then underwent a video-based sexual arousal manipulation (sexually aroused/non-aroused) and indicated their intentions to engage in condom-protected and condomless sexual acts in a standardized paradigm. The primary outcome entailed intentions to engage in condomless receptive and condomless insertive anal sex.

Results—A total of 282 MSM (141 HIV-positive; 141 HIV-negative) completed experimental procedures. MSM who received alcohol reported significantly stronger intentions to engage in condomless sex than those who received placebo alcohol or water ($F(1,274)=9.43$, $p=0.002$). The impact of alcohol did not differ between HIV-positive and HIV-negative MSM ($F(1,274)=1.86$, $p=0.174$).

Conclusions—The present investigation entailed the first risk-focused alcohol administration experiment to involve an HIV-positive sample, and results demonstrated that consuming alcohol had an independent, causal impact on intentions to engage in sexual behaviors that can result in HIV transmission. Findings strongly suggest that alcohol-focused initiatives should be incorporated into HIV prevention efforts.

Keywords

HIV/AIDS; Alcohol; Men-Who-Have-Sex-With-Men (MSM); Risk; Transmission

Introduction

Despite a recent modest decline in global HIV incidence,¹ HIV infections have been increasing among populations of men-who-have-sex-with-men (MSM) in a number of regions.²⁻⁴ Predominantly driving this persistent epidemic is the continued engagement in condomless sex between HIV-positive MSM and their non-infected counterparts.⁵ The majority of HIV-positive MSM remain sexually active after HIV diagnosis,^{6;7} and approximately 40% report engaging in condomless sex.^{8;9} While newly promoted biomedical HIV prevention strategies, entailing pre-exposure prophylaxis (PrEP)¹⁰ and “treatment-as-prevention” (TasP),¹¹ can substantially decrease the likelihood of sexually-based HIV acquisition and transmission, even when condoms are not used,¹⁰⁻¹⁵ rates of PrEP uptake among MSM are low,^{16;17} and marked gaps in the HIV treatment cascade remain, such that only an estimated 16-34% of HIV-positive MSM in the US have attained an undetectable HIV viral load.¹⁸ As a result, a sizable number of HIV-negative and HIV-positive MSM remain “biologically unprotected” from acquiring or transmitting HIV, respectively. Condoms therefore maintain a role in HIV prevention, and efforts to increase condom use among HIV-positive MSM could help reduce HIV transmission.¹⁹

Condoms are not without their challenges, however, and among the many factors purported to hinder condom use, alcohol consumption has received particular attention.²⁰⁻²⁶ It has been theorized that consuming alcohol limits cognitive processing capabilities; causing individuals to attend only to salient risk-impelling cues (e.g., feeling sexually aroused) while disregarding risk-inhibiting information (e.g., possible HIV transmission), which in turn increases the likelihood of engaging in sex without condoms.²⁷ Although results from meta-analyses and reviews are consistent with this theory,^{28;29} the evidence has largely been based

on cross-sectional studies, and findings from event-level analyses have been mixed.³⁰⁻³² Questions therefore remain as to the causal nature of the alcohol-condomless sex association. Causality is further questioned by research suggesting that alcohol in and of itself may not have a direct causal impact on condomless sex. Rather, the association may be an indirect one; stemming from risk-relevant traits that underlie both the consumption of alcohol at heavy levels and the engagement in sex without condoms, such as sexual sensation seeking,³³⁻³⁶ sexual compulsivity,³⁷⁻⁴⁰ and sex-related alcohol expectancies.^{21;23;41-46}

More recently, a small number of experiments entailing controlled alcohol administration in conjunction with assessments of participants' condom use intentions have yielded some evidence of a causal link between alcohol use and condomless sex.^{47;48} For example, in their alcohol administration experiment involving HIV-negative MSM, Maisto et al.⁴⁹ found that alcohol was linked to determinants of sexual risk behavior, and that the impact of alcohol was enhanced under conditions of sexual arousal. Importantly, however, none of these experiments involved HIV-positive individuals. This limitation is critical because condom use motivations may differ considerably between HIV-positive and HIV-negative populations, with the former potentially driven to protect their partners from HIV, versus the latter who may be driven to protect themselves.^{34;50;51} These "other" vs "self" motivations may be disparate in salience and strength, and as such, they may be differentially impactful while under the influence of alcohol. Similarly, HIV-positive individuals may possess riskier underlying traits and behavioral profiles than HIV-negative individuals,⁵² which may result in an augmented proclivity toward engaging in condomless sex when consuming alcohol. Finally, recognizing that every new case of HIV derives from someone who is already infected with the virus, it remains crucial to identify the specific role of alcohol in HIV transmission risk behavior among those living with HIV.

Objectives

The present experiment tested the hypothesis that acute alcohol consumption would directly increase HIV-positive MSM's intentions to engage in condomless sex. The experiment also assessed whether the impact of acute alcohol consumption on condomless sex intentions differed between HIV-positive and HIV-negative MSM. Finally, the experiment assessed the potential moderating role of risk-relevant personality traits and expectancies in the alcohol-condomless sex association.

Methods

Study Design

A randomized controlled experiment with blinding was conducted in which HIV-positive and HIV-negative MSM were 1) randomly assigned to receive a body weight-specified dose of alcohol (target blood alcohol concentration (BAC)=.080%), placebo alcohol (target BAC=.000%), or water (control); 2) randomly presented with video clips designed to induce either a moderate level of sexual arousal or no sexual arousal; and 3) asked to report their intentions to engage in condom-protected and condomless sexual acts with a series of hypothetical partners. Intentions to engage in condomless anal sex comprised the primary

study outcome, as 1) such intention measures have been shown to be significantly associated with one's engagement in actual sexual risk behavior;^{53;54} and 2) measuring real-world engagement in condomless sex following alcohol administration would be ethically and logistically unfeasible. Details pertaining to the study protocol have been published.⁵⁵ Procedures were approved by the Research Ethics Board at the Centre for Addiction and Mental Health in Toronto, Canada (Protocol# 034/2010-04).

Study Population

From February 2012 to March 2015, HIV-positive and HIV-negative MSM were recruited from a clinic in Toronto, Canada providing specialized care for HIV-positive patients and MSM. Eligibility criteria included 1) 19+ years of age (legal drinking age in the jurisdiction); 2) anal sex with a man during the past 6 months; 3) social drinker (i.e., consumed 5 drinks/week on average and 5 drinks in one episode during the past six months);⁵⁶ 4) no recent history (i.e., past 5 years) of problematic alcohol or substance use; and 5) no medical contraindication for consuming alcohol to a BAC of approximately .100%.

Experimental Procedures

The experiment was conducted at an addictions and mental health hospital in Toronto, Canada. Participants first completed an electronic self-administered questionnaire assessing demographics, alcohol,⁵⁷ substance use,⁵⁸ sexual history,⁵⁹ underlying risk-relevant traits (i.e., sexual compulsivity,⁶⁰ sexual sensation seeking³⁹), and sex-related alcohol expectancies.⁶¹ Participants were then taken into a specialized barroom laboratory where they were randomly assigned to one of three beverage consumption conditions – alcohol, placebo alcohol, or water (control). Alcohol condition participants received 0.7g alcohol/kg body weight through beverages formulated using a 1:3 vodka:tonic water ratio; with the total volume of fluid divided equally across three cups (target peak BAC=.080%). For placebo alcohol condition participants, beverages were based on a 1:3 mixture of flat tonic water (poured from a vodka bottle):regular tonic water, with a small amount of vodka floated on top of each cup as well as vodka rubbed around the cups' rims (target BAC=.000%). Control condition participants were provided with a volume of water that matched the total beverage volume they would have received in alcohol and placebo conditions. All beverages were consumed over a 15-minute period, followed by a 13-minute absorption period (see⁵⁵ for additional details).

Following beverage procedures, participants proceeded through a touchscreen-based study-designed program, which commenced by randomly presenting two video clips containing either sexual content (arousal condition) or no sexual content (no arousal condition); intended to induce a moderate level of sexual arousal or no sexual arousal, respectively. The program then presented 18 hypothetical “first time” sexual partner profiles in random order, each of which included 1) a photograph of a male partner who was either physically attractive or unattractive; and 2) text describing the partner's HIV-serostatus (HIV-positive/HIV-negative/HIV status unknown) and condom use preference (use/don't use/ not stated).^{55;62} For each partner, participants indicated their intentions to engage in condom-protected and condomless sexual acts (see “Primary Outcome” below).

After completing the program, alcohol condition participants remained onsite until a BAC of <.040% was achieved. HIV-positive participants completed additional questions assessing adherence to antiretroviral therapy (ART),^{63;64} and following the study, clinic medical charts were reviewed to extract HIV-pertinent data (e.g., HIV viral load). All participants received \$50 for taking part, and those in the alcohol condition received an additional \$15 per hour for the detoxification period.

Primary Outcome

For each hypothetical partner, participants were asked “Which of the following would you consider doing with this partner?” This was followed by six items, each referring to a distinct condom-protected or condomless sexual act, and accompanied by a 5-point likert scale, ranging from 1=definitely to 5=definitely not. Scores from two items, identifying intentions to engage in condomless receptive anal sex (“Anal sex – HIS penis in YOUR butt – no condom”), and condomless insertive anal sex (“Anal sex – YOUR penis in HIS butt – no condom”), were extracted and reverse coded, where higher scores reflected stronger intentions to engage in the specified condomless sexual act. The mean score of these two items for each hypothetical partner comprised the primary outcome measure.

Sample Size

A priori established sample size calculations were conducted to assure that the sample size was associated with sufficient power (>80%) to 1) test for the difference in condomless sex intentions between the alcohol condition and the two other beverage conditions; and 2) compare the effect of alcohol on condomless sex intentions between HIV-positive and HIV-negative MSM. Full details pertaining to sample size calculations are provided in the eMethods of the Supplemental Digital Content.

Randomization

Randomization sequences were generated through www.random.org.⁶⁵ Randomization to beverage consumption condition was based on a 50%-25%-25% distribution to alcohol, placebo alcohol, and water conditions, respectively. Randomization to sexual arousal condition was based on a 50%-50% distribution. Additional information regarding randomization procedures can be found in the eMethods of the Supplemental Digital Content.

Blinding

A two-Research Assistant (RA) paradigm was employed for the purposes of experimenter blinding, and participant blinding was facilitated through the inclusion of the placebo alcohol condition. Blinding procedures are described in detail in the eMethods of the Supplemental Digital Content.

Statistical Analyses

The primary analysis entailed a general linear model (GLM) based on a repeated measures full-factorial analysis of variance (ANOVA) design. Beverage consumption conditions were dichotomized into a “drink condition” factor in accordance with a priori-determined analytic

procedures, such that participants in placebo and control conditions were combined into a “non-alcohol” condition and compared to those who received alcohol (note that as anticipated, no significant differences between placebo and water conditions with respect to outcome were yielded). Drink condition (alcohol vs. non-alcohol), participant HIV serostatus (HIV-positive vs. HIV-negative), and sexual arousal condition (arousal vs. non-arousal) were entered into the full-factorial GLM as between-subjects factors, whereas partner characteristics (i.e., attractiveness, HIV-serostatus, and condom use preference) were entered as within-subjects factors.

In accordance with these analytic procedures, the main effect for drink condition was examined in the context of main effects and interactions among between- and within-subjects factors to determine whether acute alcohol consumption had a causal, independent impact on condomless sex intentions. The drink condition $\times$ participant HIV serostatus interaction term was examined to determine whether the impact of acute alcohol consumption differed between HIV-positive and HIV-negative MSM. Finally, to identify whether the link between acute alcohol consumption and condomless sex intentions was attributable to underlying risk-relevant traits and cognitive expectancies, the above-mentioned GLM was run three additional times, in which one potential moderating factor (i.e., sexual sensation seeking, sexual compulsivity, sex-related alcohol expectancies) was included as a covariate and as an interaction term with drink condition.

Results

Screening

As depicted in Figure 1, 585 MSM (264 HIV-positive, 321 HIV-negative) completed screening procedures, among whom 302 were excluded prior to randomization. Among the 302 excluded, 214 were ineligible as a result of 1) not being a social drinker ($n=60$); 2) reporting no anal sex in the past six months ($n=57$); 3) reporting concern over alcohol use ($n=4$); and 4) failing to meet two or more eligibility criteria ($n=93$). Clinic physicians additionally denied medical clearance for 20 MSM. Of the remaining 351 eligible MSM (179 HIV-positive, 172 HIV-negative), 68 did not present for a study session (e.g., due to scheduling difficulties), resulting in a sample of 283 randomized participants (142 HIV-positive, 141 HIV-negative). One participant randomized to the alcohol condition completed the survey but not the experimental procedures due to a mild allergic reaction; yielding a final sample of 282 MSM (141 HIV-positive, 141 HIV-negative) for analysis.

Sample Characteristics

Sociodemographic and other relevant characteristics are presented in Table 1. As shown in the table, participants' age ranged from 20–69 ($M=42.6$), and the majority were white (72.2%), employed (67.9%), and educated at or above the college diploma level (71.5%). Participants reported consuming an average of 9.9 drinks/week (Canadian drink size=13.7g alcohol), and mean AUDIT score was 7.8. Condomless sex during the past six months was reported by 73.2% of participants, serodiscordant condomless sex was reported by 36.7%, and roughly three quarters (76.4%) of the sample indicated multiple sexual partnerships.

Approximately 9 out of 10 (88.6%) participants indicated that they had consumed alcohol before or during sex in the past six months.

Among HIV-positive participants, 82.1% had an undetectable viral load, and mean CD4 cell count was 601.6. The majority (85.1%) were currently receiving ART, among whom roughly half (48.3%) reported <100% adherence during the past month.

Efficacy of Experimental Manipulations

All experimental manipulations worked as anticipated. For example, mean peak BAC was .086% ($SD=.012$) for alcohol condition participants. A full summary of manipulation check data is presented in eTable 1 of the Supplemental Digital Content.

Primary Outcome Analysis

As indicated in Table 2, consistent with our main hypothesis, participants in the alcohol condition reported significantly stronger intentions to engage in condomless sex than non-alcohol condition participants. Furthermore, although HIV-positive participants indicated significantly stronger condomless sex intentions than HIV-negative participants, the interaction between drink condition and participant HIV serostatus was not significant, indicating that the effect of alcohol on condomless sex intentions was similar for HIV-positive and HIV-negative participants. No significant two-way interactions were found between alcohol condition and sexual arousal, partner HIV serostatus, partner condom use preference, and partner attractiveness (all p 's $>.05$).

Results from moderator analyses are presented in Table 3. In all three models, alcohol condition participants reported significantly stronger intentions to engage in condomless sex than non-alcohol condition participants. The main effects for sexual sensation seeking, sexual compulsivity, and sex-related alcohol expectancies were also shown to be significant, such that individuals scoring higher on these dimensions reported stronger intentions to engage in condomless sex. However, the interactions between each of these factors and drink condition were not significant, suggesting that the impact of alcohol on condomless sex intentions was not moderated by these factors.

Discussion

The present investigation entailed the first risk-focused experiment to employ alcohol administration procedures with HIV-positive individuals, and it was uniquely positioned to identify alcohol's causal impact on HIV-positive MSM's condomless sex intentions within the context of potential moderating factors. Findings supported the primary hypothesis, demonstrating that acute alcohol consumption directly increased HIV-positive MSM's intentions to engage in sex without condoms. The impact of alcohol did not differ between HIV-positive and HIV-negative MSM, and it remained significant even when accounting for underlying intra-individual risk factors. As such, findings not only accord with results from cross-sectional research linking alcohol use to condomless sex^{5;24;28-30} and HIV seroconversion,⁶⁶ but they are also consistent with recent meta-analyses showing direct links between experimentally-manipulated alcohol consumption and sexual risk intentions among non-infected populations.^{47;48} Importantly, our work extends the research field by providing

evidence for a causal link between consuming alcohol and the engagement in sexual behaviors that possess an inherent risk for HIV transmission.

Interestingly, whereas results from Maisto et al.'s MSM-focused alcohol administration experiment⁴⁹ were suggestive of a synergistic effect of alcohol and sexual arousal on risk; findings from the present experiment showed no significant interaction between these two factors, whereby the impact of alcohol was consistent regardless of level of sexual arousal. It is important to note, however, that while the sexual arousal manipulation in the present experiment successfully induced significantly higher levels of sexual arousal among those receiving the arousing vs. non-arousing videos, the level of arousal among those in the former group was still relatively modest. As a result, it may not have been sufficiently robust to significantly amplify alcohol's impact on risk intentions. Additional research in which sexual arousal is manipulated to varying degrees would help clarify the possibly synergy between alcohol and arousal on risk intentions.

Study findings have direct implications for efforts to reduce HIV incidence among MSM. Recognizing alcohol's independent causal role in HIV transmission, prevention efforts should focus on providing HIV-positive MSM with alcohol-reduction interventions to diminish consumption levels and binge drinking, which could lead to corresponding declines in condomless sex.⁶⁷⁻⁷⁰ As an alternative, alcohol screening could be offered at clinic- and community-based venues, and hazardous drinkers could then be provided not only with evidence-based behavioral interventions to increase condom use (e.g.⁷¹⁻⁷⁴), but also with ART adherence promotion programs to help maintain HIV viral suppression in order to reduce viral transmissibility.¹¹ (see also⁷⁵)

In addition to clinic- and community-based HIV prevention efforts, HIV-positive MSM could benefit from interventions delivered precisely within alcohol and sexual partnership contexts. Research has suggested that when under the influence of alcohol, receiving salient, risk-inhibiting information or cues “in-the-moment” of sexual decision making can potentially reduce the likelihood of condomless sex.⁷⁶ Although the implementation of such an approach is not without its challenges, new smartphone-based technologies, including geolocation and real-time alcohol monitoring, could be employed to intuitively deliver timely HIV prevention messages that are strong, simple, and salient enough to be impactful under conditions of intoxication. For instance, tailored HIV prevention messages or cues could be sent to one's device upon entering an alcohol establishment or upon attaining a specified BAC. These initiatives would not only deliver HIV prevention when it is needed most, but they would also be efficient, seamless, and cost-effective.

Study results should be viewed in terms of potential limitations. First, the investigation involved a controlled experiment in which condomless sex intentions constituted the outcome. Even though intentions have been shown to be excellent surrogate indicators of actual condom use,^{53;54} concerns surrounding the ecological validity of our findings cannot be discounted entirely. Second, for ethical, safety, and legal reasons, MSM with a recent history of alcohol or substance abuse, as well as those under the age of 19, were excluded from participation. As these groups continue to be particularly hard-hit by the HIV epidemic,^{3;66;77} further efforts are necessary to better understand their alcohol-related risk

dynamics. Third, the study sample was primarily white, well-educated, and employed, and may not be fully reflective of MSM in general. Finally, knowledge and evidence regarding TasP and PrEP became more widely available as our study progressed. It is therefore unclear if and how perceptions surrounding these aspects affected MSM's condomless sex intentions. However, recognizing that PrEP was only approved in the study jurisdiction after all experimental sessions had been run, and given that HIV viral load suppression was unrelated to condomless sex intentions among our HIV-positive participants, it is unlikely that these aspects significantly influenced study outcomes.

These limitations, however, neither detract from the strengths and uniqueness of the experiment, nor do they diminish the importance of the findings that demonstrated a direct, causal impact of acute alcohol consumption on HIV-positive MSM's intentions to engage in condomless sex. Incorporating alcohol screening and interventions into HIV prevention initiatives could lead to reductions in HIV transmission risk behavior, which in turn could help curtail the persistent HIV epidemic among MSM on a broad scale.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Funding for this work was provided by the National Institutes of Health (NIH)/National Institute on Alcohol Abuse and Alcoholism (NIAAA) (4R21AA020236-02, PI: Shuper) and the Canadian Institutes of Health Research (HHP111404, PI: Shuper). We would like to thank study participants, research team members, and physicians and staff at the Maple Leaf Medical Clinic.

Sources of Funding: Funding for this work was provided by the National Institutes of Health (NIH)/National Institute on Alcohol Abuse and Alcoholism (NIAAA) (4R21AA020236-02, PI: Shuper) and the Canadian Institutes of Health Research (HHP111404, PI: Shuper).

Reference List

1. UNAIDS. [Accessed April 13, 2017] Global Aids Update 2016. 2016. Available at: http://www.unaids.org/sites/default/files/media_asset/global-AIDS-update-2016_en.pdf
2. Beyrer C, Baral SD, van GF, et al. Global epidemiology of HIV infection in men who have sex with men. *Lancet*. 2012; 380:367–377. [PubMed: 22819660]
3. Centers for Disease Control and Prevention (2016). [Accessed April 13, 2017] CDC FACT SHEET: HIV among Gay and Bisexual Men. 2016. Available at: <https://www.cdc.gov/nchhstp/newsroom/docs/factsheets/cdc-msm-508.pdf>
4. UNAIDS (2014). [Accessed April 13, 2017] The Gap Report. 2014. Available at: http://files.unaids.org/en/media/unaids/contentassets/documents/unaidspublication/2014/UNAIDS_Gap_report_en.pdf
5. Wolf SE, Maisto SA. Alcohol use and risk of HIV infection among men who have sex with men. *AIDS Behav*. 2009; 13:757–782. [PubMed: 18236149]
6. Centre for Disease Control and Prevention (2004). High-Risk Sexual Behavior by HIV-Positive Men Who Have Sex with Men-- 16 Sites, United States, 2000--2002. *MMWR*. Oct; 2004 53(38):891–894. [Accessed April 13, 2017] Available at: <https://www.cdc.gov/mmwr/preview/mmwrhtml/mm5338a1.htm>. [PubMed: 15457144]
7. Grewal R, Allen VG, Gardner S, et al. Serosorting and recreational drug use are risk factors for diagnosis of genital infection with chlamydia and gonorrhoea among HIV-positive men who have

- sex with men: results from a clinical cohort in Ontario, Canada. *Sex Transm Infect.* 2017; 93:71–75. [PubMed: 27154185]
8. Crepaz N, Marks G, Liao A, et al. Prevalence of unprotected anal intercourse among HIV-diagnosed MSM in the United States: a meta-analysis. *AIDS.* 2009; 23:1617–1629. [PubMed: 19584704]
 9. van Kesteren NM, Hospers HJ, Kok G. Sexual risk behavior among HIV-positive men who have sex with men: a literature review. *Patient Educ Couns.* 2007; 65:5–20. [PubMed: 17098392]
 10. Grant RM, Lama JR, Anderson PL, et al. Preexposure chemoprophylaxis for HIV prevention in men who have sex with men. *N Engl J Med.* 2010; 363:2587–2599. [PubMed: 21091279]
 11. Cohen MS, Chen YQ, McCauley M, et al. Prevention of HIV-1 infection with early antiretroviral therapy. *N Engl J Med.* 2011; 365:493–505. [PubMed: 21767103]
 12. Baeten JM, Donnell D, Ndase P, et al. Antiretroviral prophylaxis for HIV prevention in heterosexual men and women. *N Engl J Med.* 2012; 367:399–410. [PubMed: 22784037]
 13. McCormack S, Dunn DT, Desai M, et al. Pre-exposure prophylaxis to prevent the acquisition of HIV-1 infection (PROUD): effectiveness results from the pilot phase of a pragmatic open-label randomised trial. *Lancet.* 2016; 387:53–60. [PubMed: 26364263]
 14. Molina JM, Capitant C, Spire B, et al. On-Demand Preexposure Prophylaxis in Men at High Risk for HIV-1 Infection. *N Engl J Med.* 2015; 373:2237–2246. [PubMed: 26624850]
 15. Thigpen MC, Kebaabetswe PM, Paxton LA, et al. Antiretroviral Preexposure Prophylaxis for Heterosexual HIV Transmission in Botswana. *N Engl J Med.* 2012; 367:423–434. [PubMed: 22784038]
 16. Holloway I, Dougherty R, Gildner J, et al. PrEP Uptake, Adherence, and Discontinuation among California YMSM Using Geosocial Networking Applications. *J Acquir Immune Defic Syndr.* 2016
 17. Parsons JT, Rendina HJ, Lassiter JM, Whitfield TH, Starks TJ, Grov C. Uptake of HIV Pre-Exposure Prophylaxis (PrEP) in a National Cohort of Gay and Bisexual Men in the United States. *J Acquir Immune Defic Syndr.* 2017; 74:285–292. [PubMed: 28187084]
 18. Risher K, Mayer KH, Beyrer C. HIV treatment cascade in MSM, people who inject drugs, and sex workers. *Curr Opin HIV AIDS.* 2015; 10:420–429. [PubMed: 26352393]
 19. Smith DK, Herbst JH, Zhang X, Rose CE. Condom effectiveness for HIV prevention by consistency of use among men who have sex with men in the United States. *J Acquir Immune Defic Syndr.* 2015; 68:337–344. [PubMed: 25469526]
 20. Cooper ML. Alcohol use and risky sexual behavior among college students and youth: evaluating the evidence. *J Stud Alcohol Suppl.* 2002:101–117. [PubMed: 12022716]
 21. Crowe LC, George WH. Alcohol and human sexuality: review and integration. *Psychol Bull.* 1989; 105:374–386. [PubMed: 2660179]
 22. Dingle GA, Oei TP. Is alcohol a cofactor of HIV and AIDS? Evidence from immunological and behavioral studies. *Psychol Bull.* 1997; 122:56–71. [PubMed: 9204781]
 23. George WH, Stoner SA. Understanding acute alcohol effects on sexual behavior. *Annu Rev Sex Res.* 2000; 11:92–124. [PubMed: 11351836]
 24. Shuper PA, Neuman M, Kanteres F, Baliunas D, Joharchi N, Rehm J. Causal considerations on alcohol and HIV/AIDS—a systematic review. *Alcohol Alcohol.* 2010; 45:159–166. [PubMed: 20061510]
 25. Weinhardt LS, Carey MP. Does alcohol lead to sexual risk behavior? Findings from event-level research. *Annu Rev Sex Res.* 2000; 11:125–157. [PubMed: 11351830]
 26. Rehm J, Probst C, Shield KD, Shuper PA. Does alcohol use have a causal effect on HIV incidence and disease progression? A review of the literature and a modeling strategy for quantifying the effect. *Popul Health Metr.* 2017; 15:4. [PubMed: 28183309]
 27. Steele CM, Josephs RA. Alcohol myopia Its prized and dangerous effects. *Am Psychol.* 1990; 45:921–933. [PubMed: 2221564]
 28. Shuper PA, Joharchi N, Irving H, Rehm J. Alcohol as a correlate of unprotected sexual behavior among people living with HIV/AIDS: review and meta-analysis. *AIDS Behav.* 2009; 13:1021–1036. [PubMed: 19618261]
 29. Woolf-King SE, Maisto SA. Alcohol use and high-risk sexual behavior in Sub-Saharan Africa: a narrative review. *Arch Sex Behav.* 2011; 40:17–42. [PubMed: 19705274]

30. Halpern-Felsher BL, Millstein SG, Ellen JM. Relationship of alcohol use and risky sexual behavior: a review and analysis of findings. *J Adolesc Health*. 1996; 19:331–336. [PubMed: 8934293]
31. Leigh BC, Stall R. Substance use and risky sexual behavior for exposure to HIV Issues in methodology, interpretation, and prevention. *Am Psychol*. 1993; 48:1035–1045. [PubMed: 8256876]
32. Leigh BC. Alcohol and condom use: a meta-analysis of event-level studies. *Sex Transm Dis*. 2002; 29:476–482. [PubMed: 12172533]
33. Kalichman SC, Johnson JR, Adair V, Rompa D, Multhau K, Kelly JA. Sexual sensation seeking: scale development and predicting AIDS-risk behavior among homosexually active men. *J Pers Assess*. 1994; 62:385–397. [PubMed: 8027907]
34. Kalichman SC, Heckman T, Kelly JA. Sensation seeking as an explanation for the association between substance use and HIV-related risky sexual behavior. *Arch Sex Behav*. 1996; 25:141–154. [PubMed: 8740520]
35. Kalichman SC, Weinhardt L, DiFonzo K, Austin J, Luke W. Sensation seeking and alcohol use as markers of sexual transmission risk behavior in HIV-positive men. *Ann Behav Med*. 2002; 24:229–235. [PubMed: 12173680]
36. Zuckerman, M. *Biological Expression and Biological Bases of Sensation Seeking*. New York: Cambridge University Press; 1994.
37. Benotsch EG, Kalichman SC, Kelly JA. Sexual compulsivity and substance use in HIV-seropositive men who have sex with men: prevalence and predictors of high-risk behaviors. *Addict Behav*. 1999; 24:857–868. [PubMed: 10628518]
38. Benotsch EG, Kalichman SC, Pinkerton SD. Sexual compulsivity in HIV-positive men and women: Prevalence, predictors, and consequences of high-risk behaviors. *Sex Addict Compuls*. 2001; 8:83–99.
39. Kalichman SC, Rompa D. Sexual sensation seeking and Sexual Compulsivity Scales: reliability, validity, and predicting HIV risk behavior. *J Pers Assess*. 1995; 65:586–601. [PubMed: 8609589]
40. Kalichman SC, Cain D. The relationship between indicators of sexual compulsivity and high risk sexual practices among men and women receiving services from a sexually transmitted infection clinic. *J Sex Res*. 2004; 41:235–241. [PubMed: 15497052]
41. Bryan A, Ray LA, Cooper ML. Alcohol use and protective sexual behaviors among high-risk adolescents. *J Stud Alcohol Drugs*. 2007; 68:327–335. [PubMed: 17446971]
42. Gordon CM, Carey MP, Carey KB. Effects of a drinking event on behavioral skills and condom attitudes in men: implications for HIV risk from a controlled experiment. *Health Psychol*. 1997; 16:490–495. [PubMed: 9302547]
43. Hull JG, Bond CF Jr. Social and behavioral consequences of alcohol consumption and expectancy: a meta-analysis. *Psychol Bull*. 1986; 99:347–360. [PubMed: 3714923]
44. LaBrie J, Earleywine M, Schiffman J, Pedersen E, Marriot C. Effects of Alcohol, Expectancies, and Partner Type on Condom Use in College Males: Event-Level Analyses. *J Sex Res*. 2005; 42:259–266. [PubMed: 19817039]
45. Maisto SA, Carey MP, Carey KB, Gordon CM, Schum JL. Effects of alcohol and expectancies on HIV-related risk perception and behavioral skills in heterosexual women. *Exp Clin Psychopharmacol*. 2004; 12:288–297. [PubMed: 15571446]
46. Murphy ST, Monahan JL, Miller LC. Inference Under the Influence: The Impact of Alcohol and Inhibition Conflict on Women's Sexual Decision Making. *Pers Soc Psychol Bull*. 1998; 24:517–528.
47. Rehm J, Shield KD, Joharchi N, Shuper P. Alcohol consumption and the intention to engage in unprotected sex: systematic review and meta-analysis of experimental studies. *Addiction*. 2012; 107:51–59. [PubMed: 22151318]
48. Scott-Sheldon LA, Carey KB, Cunningham K, Johnson BT, Carey MP. Alcohol Use Predicts Sexual Decision-Making: A Systematic Review and Meta-Analysis of the Experimental Literature. *AIDS Behav*. 2016; 20(1):S19–S39. [PubMed: 26080689]

49. Maisto SA, Palfai T, Vanable PA, Heath J, Woolf-King SE. The effects of alcohol and sexual arousal on determinants of sexual risk in men who have sex with men. *Arch Sex Behav.* 2012; 41:971–986. [PubMed: 22009480]
50. Parsons JT, Vicioso KJ, Punzalan JC, Halkitis PN, Kutnick A, Velasquez MM. The impact of alcohol use on the sexual scripts of HIV-positive men who have sex with men. *J Sex Res.* 2004; 41:160–172. [PubMed: 15326541]
51. Parsons JT, Vicioso K, Kutnick A, Punzalan JC, Halkitis PN, Velasquez MM. Alcohol use and stigmatized sexual practices of HIV seropositive gay and bisexual men. *Addict Behav.* 2004; 29:1045–1051. [PubMed: 15219356]
52. Shuper PA, Joharchi N, Rehm J. Personality as a predictor of unprotected sexual behavior among people living with HIV/AIDS: a systematic review. *AIDS Behav.* 2014; 18:398–410. [PubMed: 23835736]
53. Sheeran P, Orbell S. Do intentions predict condom use? Meta-analysis and examination of six moderator variables. *Br J Soc Psychol.* 1998; 37:231–250. [PubMed: 9639864]
54. Sheeran P, Abraham C, Orbell S. Psychosocial correlates of heterosexual condom use: a meta-analysis. *Psychol Bull.* 1999; 125:90–132. [PubMed: 9990846]
55. Shuper PA, Joharchi N, Rehm J. Protocol for a Controlled Experiment to Identify the Causal Role of Acute Alcohol Consumption in Condomless Sex among HIV-Positive MSM: Study Procedures, Ethical Considerations, and Implications for HIV Prevention. *AIDS Behav.* 2016; 20(1):S173–S184. [PubMed: 26163147]
56. Davis KC, Hendershot CS, George WH, Norris J, Heiman JR. Alcohol's effects on sexual decision making: an integration of alcohol myopia and individual differences. *J Stud Alcohol Drugs.* 2007; 68:843–851. [PubMed: 17960302]
57. Babor, TF., Higgins-Biddle, JC., Saunders, JB., Monteiro, MG. World Health Organization. Dept. of Mental Health and Substance Dependence. The Alcohol Use Disorders Identification Test Guidelines for Use in Primary Care. 2nd. World Health Organization; Geneva: 2001.
58. National Institute on Drug Abuse. [Accessed April 13, 2017] Resource Guide: Screening for Drug Use in General Medical Settings. 2012. Available at: <https://www.drugabuse.gov/publications/resource-guide>
59. The Explore Team. Effects of a behavioural intervention to reduce acquisition of HIV infection among men who have sex with men: the EXPLORE randomised controlled study. *Lancet.* 2004; 364:41–50. [PubMed: 15234855]
60. Kalichman SC, Rompa D. The Sexual Compulsivity Scale: further development and use with HIV-positive persons. *J Pers Assess.* 2001; 76:379–395. [PubMed: 11499453]
61. Leigh BC. Alcohol expectancies and reasons for drinking: Comments from a study of sexual experiences. *Psychol Addict Behav.* 1990; 4:91–96.
62. Shuper PA, Fisher WA. The role of sexual arousal and sexual partner characteristics in HIV+ MSM's intentions to engage in unprotected sexual intercourse. *Health Psychol.* 2008; 27:445–454. [PubMed: 18643002]
63. Chesney MA, Ickovics JR, Chambers DB, et al. Self-reported adherence to antiretroviral medications among participants in HIV clinical trials: the AACTG adherence instruments Patient Care Committee & Adherence Working Group of the Outcomes Committee of the Adult AIDS Clinical Trials Group (AACTG). *AIDS Care.* 2000; 12:255–266. [PubMed: 10928201]
64. Walsh JC, Mandalia S, Gazzard BG. Responses to a 1 month self-report on adherence to antiretroviral therapy are consistent with electronic data and virological treatment outcome. *AIDS.* 2002; 16:269–277. [PubMed: 11807312]
65. Haahr, M. List randomiser. 1998. Available from: www.random.org/lists/
66. Baliunas D, Rehm J, Irving H, Shuper P. Alcohol consumption and risk of incident human immunodeficiency virus infection: a meta-analysis. *Int J Public Health.* 2010; 55:159–166. [PubMed: 19949966]
67. Kalichman SC, Simbayi LC, Vermaak R, et al. Randomized trial of a community-based alcohol-related HIV risk-reduction intervention for men and women in Cape Town South Africa. *Ann Behav Med.* 2008; 36:270–279. [PubMed: 18836789]

68. Monti PM, Mastroleo NR, Barnett NP, Colby SM, Kahler CW, Operario D. Brief motivational intervention to reduce alcohol and HIV/sexual risk behavior in emergency department patients: A randomized controlled trial. *J Consult Clin Psychol*. 2016; 84:580–591. [PubMed: 26985726]
69. Samet JH, Walley AY. Interventions Targeting HIV Infected Risky Drinkers: Drops in a Bottle. *Alcohol Res Health*. 2010; 33:267–279. [PubMed: 23584068]
70. Velasquez MM, von SK, Johnson DH, Green C, Carbonari JP, Parsons JT. Reducing sexual risk behaviors and alcohol use among HIV-positive men who have sex with men: a randomized clinical trial. *J Consult Clin Psychol*. 2009; 77:657–667. [PubMed: 19634959]
71. Noar SM. Behavioral interventions to reduce HIV-related sexual risk behavior: review and synthesis of meta-analytic evidence. *AIDS Behav*. 2008; 12:335–353. [PubMed: 17896176]
72. Johnson WD, Hedges LV, Diaz RM. Interventions to modify sexual risk behaviors for preventing HIV infection in men who have sex with men. *Cochrane Database Syst Rev*. 2003:CD001230. [PubMed: 12535405]
73. Johnson WD, Diaz RM, Flanders WD, et al. Behavioral interventions to reduce risk for sexual transmission of HIV among men who have sex with men. *Cochrane Database Syst Rev*. 2008:CD001230. [PubMed: 18646068]
74. Covey J, Rosenthal-Stott HE, Howell SJ. A synthesis of meta-analytic evidence of behavioral interventions to reduce HIV/STIs. *J Behav Med*. 2016; 39:371–385. [PubMed: 26831053]
75. Crepaz N, Baack BN, Higa DH, Mullins MM. Effects of integrated interventions on transmission risk and care continuum outcomes in persons living with HIV: meta-analysis, 1996-2014. *AIDS*. 2015; 29:2371–2383. [PubMed: 26372483]
76. Dal Cin S, MacDonald TK, Fong GT, Zanna MP, Elton-Marshall TE. Remembering the message: the use of a reminder cue to increase condom use following a safer sex intervention. *Health Psychol*. 2006; 25:438–443. [PubMed: 16719617]
77. Sander PM, Cole SR, Stall RD, et al. Joint effects of alcohol consumption and high-risk sexual behavior on HIV seroconversion among men who have sex with men. *AIDS*. 2013; 27:815–823. [PubMed: 23719351]

Figure 1.
Participant flow.

Table 1

Sample characteristics.

	Total N=282	HIV-Positive n=141	HIV-Negative n=141	p*
Demographics				
Age: M (SD)	42.6 (10.4)	40.9 (10.6)	44.3 (9.9)	.005
Range	20-69	20-69	20-69	
Race: n (%)				
White	203 (72.2)	92 (65.2)	111 (79.3)	.009
Latin American	20 (7.1)	13 (9.2)	7 (5.0)	.169
Black	12 (4.3)	12 (8.5)	0 (0)	.000
Chinese	7 (2.5)	3 (2.1)	4 (2.9)	.496
South-Asian	4 (1.4)	3 (2.1)	1 (0.7)	.314
Filipino	2 (0.7)	2 (1.4)	0 (0)	.251
Arab	1 (0.4)	0 (0)	1 (0.7)	.498
West Asian	1 (0.4)	0 (0)	1 (0.7)	.498
Multi-Race	31 (11.0)	16 (11.3)	15 (10.7)	.865
Identify as "Gay": n (%)	275 (97.9)	138 (97.9)	137 (97.9)	.655
Education=college diploma or higher: n (%)	201 (71.5)	84 (59.6)	117 (83.6)	.000
Employment=full/part-time: n (%)	190 (67.9)	76 (53.9)	114 (82.0)	.000
Currently in a steady partnership: n (%)	155 (56.4)	68 (48.2)	87 (64.9)	.005
Alcohol consumption				
AUDIT total: M (SD)	7.8 (3.6)	8.1 (4.0)	7.4 (3.1)	.122
Drinks/week: M (SD)	9.9 (5.4)	10.0 (5.8)	9.8 (5.1)	.801
No. of binge episodes: past 6 months: M (SD)	14.6 (16.8)	15.6 (18.5)	13.5 (15.0)	.305
Sexual behavior – past 6 months				
Engaged in condomless sex: n (%)	202 (73.2)	112 (81.8)	90 (64.7)	.001
Engaged in condomless sex with an HIV serodiscordant/HIV status unknown partner: n (%)	101 (36.7)	61 (44.5)	40 (29.0)	.008
Had multiple sex partners: n (%)	214 (76.4)	116 (82.9)	98 (70.0)	.011
Consumed alcohol before/during sex: n (%)	248 (88.6)	122 (87.1)	126 (90.0)	.452
Risk-relevant traits and alcohol expectancies				
Sexual sensation seeking: M (SD)	2.47 (0.54)	2.54 (0.57)	2.39 (0.49)	.014
Sexual compulsivity: M (SD)	1.55 (0.49)	1.62 (0.57)	1.48 (0.39)	.013
Sex-related alcohol Expectancies: M (SD)	2.45 (0.68)	2.55 (0.69)	2.36 (0.65)	.024
HIV-related factors				
Year of HIV diagnosis: M (SD)		2003 (8.4)		
Range		1980-2015		
CD4 count (cells/mm ³): M (SD)		601.6 (205.4)		
HIV viral load = undetectable: n (%)		115 (82.1)		
Currently on ART: n (%)		120 (85.1)		
4-day ART adherence = 100%: n (%)		105 (89.0)		
4-week adherence = 100%: n (%)		62 (51.7)		

Note: Percentages are based on the number of participants who indicated a specific response divided by the number of participants who responded to the item in question.

* p values are based on t -tests and χ^2 tests assessing differences between HIV-positive and HIV-negative MSM for continuous and dichotomous variables, respectively.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 2

Tests of experimental hypotheses: GLM repeated measures analysis of variance (ANOVA).

Effect*	Condition	Sub-Condition	Risk Intention (M)	F	p
Drink Condition	Non-Alcohol		1.61	9.43	.002
	Alcohol		1.86		
Participant HIV Serostatus	HIV-negative		1.41	63.22	.000
	HIV-positive		2.06		
Drink Condition × Participant HIV Serostatus	Non-Alcohol	HIV-negative	1.23	1.86	.174
		HIV-positive	1.99		
	Alcohol	HIV-negative	1.56		
		HIV-positive	2.13		

* Results are based on a full factorial GLM that included the following factors: *Between subjects*: Drink Condition, Participant HIV Status, Sexual Arousal; *Within subjects*: Partner Attractiveness, Partner Serostatus, Partner Condom Use Preference. Only those effects pertaining to the study objectives are presented.

Table 3

Tests of moderating effects of risk-relevant traits and sex-related alcohol expectancies on the alcohol-condomless sex association: GLM repeated-measures ANOVAs.

Effect*	<i>F</i>	<i>p</i>
Moderator: Sexual Sensation Seeking		
Drink Condition	13.30	.000
Sexual Sensation Seeking	46.15	.000
Sexual Sensation Seeking × Drink Condition	1.62	.205
Moderator: Sexual Compulsivity		
Drink Condition	8.48	.004
Sexual Compulsivity	7.78	.006
Sexual Compulsivity × Drink Condition	0.01	.907
Moderator: Sex-Related Alcohol Expectancies		
Drink Condition	11.45	.001
Sex-Related Alcohol Expectancies	21.23	.000
Sex-Related Alcohol Expectancies × Drink condition	2.21	.139

* Results are based on a three separate full factorial GLMs, with each containing only one of the above-indicated potential moderators. Each GLM included all of the following factors: *Between subjects*: Drink Condition, Participant HIV Status, Sexual Arousal; *Within subjects*: Partner Attractiveness, Partner Serostatus, Partner Condom Use Preference. For each GLM, only those effects pertaining to the study objectives are presented.