Published in final edited form as:

Curr Pharm Des. 2017 ; 23(41): 6347–6357. doi:10.2174/1381612823666170510124348.

Modulation of Inflammatory Response to Implanted Biomaterials Using Natural Compounds

Maria Yanez¹, James Blanchette², and Ehsan Jabbarzadeh^{1,2,3,*}

¹Department of Chemical Engineering, University of South Carolina, Columbia, SC 29208, USA

²Biomedical Engineering Program, University of South Carolina, Columbia, SC 29208, USA

³Department of Orthopedic Surgery, University of South Carolina School of Medicine, Columbia SC, 29209, USA

Abstract

Tissue engineering offers a promising strategy to restore injuries resulting from trauma, infection, tumor resection, or other diseases. In spite of significant progress, the field faces a significant bottleneck; the critical need to understand and exploit the interdependencies of tissue healing, angiogenesis, and inflammation. Inherently, the balance of these interacting processes is affected by a number of injury site conditions that represent a departure from physiological environment, including reduced pH, increased concentration of free radicals, hypoglycemia, and hypoxia. Efforts to harness the potential of immune response as a therapeutic strategy to promote tissue repair have led to the identification of natural compounds with significant anti-inflammatory properties. This article provides a concise review of the body's inflammatory response to biomaterials and describes the role of oxygen as a physiological cue in this process. We proceed to highlight the potential of natural compounds to mediate inflammatory response and improve host-graft integration. Herein, we discuss the use of natural compounds to map signaling molecules and checkpoints that regulate the cross-linkage of immune response and skeletal repair.

Keywords

Inflammatory response; biomaterials; angiogenesis; tissue engineering; macrophages; natural compounds

1. INTRODUCTION

The field of tissue engineering has utilized biocompatible materials to synthesize scaffolds with optimal chemical and mechanical properties, in addition to low-level cytotoxicity [1–4]. However, these efforts have been confounded by the host reaction after implantation. Most implanted biomaterials trigger an initial inflammatory response due to the host tissue

*Address correspondence to this author at the Associate Professor of Chemical Engineering, Associate Professor of Biomedical Engineering, Associate Professor of Orthopedic Surgery. University of South Carolina, Columbia, SC 29208; Tel: 001-803-777-3297; Fax: 001-803-777-8265; ehsan@sc.edu.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

reaction [5]. Inflammation is the body's response to injury or foreign materials, and is present in numerous diseases, such as rheumatoid arthritis, infection, and cancer [6–8]. As demonstrated in Figure 1A, following the implantation of a material, the wound healing process begins with the recruitment of neutrophils (polymorphonuclear leukocytes) to the injured site to remove bacteria and foreign bodies by phagocytosis [9–11]. To accelerate the phagocytosis process, neutrophils generate reactive oxygen species (ROS) which sometimes can cause tissue damage [11, 12]. These cells are responsible for producing inflammatory mediators that later recruit and differentiate monocyte to M0 macrophages followed by polarization to M1 pro-inflammatory and M2 pro-healing macrophages. (Figure 1B). If the host integration is not properly orchestrated, chronic inflammation, inadequate angiogenesis, and biomaterial/device fibrous encapsulation can lead to transplant failure (Figure 1C). Therefore, for tissue engineering strategies to be successful, it is vital to control biomaterial-host tissue processes interaction [2, 13, 14]. This requires a thorough understanding of the mechanisms by which physiological cues regulate the interaction of inflammatory cells to biomaterials [13–15].

Inflammatory response progresses through adherence of macrophages to implanted materials secreting growth factors, such as fibroblast growth factor (FGF), vascular endothelial growth factor (VEGF), cytokines, including tumor necrosis factor alpha (TNF- α), interleukin 6 (IL-6), granulocyte-colony stimulating factor (G-CSF), and granulocyte macrophage colony stimulating factor (GM-CSF) [16–18]. If acute inflammation is not resolved, biomaterial adherent M1 macrophages will begin to form foreign body giant cells, and the acute inflammation becomes chronic inflammation [18].

It is generally believed that three types of macrophages (M0, M1, and M2) are involved during different phases of wound healing [19]. These macrophages can be polarized by cytokines to differentiate from one phenotype to another [2]. M1 macrophages (pro-inflammatory) are present in the early stage of inflammation and are activated by foreign agents, such as microbes, necrosis, and/or tissue injury [13, 17]. It is hypothesized that M1 macrophages are subsequently differentiated to M2 macrophages (anti-inflammatory) to continue the wound-healing process [20, 21]. M1 macrophages express a number of pro-inflammatory cytokines, including TNF- α , IL-1 β , IL-6, IL-12, and IL-23, as well as chemokines such as CXC ligand motif 2 (CXL2), -9, -10, -11, -12, CC-motif ligand 2 (CCL2), -3, -4, and -5, and inducible nitric oxide synthase (iNOS) [22]. In contrast, M2 macrophages express anti-inflammatory agents including IL-10, along with scavenger, mannose, and galactose receptors [23]. There are multiple sub-types of M2 macrophages, which are distinguished from each other according to their function: M2a macrophages (alternatively activated macrophages) are activated by IL-4, whereas M2c macrophages (activated regulatory macrophages) are activated by immune complexes, glucocorticoids, prostaglandins, and IL-10 [24–26].

Inflammation is mediated by various enzymes; amongst which, the cyclo-oxygenase (COX) is the most extensively studied. COX-1 and COX-2 play critical roles during the transformation of arachidonic acid in prostaglandins [27, 28]. The overexpression of COX2 and prostaglandin is generated by several pro-inflammatory mediators, such as IL-1, IL-6, and TNF- α , and is related to fibrosis encapsulation, scar formation, and delayed wound

healing [29, 30]. Due to its role, COX2 has been used in drug discovery studies aimed at the modulation of inflammation. Currently, there are different non-steroidal anti-inflammatory drugs such as celecoxib and indomethacin that block COX2 and create an anti-inflammatory effect [31, 32]. In addition to this, other natural compounds, such as flavonoids, have been tested to reduce cytokine production and inhibit COX2 overproduction [28].

Inflammation is associated with a number of conditions that represent a departure from physiological environment, including reduced pH, an increased concentration of free radicals, hypoglycemia, and hypoxia. The relationship between oxygen partial pressure and inflammation is difficult to characterize due to the complex interplay of numerous signaling pathways. A departure from physiological oxygen tension in either direction (hypoxia or hyperoxia) can alter gene expression and impact cellular behavior with the consequences of initiating or augmenting inflammation. Additional information regarding the impact of hyperoxia on inflammation can be found in a number of recent reviews [33, 34]. In the context of tissue repair, reduced oxygen levels are more relevant due to the damage to the vasculature at an injury site. In addition, because the establishment of a mature vascular network that can return the local oxygen tension to physiological levels is a process that takes weeks, the impact of the hypoxic conditions on the inflammatory response will be chronic rather than acute.

Alterations in local oxygen tension can be translated into a change in cell phenotype through numerous mechanisms, but the hypoxia-inducible factor (HIF) family of transcription factors is often referred to as the master regulator of this process [35, 36]. HIFs are heterodimers formed between oxygen-sensitive alpha subunits (HIF-1 α and HIF-2 α) and a constitutively expressed beta subunit (HIF-1 β , also referred to as ARNT). HIF-1 α degradation is tied to cellular oxygen tension. As the oxygen levels drop below 6% (~47 mm Hg), HIF-1 α expression increases and its proteasomal degradation decreases. This change allows dimerization with the beta subunit, interaction with the coactivator CREB-binding protein, and regulation of the transcription of genes with hypoxia-responsive elements in their promoter regions [37].

HIF activity impacts immune cell phenotypes by triggering the expression of IL-1, IL-6, iNOS, TNF- α , and a host of other pro-inflammatory cytokines. Regulatory T cells (T_{reg}) require HIF activity for function, but when HIF activity is elevated by genetic modification, the T_{reg} function in a colitis model is impaired [38]. Similarly, HIF activity is essential for cytotoxic T lymphocytes (CTL) [39]. HIF activity mediates their elevated glycolysis, differentiation, and enhanced granzyme B levels. These results indicate that any perturbation from baseline activity can initiate a cascade with wide-ranging consequences.

The impacts of hypoxia and elevated HIF activity on the phenotypes of macrophages are complex due to the interplay of numerous pathways. Conclusions regarding the impact of hypoxia on macrophage polarity remain somewhat muddled. While oxygen levels can be manipulated easily outside the body, hypoxia is often accompanied by reduced pH, increased concentrations of free radicals, and hypoglycemia *in vivo*. Hypoxia is also used to describe a wide range of oxygen levels; it is expected that macrophages, like other cell types, will show different behaviors in 5, 1 and 0.1% oxygen environments, which is further

complicating the goal of reaching consensus [40]. The majority of studies have indicated that hypoxia and HIF activity promote the M1 to M2 transition, which has been summarized in previous review articles [41, 42]. Macrophages have the capacity to switch to anaerobic respiration in low-oxygen environments. In a study that investigated glioblastoma multiforme, hypoxic conditions led to elevated macrophage migration and differentiation of both M0 and M1 macrophages to the M2 phenotype, as indicated by iNOS and arginase 1 (Arg1) staining [41]. A study of intermittent hypoxia (IH) found that resveratrol could reduce the macrophage migration triggered by low oxygen [43]. It is worth noting that this study also showed a shift in the macrophage phenotype towards the M1 form, although this shift could have been the result of differences between IH and chronic hypoxia.

Reactive oxygen species (ROS) are generated as part of the inflammatory response and levels are altered depending on local oxygen tension [44, 45]. ROS can react with, and alter, essential macromolecules in cells such as proteins and nucleic acids. While generation of ROS is the result of oxidative phosphorylation, and therefore occurring in mitochondria of healthy cells, departures from physiological levels can overwhelm antioxidant capacity. This can cause loss of cells, and in turn impaired organ function, due to apoptosis or even initiate oncogenesis.

Hyperoxia can elevate levels of ROS, and this has been studied in neonatal infants placed in high oxygen incubators and in the epithelial lining of the lung during hyperoxia [46]. Such conditions are less relevant in the immediate vicinity of implanted biomaterials. Hypoxia is also linked with ROS generation, in which mitochondrial function plays an essential role. Numerous studies have shown the necessity of mitochondrial activity for ROS-mediated HIF-1 α stabilization [47, 48]. This causes elevated expression of HIF target genes and a further increase in ROS levels. These studies also identify potential therapeutic targets to reduce the impact of inflammation on HIF stabilization through ROS. A recent review summarizes many of these phenomena [49, 50].

2. CONVENTIONAL SYNTHETIC DRUGS TO SUPPRESS INFLAMMATORY RESPONSE

The interest in effective mitigation of fibrous capsul formation has led to development of steroidal and nonsteroidal classes of anti-inflammatory drugs. In this context, corticosteroids (e.g. dexamethasone) or glucocorticoids are most commonly used in clinical studies to mimic the functions of anti-inflammatory hormones [51]. Recent *in vivo* experiments demonstrated that augmenting poly(lactic-co-glycolic acid) (PLGA) scaffolds with dexamethasone improved bone regeneration and reduced inflammation. However, other studies have found that this improvement was sustained for only 4 days and corticosteroids induced a negative side-effect on bone homeostasis and density as well as metabolic system interference [13, 52].

A number of antibodies and biologic agents (e.g. infliximab, adalimumab, etanercept, golimumab, certolizumab-pegol, anakinra, and tocilizumab, which antagonize IL-1 and IL-6) have been used to treat chronic inflammation by targeting and inhibiting inflammatory

cytokines including TNF- α , IL-1, and IL-6. A drawback with the use of these inhibitors is the increased risk of infection associated with orthopedic surgical implantation [52].

Non-steroidal anti-inflammatory drugs (NSAIDs) have also been used to modulate inflammatory response. Examples of such biological agents include ibuprofen, aspirin, naproxen sodium, celecoxib, and COX2 inhibitors. These types of drugs often target the cyclooxygenase enzymes [52]. In general, COX inhibitors are responsible for the conversion of arachidonic acid to prostaglandins. However, studies performed in animal models revealed that these medications affect bone healing by reducing the bone mineral density and mechanical strength in the injured area. Despite limited success, NSAIDs provide short-term relief and do not target the real problem causing the inflammation. [51, 53].

3. NATURAL ANTI-INFLAMMATORY COMPOUNDS TO MODULATE INFLAMMATORY RESPONSE

Over the years, natural supplements have been commonly used to improve health and to reduce effects associated with aging. Natural products can serve as multi-target drugs and thereby improve more than one health problem. These supplements are extracted from various plants, fruits, and herbs and have been used for many years as remedies. Lately, these types of compounds are becoming more popular in modern medicine due to their ability to block or activate multiple pathways. These extracts include flavonoids, triterpenes, and polyphenols, to name a few. In the following, we provide a brief review of *in vitro* and *in vivo* investigations that used different forms of natural compounds to combat inflammatory diseases.

3.1. Polyphenols

Polyphenols have garnered a great deal of interest due to their ability to reduce arthritic damage through their anti-inflammatory and anti-oxidant properties [54, 55]. Recent *in vitro* and *in vivo* studies have shown that polyphenols down-regulate pro-inflammatory cytokines, stimulate anti-inflammatory agents, increase forkhead box P3 (Foxp3)-expression in CD4+ regulatory T-cells, and prevent bone loss [56]. In the following, we review the promising application of several polyphenols in studies aimed at modulation of inflammation.

3.1.1. Resveratrol—Resveratrol is a natural polyphenol found in various plants including grapes, nuts, and berries among which the skin of red grapes have the highest concentration [57]. Its chemical structure has two phenolic rings bonded together by a double styrene bond, which is responsible for the isometric *cis*- and *trans*-forms of resveratrol [58]. Resveratrol has attracted much attention over the years for its ability to modulate various pathways associated with cell growth, apoptosis, cancer, and inflammation [59]. In addition, recent studies have demonstrated resveratrol to be effective in anti-aging, anti-oxidant, anti-tumor, anti-platelet aggregation, and anti-atherogenic studies [7, 60–64].

Resveratrol has anti-inflammatory properties due to its inhibitory effect on the expression of mediators including IL-1 β , matrix metalloproteinase 13 (MMP-13), and COX2 [13, 60, 62]. The mechanism of action of resveratrol is not completely understood, but current research

suggests that resveratrol activates the silent information regulator T1 (SIRT1) protein in the cells which helps to reduce pro-inflammatory cytokines by cutting down nuclear factor kappa B (NF- κ B).

The tissue engineering community has utilized resveratrol to modulate inflammatory response to biomaterials. Resveratrol encapsulated in specific scaffolds for local delivery potently down-regulated TNF- α and IL-1 β and suppressed COX2 while blocking NF- κ B activation [62, 65]. Cellular analysis revealed that resveratrol encapsulated in PLGA nanoparticles reduced the expression of pro-inflammatory agents (IL-6 and TNF- α), whereas it increased the expression of anti-inflammatory genes including IL-10 and vascular endothelial growth factor (VEGF), which promoted osteogenesis as demonstrated by alkaline phosphatase (ALP) expression in mesenchymal stem cells (MSC). In a recent discovery, resveratrol-polyacrylic acid particles were utilized in an atelocollagen matrix to enhance cyto-compatibility, proliferation and cell viability of marrow stromal cells. In addition, the authors found that these scaffolds down-regulated IL-1 β , COX2, and MMP-13 [65]. The *in vivo* data obtained in a rabbit osteochondral defect model showed that a resveratrol-poly(acrylic acid)-collagen hydrogel increased bone and cartilage gene expression. In another study, Elmali *et al.* evaluated the effect of resveratrol in rabbits with induced arthritis by injecting resveratrol in the affected area. The authors observed that resveratrol decreased the cartilage damage and inflammation [62]. These studies demonstrate the potential of resveratrol to be used as an anti-inflammatory natural molecule which can also enhance tissue integration.

Resveratrol has demonstrated a dose response effect in various clinical, *in vitro*, and *in vivo* studies. While published data have not reported adverse effects of resveratrol in humans, in some animal studies, high dosages (3 g/kg/day in rats) has led to death due to nephrotoxicity [66]. In contrast, several *in vitro* and *in vivo* studies have reported that resveratrol is well-tolerated and relatively non-toxic [67, 68]. Further studies are warranted to elucidate the mechanism of action, as well as the dose response safety, of resveratrol in clinical applications aimed at modulation of inflammation. Another major challenge is the low bioavailability due to rapid and extensive metabolism of resveratrol. Approaches that utilize nanotechnological formulations to control release of resveratrol may help expedite the path of resveratrol to clinical trials.

3.1.2. Curcumin—Curcumin is a polyphenol found in the root of the turmeric plant, *Curcuma longa*. Curcumin exhibits anti-infectious, chemopreventive, pro-apoptotic, anti-angiogenic, and anti-inflammatory properties [8, 69]. Its anti-oxidant properties are associated with the inhibition of the nitrite-induced oxidation of hemoglobin [70]. The antioxidant effect of curcumin is highly potent due to its multiple functional groups (B-diketo group, carbon-carbon double bonds, and phenyl rings containing various amounts of hydroxyl and methoxy substituents) [71].

The mechanism of action of curcumin during inflammation is related to the suppression of pro-inflammatory pathways (nuclear factor erythroid-2 related factor-1, Nrf2-keap 1, and NF- κ B) and the down-regulation of several cytokines including TNF- α , IL-1, IL-6, IL-8, and IL-12 in various cell types [70–75]. Studies performed in rats that were positive for

inflammatory markers showed significant suppression of TNF- α , C-reactive protein (CRP), and COX2 [76]. Consistent with these findings, studies performed with colon cancer cells (HT29 and SW480) showed a down-regulation of COX2 and inhibition of prostaglandin-E2 (PEG2) when exposed to curcumin [77]. The *in vivo* studies by Miao *et al.* showed that in rats subjected to an induced middle cerebral artery occlusion, curcumin administration by intraperitoneal injections for 5 days led to TNF- α and IL-6 down-regulation [78]. In addition, the oral administration of curcumin has led to reduced inflammation in patients with ulcerative proctitis and Crohn's disease as well as accelerated wound healing [79].

Curcumin blocks the adhesion of monocytes to endothelial cells which suppresses tumor vascularization, a process essential for tumor metastasis [70]. Clinical research with osteoarthritis patients has shown that a curcumin-phosphatidylcholine complex reduces the oxidative stress, down-regulates IL-1 β , IL-6, and improves pain and various osteoarthritis symptoms [72]. Another interesting study was conducted by Sun *et al.* to encapsulate the curcumin into exosome nanoparticles in order to increase its solubility and induce lipopolysaccharide (LPS) [8, 69, 79]. Curcumin has been also encapsulated in pH-sensitive nanoparticles to decrease neutrophil infiltration and reduce TNF- α [80].

Although more research is called for, the immune modulatory property of curcumin renders it as an inexpensive, natural choice in tissue engineering applications [81–83]. Nevertheless, its short biological life, low absorption, and fast metabolism have hampered the enthusiasm for its use in clinical applications [84]. In spite of these limitations, further explorations to design novel curcumin formation derivatives to enhance systemic bioavailability and efficacy are needed for future human use.

3.1.3. Green tea (*Camellia sinensis*)—Green tea (*Camellia sinensis*) contains the major polyphenolic compounds epigallocatechin-3 galate (EGCG) and flavonoids. Evidence in literature suggest that the bioactive compounds of green tea prevent cardiovascular problems, cancer, inflammation, and neurodegenerative problems due to its anti-oxidant and anti-inflammatory properties [85–89]. Several studies have examined the mechanism of action of the green tea components in the improvement of bone formation [25]. *In vitro* studies utilizing cancer cell lines revealed that EGCG induces apoptosis and promotes cell growth arrest by regulating the cell cycle regulatory proteins, activating killer caspases, and down regulating IL-1 β and NF- κ B [57, 90]. However, cell apoptosis was not observed in healthy cells. Studies performed with osteoblastic cells revealed that green tea increased osteoblast survival and decreased the cell apoptosis due to TNF- α and IL-6 inhibition [91]. Additional findings suggest that EGCG inactivates additional pro-inflammatory agents, such as PEG2, nitric oxide (NO), and COX2 [86]. Consistent with these results, studies performed in rats with induced chronic inflammation demonstrated that the green tea polyphenolic compounds reduced chronic inflammation while preserving the bone mass and microarchitecture [87]. For example, Leong *et al.* found that green tea reduced the expression of pro-inflammatory agents (IL-1 β , TNF- α , and IL-6) and inflammatory cell infiltration in osteoarthritis mouse model [92].

The toxicity of compounds isolated from green tea has been assessed in animal models to determine the lethal dose as well as hematologic and biochemical abnormalities. In one

study, it was found that the median lethal dose was above 3–5 g/kg in rats [93]. The modulatory effect of EGCG on bone mineralization and VEGF activation combined with its effect on down-regulation of inflammatory cytokines reaffirms the significant potential of green tea extracts in tissue engineering applications.

3.1.4. Omega 3—Omega 3 is a natural anti-inflammatory compound mainly found in fish oil with active ingredients of eicosapentaenoic acid (EPA), docosahexanoic acid (DHA), and poly-unsaturated fatty acids (PUFAs) [94, 95]. Most of the studies of omega 3 have traditionally focused on cardiovascular diseases, obesity, and diabetes showing the potential of EPA and DHA to augment the production of PEG3 and inhibit the transformation of arachidonic acid to PEG2 [23, 57]. Studies performed in LPS-stimulated mouse microglial cells assessed the mechanism of action of EPA, DHA, and PUFAs. The data demonstrated that these three active ingredients of Omega 3 activate the SIRT1 pathway by inhibiting NF- κ B and reducing IL-6 and TNF- α expression [96]. Clinical investigations have revealed that patients who receive an omega 3 treatment demonstrated a suppression in arachidonic acid [97, 98]. In addition, it was shown that omega 6 activates the choline and ethanolamine phosphoglycerides and inhibited NF- κ B activation [97, 98]. In another important study, Turnen et al. found that TNF- α and IL-6 expression in serum decreased in patients who were treated with omega 3 [95]. Further evidence was demonstrated in a recent *in vitro* study showing that PUFA inhibited LPS effect in murine RAW macrophages leading to a reduction in the numbers of M1 macrophages and an increase of M2 macrophages in the adipose tissue in a mouse model [99].

A recent report coming out of Mayo Clinic has provided evidence that omega 3 is not toxic in low and moderate dosages [100]. In contrast, Sijben and Cader revealed that an n-3 PUFA supplement depressed immune system due to an alteration in cytokine production, T-cell proliferation, and T cell-mediated cytotoxicity [101]. In another study, Woodworth et al. pre-treated mice with a 3 g/kg EPA + 32.4 g/kg DHA diet for 6–8 weeks before inducing colitis. The histopathology results revealed that EPA and DHA pre-treated animals had a higher degree of inflammation and rate of mortality as compared to the control group (regular diet enriched with corn oil or safflower oil). These observations can be related to immune system alteration [102].

3.1.5. Proanthocyanidins—Proanthocyanidins are a class of polyphenolic compounds present in the fruits, bark, leaves, and seeds of various plants, which possess anti-bacterial, anti-viral, anti-inflammatory, and vasodilatory properties [26, 103, 104]. The treatment of murine macrophages with a proanthocyanidin has recently been shown to down-regulate a number of inflammatory cytokines including IL-1 β , COX2, IL-6, and TNF- α [105]. In addition, *in vivo* studies have revealed that proanthocyanidin augments the expression of IL-4, IL-10, and transforming growth factor β (TGF- β 1) while dampening the expression of IL-1 β , macrophage inflammatory protein 1 (MIP-1), and monocyte chemoattract protein-1 (MCP-1) in a mouse model of induced arthritis [26, 103].

Proanthocyanidin has a great potential to combat venous and capillary diseases as well as tumor formation due to its anti-thrombotic and anti-inflammatory effects. In a recent study in rats with cardiovascular conditions, the oral administration of oligomeric

proanthocyanidin was shown to reduce atherosclerosis [106]. The macrophage differentiation was controlled through down-regulation of vascular cell adhesion protein 1 (VCAM-I) and monocyte chemoattractant protein-1. *In vitro* studies on exposing two types of pancreatic cells (Miapaca-2 and AsPC-1) to proanthocyanadin extract for 48 hour found a significant reduction in NF- κ B and I κ B kinase- α (IKK- α) in addition to reversal of the epithelial to mesenchymal transition process [107, 108]. Published data utilizing cardiomyocytes have shown that at low concentrations (5–10 μ g/ml), proanthocyanidin possesses antioxidant properties, while at high concentrations, (100–500 μ g/ml) proanthocyanidin becomes a pro-oxidant compound, inducing ROS production and causing cell death [109].

3.1.6. Uncaria tomentosa—Uncaria tomentosa is a Peruvian woody vine known as cat's claw. The active compounds of this source include diverse arrays of polyphenols. Uncaria tomentosa has been shown to be effective in the treatment of various diseases including rheumatism, irregular menstruation, tumorigenesis, and kidney problems [110, 111]. Several studies have revealed that this natural compound has anti-inflammatory properties on the basis that it inhibits TNF- α expression and NO synthase and suppresses the activation of NF- κ B [57, 110]. This observation was consistent with an *in vivo* study demonstrating that an oral dose of Uncaria tomentosa suppressed TNF- α , IL-1 α , and IL-1 β , IL-4, and IL-17 [111]. Furthermore, the administration of Uncaria tomentosa for 8 days lowered the expression of pro-inflammatory markers including NF- κ B, COX1, and COX2 [112–115]. Other studies have shown that Uncaria tomentosa inhibits proliferation of tumor cells without causing cell death and improving cells' DNA repair [116]. Despite its significant potential, the use of uncaria tomentosa at higher doses (320 μ g/ml) is not recommended due to cytotoxic effects (blocking cell proliferation) and safety issues [110, 113].

3.2. Flavonoids

Flavonoids are water soluble metabolites abundantly found in plants which have shown many health benefits. They are most famous for their antioxidant, anti-inflammatory, and anti-carcinogenic properties [56, 117]. In the following, we provide a review of four promising flavonoids with potential to mediate inflammatory response to biomaterials.

3.2.1. 6-Shogaol—6-shogaol is one of the main compounds extracted from ginger which has been shown to possess anti-inflammatory, anti-carcinogenic, and anti-oxidant properties due to its inactivation of NF- κ B and down-regulation of IL-1 β in a mouse model [56, 118–120]. It has been used to treat neurodegenerative diseases such as Alzheimer's and Parkinson's due to its anti-neuroinflammatory and anti-oxidant properties [121]. In a recent study it was found that LPS-induced rat astrocytes treated with 6-shogaol showed an improvement in neuro-inflammation due to reduction in the levels of IL-1 β and IL-6, down-regulation of iNOS and COX2, and inactivation of NF- κ B pathway [122]. Furthermore, it has been demonstrated that 6-shogaol blocks the expression of iNOS and COX2 and inhibits the transcription of NF- κ B in LPS-induced macrophages and in a mouse model [119, 123]. The applicability of this compound is not limited to neuro-inflammation. In fact, 6-shogaol treatment of rats, in which complete Freund's adjuvant (CFA) had been administered to the synovial cavity of the knee, led to a significant reduction in the knee swelling and

inflammatory cell infiltration in the affected area [53]. In the context of cancer therapeutics, Shogaol compounds (6-shogaol and 3-phenyl-3-shogaol) down-regulated NF- κ B and the associated MMP-9 expression in breast cancer cells while suppressing IL-1 β and TNF- α in LPS-induced microglia cells [123, 124]. Studies performed using breast cancer cells (MCF-7, dosed with 6-shogaol 39.52 μ M), revealed that 6-shogaol reduces the expression of cancer stem cell markers (CD44⁺, and CD24^{-/low}) due to the mitotic block and interference with the self-renewal pathway. Higher concentrations of 6-shogaol have been shown to be safe for noncancerous cells (HEK 293 and HaCaT cells treated with 69.97 and 103.84 μ M, respectively) [125]. Further *in vivo* studies will be needed to obtain a definitive evaluation of toxic effects and clinical potential of 6-shogaol.

3.2.2. Salicylate (Salicylic Acid or Salicin)—Salicylate (salicylic acid or salicin), the main component of aspirin, is a natural compound extracted from white willow bark which is and has been used to treat pain, inflammation, fever, and recently, osteoarthritis and lower back pain [57, 126, 127]. Salicylate is a non-selective inhibitor of COX1 and COX2 and induces an anti-inflammatory effect in a variety of diseases [127, 128]. A recent *in vitro* study has shown that administration of D(–)-salicin down-regulates TNF- α , IL-1 β , and IL-6 in LPS-induced inflammatory macrophages [129]. Consistent with this finding, the intraperitoneal injection of D(–)-salicin reduced TNF- α , IL-1 β , and IL-6 levels and significantly increased IL-10 levels in mice with LPS-induced inflammation [129]. In addition, D (–)-salicin was shown to inhibit the infiltration of neutrophils and macrophages into LPS-induced lung tissue, improve the accumulation of nuclear factor erythroid 2-related factor-2 (NRF-2), and reduce oxidative stress [129]. Due to these anti-inflammatory properties, salicylate has been used to treat a number of inflammatory diseases. For example, in a recent study colitis was treated with salicylate by intraperitoneal injection for 7 days leading to suppressed edema, mucosal damage, and gross bleeding [130]. Another relevant application is the development of fully bio-absorbable salicylate-base sirolimus-eluting stent, which was implanted in a porcine coronary artery. The authors reported a positive device incorporation without causing excessive thrombotic reaction [131]. The therapeutic use of salicylate, however, faces challenges due to side effects including gastrointestinal irritation [128, 132, 133]. Studies performed in rats showed evidence that salicylate (10–100 mM) intestinal irritation is due to the intestinal epithelial cell permeability [134]. There have also been isolated reports of toxic side effects associated with the use of salicylates. As such, future investigations should determine the efficacy of low-dose therapies.

3.2.3. Naringin—Naringin is a bioflavonoid mainly found in grapefruits and other citrus fruit. Naringin possesses anti-carcinogenic, anti-oxidant, anti-inflammatory, lipid-lowering, superoxide scavenging, anti-atherogenic, and metal-chelating effects [117, 135]. Recent investigations have revealed that naringin administration suppressed NF- κ B activation and inhibited NO production in both LPS-induced macrophages and LPS-induced mice [136, 137]. A number of clinical studies have used naringin to treat inflammatory diseases *in vivo*. For example, using a rat model, investigators have shown that naringin down-regulates TNF- α and NF- κ B levels in cis-induced striatal injury [135] and kidney injury (gentamicin nephrotoxicity) [138]. In another study, a two-week treatment with naringin improved swelling, decreased the T-cell numbers, down-regulated IL-17A and IL-1 β , increased the

Treg population, up-regulated Foxp3 and trans-acting T-cell-specific transcription factor (GATA-3), and activated IL-10 and IL-4 in mice with induced polyarthritis [117]. The oral administration of naringin has also been demonstrated to inhibit NF- κ B in bronchoalveolar lavage fluid and reduce neutrophil infiltration into the lungs of LPS-induced acute lung injury as well as suppress MMP-9 expression in paraquat-induced pulmonary fibrosis in mice [139, 140]. Despite the significant benefits of Naringin, this natural compound has shown to cause several side effects including nausea, anorexia, and vomiting due to the inappropriate diffusion in the brain [141].

3.2.4. Quercetin—Quercetin is a polyphenolic flavonoid mainly found in fruits and vegetables. Quercetin possesses anti-oxidant, anti-carcinogenic, anti-inflammatory [142], and mitochondria protective effects [143]. Recent studies have revealed that quercetin administration prevented the increase of inflammatory cytokines including IL-1 β , TNF- α , and IFN- γ in addition to suppressing the activation of NF- κ B both *in vivo* and *in vitro* [143, 144]. The administration of quercetin induced the activation of nuclear factor NRF2 in LPS-induced human aortic endothelial cells [145]. It has been discovered that NRF2 protects against inflammation by down-regulating pro-inflammatory cytokines [146]. Consistent with this finding, human intestinal biopsies from inflamed areas were treated with quercetin, and the results revealed that quercetin up-regulated secretory leukocyte protease inhibitor (Slpi) expression and suppressed TNF- α [145]. More recent clinical studies revealed that quercetin oral administration of quercetin to patients with rheumatoid arthritis reduced TNF- α levels in plasma [147].

The application of quercetin is hampered by deglycosylation limiting the lifespan of this flavonoid. To overcome this limitation, researchers encapsulated quercetin in poly(L-lactic acid) scaffolds to preserve the antioxidant's stability properties and more effectively deliver the compound [148]. In light of advances in drug delivery approaches and given the striking potential of this herb in modulating inflammation, further studies are warranted to assess its clinical utility in combination with scaffolds in regenerative medicine.

3.3. Triterpenes

Triterpenes are a class of active natural compounds that are found in leaves, stem bark, fruits, and roots which have been the focus of phytochemical investigations. In the following, we review recent *in vitro* and *in vivo* studies that used three types of triterpenes to combat inflammatory diseases.

3.3.1. Tinospora cordifolia (Guduchi)—*Tinospora cordifolia* (Guduchi) belongs to the Menispermaceae family and is found in Southeast Asia. Various active ingredients including alkaloids, steroids, diterpenoids, lactones, aliphatics, and glycosides can be extracted from different parts of this plant [149]. Published data have shown the potential of *tinospora cordifolia* to treat various diseases due to its anti-diabetic, anti-malarial, anti-oxidant, anti-allergic, and anti-inflammatory properties [24, 149–153]. Diterpenoids, lactones, and aliphatic compounds are responsible for the anti-inflammatory effects of this plant [149]. *Tinospora cordifolia* regulates both pro- and anti-inflammatory pathways by the activation of various cytokines [153]. In one study, Sannegowda *et al.* found that when rats with an

induced form of arthritis were treated with *tinofora cordifolia* extract, it down-regulated several pro-inflammatory cytokines including IL-1 β , IL-6, IL-23, TNF- α and MIP-1 [154]. In another recent study, oral delivery of *tinofora cordifolia* led to significant down-regulation of iNOS, COX2, and intercellular adhesion molecule-1 (ICAM-1) in asthmatic mice [155]. *Tinospora cordifolia* extract (octacosanol) inhibited tumor cell proliferation and the angiogenesis of Ehrlich tumor cells implanted in mice through inhibition of NF- κ B activation [156].

Tinospora cordifolia extract has shown to be pro-osteogenic. Human osteoblast-like cells and primary rat osteoblastic cells proliferated at a higher rate and accelerated mineral content deposition when exposed to a *Tinospora cordifolia* extract [150, 157]. A recent study revealed that osteoblasts enhance mineral deposition when exposed to 12.5–25 μ g/ml of *tinofora cordifolia*. This effect was measured by monitoring alkaline phosphatase and calcium deposition in rat primary osteoblast as well as human osteoblast-like MG-63 cells in the presence of *Tinospora cordifolia* [158]. These results all together demonstrate the potential of *tinofora cordifolia* to be used in tissue engineering applications to concurrently enhance bone repair and alleviate inflammatory response.

3.3.2. Boswellic acid—Boswellic acid is a pentacyclic triperpenoid that is extracted from the resin of the *Boswellia serrata* tree [159]. Boswellic acid is a combination of the triterpenoids β -boswellic acid, 3-acetyl- β -boswellic acid, 11-keto- β -boswellic acid, and 3-acetyl-11-keto- β -boswellic acid. It has been used to treat chronic inflammation in rheumatoid arthritis, asthma, ulcerative colitis, and Crohn's disease on the basis that it targets NF- κ B, signal transducer and activator of transcription 3 (STAT3), androgen receptor (AR), p21, death receptor 5 (DR5), and caspase-3 and -8 [159–161]. It has been reported that this plant has anti-cancer properties due to its capacity to inhibit growth and to induce apoptosis [159, 162]. *In vitro* studies have reported that boswellic acid inhibits iNOS activity in LPS-induced macrophages [163]. Other studies have revealed that boswellic acid suppresses NF- κ B and STAT in several cancer cell lines, including HT-29 colon, SW-620, Colo-205, Hep-2 larynx, DU-145 prostate, and PC-3, confirming the anti-cancer activity of boswellic acid [162]. Animal studies have shown that boswellic acid leads to reduction in the weight of the tumor-like cells, inhibition of fibrovascular tissue, and a down-regulation tumor angiogenesis.

Arthritis is another inflammatory disease for which boswellic acid has demonstrated significant therapeutic promise. *In vivo* studies in mice with an induced form of steoarthritis demonstrated significant improvement when treated with boswellic acid for 12 weeks. Safranin-O-staining analysis showed that the boswellic acid treatment reduced the articular cartilage erosion, knee synovitis, and osteophyte formation and inhibited IL-1 β and Toll-like receptor 4 (TLR4, an innate immune system activator) in the osteoarthritic synovial explanted tissue [164]. Most recently boswellic acid was declared as a new class of safe and high tolerance NSAID [165].

3.3.3. Celastrol—Celastrol is a compound found in the root of *Tripterygium wilfordii*, which is better known as “Thunder God Vine” [166]. Celastrol has been studied for the treatment of asthma, skin inflammation, arthritis, systemic lupus erythematosus,

hypertension, and neurodegenerative disorders [167]. Recent data has shown that celastrol inhibits NF- κ B pathway leading to a significant reduction in the expression of IL-6 and IL-1 β in myeloid cells [168]. Clinical investigation of this natural compound has identified a number of health benefits including anti-oxidant, anti-cancer, and anti-inflammatory properties [169]. In a recent study, Cascão *et al.* found that celastrol, injected intraperitoneally for 19 days, down-regulated the expression of IL-1 β and TNF- α and suppressed NF- κ B and caspase-1 activation in rats with adjuvant-induced arthritis [166]. The underlying mechanism for this effect was the ability of celastrol to induce heme oxygenase-1 (HO-1), inhibit the effect of interferon gamma (IFN γ)-induced monocyte adhesion, and down-regulate ICAM-1 [166]. A similar effect on inflammatory cytokine expression in human THP-1 and T cells was noted in other studies [170–172]. Toxicological analysis performed on zebra fish have shown that 1 μ M celastrol is toxic for this animal model as a high rate of mortality was observed 48 hours after exposure. In addition, this study revealed that lower concentrations (0.5 μ M) generated animal tail malformation [173]. A comprehensive study of Celastrol's pharmacokinetics profile, bioavailability, and toxicological effects in animal models is warranted to expedite its potential move to clinical trials.

CONCLUSION

The emerging insight into the role of inflammation in tissue remodeling has led to a paradigm shift in tissue engineering approaches; rather than seeking means to suppress immune response, current efforts focus on identifying ways to harness the potential of inflammatory cells to encourage vascularization and osteogenicity. The limited success of above strategies in recapitulating the temporal aspects of the crosstalk between various cellular constituents is inherent in the design principle to target a single biological event. Having developed over an evolutionary timescale, complex pathologies such as tissue regeneration embrace a plethora of signaling pathways and inflammatory mediators acting together. The pathogenesis of these biological processes are rather multi-factorial in nature and not due to a single cytokine or cellular phenomenon. The remarkable capability of natural compounds to bind to various sites of multiple interacting molecular targets offers an innovative approach to identify and modulate targets in interdependent biological processes such as inflammation, angiogenesis, and tissue repair. Exploiting the potential of pharmacologically active natural compounds may offer a safe and robust approach to regulate the complex signaling network that governs these pathologies.

Acknowledgments

We gratefully acknowledge support from the National Institutes of Health (NIH P20 GM103641), (NIH AR063338), and National Science Foundation (NSF 1631439).

References

1. Spiller K, Freytes D, Vunjak-Novakovic G. Macrophages Modulate Engineered Human Tissues for Enhanced Vascularization and Healing. *Annals of Biomedical Engineering*. 2015; 43(3):616–627. [PubMed: 25331098]
2. Kzhyshkowska J, Gudima A, Riabov V, Dollinger C, Lavalle P, Vrana NE. Macrophage responses to implants: prospects for personalized medicine. *Journal of Leukocyte Biology*. 2015; 98

3. Jiang T, Nukavarapu SP, Deng M, Jabbarzadeh E, Kofron MD, Doty SB, Abdel-Fattah WI, Laurencin CT. Chitosan-poly(lactide-co-glycolide) microsphere-based scaffolds for bone tissue engineering: in vitro degradation and in vivo bone regeneration studies. *Acta Biomater.* Sep; 2010 6(9):3457–70. [PubMed: 20307694]
4. Mikos AG, Herring SW, Ochareon P, Elisseeff J, Lu HH, Kandel R, Schoen FJ, Toner M, Mooney D, Atala A, Van Dyke ME, Kaplan D, Vunjak-Novakovic G. Engineering complex tissues. *Tissue Eng.* Dec; 2006 12(12):3307–39. [PubMed: 17518671]
5. Aamodt JM, Grainger DW. Extracellular matrix-based biomaterial scaffolds and the host response. *Biomaterials.* Apr.2016 86:68–82. [PubMed: 26890039]
6. Schmidt-Bleek K, Kwee BJ, Mooney DJ, Duda GN. Boon and Bane of Inflammation in Bone Tissue Regeneration and Its Link with Angiogenesis. *TISSUE ENGINEERING: Part B.* 2015; 21(4):354–364.
7. Baur JA, Sinclair DA. Therapeutic Potential of Resveratrol: The in vivo evidence. *Drug Discovery.* 2006; 5:493–506.
8. Koeberle A, Werz O. Multi-target approach for natural products in inflammation. *Drug Discovery Today.* 2014; 19(12):1871–1882. [PubMed: 25172801]
9. Cote CK, Van Rooijen N, Welkos SL. Roles of macrophages and neutrophils in the early host response to *Bacillus anthracis* spores in a mouse model of infection. *Infect Immun.* Jan; 2006 74(1):469–80. [PubMed: 16369003]
10. Kruger P, Saffarzadeh M, Weber AN, Rieber N, Radsak M, von Bernuth H, Benarafa C, Roos D, Skokowa J, Hartl D. Neutrophils: Between host defence, immune modulation, and tissue injury. *PLoS Pathog.* Mar.2015 11(3):e1004651. [PubMed: 25764063]
11. Wright HL, Moots RJ, Bucknall RC, Edwards SW. Neutrophil function in inflammation and inflammatory diseases. *Rheumatology (Oxford).* Sep; 2010 49(9):1618–31. [PubMed: 20338884]
12. Medzhitov R. Origin and physiological roles of inflammation. *Nature.* Jul 24; 2008 454(7203):428–35. [PubMed: 18650913]
13. Browne S, Pandit A. Biomaterial-mediated modification of the local inflammatory environment. *Frontiers in Bioengineering and Biotechnology.* 2015; 3(67):1–14. [PubMed: 25654078]
14. Rutledge K, Cheng Q, Pryzhkova M, Harris GM, Jabbarzadeh E. Enhanced differentiation of human embryonic stem cells on extracellular matrix-containing osteomimetic scaffolds for bone tissue engineering. *Tissue Eng Part C Methods.* Nov; 2014 20(11):865–74. [PubMed: 24634988]
15. Harris GM, Rutledge K, Cheng Q, Blanchette J, Jabbarzadeh E. Strategies to direct angiogenesis within scaffolds for bone tissue engineering. *Curr Pharm Des.* 2013; 19(19):3456–65. [PubMed: 23432671]
16. Spiller KL, Anfang R, Spiller KJ, Ng J, Nakazawa KR, Daulton JW, Vunjak-Novakovic G. The Role of Macrophage Phenotype in Vascularization of Tissue Engineering Scaffolds. *Biomaterials.* 2014; 35(15):4477–4488. [PubMed: 24589361]
17. Jetten N, Verbruggen S, Gijbels MJ, Post MJ, De Winther MPJ, Donners MMPC. Anti-inflammatory M2, but not pro-inflammatory M1 macrophages promote angiogenesis in vivo. *Angiogenesis.* 2014; 17:109–118. [PubMed: 24013945]
18. Anderson JM, Rodriguez A, Chang DT. Foreign body reaction to biomaterials. *Semin Immunol.* Apr; 2008 20(2):86–100. [PubMed: 18162407]
19. Schmidt-Bleek K, Kwee BJ, Mooney DJ, Duda GN. Boon and Bane of Inflammation in Bone Tissue Regeneration and Its Link with Angiogenesis. *Tissue Eng Part B Rev.* Aug; 2015 21(4):354–64. [PubMed: 25742724]
20. Spiller K, Nassiri S, Witherel CE, Anfang RR, Ng J, Nakazawa KR, Yu T, Vunjak-Novakovic G. Sequential delivery of immunomodulatory cytokines to facilitate the M1-to-M2 transition of macrophages and enhance vascularization of bone scaffolds. *Biomaterials.* 2015; 37:197–207.
21. Lawrence T, Gilroy DW. Chronic inflammation: a failure of resolution? *Int J Exp Pathol.* Apr; 2007 88(2):85–94. [PubMed: 17408451]
22. Koh TJ, DiPietro LA. Inflammation and wound healing: the role of the macrophage. *Expert Rev Mol Med.* Jul 11.2011 13:e23. [PubMed: 21740602]
23. Im D-S. Functions of omega-3 fatty acids and FFA4 (GPR120) in macrophages. *European Journal of Pharmacology.* 2015

24. Shaikh R, Pund M, Dawane A, Iliyas S. Evaluation of Anticancer, Antioxidant, and Possible Anti-inflammatory Properties of Selected Medicinal Plants Used in Indian Traditional Medication. *Journal of Traditional and Complementary Medicine*. 2014; 4(4):253–257. [PubMed: 25379467]
25. Shen C-L, Yeh JK, Cao J, Wang J-S. Green Tea and Bone metabolism. *Nutr Res*. 2009; 29(7):437–456. [PubMed: 19700031]
26. Ahmad SF, Zoheir KMA, Abdel-Hamied HE, Ashour AE, Bakheet SA, Attia SM, Abd-Allah ARA. Grape seed proanthocyanidin extract has potent anti-arthritis effects on collagen-induced arthritis by modifying the T cell balance. *International Immunopharmacology*. 2013; 17:79–87. [PubMed: 23759204]
27. Oberyszyn TM. Inflammation and wound healing. *Front Biosci*. May 01.2007 12:2993–9. [PubMed: 17485276]
28. Ribeiro D, Freitas M, Tome SM, Silva AM, Laufer S, Lima JL, Fernandes E. Flavonoids inhibit COX-1 and COX-2 enzymes and cytokine/chemokine production in human whole blood. *Inflammation*. Apr; 2015 38(2):858–70. [PubMed: 25139581]
29. Futagami A, Ishizaki M, Fukuda Y, Kawana S, Yamanaka N. Wound healing involves induction of cyclooxygenase-2 expression in rat skin. *Lab Invest*. Nov; 2002 82(11):1503–13. [PubMed: 12429810]
30. Durant S, Duval D, Homo-Delarche F. Effect of exogenous prostaglandins and nonsteroidal anti-inflammatory agents on prostaglandin secretion and proliferation of mouse embryo fibroblasts in culture. *Prostaglandins Leukot Essent Fatty Acids*. Oct; 1989 38(1):1–8. [PubMed: 2608699]
31. Rouzer CA, Marnett LJ. Cyclooxygenases: structural and functional insights. *J Lipid Res*. Apr; 2009 50(Suppl):S29–34. [PubMed: 18952571]
32. Ricciotti E, FitzGerald GA. Prostaglandins and inflammation. *Arterioscler Thromb Vasc Biol*. May; 2011 31(5):986–1000. [PubMed: 21508345]
33. Wang L, Cui S, Ma L, Kong L, Geng X. Current advances in the novel functions of hypoxia-inducible factor and prolyl hydroxylase in invertebrates. *Insect Mol Biol*. Dec; 2015 24(6):634–48. [PubMed: 26387499]
34. Gonzalez-Muniesa P, Lopez-Pascual A, de Andres J, Lasa A, Portillo MP, Aros F, Duran J, Egea CJ, Martinez JA. Impact of intermittent hypoxia and exercise on blood pressure and metabolic features from obese subjects suffering sleep apnea-hypopnea syndrome. *J Physiol Biochem*. Sep; 2015 71(3):589–99. [PubMed: 25913417]
35. Semenza GL, Agani F, Iyer N, Jiang BH, Leung S, Wiener C, Yu A. Hypoxia-inducible factor 1: from molecular biology to cardiopulmonary physiology. *Chest*. Jul; 1998 114(1 Suppl):40S–45S. [PubMed: 9676622]
36. Semenza G. Signal transduction to hypoxia-inducible factor 1. *Biochem Pharmacol*. Sep; 2002 64(5–6):993–8. [PubMed: 12213597]
37. McNamee EN, Vohwinkel C, Eltzschig HK. Hydroxylation-independent HIF-1 α stabilization through PKA: A new paradigm for hypoxia signaling. *Sci Signal*. 2016; 9(430):fs11. [PubMed: 27245611]
38. Fluck K, Breves G, Fandrey J, Winning S. Hypoxia-inducible factor 1 in dendritic cells is crucial for the activation of protective regulatory T cells in murine colitis. *Mucosal Immunol*. Mar; 2016 9(2):379–90. [PubMed: 26220168]
39. Noman MZ, Buart S, Van Pelt J, Richon C, Hasmmim M, Leleu N, Suchorska WM, Jalil A, Lecluse Y, El Hage F, Giuliani M, Pichon C, Azzarone B, Mazure N, Romero P, Mami-Chouaib F, Chouaib S. The cooperative induction of hypoxia-inducible factor-1 α and STAT3 during hypoxia induced an impairment of tumor susceptibility to CTL-mediated cell lysis. *J Immunol*. Mar 15; 2009 182(6):3510–21. [PubMed: 19265129]
40. Ivanovic Z. Hypoxia or in situ normoxia: The stem cell paradigm. *J Cell Physiol*. May; 2009 219(2):271–5. [PubMed: 19160417]
41. Leblond MM, Gerault AN, Corroyer-Dulmont A, MacKenzie ET, Petit E, Bernaudin M, Valable S. Hypoxia induces macrophage polarization and re-education toward an M2 phenotype in U87 and U251 glioblastoma models. *Oncoimmunol*. 2016; 5(1):e1056442.
42. Fumagalli S, Perego C, Pischitta F, Zanier ER, De Simoni MG. The ischemic environment drives microglia and macrophage function. *Front Neurol*. 2015; 6:81. [PubMed: 25904895]

43. Carreras A, Zhang SX, Almendros I, Wang Y, Peris E, Qiao Z, Gozal D. Resveratrol attenuates intermittent hypoxia-induced macrophage migration to visceral white adipose tissue and insulin resistance in male mice. *Endocrinology*. Feb; 2015 156(2):437–43. [PubMed: 25406018]
44. Perrone S, Bracciali C, Di Virgilio N, Buonocore G. Oxygen Use in Neonatal Care: A Two-edged Sword. *Front Pediatr*. 2016; 4:143. [PubMed: 28119904]
45. Kaelin WG Jr. ROS: really involved in oxygen sensing. *Cell Metab*. Jun; 2005 1(6):357–8. [PubMed: 16054083]
46. Brueckl C, Kaestle S, Kerem A, Habazettl H, Krombach F, Kuppe H, Kuebler WM. Hyperoxia-induced reactive oxygen species formation in pulmonary capillary endothelial cells in situ. *Am J Respir Cell Mol Biol*. Apr; 2006 34(4):453–63. [PubMed: 16357365]
47. Guzy RD, Schumacker PT. Oxygen sensing by mitochondria at complex III: the paradox of increased reactive oxygen species during hypoxia. *Exp Physiol*. Sep; 2006 91(5):807–19. [PubMed: 16857720]
48. Sato H, Sato M, Kanai H, Uchiyama T, Iso T, Ohyama Y, Sakamoto H, Tamura J, Nagai R, Kurabayashi M. Mitochondrial reactive oxygen species and c-Src play a critical role in hypoxic response in vascular smooth muscle cells. *Cardiovasc Res*. Sep 01; 2005 67(4):714–22. [PubMed: 15913578]
49. Hielscher A, Gerecht S. Hypoxia and free radicals: role in tumor progression and the use of engineering-based platforms to address these relationships. *Free Radic Biol Med*. Feb.2015 79:281–91. [PubMed: 25257256]
50. Skiles ML, Hanna B, Rucker L, Tipton A, Brougham-Cook A, Jabbarzadeh E, Blanchette JO. ASC spheroid geometry and culture oxygenation differentially impact induction of preangiogenic behaviors in endothelial cells. *Cell Transplant*. 2015; 24(11):2323–35. [PubMed: 25197983]
51. Challem J. The Inflammation Syndrome: The Complete Nutritional Program to Prevent and Reverse Heart Disease, Arthritis, Diabetes, Allergies and Asthma. 2003
52. Mountziaris PM, Spicer PP, Kasper FK, Mikos AG. Harnessing and Modulating Inflammation in Strategies for Bone Regeneration. *TISSUE ENGINEERING: Part B*. 2011; 17(6):393–402.
53. Levy AS, Simon O, Shelly J, Gardener M. 6-Shogaol reduced chronic inflammatory response in the knees of rats treated with complete Freund's adjuvant. *BMC Pharmacology*. 2006; 6(12)
54. Shen C-L, Yeh JK, Cao JJ, Chyu M-C, Wang J-S. Green Tea and Bone Health: Evidence from Laboratory Studies. *Pharmacol Res*. 2011; 64(2):155–161. [PubMed: 21473914]
55. Mahendran G, Manoj M, Rajendra Prasad KJ, Narmatha Bai V. Antioxidants, anti-proliferative, anti-inflammatory, anti-diabetic and anti-microbial effects of isolated compounds from *Swertia corymbosa* (Grieb.) Wight ex C.B. Clark – An in vitro approach. *Food Science and Human Wellness*. 2015
56. Nanjundaiah SM, Astry B, Moudgil KD. Mediators of Inflammation-Induced Bone Damage in Arthritis and Their Control by Herbal Products. *Evidence-Based Complementary and Alternative Medicine*. 2013
57. Maroon JC, Bost JW, Maroon A. Natural anti-inflammatory agents for pain relief. *Surgical Neurology International*. 2010; 1(80)
58. Gambini J, Ingles M, Olaso G, Lopez-Grueso R, Bonet-Costa V, Gimeno-Mallench L, Mas-Bargues C, Abdelaziz KM, Gomez-Cabrera MC, Vina J, Borrás C. Properties of Resveratrol: In Vitro and In Vivo Studies about Metabolism, Bioavailability, and Biological Effects in Animal Models and Humans. *Oxid Med Cell Longev*. 2015; 2015:837042. [PubMed: 26221416]
59. Athar M, Back JH, Tang X, Kim KH, Kopelovich L, Bickers DR, Kim AL. Resveratrol: a review of preclinical studies for human cancer prevention. *Toxicol Appl Pharmacol*. Nov 01; 2007 224(3): 274–83. [PubMed: 17306316]
60. Das S, Das D. Anti-inflammatory responses of resveratrol. *Inflamm Allergy Drug Targets*. 2007; 6(3):168–173. [PubMed: 17897053]
61. Bereswill S, Muñoz M, Fischer A, Plickert R, Haag L-M, Otto B, Kuhl AA, Loddenkemper C, Gobel UB, Heimesaat MM. Anti-Inflammatory Effects of Resveratrol, Curcumin and Simvastatin in Acute Small Intestinal Inflammation. *PLoS ONE*. 2010; 5(12):1–11.
62. Elmali M, Baysal O, Harma A, Esenkaya I, Mizrak B. Effects of Resveratrol in Inflammatory Arthritis. *Inflammation*. 2007; 30(1–2)

63. Rutledge K, Cheng Q, Jabbarzadeh E. Modulation of Inflammatory Response and Induction of Bone Formation Based on Combinatorial Effects of Resveratrol. *Journal of Nanomedicine and Nanotechnology*. 2016; 7(1)
64. Soto BL, Hank JA, Darjatmoko SR, Polans AS, Yanke EM, Rakhmilevich AL, Seo S, Kim K, Reisfeld RA, Gillies SD, Sondel PM. Anti-tumor and immunomodulatory activity of resveratrol in vitro and its potential for combining with cancer immunotherapy. *Int Immunopharmacol*. 2011; 11(11):1877–1886. [PubMed: 21854876]
65. Wei W, Sun L, Zhang P, Song J, Liu W. An anti-inflammatory cell-free collagen/resveratrol scaffold for repairing osteochondral defects in rabbits. *Acta Biomaterialia*. 2014; 10:4983–4995. [PubMed: 25169257]
66. Cottart CH, Nivet-Antoine V, Laguillier-Morizot C, Beaudeau JL. Resveratrol bioavailability and toxicity in humans. *Mol Nutr Food Res*. Jan; 2010 54(1):7–16. [PubMed: 20013887]
67. Carsten RE, Bachand AM, Bailey SM, Ullrich RL. Resveratrol reduces radiation-induced chromosome aberration frequencies in mouse bone marrow cells. *Radiat Res*. Jun; 2008 169(6): 633–8. [PubMed: 18494544]
68. Williams LD, Burdock GA, Edwards JA, Beck M, Bausch J. Safety studies conducted on high-purity trans-resveratrol in experimental animals. *Food Chem Toxicol*. Sep; 2009 47(9):2170–82. [PubMed: 19505523]
69. Sun D, Zhuang X, Xiang X, Liu Y, Zhang S, Liu C, Barnes S, Grizzle W, Miller D, Zhang H-G. A Novel Nanoparticle Drug Delivery System: The Anti-inflammatory Activity of Curcu-minIs Enhanced When Encapsulated in Exosomes. *The American Society of Gene & Cell Therapy*. 2010; 18(9):1606–1614.
70. Sethi, G., Sung, B., Aggarwal, BB. The Role of Curcumin in Modern Medicine. In: Ramawat, KG., editor. *Herbal Drugs: Ethnomedicine to Modern Medicine*. 2009. p. 92-112.
71. Menon, VP., Sudheer, AR. The Molecular Targets and Therapeutic Uses of Curcumin in Health and Disease. New York: Springer Science + Business Media, LLC; 2007. Antioxidant and Anti-inflammatory Properties of Curcumin; p. 105-125.
72. Belcaro G, Cesarone MR, Dugall M, Pellegrini L, Ledda A, Grossi MG, Togni S, Appendino G. Efficacy and Safety of Meriva®, a Curcumin-phosphatidylcholine Complex, during Extended Administration in Osteoarthritis Patients. *Alternative Medicine Review*. 2010; 15(4):337–344. [PubMed: 21194249]
73. Ghosh S, Banerjee S, Sil PC. The beneficial role of curcumin on inflammation, diabetes and neurodegenerative disease: A recent update. *Food and Chemical Toxicology*. 2015; 83:111–124. [PubMed: 26066364]
74. Jurenka JS. Anti-inflammatory Properties of Curcumin, a Major Constituent of Curcuma longa: A Review of Preclinical and Clinical Research. *Alternative Medicine Review*. 2009; 14(2):141–153. [PubMed: 19594223]
75. He Y, Yue Y, Zheng X, Zhang K, Chen S, Du Z. Curcumin, inflammation, and chronic diseases: how are they linked? *Molecules*. May 20; 2015 20(5):9183–213. [PubMed: 26007179]
76. Maithilikarpagaselvi N, Sridhar MG, Swaminathan RP, Zachariah B. Curcumin prevents inflammatory response, oxidative stress and insulin resistance in high fructose fed male Wistar rats: Potential role of serine kinases. *Chemico-Biological Interactions*. 2016; 244:187–194. [PubMed: 26713546]
77. Levi-Ari S, Maimon Y, Strier L, Kazanov D, Arber N. Down-regulation of prostaglandin E2 by curcumin is correlated with inhibition of cell growth and induction of apoptosis in human colon carcinoma cell lines. *J Soc Integr Oncol*. 2006; 4(1):21–6. [PubMed: 16737669]
78. Miao Y, Zhao S, Gao Y, Wang R, Wu Q, Wu H, Luo T. Curcumin pretreatment attenuates inflammation and mitochondrial dysfunction in experimental stroke: The possible role of Sirt1 signaling. *Brain Research Bulletin*. 2016; 121:9–15. [PubMed: 26639783]
79. Basnet P, Skalko-Basnet N. Curcumin: An Anti-Inflammatory Molecule from a Curry Spice on the Path to Cancer Treatment. *Molecules*. 2011; 16:4567–4598. [PubMed: 21642934]
80. Belouqui A, Coco R, Memvanga PB, Ucakar B, des Rieux A, Preat V. pH-sensitive nanoparticles for colonic delivery of curcumin in inflammatory bowel disease. *Int J Pharm*. Oct 01; 2014 473(1–2): 203–12. [PubMed: 25014369]

81. Deodhar SD, Sethi R, Srimal RC. Preliminary study on antirheumatic activity of curcumin (diferuloyl methane). *Indian J Med Res.* Apr.1980 71:632–4. [PubMed: 7390600]
82. Hanai H, Iida T, Takeuchi K, Watanabe F, Maruyama Y, Andoh A, Tsujikawa T, Fujiyama Y, Mitsuyama K, Sata M, Yamada M, Iwaoka Y, Kanke K, Hiraishi H, Hirayama K, Arai H, Yoshii S, Uchijima M, Nagata T, Koide Y. Curcumin maintenance therapy for ulcerative colitis: randomized, multicenter, double-blind, placebo-controlled trial. *Clin Gastroenterol Hepa-tol.* Dec; 2006 4(12): 1502–6.
83. Cole GM, Teter B, Frautschy SA. Neuroprotective effects of curcumin. *Adv Exp Med Biol.* 2007; 595:197–212. [PubMed: 17569212]
84. Anand P, Kunnumakkara AB, Newman RA, Aggarwal BB. Bioavailability of curcumin: problems and promises. *Mol Pharm.* Nov-Dec;2007 4(6):807–18. [PubMed: 17999464]
85. Chatterjee P, Chandra S, Dey P, Bhattacharya S. Evaluation of anti-inflammatory effects of green tea and black tea: A comparative in vitro study. *Journal of Advance Pharmaceutical Technology & Research.* 2012; 3(2):136–138.
86. Leong DJ, Choudhury M, Hanstein R, Hirsh DM, Kim SJ, Majeska RJ, Schaffler MB, Hardin JA, Spray DC, Goldring MB, Cobelli NJ, Sun HB. Green tea polyphenol treatment is chondroprotective, anti-inflammatory and palliative in a mouse posttraumatic osteoarthritis model. *Arthritis Research Therapy.* 2014; 16(6)
87. Shen CL, Yeh Jk, Samathanam C, Cao JJ, Stoecker BJ, Dagda RY, Chyu MC, Dunn DM, Wang JS. Green tea polyphenols attenuate deterioration of bone microarchitecture in female rats with systemic chronic inflammation. *International Osteoporosis.* 2009; 22:327–337.
88. Gennaro G, Claudino M, Cestar TM, Ceolin D, Germino P, Garlet GP, Francisco de Assis G. Green Tea Modulates Cytokine Expression in the Periodontium and Attenuates Alveolar Bone Resorption in Type 1 Diabetic Rats. *PLOS ONE.* 2015; 10(8)
89. Weinreb O, Mandel S, Amit T, Youdim MB. Neurological mechanisms of green tea polyphenols in Alzheimer's and Parkinson's diseases. *J Nutr Biochem.* Sep; 2004 15(9):506–16. [PubMed: 15350981]
90. Singh BN, Shankar S, Srivastava RK. Green tea catechin, epigallocatechin-3-gallate (EGCG): mechanisms, perspectives and clinical applications. *Biochem Pharmacol.* Dec 15; 2011 82(12): 1807–21. [PubMed: 21827739]
91. Choi EM, Hwang JK. Effects of (+)-catechin on the function of osteoblastic cells. *Biol Pharm Bull.* 2003; 26(4):523–526. [PubMed: 12673036]
92. Choi EM, Hwang JK. Effects of (+)-catechin on the function of osteoblastic cells. *Biol Pharm Bull.* 2003; 118(1):523–526.
93. Yamane T, Nakatani H, Kikuoka N, Matsumoto H, Iwata Y, Kitao Y, Oya K, Takahashi T. Inhibitory effects and toxicity of green tea polyphenols for gastrointestinal carcinogenesis. *Cancer.* Apr 15; 1996 77(8 Suppl):1662–7. [PubMed: 8608559]
94. Skulas-Ray AC. Omega-3 fatty acids and inflammation: A perspective on the challenges of evaluating efficacy in clinical re-search. *Prostaglandins and Other Lipid Mediators.* 2015:104–111. [PubMed: 25698680]
95. Turunen AW, Jula A, Suominen AL, Mannisto S, Marniemi J, Kiviranta H, Tiittanen P, Karanko H, Moilanen L, Nieminen MS, Kesaniemi YA, Kahonen M, Verkasalo PK. Fish consumption, omega-3 fatty acids, and environmental contaminants in relation to low-grade inflammation and early atherosclerosis. *Environmental Research.* 2013; 120:43–54. [PubMed: 23089109]
96. Inoue T, Tanaka M, Masuda S, Ohue-Kitano R, Yamakage H, Muranaka K, Wada H, Kusakabe T, Shimatsu A, Hasegawa K, Satoh-Asahara N. Omega-3 polyunsaturated fatty acids suppress the inflammatory responses of lipopolysaccharide-stimulated mouse microglia by activating SIRT1 pathways. *Biochim Biophys Acta.* Feb 22; 2017 1862(5):552–560. [PubMed: 28254441]
97. Dasilva G, Pazos M, García-Egido E, Gallardo JM, Rodríguez I, Cela R, Medina I. Healthy effect of different proportions of marine ω -3 PUFAs EPA and DHA supplementation in Wistar rats: Lipidomic biomarkers of oxidative stress and inflammation. *Journal of Nutritional Biochemistry.* 2015; 26:1385–1392. [PubMed: 26320676]
98. Daak AA, Elderderly AY, Elbashir LM, Mariniello K, Mills J, Scarlett G, Elbashir MI, Ghebremeskel K. Omega 3 (n-3) fatty acids down-regulate nuclear factor-kappa B (NF- κ B) gene

and blood cell adhesion molecule expression in patients with homozygous sickle cell disease. *Blood Cells, Molecules and Diseases*. 2015; 55:48–55.

99. Oh DY, Talukdar S, Bae EJ, Imamura T, Morinaga H, Fan W, Li P, Lu WJ, Watkins SM, Olefsky JM. GPR120 is an Omega-3 Fatty Acid Receptor Mediating Potent Anti-Inflammatory and Insulin Sensitizing Effects. *Cell*. 2010;687–698. [PubMed: 20813258]
100. Clinic M. Omega-3 fatty acids, fish oil, alpha-linolenic acid. 2013
101. Sijben JW, Calder PC. Differential immunomodulation with long-chain n-3 PUFA in health and chronic disease. *Proc Nutr Soc*. May; 2007 66(2):237–59. [PubMed: 17466105]
102. Woodworth HL, McCaskey SJ, Duriancik DM, Clinthorne JF, Langohr IM, Gardner EM, Fenton JJ. Dietary fish oil alters T lymphocyte cell populations and exacerbates disease in a mouse model of inflammatory colitis. *Cancer Res*. Oct 15; 2010 70(20):7960–9. [PubMed: 20798218]
103. Cho M-L, Heo Y-J, Park M-K, Oh H-J, Park J-S, Woo Y-J, Ju J-H, Park S-H, Kim H-Y, Min J-K. Grape seed proanthocyanidin extract (GSPE) attenuates collagen-induced arthritis. *Immunology Letters*. 2009; 124:102–110. [PubMed: 19446580]
104. Park M-K, Park J-S, Cho M-L, Oh H-J, Heo Y-J, Woo Y-J, Heo Y-M, Park M-J, Park H-S, Park S-H, Kim H-Y, Min J-K. Grape seed proanthocyanidin extract (GSPE) differentially regulates Foxp3+ regulatory and IL-17+ pathogenic T cell in autoimmune arthritis. *Immunology Letters*. 2011; 135:50–58. [PubMed: 20933009]
105. Chu H, Tang Q, Huang H, Hao W, Wei X. Grape-seed proanthocyanidins inhibit the mRNA expression of inflammatory mediators by suppressing MAPK and nuclear factor- κ B signal pathways in lipopolysaccharide-induced RAW 264.7 macrophages. *Environmental Toxicology and Pharmacology*. 2015
106. Mohana T, Navin AV, Jamuna S, Sadullah MSS, Devaraj SN. Inhibition of differentiation of monocyte to macrophages in atherosclerosis by oligomeric proanthocyanidins –In-vivo and in-vitro study. *Food and Chemical Toxicology*. 2015; 82:96–105. [PubMed: 25981678]
107. Prasad R, Katiyar SK. Grape seed proanthocyanidins inhibit migration potential of pancreatic cancer cells by promoting mesenchymal-to-epithelial transition and targeting NF- κ B. *Cancer Letters*. 2013; 334:118–126. [PubMed: 22902508]
108. Chen K-Y, Shyu P-C, Dong G-C, Chen Y-S, Kuo W-W, Yao C-H. Reconstruction of calvarial defect using a tricalcium phosphate-oligomeric proanthocyanidins cross-linked gelatin composite. *Biomaterials*. 2009; 30:1682–1688. [PubMed: 19136152]
109. Shao ZH, Vanden Hoek TL, Xie J, Wojcik K, Chan KC, Li CQ, Hamann K, Qin Y, Schumacker PT, Becker LB, Yuan CS. Grape seed proanthocyanidins induce pro-oxidant toxicity in cardiomyocytes. *Cardiovasc Toxicol*. 2003; 3(4):331–9. [PubMed: 14734830]
110. Akesson C, Lindgren H, Pero RW, Leanderson T, Ivars F. An extract of *Uncaria tomentosa* inhibiting cell division and NF- κ B activity without inducing cell death. *International Immunopharmacology*. 2003; 3:1889–1900. [PubMed: 14636838]
111. Rojas-Duran R, Gonzalez-Aspajo G, Ruiz-Martel C, Bourdy G, Doroteo-Ortega VH, Alban-Castillo J, Robert G, Auberger P, Deharo E. Anti-inflammatory activity of Mitraphylline isolated from *Uncaria tomentosa* bark. *Journal of Ethnopharmacology*. 2012; 143:801–804. [PubMed: 22846434]
112. Aguilar JL, Rojas P, Marcelo A, Plaza A, Bauer R, Reininger E, Klaas CA, Merfort I. Anti-inflammatory activity of two different extracts of *Uncaria tomentosa* (Rubiaceae). *Journal of Ethnopharmacology*. 2002; 81:271–276. [PubMed: 12065162]
113. Allen-Hall L, Arnason JT, Cano P, Lafrenie RM. *Uncaria tomentosa* acts as a potent TNF-inhibitor through NF- κ B. *Journal of Ethnopharmacology*. 2010; 127:685–693. [PubMed: 19995599]
114. Allen-Hall L, Cano P, Arnason JT, Rojas R, Lock O, Lafrenie RM. Treatment of THP-1 cells with *Uncaria tomentosa* extracts differentially regulates the expression of IL-1 β and TNF- α . *Journal of Ethnopharmacology*. 2007; 109:312–317. [PubMed: 16959454]
115. Cisneros FJ, Jayo M, Niedziela L. An *Uncaria tomentosa* (cat's claw) extract protects mice against ozone-induced lung inflammation. *Journal of Ethnopharmacology*. 2005; 96:355–364. [PubMed: 15619552]

116. Akesson C, Lindgren H, Pero RW, Leanderson T, Ivars F. An extract of *Uncaria tomentosa* inhibiting cell division and NF-kappa B activity without inducing cell death. *Int Immunopharmacol.* Dec; 2003 3(13–14):1889–900. [PubMed: 14636838]
117. Ahmad SF, Zoheir KMA, Abdel-Hamied HE, Ashour AE, Bakheet SA, Attia SM, Abd-Allah ARA. Amelioration of autoimmune arthritis by naringin through modulation of T regulatory cells and Th1/Th2 cytokines. *Cellular Immunology.* 2014; 287:112–120. [PubMed: 24487035]
118. Debnath T, Kim DH, Lim BO. Natural Products as a Source of Anti-Inflammatory Agents Associated with Inflammatory Bowel Disease. *Molecules.* 2013; 18:7253–7270. [PubMed: 23783459]
119. Pan M, Hsieh M, Hsu P, Ho S, Lai C, Wu H, Sang S, Ho C. 6-Shogaol suppressed lipopolysaccharide-induced up-expression of iNOS and COX-2 in murine macrophages. *Molecular Nutrition & Food Research.* 2008; 52(12):1467–1477.
120. Gan F-F, Ling H, Ang X, Reddy SA, Lee SS-H, Yang H, Tan S-H, Hayes JD, Chui W-K, Chew E-H. A novel shogaol analog suppresses cancer cell invasion and inflammation, and displays cytoprotective effects through modulation of NF- κ B and Nrf2-Keap1 signaling pathways. *Toxicology and Applied Pharmacology.* 2013; 272:852–862. [PubMed: 23899529]
121. Moon M, Kim HG, Choi JG, Oh H, Lee PK, Ha SK, Kim SY, Park Y, Huh Y, Oh MS. 6-Shogaol, an active constituent of ginger, attenuates neuroinflammation and cognitive deficits in animal models of dementia. *Biochemical and Biophysical Research Communications* vol. 2014; 449:8–13.
122. Shim S, Kim S, Choi D-S, Kwon Y-B, Kwon J. Anti-inflammatory effects of [6]-shogaol: Potential roles of HDAC inhibition and HSP70 induction. *Food and Chemical Toxicology.* 2011; 49:2734–2740. [PubMed: 21864631]
123. Ha SK, Moon E, Ju MS, Kim DH, Ryu JH, Oh MS, Kim SY. 6-Shogaol, a ginger product, modulates neuroinflammation: A new approach to neuroprotection. *Neuropharmacology.* 2012; 63:211–223. [PubMed: 22465818]
124. Sabina EP, Rasool M, Mathew L, EzilRani P, Indu H. 6-Shogaol inhibits monosodium urate crystal-induced inflammation – An in vivo and in vitro study. *Food and Chemical Toxicology.* 2010; 48:229–235. [PubMed: 19819286]
125. Ray A, Vasudevan S, Sengupta S. 6-Shogaol Inhibits Breast Cancer Cells and Stem Cell-Like Spheroids by Modulation of Notch Signaling Pathway and Induction of Autophagic Cell Death. *PLoS One.* 2015; 10(9):e0137614. [PubMed: 26355461]
126. Beg S, Swain S, Hasan H, Barkat A, Hussain S. Systematic review of herbals as potential anti-inflammatory agents: Recent advances, current clinical status and future perspectives. *Pharmacogn Rev.* 2011; 5(10):120–137. [PubMed: 22279370]
127. Ishikado A, Sono Y, Matsumoto M, Robida-Stubbs S, Okuno A, Goto M, King GL, Blackwell TK, Makino T. Willow bark extract increases antioxidant enzymes and reduces oxidative stress through activation of Nrf2 in vascular endothelial cells and *Caenorhabditis elegans*. *Radic Biol Med.* 2013; 65
128. Bucolo C, Marrazzo G, Platania CBM, Drago F, Leggio GM, Salomone S. Fortified Extract of Red Berry, Ginkgo biloba, and White Willow Bark in Experimental Early Diabetic Retinopathy. *J Diabetes Res.* 2013
129. Li Y, Wu Q, Deng Y, Lv H, Qiu J, Chi G, Feng H. D(–)-Salicin inhibits the LPS-induced inflammation in RAW264.7 cells and mouse models. *International Immunopharmacology.* 2015; 26:286–294. [PubMed: 25907238]
130. Verma NVN, Pau JPJ. Salicin ameliorates the inflammation and prevents the loss of gut flora in DSS induced mice model of colitis.
131. Jabara R, Pendyala L, Geva S, Chen J, Chronos N, Robinson K. Novel fully bioabsorbable salicylate-based sirolimuseluting stent. *EuroIntervention.* Dec 15; 2009 5(Suppl F):F58–64. [PubMed: 22100678]
132. Mahdi JG. Medicinal potential of willow: A chemical perspective of aspirin discovery. *Journal of Saudi Chemical Society.* 2010; 14:317–322.

133. Shakibaei M, Allaway D, Nebrich S, Mobasheri A. Botanical Extracts from Rosehip (*Rosa canina*), Willow Bark (*Salix alba*), and Nettle Leaf (*Urtica dioica*) Suppress IL-1 β -Induced NF- κ B Activation in Canine Articular Chondrocytes. *Evid Based Complement Alternat Med*. 2012
134. Fukui A, Naito Y, Handa O, Kugai M, Tsuji T, Yoriki H, Qin Y, Adachi S, Higashimura Y, Mizushima K, Kamada K, Katada K, Uchiyama K, Ishikawa T, Takagi T, Yagi N, Kokura S, Yoshikawa T. Acetyl salicylic acid induces damage to intestinal epithelial cells by oxidation-related modifications of ZO-1. *Am J Physiol Gastrointest Liver Physiol*. Oct 15; 2012 303(8):G927–36. [PubMed: 22917627]
135. Chtourou Y, Aouey B, Kebieche M, Fetoui H. Protective role of naringin against cisplatin induced oxidative stress, inflammatory response and apoptosis in rat striatum via suppressing ROS-mediated NF- κ B and P53 signaling pathways. *Chemico-Biological Interactions*. 2015; 239:76–86. [PubMed: 26120027]
136. Golechha M, Sarangal V, Bhatia J, Chaudhry U, Saluja D, Arya DS. Naringin ameliorates pentylenetetrazol-induced seizures and associated oxidative stress, inflammation, and cognitive impairment in rats: Possible mechanisms of neuroprotection. *Epilepsy & Behavior*. 2014; 41:98–102. [PubMed: 25461197]
137. Kanno, S-i, Shouji, A., Tomizawa, A., Hiura, T., Osanai, Y., Ujibe, M., Obara, Y., Nakahata, N., Ishikawa, M. Inhibitory effect of naringin on lipopolysaccharide (LPS)-induced endotoxin shock in mice and nitric oxide production in RAW 264.7 macrophages. *Life Sciences*. 2006; 78:673–681. [PubMed: 16137700]
138. Sahu BD, Tatireddy S, Koneru M, Borkar RM, Kumar JM, Kuncha M, R S, R SS, Sistla R. Naringin ameliorates gentamicin-induced nephrotoxicity and associated mitochondrial dysfunction, apoptosis and inflammation in rats: Possible mechanism of nephroprotection. *Toxicology and Applied Pharmacology*. 2014; 277:8–20. [PubMed: 24637089]
139. Liu Y, Wu H, Nie Y-c, Chen J-l, Su W-w, Li P-b. Naringin attenuates acute lung injury in LPS-treated mice by inhibiting NF- κ B pathway. *International Immunopharmacology*. 2011; 11:1606–1612. [PubMed: 21640201]
140. Chen Y, Nie Y-c, Luo Y-l, Lin F, Zheng Y-f, Cheng G-h, Wu H, Zhang K-j, Su W-w, Shen J-g, Li P-b. Protective effects of naringin against paraquat-induced acute lung injury and pulmonary fibrosis in mice. *Food and Chemical Toxicology*. 2013; 58:133–140. [PubMed: 23603004]
141. Jung UJ, Kim SR. Effects of naringin, a flavanone glycoside in grapefruits and citrus fruits, on the nigrostriatal dopaminergic projection in the adult brain. *Neural Regen Res*. Aug 15; 2014 9(16): 1514–7. [PubMed: 25317167]
142. Li Y, Yao J, Han C, Yang J, Chaudhry MT, Wang S, Liu H, Yin Y. Quercetin, Inflammation and Immunity. *Nutrients*. Mar 15.2016 8(3):167. [PubMed: 26999194]
143. Carrasco-Pozo C, Tan KN, Reyes-Farias M, De La Jara N, Ngo ST, Garcia-Diaz DF, Llanos P, Cires MJ, Borges K. The deleterious effect of cholesterol and protection by quercetin on mitochondrial bioenergetics of pancreatic beta-cells, glycemic control and inflammation: In vitro and in vivo studies. *Redox Biol*. Oct.2016 9:229–243. [PubMed: 27591402]
144. Overman A, Chuang CC, McIntosh M. Quercetin attenuates inflammation in human macrophages and adipocytes exposed to macrophage-conditioned media. *Int J Obes (Lond)*. Sep; 2011 35(9): 1165–72. [PubMed: 21224828]
145. K D, De Santis S, Serino G, Galleggiante V, Caruso ML, Mastronardi M, Cavalcanti E, Ranson N, Pinto A, Campiglia P, Santino A, Eri R, Chieppa M. Secretory leukoprotease inhibitor is required for efficient quercetin-mediated suppression of TNF α secretion. *Oncotarget*. 2016
146. Khor TO, Huang MT, Kwon KH, Chan JY, Reddy BS, Kong AN. Nrf2-deficient mice have an increased susceptibility to dextran sulfate sodium-induced colitis. *Cancer Res*. Dec 15; 2006 66(24):11580–4. [PubMed: 17178849]
147. Javadi F, Ahmadzadeh A, Eghtesadi S, Aryaeian N, Zabihyeganeh M, Rahimi Foroushani A, Jazayeri S. The Effect of Quercetin on Inflammatory Factors and Clinical Symptoms in Women with Rheumatoid Arthritis: A Double-Blind, Randomized Controlled Trial. *J Am Coll Nutr*. Oct 6.2016 :1–7. [PubMed: 25932956]
148. Cruz-Zuniga JM, Soto-Valdez H, Peralta E, Mendoza-Wilson AM, Robles-Burgueno MR, Auras R, Gamez-Meza N. Development of an antioxidant biomaterial by promoting the deglycosylation

of rutin to isoquercetin and quercetin. Food Chem. Aug 01.2016 204:420–6. [PubMed: 26988520]

149. Saha S, Ghosh S. *Tinospora cordifolia*: One plant, many roles. Ancient Science of Life. 2012; 31(4):151–159. [PubMed: 23661861]
150. Abiramasundari G, Sumalatha KR, Sreepriya M. Effects of *Tinospora cordifolia* (Menispermaceae) on the proliferation, osteogenic differentiation and mineralization of osteoblast model systems in vitro. Journal of Ethnopharmacology. 2012; 141:474–480. [PubMed: 22449439]
151. Sharma U, Bala M, Kumar N, Singh B, Munshi RK, Bhalerao S. Immunomodulatory active compounds from *Tinospora cordifolia*. Journal of Ethnopharmacology. 2012; 141:918–926. [PubMed: 22472109]
152. Patgiri B, Umretia BL, Vaishnav PU, Prajapati PK, Shukla VJ, Ravishankar B. Anti-inflammatory activity of Guduchi Ghana (aqueous extract of *Tinospora Cordifolia* Miers). AYU. 2014; 35(1): 108–110. [PubMed: 25364210]
153. Koppada R, Norozian FM, Torbati D, Kalomiris S, Ramachandran C, Totapally BR. Physiological Effects of a Novel Immune Stimulator Drug, (1,4)- α -D-Glucan, in Rats. Basic & Clinical Pharmacology & Toxicology. 2009; 105:217–221. [PubMed: 19389049]
154. Sannegowda KM, Venkatesha SH, Moudgil KD. *Tinospora cordifolia* inhibits autoimmune arthritis by regulating key immune mediators of inflammation and bone damage. International Journal of Immunopathology and Pharmacology. 2015; 28(4):251–531.
155. Sannegowda KM, Venkatesha SH, Moudgil KD. *Tinospora cordifolia* inhibits autoimmune arthritis by regulating key immune mediators of inflammation and bone damage. Int J Immunopathol Pharmacol. Dec; 2015 28(4):521–31. [PubMed: 26467057]
156. Thippeswamy G, Sheela ML, Salimath BP. Octacosanol isolated from *Tinospora cordifolia* downregulates VEGF gene expression by inhibiting nuclear translocation of NF- κ B and its DNA binding activity. European Journal of Pharmacology. 2008; 588:141–150. [PubMed: 18513715]
157. Saha S, Ghosh S. *Tinospora cordifolia*: One plant, many roles. Anc Sci Life. Apr; 2012 31(4): 151–9. [PubMed: 23661861]
158. Abiramasundari G, Sumalatha KR, Sreepriya M. Effects of *Tinospora cordifolia* (Menispermaceae) on the proliferation, osteogenic differentiation and mineralization of osteoblast model systems in vitro. J Ethnopharmacol. May 07; 2012 141(1):474–80. [PubMed: 22449439]
159. Sharma S, Gupta S, Khajuria V, Bhagat A, Ahmed Z, Shah BA. Analogues of boswellic acids as inhibitors of pro-inflammatory cytokines TNF- α and IL-6. Bioorganic & Medicinal Chemistry Letters. 2016; 26:695–698. [PubMed: 26711891]
160. Yadav VR, Prasad S, Sung B, Kannappan R, Aggarwal BB. Targeting Inflammatory Pathways by Triterpenoids for Prevention and Treatment of Cancer. Toxins. 2010; 2:2428–2466. [PubMed: 22069560]
161. Saraswati S, Pandey M, Mathur R, Agrawal SS. Boswellic acid inhibits inflammatory angiogenesis in a murine sponge model. Microvascular Research. 2011; 82:263–268. [PubMed: 21864545]
162. Kumar A, Shah BA, Singh S, Hamid A, Singh SK, Sethi VK, Saxena AK, Singh J, Taneja SC. Acyl derivatives of boswellic acids as inhibitors of NF- κ B and STATs. Bioorganic & Medicinal Chemistry Letters. 2012; 22:431–435. [PubMed: 22123322]
163. Henkel A, Kather N, Monch B, Northoff H, Jauch J, Werz O. Boswellic acids from frankincense inhibit lipopolysaccharide functionality through direct molecular interference. Biochemical Pharmacology. 2012; 83:115–121. [PubMed: 22001311]
164. Wang Q, Pan X, Wong H, Wagner CA, Lahey LJ, Robinson WH, Sokolove J. Oral and topical boswellic acid attenuates mouse osteoarthritis. Oral and topical boswellic acid attenuates mouse osteoarthritis. 2014; 22:128–132.
165. Singh GB, Bani S, Singh S. Toxicity and safety evaluation of boswellic acids. Phytomedicine. May; 1996 3(1):87–90. [PubMed: 23194869]

166. Seo WY, Ju SM, Song HY, Goh AR, Jun J-G, Kang Y-H, Choi SY, Park J. Celastrol suppresses IFN-gamma-induced ICAM-1 expression and subsequent monocyte adhesiveness via the induction of heme oxygenase-1 in the HaCaT cells. *Biochemical and Biophysical Research Communications*. 2010; 398:140–145. [PubMed: 20599745]
167. Shaker ME, Ashamalla SA, Houssen ME. Celastrol ameliorates murine colitis via modulating oxidative stress, inflammatory cytokines and intestinal homeostasis. *Chemico-Biological Interactions*. 2014; 210:26–33. [PubMed: 24384223]
168. Astry B, Venkatesha SH, Laurence A, Christensen-Quick A, Garzino-Demo A, Frieman MB, O'Shea JJ, Moudgil KD. Celastrol, a Chinese herbal compound, controls autoimmune inflammation by altering the balance of pathogenic and regulatory T cells in the target organ. *Clin Immunol*. Apr; 2015 157(2):228–38. [PubMed: 25660987]
169. Kannaiyan R, Shanmugam MK, Sethi G. Molecular targets of celastrol derived from Thunder of God Vine: Potential role in the treatment of inflammatory disorders and cancer. *Cancer Letters*. 2011; 303:9–20. [PubMed: 21168266]
170. Cascão R, Vidal B, Raquel H, Neves-Costa A, Figueiredo N, Gupta V, Fonseca JE, Moita LF. Effective treatment of rat adjuvant-induced arthritis by celastrol. *Autoimmunity Reviews*. 2012; 11:856–862. [PubMed: 22415021]
171. Astry B, Venkatesha SH, Laurence A, Christensen-Quick A, Garzino-demo A, Frieman MB, O'Shea JJ, Moudgil KD. Celastrol, a Chinese herbal compound, controls autoimmune inflammation by altering the balance of pathogenic and regulatory T cells in the target organ. *Clinical Immunology*. 2015; 157:228–238. [PubMed: 25660987]
172. Li, G-q, Zhang, Y., Liu, D., Qian, Y-y, Zhang, H., Guo, S-y, Sunagawa, M., Hisamitsu, T., Liu, Y-q. Celastrol inhibits interleukin-17A-stimulated rheumatoid fibroblast-like synoviocyte migration and invasion through suppression of NF- κ B-mediated matrix metalloproteinase-9 expression. *International Immunopharmacology*. 2012; 14:422–431. [PubMed: 22954486]
173. Wang S, Liu K, Wang X, He Q, Chen X. Toxic effects of celastrol on embryonic development of zebrafish (*Danio rerio*). *Drug Chem Toxicol*. Jan; 2011 34(1):61–5. [PubMed: 20954803]

Biomaterial - Host Interaction

Fig. 1.

Biomaterial host interaction. A) Host reaction after biomaterial implantation where neutrophils migrate to the wound site, and monocytes differentiate into M1 macrophages (acute inflammation). Hypoxia can trigger the expression of pro-inflammatory cytokines helping the transition from M1 to M2 macrophages. B) Pro-healing, normal wound healing process where monocytes are polarized to M2 macrophages. The addition on natural compounds (NC) help to improve the inflammation process when they target pro-inflammatory mediators including IFN- γ , IL-6, TNF- α , and IL-1 β . C) Chronic

inflammation, normal inflammatory response fails, and pro-inflammatory mediators are over expressed. M0 macrophages are polarized to M1 macrophages and fibroblasts migrate to the device generating fibrosis.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript