Published in final edited form as:

Qual Life Res. 2017 December ; 26(12): 3239–3250. doi:10.1007/s11136-017-1647-9.

Internal resources among informal caregivers: Trajectories and associations with well-being

Kristin Litzelman^{1,*}, Gina Tesaro², and Rebecca Ferrer²

¹University of Wisconsin

²National Cancer Institute

Abstract

Purpose—The experiences, skills, and internal resources that informal caregivers bring into their role may play a critical part in their mental health and well-being. This study examined how caregiver internal resources changed over a 10-year period, and how this was related to caregivers' well-being.

Methods—Data are from the Midlife in the United States (MIDUS) study, a national sample of adults, at two time points: 1995–1996 (T1), and 2004–2006 (T2). We identified subjects who reported being a caregiver at T2 and starting care after T1 (mean age=56; 65% female). We examined internal resources – sense of control (personal mastery); primary and secondary control strategies (persistence in goal striving; positive reappraisal; lowering expectations), and social support seeking – and psychological and subjective well-being. We evaluated how internal resources changed over time, and how these trajectories were associated with well-being at T2 using multivariable linear regressions.

Results—Most caregivers had stable levels of internal resources (between 4% and 13% showed an increase or decrease). Caregivers with increasing or high-stable levels of personal mastery had significantly better well-being on 6 out of 8 subscales compared to low-stable levels (effect sizes [ES] between 0.39 and 0.79). Increasing persistence was associated with better personal growth and environmental mastery (ES=0.96 and 0.91), and increasing and high-stable positive reappraisal were associated with better affect (ES=0.63 and 0.48) compared to low-stable levels. Lowering aspirations and support seeking were not associated with well-being outcomes.

Conclusions—Practices or interventions that support or improve internal resources could potentially improve caregiver well-being.

Keywords

Caregiver; MIDUS; resources; mastery; control; social support seeking; longitudinal

*Corresponding author: 4115 Nancy Nicholas Hall, 1300 Linden Drive, Madison, WI 53706, Phone: 608-262-3314, litzelman@wisc.edu.

ORCID: 0000-0002-5542-6434

INTRODUCTION

Informal caregiving (providing unpaid support to a family member or a friend with an illness or a disability) has both positive and negative consequences for caregivers. While many caregivers find benefits in the experience, such as a sense of purpose [1; 2], many also report feeling stressed or burdened by their role [3]. Moreover, some caregivers experience serious problems like depression [4].

The *internal resources* that caregivers bring into their role – experiences, strengths, personality characteristics, coping skills, social background, and other psychological resources that support individuals' resilience or coping – may play a critical part in how caregiving influences emotional health, psychological well-being, or life satisfaction. Internal resources and related concepts (e.g., “intrapersonal resources” [5], “psychological capital” [6], and “resilience resources” [7]) are defined diversely in the literature, but often encompass personality characteristics such as self-esteem and optimism, as well as skills such as self-efficacy or perceived control [5; 8]. One general definition refers to “characteristics which are internal...When present, these characteristics result in better coping.” [9]

Theory and limited empirical research support the notion that caregivers with certain internal resources may be more resilient. Theories of stress process and caregiving suggest that antecedent factors such as the caregiver's characteristics, caregiving history, or family network play an important role in the development of caregiver stress and subsequent outcomes [10]. In particular, Goode's longitudinal adaptation of the stress process model [11] suggests that changes in caregivers' mental and physical health are influenced by their initial psychological resources, changes in stressors, and changes in psychological resources. Furthermore, psychological homeostasis theory [5] suggests that psychological resources play a critical role in the maintenance of well-being during stress (such as caregiving). Research among caregivers of patients with dementia has shown the fundamental role of psychosocial resources (including appraisals, coping, social support, and dispositional resilience) as predictors of stable caregiver mental health and resilience outcomes overtime [11; 12]. Little research, however, has addressed how such attributes may change after the onset of caregiving, and how this may be related to caregiver outcomes. In a recent longitudinal study, Hajek and Konig [13] reported that self-efficacy moderated the adverse impact of informal caregiving on subjective well-being, in particular mental health. Other internal resources may too be associated with caregiver well-being outcomes.

In order to better understand the role that internal resources play in caregiver well-being, this study sought to 1) evaluate how caregivers' internal resources change over time, from prior to the commencement of caregiver to during or shortly after the caregiving role; and 2) determine how these internal resource trajectories are associated with caregiver well-being over time. In particular, we examine both psychological (eudaimonic) and subjective (hedonic) well-being [14; 15], embracing the World Health Organization's definition of health as not merely the absence of disease and the biopsychosocial impact of well-being on other physical and mental health outcomes [16; 17]. The findings stand to improve our general understanding of the caregiving experience, and highlight novel targets for

intervention to support caregiver well-being and mitigate the adverse impacts of caregiver stress and burden.

METHODS

This study was a secondary analysis of data collected as part of Mid-life in the United States (MIDUS) study [18]. Briefly, MIDUS is a national study of health and aging that follows a cohort of adults ages 25–74 at baseline. The baseline sample was selected via national RDD (random digit dialing), with oversampling from five metropolitan areas in the US. In addition, siblings of individuals from the RDD sample and a national RDD sample of twin pairs were included. Non-institutionalized, English-speaking adults in the contiguous United States were eligible. Data collection consisted of a phone interview and self-administered questionnaires. Baseline response rates were 70% for the phone survey and 87% for the self-administered questionnaire for the RDD sample; 64% and 81% respectively for the sibling sample, and 60% and 92% respectively for the twin sample [19]. The overall longitudinal response rate was 75% [19]. We used data from waves one (1995–1996; “Time 1”) and two (2004–2005; “Time 2”). Additional information about MIDUS is available at <http://midus.wisc.edu/>.

Identification of caregivers

In the second wave of MIDUS, subjects were asked “During the last 12 months have you, yourself, given personal care for a period of one month or more to a family member or friend because of a physical or mental condition, illness, or disability?” Subjects who responded affirmatively and indicated that the caregiving began after 1996 (Time 1) were included in this study.

Of the 4,959 MIDUS II participants who responded to this item, 629 (12.7%) reported caregiving the past 12 months. Of these, 117 were missing data on the outcome variables and 145 were missing data on one or more other key variables, resulting in a final complete-case sample of 367 caregivers. Caregivers who were dropped due to missing data were younger; more likely to be non-white or have unreported race; and less likely to be college graduates; and had more chronic conditions at Time 2. They did not differ on other sociodemographic or caregiving factors.

Key variables

Dependent variables

Subjective well-being: As in Persoskie et. al [20], subjective well-being (SWB) was assessed in terms of 1) affect and 2) life satisfaction [21]. Specifically, participants reported their frequency of experiencing six positive (e.g., feeling cheerful) and six negative (e.g., feeling nervous) affective states during the past 30 days (5-point Likert scale for none of the time (1) to all the time (5)). Negative items were reverse coded and the mean of all 12 items was used as a composite score, as has been used previously (e.g., ref [20]). Cronbach’s alphas in the MIDUS sample at Time 1 [22] and Time 2 [23] respectively were 0.87 and 0.85 for negative affect and 0.91 and 0.90 for positive affect. Overall life satisfaction was assessed using participants’ reports of their quality of life in five domains: work, health,

their relationship with their spouse or partner, their relationship with their children, and life overall (10-point scale ranging from worst possible (1) to best possible (10)). This evaluation is consistent with the definition of life satisfaction offered by Luhman et. al: “evaluation of life overall...as well as of specific life domains (e.g., job satisfaction or marital satisfaction)” [21]. The items for relationship with spouse/partner and relationship with children were averaged to create single “relationships” item, and this was then averaged with the remaining three items to create an overall score. For participants with missing items (e.g., no spouse or partner), the overall score was created using the remaining items. This composite measure was calculated by MIDUS researchers [24] and has been used successfully as an assessment of overall life satisfaction in studies using MIDUS data (e.g., ref [20]). Cronbach’s alpha in the MIDUS sample was 0.67 at both timepoints [22; 23].

Psychological well-being: Psychological well-being (PWB) was measured using Ryff’s Psychological Well-Being Scale [25]. This well-validated scale consists of six, 3-item subscales: positive relations with others, self-acceptance, autonomy, personal growth, environmental mastery, and purpose in life. Respondents indicated how much they agreed or disagreed with each item (7-point Likert scale from strongly agree (1) to strongly disagree (7)). Negative items were reverse-coded and summed scores were created for all scales. Higher scores reflect more positive appraisals. Cronbach’s alphas in the MIDUS sample were 0.58, 0.59, 0.48, 0.55, 0.52, and 0.36 at Time 1 [22] and 0.63, 0.66, 0.45, 0.54, 0.54, and 0.29 at Time 2 [23].

Independent variables (Internal resources)—We evaluated five factors capturing an individual’s internal resources: sense of control (personal mastery), primary and secondary control strategies (persistence in goal setting, positive reappraisal, lowering aspirations), and social support seeking. These resources were selected align with our theoretical conceptualization of internal resources described above and were included in both the baseline and follow-up MIDUS questionnaires. *Personal mastery* (i.e., one’s sense of efficacy or effectiveness in carrying out goals) [26] was measured using 4 items assessing subjects’ sense of control. All items were measured on a 7-point Likert scale (strongly agree (1) to strongly agree (7)). Scales scores were calculated as the mean across each set of items for those who had valid values for at least half of the items. Higher scores indicated higher standing in each dimension. Cronbach’s alphas in the MIDUS sample were 0.70 at Time 1 [22] and 0.73 at Time 2 [23]. *Persistence in goal striving* (primary control; 5 items), *positive reappraisals* (secondary control; 4 items) and *lowering aspirations* (secondary; 5 items) were measured by asking participants how often they used various primary and secondary control strategies (e.g., “When I encounter problems, I don’t give up until I solve them”). All items were measured on a 4-point Likert scale (a lot (1) to not at all (4)). Scale scores were calculated as the mean across each set of items for those who had valid values for at least half of the items. Higher scores indicated higher standing in each dimension; Cronbach’s alphas were 0.77, 0.78 and 0.63 at Time 1 [22] and 0.78, 0.78, and 0.61 at Time 2 [23] respectively. *Social support seeking* was measured using two items (“I don’t like to ask others for help unless I have to” and “Asking others for help comes naturally to me” [reverse coded]) scored on a 4-point Likert scale (a lot (1) to not at all (4)). The scale score (mean) was calculated for those with data for at least one item [27]. Higher scores indicate greater

support seeking; Cronbach's alphas in the MIDUS sample were 0.68 at Time 1 [22] and 0.62 at Time 2 [23].

Internal resources were measured at baseline and follow-up. Following Ryff 2015 [16], we then categorized subjects into trajectory subgroups: increasing, decreasing, high-stable, mid-stable, or low stable by cross-tabulating quartiles for each scale at both baseline and followup. Figure 1 provides a visual depiction of how subjects were categorized. Social support seeking was coded slightly differently due to a floor effect: scores ≤ 1.5 (the median value) were considered "low" and all other scores were considered "high", and trajectories were categorized accordingly with the "mid-stable" category removed.

Covariates—Sociodemographic characteristics included age (years), gender, race (white vs nonwhite/unreported race), educational level (high school graduate or less, some college, or college graduate). In addition, subjects reported on a number of caregiving characteristics including: their relationship to the care recipient (care recipient is spouse, child, parent/in-law, or other); care recipient gender; whether they were still providing care at the time of the survey; whether they reside with the care recipient (co-resident: yes vs no/some of the time); number of caregiving tasks they performed (bathing, dressing, eating or going to the bathroom; getting around inside the house or going outside; shopping, cooking, housework or laundry; managing money, making phone calls, or taking medications; range 0–4), duration for which they had been caring for the care recipient (weeks), and hours per week of care. Personality characteristics (neuroticism, extraversion, openness to experience, conscientiousness, agreeableness, and agency measured at T1 using subjects' responses about how well 30 self-descriptive adjectives described them [27]; Cronbach's alpha on subscales ranged from 0.58–0.80 in the MIDUS RDD sample) were also included as covariates, as suggested by Ferrer-i-Carbonell and Frijters [28].

Statistical approach

Descriptive statistics were used to assess the sample (percentages; mean/standard deviation; histograms). Multi-collinearity was evaluated among the trajectory variables using Cramer's V scores from chi-squared tests, and among the continuous internal resource and well-being variables using Pearson's correlation. There was no collinearity among the internal resources (Cramer's V ranged from 0.09 to 0.31), among the well-being variables (r ranged from 0.23 to 0.61), or between the continuous internal resources variables and the well-being variables (r ranged from 0.01 to 0.45). The core multivariable linear regression models regressed each PWB and SWB outcome on the sociodemographic and caregiving-related covariates, and the six personality traits [28]; models used robust standard errors. Each internal resource was then added to these core models, for a total of 40 final regressions. We also tested including caregiver chronic health conditions as a control variable; as this did not impact the results, this variable was not included in the presented models. To assess the possibility of regression to the mean, dependent variables were examined with scatter plots of change score (T2 minus T1) by T1 score. If possible regression to the mean was observed (i.e., if subjects with particularly low scores at T1 were more likely to have an increase, and subjects with particularly high scores were more likely to have a decrease between T1 and T2), the subject's baseline value less the sample mean at baseline was included in the regression [29].

Effect sizes (ES) were evaluated using Cohen's d [30]. Given the large number of tests (40 models $\times$ 4 contrasts in each model), a Bonferroni correction was applied ($p = 0.0003$) when interpreting the p -values in the final regressions.

Sensitivity analyses—In order to assess whether unobserved heterogeneity may have impacted our results [28], fixed effects models were also run regression the difference score (T2 minus T1) in the dependent variables on the different score for the independent variables, and controlling for age and caregiving characteristics. P -values were interpreted after a Bonferroni correction ($p < 0.00125$).

RESULTS

Table 1 depicts the sociodemographic and caregiving characteristics of the informal caregivers. On average, caregivers were 56 years of age. The majority were female (65%), white (92%), and had at least some college education (70%). Almost half were caring for a parent or parent-in-law (46%) and lived with the care recipient (44%). On average, caregivers provided more than 20 hours of care per week, and had been providing care for nearly half a year.

Most caregivers reported stable internal resources from the pre-caregiving time-point to 10 years later (Figure 2). PWB and SWB were also largely stable over time (Table 2; Supplemental Figure 1): for example, only 8% reported a decline in life satisfaction and 6% reported a decline in affect, while almost 8% reported an increase in both SWB outcomes. The largest proportion reporting a decline was seen for autonomy (13%); while the largest group reporting an increase was for personal growth (10%).

Multivariable analyses indicated that increasing or high-stable levels of internal resources were strong predictors of PWB and SWB at time 2 (Table 3; Supplemental Figure 2). For *personal mastery*, increasing and/or high-stable levels were associated with 6 out of 8 well-being outcomes (positive relationships with others; self-acceptance; autonomy; personal growth; environmental mastery; affect) compared to low-stable. The effect sizes were moderate to large, ranging from 0.39 to 0.79. For *persistence*, increasing persistence was associated with better personal growth and environmental mastery ($ES = 0.96$ and 0.91) compared to low-stable. For *positive reappraisal*, mid-stable positive reappraisal was associated with better relationships with others ($ES = 0.54$) compared with low-stable levels. Increasing and high-stable levels of positive reappraisal were also associated with better affect ($ES = 0.63$ and 0.48) compared to low-stable levels. *Lowering aspirations* and *social support seeking* were not significantly associated with PWB or SWB outcomes after Bonferroni corrections.

Sensitivity Analyses

When we ran the analyses using fixed effects models (Table 4), an increase in *personal mastery* was associated with an increase in 6 out of the 8 well-being subscales (positive relationships with others, self-acceptance, autonomy, personal growth, environmental mastery and life satisfaction, ES for a one-point increasing in personal mastery ranging from 0.14 to 0.25). An increase in *persistence* was associated with an increase in 3 well-being

subscales (self-acceptance, autonomy, and environmental mastery, ES ranging from 0.37 to 0.39). An increase in *positive reappraisal* was associated with 5 well-being subscales (autonomy, personal growth, environmental mastery, affect and life satisfaction, ES ranging from 0.32 to 0.44). Changes in *lowering aspirations* and *support seeking* were not associated with changes in well-being.

DISCUSSION

Using a national dataset, this study assessed how internal resources (personal mastery; control strategies; social support seeking) changed over time for informal caregivers, and how such trajectories were associated with psychological and subjective well-being. While most caregivers showed stable internal resources over time, those with increasing and/or levels of personal mastery had better well-being (6 out of 8 subscales) than those with consistently low levels of internal resources. Those with increasing levels of persistence also had better well-being on two subscales (personal growth and mastery), and those with increasing or high-stable levels of positive reappraisal had better affect, compared to those with low-stable levels. This suggests that approaches for increasing or sustaining high levels of internal psychological resources throughout the course of (or even before) informal caregiving may be valuable tools for improving mental and emotional health outcomes of caregivers.

Research suggests that certain internal psychological resources play a critical role in buffering the negative effects of stress and life challenges, promoting resilience [5; 31]. For example, the ability to productively engage with goals – including persistence in goal striving as well as adjustment or disengagement with goals that are unattainable or not aligned with personal values – is associated with better psychological outcomes [32–35]. Self-efficacy has also been found to moderate the association between caregiving and poor mental health [13]. Our findings provide evidence that caregivers who experience increases in these resources, as well as those who have high levels both before and during caregiving, may have better well-being outcomes over time.

It is particularly interesting to note that most caregivers did not report changes in internal resources from before caregiving began. Furthermore, the fact that both increasing levels and high-stable levels of some internal resources were associated with some well-being scores suggests that where caregivers end up in their levels of internal resources may be more important than where they start. The start of caregiving, therefore, may be a window of opportunity for helping caregivers to acknowledge, utilize, and/or bolster their internal psychological resources. The stability of internal resources over time, however, suggests that sustainable changes in these factors do not often occur spontaneously, but rather likely require intentional effort. Mindfulness-based interventions, in particular, may increase positive reappraisal [36–38] and/or sense of control [39] and may plausibly bolster other internal resources as well.

Caregivers respond to their role in a variety of ways. One qualitative study of caregivers of seniors with chronic conditions identified both barriers and facilitators of resilience, with caregivers' sense of control, mastery, and support (or lack thereof) playing a role in both

categories [40]. Importantly, many caregivers report positive changes or growth due to caregiving, even while reporting high levels of burden [1–3]. Helping caregiver to bolster their internal resources, or leverage their existing strengths, may improve resilience or benefit finding, or otherwise mitigate some of the potentially deleterious effects of caregiving stress.

Despite the well-researched connection between social support and emotional outcomes for informal caregivers, this study did not find that a willingness to seek out social support was associated with well-being outcomes. This was contrary to our expectations. Caregivers may experience a desire to be (or appear) strong and self-sufficient, balancing the effects of a willingness to seek out help and support; although we theorized that such stoicism would be deleterious to caregivers, it may in some cases facilitate self-management [41]. It is also possible that the measure used in this study did not adequately capture the construct of interest. Indeed, the hypothesis driving this concept would naturally be moderated by the receipt of actual and useful social support (that is, caregivers who are inclined to seek out social support – rather than attempting to undertake everything on their own – would potentially have better outcomes precisely because they would be more likely to receive help and support); we could not fully investigate those effects here. Future research may wish to pursue this question more fully.

Importantly, targeting internal resources may be an effective way to improve well-being [42; 43]. For example, a review of strengths interventions concluded that they “demonstrate small but consistent well-being effects” [44]. Internal resources and strengths, however, have not been well-studied among informal caregivers despite theoretic support for their importance, particularly in a longitudinal context. Future research will be needed to determine whether helping caregivers bolster and best utilize their existing strengths, skills, and other internal resources can have a tangible impact on family outcomes. This information will be particularly relevant for clinical psychologists and other mental health professionals working with family caregivers, who may have the opportunity to work with caregivers on maintaining or enhancing their internal resources. In addition, the care recipient’s clinical care team may have an important role in supporting caregivers: one recent study indicated that providing caregivers with needed training on medical tasks, for example, has been shown to improve caregivers’ confidence [45], and may also positively impact self-efficacy, personal mastery, or other internal resources.

This study should be interpreted in the context of several potential limitations. First, informal caregivers were identified by self-report, and included all adults who reported being a caregiver in the past 12 months. Although this study was longitudinal, the findings are based on two time points approximately ten years apart, and no intervening measurements of internal resources or well-being outcomes were available. Response rates were good at both baseline and follow-up, but subjects with the greatest burden or the lowest levels of internal resources or well-being may have chosen not to participate, potentially contributing to selection bias. We conducted a complete case analysis, and subjects dropped due to missing data may represent a particularly vulnerable subset of caregivers. The sample size is also a limitation, especially in light of the multiple exploratory tests. We applied a Bonferroni correction in order to reduce the possibility of Type I error (i.e., false positive

findings); however, the small sample size and resultant limitations to statistical power leave open the possibility of Type II error (i.e., false negative findings). A post-hoc power analysis revealed that we had 80% power to detect small-medium effect sizes ($f^2=0.05$); nevertheless, non-significant findings should be interpreted with caution and all findings should be replicated in other, larger samples. We also include a lagged term in the models to control for the possibility of regression to the mean [29]. However, this could result in Nickell-Bias [46] and should be interpreted accordingly; models using the lagged term produced more conservative results (i.e., estimates closer to the null and larger p-values) than those without the lagged term. Finally, we were not able to account for some potentially relevant factors such as perceived stress and care recipient health condition. Nevertheless, this study provides novel evidence of how individuals change after informal caregiving begins, and how such changes are associated with well-being outcomes.

In conclusion, this study provides evidence that greater (and increasing) psychological resources, particularly personal master, may be associated with better well-being outcomes among informal caregivers. Informal caregivers may therefore be a prime target for testing, dissemination, and implementation of existing interventions targeting psychological resources. Future observational resource should confirm these findings, and randomized controlled trials should test whether existing or adapted interventions or trainings improve positive or prevent/mitigate negative caregiver outcomes, as well as assess the acceptability, feasibility, and sustainability of such approaches. Further targeted observational research should seek to understand how other elements of positive psychology may be related to better caregiver outcomes.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

The MIDUS 1 study (Midlife in the U.S.) was supported by the John D. and Catherine T. MacArthur Foundation Research Network on Successful Midlife Development. The MIDUS 2 research was supported by a grant from the National Institute on Aging (P01-AG020166) to conduct a longitudinal follow-up of the MIDUS 1 investigation.

BIBLIOGRAPHY

1. Lloyd J, Patterson T, Muers J. The positive aspects of caregiving in dementia: A critical review of the qualitative literature. *Dementia (London)*. 2016; 15(6):1534–1561. [PubMed: 25547208]
2. Cohen CA, Colantonio A, Vernich L. Positive aspects of caregiving: rounding out the caregiver experience. *Int J Geriatr Psychiatry*. 2002; 17(2):184–188. [PubMed: 11813283]
3. Litzelman K, Skinner HG, Gangnon RE, Nieto FJ, Malecki K, Witt WP. Role of global stress in the health-related quality of life of caregivers: evidence from the Survey of the Health of Wisconsin. *Qual Life Res*. 2014; 23(5):1569–1578. [PubMed: 24322907]
4. Pinquart M, Sorensen S. Differences between caregivers and noncaregivers in psychological health and physical health: a meta-analysis. *Psychol Aging*. 2003; 18(2):250–267. [PubMed: 12825775]
5. Cummins RA, Wooden M. Personal Resilience in Times of Crisis: The Implications of SWB Homeostasis and Set-Points. *J Happiness Stud*. 2014; 15(1):223–235.
6. Luthans F, Avolio BJ, Avey JB, Norman SM. Positive psychological capital: Measurement and relationship with performance and satisfaction. *Personnel Psychology*. 2007; 60(3):541–572.

7. Yi-Frazier JP, Smith RE, Vitaliano PP, Yi JC, Mai S, Hillman M, Weinger K. A Person-Focused Analysis of Resilience Resources and Coping in Diabetes Patients. *Stress Health*. 2010; 26(1):51–60. [PubMed: 20526415]
8. Masten AS. Resilience in developing systems: Progress and promise as the fourth wave rises. *Development and Psychopathology*. 2007; 19(3):921–930. [PubMed: 17705908]
9. Andrykowski MA, Lykins E, Floyd A. Psychological health in cancer survivors. *Semin Oncol Nurs*. 2008; 24(3):193–201. [PubMed: 18687265]
10. Pearlin LI, Mullan JT, Semple SJ, Skaff MM. Caregiving and the Stress Process – an Overview of Concepts and Their Measures. *Gerontologist*. 1990; 30(5):583–594. [PubMed: 2276631]
11. Goode KT, Haley WE, Roth DL, Ford GR. Predicting longitudinal changes in caregiver physical and mental health: a stress process model. *Health Psychol*. 1998; 17(2):190–198. [PubMed: 9548710]
12. O'Rourke N, Kupferschmidt AL, Claxton A, Smith JZ, Chappell N, Beattie BL. Psychological resilience predicts depressive symptoms among spouses of persons with Alzheimer disease over time. *Aging & Mental Health*. 2010; 14(8):984–993. [PubMed: 21069604]
13. Hajek A, Konig HH. Informal Caregiving and Subjective Well-Being: Evidence of a Population-Based Longitudinal Study of Older Adults in Germany. *J Am Med Dir Assoc*. 2016; 17(4):300–305. [PubMed: 26705001]
14. Joshanloo M. Revisiting the Empirical Distinction Between Hedonic and Eudaimonic Aspects of Well-Being Using Exploratory Structural Equation Modeling. *J Happiness Stud*. 2016; 17(5): 2023–2036.
15. Henderson LW, Knight T. Integrating the hedonic and eudaimonic perspectives to more comprehensively understand wellbeing and pathways to wellbeing. *International Journal of Wellbeing*. 2012; 2(3)
16. Ryff CD, Radler BT, Friedman EM. Persistent Psychological Well-being Predicts Improved Self-Rated Health Over 9–10 Years: Longitudinal Evidence from MIDUS. *Health Psychol Open*. 2015; 2(2) 2055102915601582.
17. Friedman EM, Ryff CD. Living well with medical comorbidities: a biopsychosocial perspective. *J Gerontol B Psychol Sci Soc Sci*. 2012; 67(5):535–544. [PubMed: 22377799]
18. Brim, OG., Ryff, CD., Kessler, RC. *The MIDUS National Survey: An Overview*. University of Chicago Press; 2004.
19. MIDUS. *Mid-life in the United States: A National Study of Health and Well-being*. UW Institute on Aging; 2007.
20. Persoskie A, Ferrer RA, Nelson WL, Klein WM. Precancer risk perceptions predict postcancer subjective well-being. *Health Psychol*. 2014; 33(9):1023–1032. [PubMed: 24588629]
21. Luhmann M, Hofmann W, Eid M, Lucas RE. Subjective well-being and adaptation to life events: a meta-analysis. *J Pers Soc Psychol*. 2012; 102(3):592–615. [PubMed: 22059843]
22. UW-Madison Institute on Aging. *National Survey of Midlife Development in the United States (MIDUS), 1995–1996: Documentation of Scales and Constructed Variables in MIDUS 1*. Ann Arbor, MI: Inter-University Consortium for Political and Social Research; 2009.
23. UW-Madison Institute on Aging. *Documentation of Psychosocial Constructs and Composite Variables in MIDUS II Project 1*. Ann Arbor, MI: Inter-University Consortium for Political and Social Research; 2010.
24. Brim, O., Baltes, P., Bumpass, L., Cleary, P., Featherman, D., Hazzard, W., Shweder, RA. *National Survey of Midlife Development in the United States (MIDUS), 1995–1996*. ICPSR02760–v8. Ann Arbor, MI: Inter-university Consortium for Political and Social Research [distributor]; 2011.
25. Ryff CD. Happiness is everything, or is it? Explorations on the meaning of psychological well-being. *Journal of personality and social psychology*. 1989; 57(6):1069.
26. Lachman ME, Weaver SL. The sense of control as a moderator of social class differences in health and well-being. *Journal of Personality and Social Psychology*. 1998; 74(3):763–773. [PubMed: 9523418]
27. Lachman, ME., Weaver, SL. *The Midlife Development Inventory (MIDI) personality scales: Scale construction and scoring*. Waltham, MA: Brandeis University; 1997.

28. Ferrer-i-Carbonell A, Frijters P. How important is methodology for the estimates of the determinants of happiness? *The Economic Journal*. 2004; 114(497):641–659.
29. Barnett AG, van der Pols JC, Dobson AJ. Regression to the mean: what it is and how to deal with it. *Int J Epidemiol*. 2005; 34(1):215–220. [PubMed: 15333621]
30. Cohen, J. *Statistical power analysis for the behavioral sciences*. Hillsdale, NJ: Lawrence Erlbaum Associates, Inc; 1988.
31. Ryff CD. Psychological well-being revisited: advances in the science and practice of eudaimonia. *Psychother Psychosom*. 2014; 83(1):10–28. [PubMed: 24281296]
32. Wrosch C, Scheier MF, Miller GE, Schulz R, Carver CS. Adaptive self-regulation of unattainable goals: goal disengagement, goal reengagement, and subjective well-being. *Pers Soc Psychol Bull*. 2003; 29(12):1494–1508. [PubMed: 15018681]
33. Wrosch C, Scheier MF, Carver CS, Schulz R. The Importance of Goal Disengagement in Adaptive Self-Regulation: When Giving Up is Beneficial. *Self and Identity*. 2003; 2(1):1–20.
34. Rasmussen HN, Wrosch C, Scheier MF, Carver CS. Self-regulation processes and health: the importance of optimism and goal adjustment. *J Pers*. 2006; 74(6):1721–1747. [PubMed: 17083664]
35. Wrosch C, Heckhausen J, Lachman ME. Primary and secondary control strategies for managing health and financial stress across adulthood. *Psychol Aging*. 2000; 15(3):387–399. [PubMed: 11014704]
36. Rayan A, Ahmad M. Effectiveness of mindfulness-based interventions on quality of life and positive reappraisal coping among parents of children with autism spectrum disorder. *Research in Developmental Disabilities*. 2016; 55:185–196. [PubMed: 27107368]
37. Gerzina HA, Porfeli EJ. Mindfulness as a Predictor of Positive Reappraisal and Burnout in Standardized Patients. *Teaching and Learning in Medicine*. 2012; 24(4):309–314. [PubMed: 23035997]
38. Garland E, Gaylord S, Park J. The Role of Mindfulness in Positive Reappraisal. *Explore-NY*. 2009; 5(1):37–44. [PubMed: 19114262]
39. Astin JA. Stress reduction through mindfulness meditation – Effects on psychological symptomatology, sense of control, and spiritual experiences. *Psychother Psychosom*. 1997; 66(2): 97–106. [PubMed: 9097338]
40. Bremault-Phillips S, Parmar J, Johnson M, Huhn A, Mann A, Tian V, Sacrey LAR. The voices of family caregivers of seniors with chronic conditions: a window into their experience using a qualitative design. *Springerplus*. 2016; 5(1):1–11. [PubMed: 26759740]
41. Toms GR, Quinn C, Anderson DE, Clare L. Help Yourself: Perspectives on Self-Management From People With Dementia and Their Caregivers. *Qual Health Res*. 2015; 25(1):87–98. [PubMed: 25192761]
42. Bolier L, Haverman M, Westerhof GJ, Riper H, Smit F, Bohlmeijer E. Positive psychology interventions: a meta-analysis of randomized controlled studies. *BMC Public Health*. 2013; 13(1): 119. [PubMed: 23390882]
43. Friedman EM, Ruini C, Foy R, Jaros L, Sampson H, Ryff CD. Lighten UP! A community-based group intervention to promote psychological well-being in older adults. *Aging Ment Health*. 2015:1–7.
44. Quinlan D, Swain N, Vella-Brodrick DA. Character Strengths Interventions: Building on What We Know for Improved Outcomes. *J Happiness Stud*. 2012; 13(6):1145–1163.
45. Mollica, M., Litzelman, K., Rowland, J., Kent, E. Training informal cancer caregivers in medical tasks is associated with improved confidence and reduced burden. U. Review. , editor. 2017.
46. Nickell S. Biases in dynamic models with fixed effects. *Econometrica: Journal of the Econometric Society*. 1981:1417–1426.

Figure 1.

Categorization of trajectories from Time 1 to Time 2.

Note: Q1–Q4 refers to quartiles of each variable, from lowest to highest.

Figure 2.

Internal resource trajectories among family caregivers.

*No mid-stable category due to floor effect.

The bars indicate the proportion of caregivers who experienced decreasing, low stable, mid stable, high stable, or increasing levels of five internal resources. Data are from the Midlife in the US Survey (MIDUS).

Table 1

Characteristics of caregivers in the MIDUS

	Current Caregiver ^a n=367 Mean (SD) or %
<i>Sociodemographics</i>	
Age	56.19 (11.13)
Gender	
Male	34.6
Female	65.4
Race	
White	91.6
Non-white/unreported	8.5
Education	
High school grad or less	30.25
Some college	27.79
College graduate	41.96
<i>Caregiving Characteristics</i>	
Relationship to CR	
spouse	16.89
child	8.17
parent/in-law	46.32
other	28.61
CR Gender	
Male	38.96
Female	61.04
Still giving care	
Yes	53.68
No	46.32
Cores ident status	
Yes	44.14
No/some of the time	55.86
Caregiving tasks	
0–1	17.71
2	25.61
3	23.98
4	32.7
Duration (weeks)	24.9 (19.7)
Hours/week	22.9 (25.7)

^aReporting caregiving at Time 2 (2004–2005) that began after 1996

Table 2

Distribution of psychological and subjective well-being among caregivers in the MIDUS

	Current Caregiver^a n=367 Mean (SD) or %
<i>Psychological (Eudaimonic) Well-Being</i>	
Positive Relationship to Others	
T1	16.57 (3.88)
T2	17.29 (3.83)
Self Acceptance	
T1	16.54 (3.48)
T2	16.36 (3.89)
Autonomy	
T1	16.34 (3.31)
T2	16.68 (3.17)
Personal Growth	
T1	18.15 (3.07)
T2	17.46 (3.28)
Environmental Mastery	
T1	16.19 (3.57)
T2	16.63 (3.36)
Purpose in Life	
T1	16.75 (3.46)
T2	16.27 (3.35)
<i>Subjective (Hedonic) Well-being</i>	
Life Satisfaction	
T1	7.72 (1.23)
T2	7.74 (1.25)
Affect	
T1	3.86 (0.64)
T2	3.9 (0.58)

T1: Time 1 (1995–1996); T2: Time 2 (2004–2005)

Table 3
Multivariable regressions of the associations between internal resources and well-being among informal caregivers

	Psychological (Eudaimonic) Well-being										Subjective (Hedonic) Well-being									
	Positive Relationships with Others					Personal Growth					Environmental Mastery					Purpose				
	Beta	SE	ES	p-value	R ²	Beta	SE	ES	p-value	R ²	Beta	SE	ES	p-value	R ²	Beta	SE	ES	p-value	R ²
<i>Internal Resources</i>																				
Personal Mastery					0.41					0.39					0.33					0.25
Low Stable																				
Decreasing	0.33	0.66	0.09	0.6155																
Increasing	<i>3.03</i> [#]	0.53	0.79	<.0001																
Mid Stable	1.44	0.42	0.38	0.0008																
High Stable	1.37	0.47	0.36	0.0037																
Persistence					0.37					0.33					0.25					0.38
Low Stable																				
Decreasing	0.00	0.85	0.00	0.9978																
Increasing	1.85	0.75	0.48	0.0141																
Mid Stable	0.75	0.42	0.20	0.0731																
High Stable	0.91	0.54	0.24	0.0926																
Positive Reappraisal					0.39					0.34					0.26					0.37
Low Stable																				
Decreasing	0.26	0.65	0.07	0.692																
Increasing	1.50	0.63	0.39	0.0175																
Mid Stable	<i>2.07</i> [#]	0.45	0.54	<.0001																
High Stable	1.30	0.55	0.34	0.0175																
Lower Aspirations					0.37					0.31					0.25					0.36
Low Stable																				
Decreasing	-0.03	0.64	-0.01	0.961																
Increasing	-1.61	0.67	-0.42	0.017																
Mid Stable	-0.33	0.39	-0.09	0.3965																
High Stable	-1.24	0.45	-0.32	0.0059																
Seeking Social Support ^a					0.37					0.32					0.23					0.36
Low Stable																				
Decreasing	0.30	0.50	0.08	0.5499																
Increasing	0.68	0.49	0.18	0.1708																
High Stable	0.97	0.39	0.25	0.0144																

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

SE: Standard Error; ES: Effect size (Cohen’s d);

Models control for: age, gender, race, education, relationship to the care recipient, care recipient gender, current caregiving status, co-resident status, number of caregiving tasks, duration of caregiving(weeks), number of hours of care per week, time 1 value of outcome less the sample mean at time 1, and personality traits.

^aSelf-sufficiency coded without “mid-stable” category due to ceiling effect

* statistically different from low-stable after Bonferroni correction (p = 0.0003)

Table 4

Associations between internal resources and well-being among informal caregivers, fixed effects models

	Psychological (eudaimonic) well-being										Subjective (hedonic) well-being																													
	Positive relationships with others					Self-acceptance					Autonomy					Personal Growth					Environmental mastery					Purpose					Affect					Life satisfaction				
	Beta	SE	ES	p value	R ²	Beta	SE	ES	p value	R ²	Beta	SE	ES	p value	R ²	Beta	SE	ES	p value	R ²	Beta	SE	ES	p value	R ²	Beta	SE	ES	p value	R ²	Beta	SE	ES	p value	R ²					
Personal mastery	0.60*	0.17	0.16	0.0003	0.04	0.77*	0.17	0.20	<0.0001	0.08	0.80	0.16*	0.25	<0.0001	0.06	0.88*	0.14	0.27	<0.0001	0.08	0.96*	0.16	0.29	0.08	0.1081	0.04	0.08	0.02	0.14	0.0013	0.05	0.29*	0.05	0.24	<0.0001	0.10				
Persistence	0.98	0.39	0.26	0.0129	0.03	1.35*	0.39	0.35	0.0006	0.06	1.31	0.37*	0.41	0.0005	0.04	1.01	0.33	0.31	0.0021	0.04	1.50*	0.38	0.45	<0.0001	0.05	0.73	0.39	0.22	0.0582	0.04	0.13	0.06	0.22	0.0212	0.02	0.33	0.13	0.27	0.0087	0.05
Positive reappraisal	0.60	0.33	0.16	0.0664	0.03	0.91	0.32	0.23	0.0048	0.05	1.01	0.31*	0.32	0.0010	0.04	1.37*	0.27	0.42	<0.0001	0.06	1.47*	0.31	0.44	<0.0001	0.07	0.44	0.32	0.13	0.1698	0.04	0.19*	0.05	0.33	<0.0001	0.06	0.47*	0.10	0.38	<0.0001	0.08
Lowering aspirations	-0.55	0.33	-0.14	0.0908	0.02	-0.52	0.33	-0.13	0.1173	0.04	-0.66	0.31	-0.21	0.0367	0.02	-0.55	0.27	-0.17	0.0445	0.02	-0.87	0.31	-0.26	0.0059	0.04	-0.11	0.32	-0.03	0.7372	0.03	-0.11	0.05	-0.20	0.0125	0.04	-0.17	0.11	-0.13	0.1117	0.04
Support seeking	0.43	0.23	0.11	0.0653	0.03	0.43	0.24	0.11	0.0694	0.04	0.24	0.23	0.08	0.2909	0.02	0.28	0.20	0.09	0.1462	0.02	0.32	0.23	0.10	0.1617	0.03	0.21	0.23	0.06	0.3541	0.03	0.02	0.03	0.04	0.4912	0.03	0.09	0.08	0.07	0.2377	0.04

Fixed effects regression controls for time-invariant heterogeneity; Models also controlled for age (years, T2-T1), relationship to the care recipient, care recipient gender, current caregiving status, co-resident status, number of caregiving tasks, duration of caregiving (weeks), and number of hours of care per week

* Statistically significant association after Bonferroni correction ($p < 0.00125$)