Neurodevelopment: The Impact of Nutrition and Inflammation During Early to Middle Childhood in Low-Resource Settings

Chandy C. John, MD, MS,^a Maureen M. Black, PhD,^{b,c,d} Charles A. Nelson III, PhD^{e,f,g}

abstract

The early to middle childhood years are a critical period for child neurodevelopment. Nutritional deficiencies, infection, and inflammation are major contributors to impaired child neurodevelopment in these years, particularly in low-resource settings. This review identifies global research priorities relating to nutrition, infection, and inflammation in early to middle childhood neurodevelopment. The research priority areas identified include: (1) assessment of how nutrition, infection, or inflammation in the preconception, prenatal, and infancy periods (or interventions in these periods) affect function in early to middle childhood; (2) assessment of whether effects of nutritional interventions vary by poverty or inflammation; (3) determination of the feasibility of preschool- and school-based integrated nutritional interventions; (4) improved assessment of the epidemiology of infection- and inflammation-related neurodevelopmental impairment (NDI); (5) identification of mechanisms through which infection causes NDI; (6) identification of noninfectious causes of inflammation-related NDI and interventions for causes already identified (eg, environmental factors); and (7) studies on the effects of interactions between nutritional, infectious, and inflammatory factors on neurodevelopment in early to middle childhood. Areas of emerging importance that require additional study include the effects of maternal Zika virus infection, childhood environmental enteropathy, and alterations in the child's microbiome on neurodevelopment in early to middle childhood. Research in these key areas will be critical to the development of interventions to optimize the neurodevelopmental potential of children worldwide in the early to middle childhood years.

^aRyan White Center for Pediatric Infectious Disease and Global Health, Indiana University School of Medicine, Indianapolis, Indiana; ^bDepartments of ^bPediatrics and ^cDepartment of Epidemiology, University of Maryland School of Medicine, Baltimore, Maryland; ^dRTI International, Research Triangle Park, North Carolina; ^eLaboratories of Cognitive Neuroscience, Boston Children's Hospital, Boston, Massachusetts; ^fDepartment of Pediatrics, Harvard Medical School, Boston, Massachusetts; and ^gDepartment of Human Development, Harvard Graduate School of Education, Cambridge, Massachusetts

Dr John was a panelist at the original *Eunice Kennedy Shriver* National Institute of Child Health and Human Development scientific meeting, served as the lead author for the paper, organized the writing team, drafted the initial manuscript, incorporated edits from the additional authors and editors, and finalized the manuscript; Drs Black and Nelson were panelists at the original NICHD scientific meeting, contributed to the writing of the initial manuscript, and reviewed and revised subsequent versions of the manuscript; and all authors approved the final manuscript as submitted and are accountable for all aspects of the work.

DOI: 10.1542/peds.2016-2828H

Accepted for publication Dec 21, 2016

Address correspondence to Chandy C. John, MD, MS, Ryan White Center for Pediatric Infectious Disease and Global Health, Indiana University School of Medicine, 1044 W. Walnut St, R4 402D, Indianapolis, IN 46202. E-mail: chjohn@iu.edu

PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275).

Copyright © 2017 by the American Academy of Pediatrics

FINANCIAL DISCLOSURE: The authors have indicated they have no financial relationships relevant to this article to disclose.

FUNDING: This supplement was supported by the *Eunice Kennedy Shriver* National Institute of Child Health and Human Development (NICHD) at the United States National Institutes of Health (NIH).

POTENTIAL CONFLICT OF INTEREST: The authors have indicated they have no potential conflicts of interest to disclose.

The foundations of health and well-being stem from the interplay between genes and environment that begin as early as conception and extend through early and middle childhood (ages 3–12 years).^{1,2} Children with adequate opportunities for early care and learning, nutrition, and protection from infectious and noninfectious causes of inflammation have the best chances of thriving. In contrast, children raised in adverse conditions, characterized by poverty, limited access to opportunities for early learning and responsive caregiving, nutritional deprivation, and infectious and noninfectious threats, are at risk for negative health and social outcomes throughout their life course, beginning with neurodevelopmental delays and extending to poor academic functioning, chronic diseases, mental illness, and lack of economic productivity.²

Global estimates are that over one-third of children <5 years of age in low- and middle-income countries (LMIC) are at risk for not reaching their developmental potential, based on poverty and stunting.³ However, without consideration of infectious and noninfectious causes of inflammation or nutritional deficiencies that do not result in stunting, the problem is likely to be much larger. Children who lag behind developmental expectations before age 5 are at risk for subsequent academic and socioemotional problems as they approach middle childhood and primary school. Primary school prepares children for higher education and for the economic, interpersonal, social, and civic responsibilities of adulthood. Universal primary education is a central global goal, as illustrated by its inclusion in the Millennium Development Goals (Goal 2) and in the current Sustainable Development Goals (Goal 4).

Primary school enrollment has increased dramatically over the

past 2 decades, especially in LMICs.⁴ Based on World Bank data, for the lowest-income countries, the primary school gross enrollment rate (ie, the number of children enrolled as a percentage of the eligible population) for both boys and girls increased from 21% between 1996 and 2000 to 67% between 2011 and 2015.⁵ However, access and retention continue to be concerns, and the 2015 Global Monitoring Report indicates that 58 million primary school-age children were out of school in 2015.⁴ In addition to low enrollment and attendance related to crisis and conflict, distance, and denial of access, problems related to children's health and nutrition interfere not only with children's participation in primary school, but also their ability to learn.

KEY NEURODEVELOPMENTAL CONSIDERATIONS DURING EARLY TO MIDDLE CHILDHOOD

Although the groundwork for brain development begins just a few weeks after conception and continues through the first postnatal years of life, experiences during middle and late childhood can still exert a tremendous influence on changes in synapse number and myelin integrity, because both processes continue to develop well beyond the first years of life.⁶ It is during this time that nutrition and inflammation, for example, can have a large impact on myelination and that exposure to multiple forms of adversity can impact the development of learning and memory circuits that reside in the medial temporal lobe (eg, hippocampus) and the development of executive functions (subserved by a distributed network of regions in the prefrontal cortex). Importantly, the biggest shift in both gray and white matter occurs as children make the transition to puberty.⁷ During this time, gray matter declines first followed by white matter, a process that starts right before puberty and

may continue for another decade. This process is advanced in girls as compared with boys by 1 to 2 years; consequently, girls enter these events earlier and conclude sooner than do boys.^{8,9} Thus, even if children traverse the infancy period, a critical time for many aspects of brain development, various environmental factors can still impact brain development during later periods (see Fig 1).

Furthermore, our understanding of the developmental sequence, as well as the genetic and experiential contributions to brain development, have grown tremendously over the past few decades. So, too, has our knowledge of more functional aspects of brain development in the first 2+ years of life, largely due to advances in imaging tools suitable for young infants, such as EEG, event-related potential, and functional near-infrared spectroscopy.^{10,11} However, there are enormous gaps in our knowledge of structural and functional brain development during the early and middle childhood periods, for several reasons.

First, with the exception of the increasingly rare primate models of development, rodent models do a relatively poor job of simulating human brain development during the preschool and elementary school years. Second, the use of EEG and event-related potential methods (functional near-infrared spectroscopy is still a relatively recent addition to our imaging armamentarium) remains stubbornly difficult in younger children, mostly due to their inherent resistance to sitting still for long periods of time coupled with motor limitations that make overt responding difficult. Similarly, structural and functional MRI and, to a limited extent, magnetoencephalography have led to important new discoveries about brain development during middle childhood, whereas these tools are enormously difficult to use

FIGURE 1

Relationships among individual and environmental risk factors, inflammation, nutrition, and neurodevelopment for school-age children in low-resource settings. TB, tuberculosis.

in children <5 years of age. Finally, significant attention has been paid to malnutrition early in life, with far less attention paid to inflammatory disease and the effects these 2 forms of adversity exert on older children, particularly those with an early developmental history of malnutrition or illness.

THE IMPACT OF NUTRITION ON NEURODEVELOPMENT DURING EARLY TO MIDDLE CHILDHOOD

Food insecurity, or inconsistent access and availability to a diverse, safe, and nutritionally adequate diet for an active lifestyle, a primary cause of nutritional deprivation, is associated with poverty and may impact both the quantity

and quality of available food, contributing to lack of macro- and micronutrients. Macronutrients include energy, carbohydrates, and fat, whereas micronutrients include small quantities of vitamins and minerals, such as iron, zinc, and vitamin B₁₂, required for specific physiologic functions. At least 4 micronutrients have been associated with neurodevelopment during early to middle childhood: iodine, zinc, vitamin B₁₂, and iron.

School-age children in food-insecure households are at risk for academic and behavioral problems, which may be related to a lack of nutrients and to stress associated with an inconsistent food supply. Neuroscientific evidence has documented the adverse and

pervasive role of poverty on early brain development, specifically on psychoimmunological functioning and self-regulation,¹² but there has been little attention to the combined effects of poverty and nutritional deficiencies on neurodevelopment.

Childhood stunting, closely associated with poverty and a major threat to child development, is often used as a population-based indicator to compare nutritional adequacy across countries. A recent meta-analysis on the association between linear growth and children's development included 68 reports from 29 LMICs¹³ and found that early growth restriction is associated with lower cognitive scores throughout childhood. Findings related to

socioemotional development were less clear, primarily due to the small number of studies and differences in the measurement of socioemotional development across differing ages.

At least 3 mechanisms may link early stunting to development during early to middle childhood: (1) biological insults that disrupt early brain development, (2) delayed motor skills that may disrupt the exploration associated with cognitive development, and (3) reduced expectations from parents and peers, based on short stature. The long-term consequences of stunting extend beyond childhood into adulthood and include lower height, less schooling, and reduced economic productivity.^{14,15} Studies into the next generation have shown associations between first-generation stunting and offspring birth size¹⁵ and performance on cognitive assessments.¹⁶

Although most of the research linking nutrition with inflammation and neurodevelopment focuses on undernutrition, overnutrition, such as childhood obesity, has also become a global concern. Obesity begins early in life, increases during childhood and adolescence, and has been associated with impaired cognition. Possible mechanisms include inflammation, oxidative stress, decreased motor performance associated with a degraded musculoskeletal system, and alterations in brain structure, leptin/insulin regulation, cerebrovascular function, and/or blood-barrier function.¹⁷

Micronutrients

Iodine

Iodine deficiency disrupts production of thyroid hormones, thyroxine, and triiodothyronine, which are necessary for neurogenesis, neuronal migration, synaptogenesis, and myelination.^{18,19} Severe deficiency can cause goiter and intellectual disability and even mild/moderate

deficiencies are associated with intellectual delays that can disrupt academic functioning.¹⁸ Iodine supplementation trials appear to partially reduce the negative effects of iodine deficiency, but ensuring adequate maternal iodine status prenatally can prevent iodine deficiency during the period of rapid brain development.¹⁸

Zinc

Zinc plays a critical role in central nervous system (CNS) development, specifically neuron formation, migration, and synapse generation.²⁰ Zinc is found in high concentrations in the hippocampus, cerebellum, prefrontal cortex, cortex, and limbic system; in addition, animal studies have documented zinc's role in neurodevelopment.^{19,21} Evidence from human supplementation trials has found positive associations between prenatal or infant zinc supplementation and motor development, including processing speed and motor aspects of attention, but not with measures of cognitive processing.²¹ One supplementation trial conducted among school-age children in China found beneficial effects of zinc supplementation on neuropsychological functioning when zinc was combined with other micronutrients,²² but trials of zinc supplementation alone among school-age children conducted in Canada and Guatemala found no effects on cognition or academic performance.²³

Vitamin B₁₂

Vitamin B₁₂ plays an important role in multiple fetal developmental processes, including neurodevelopment through DNA methylation and epinephrine synthesis, along with methionine synthesis.²⁴ The fetus receives vitamin B₁₂ through the placenta from maternal vitamin B₁₂.²⁵ After birth, vitamin B₁₂ is supplied through animal source food. Vitamin B₁₂ deficiency has been associated with

demyelination, which can result in delayed cognitive development, and with gastric inflammatory states, possibly indicating an inflammation-induced autoimmune process blocking intrinsic factor, and thus preventing vitamin B₁₂ absorption.²⁶

Although longitudinal studies have shown a relationship between prenatal vitamin B₁₂ deficiency and school-age cognitive functioning involving the frontal lobe (perceptual tracking and simple sequencing tasks) and temporal lobe (short-term memory),²⁵ more research is needed to understand the role of vitamin B₁₂ in neurodevelopment and links to academic performance.

Iron

Iron deficiency is the most common nutrient deficiency globally; because of its role in hemoglobin synthesis, adequate iron is essential for oxygen delivery to all tissues, especially the brain. Iron is necessary for myelination, frontal cortex, and basal ganglia development.²⁷ In toddlerhood, iron deficiency has been associated with impaired socioemotional behavior, including shyness, wariness, and low responsivity.²⁸ Iron-deficiency anemia in early infancy is a risk factor for impaired mental and motor development and has been associated with long-term negative functional consequences.¹⁹

Although the associations between iron deficiency and infant and child development are strong, nutritional interventions early in life, when children's rate of growth is rapid and nutritional demands are high, have been met with limited and inconsistent success either in alleviating nutritional deficiencies or in promoting early development.^{29,30} Possible explanations for the limited findings are: (1) the origins of iron deficiency may occur before conception or prenatally, and postnatal interventions are too late; (2) interventions may be most

TABLE 1 Gaps in Knowledge Related to Nutrition and Neurodevelopment During Early to Middle Childhood

Problem or Question	Studies Needed
What is the impact of early nutritional deficiencies on neurodevelopment and brain function during early to middle childhood?	<ol style="list-style-type: none"> 1. Identify sensitive periods in terms of timing of intervention. 2. Determine how much plasticity there is in recovery from early nutritional deficiencies and whether interventions can be developed for implementation after a sensitive period has closed.
Individual differences in timing of nutritional intervention and recovery from early deprivation.	Identify genes that may play a role in individual differences; develop new intervention strategies.
Are there lasting effects of nutritional interventions implemented during the preconception, prenatal, infancy, and preschool periods on functioning during early to middle childhood?	<ol style="list-style-type: none"> 1. Longitudinal follow-up through middle childhood of interventions delivered at earlier time periods. 2. Measure nutritional status, brain structure and function, and academic performance.
Do the effects of early nutritional interventions vary by poverty or inflammation?	Examine how poverty and inflammation modify the above relationships.
Are preschool-/school-based integrated interventions that include nutrition and early child development (eg, self-regulation) feasible and effective?	Conduct a trial of integrated interventions in preschool/school settings and measure impact on nutritional status, executive function, and academic performance.
Can school feeding programs be implemented that are effective in promoting school performance while not increasing stigma?	Evaluate strategies to introduce school feeding programs that are effective and examine perceptions of stigma.

effective when targeted to deficient populations; or (3) iron deficiency often occurs in the context of other micronutrient deficiencies, leading to recommendations to focus on multiple micronutrient deficiencies.³¹

Multiple Micronutrients

Micronutrient deficiencies often cooccur, especially when micronutrients are derived from common sources, such as animal source foods. Recent evidence suggests that interventions supplementing multiple micronutrients are more beneficial than single micronutrient trials for child development.¹⁹ Reviews of multiple micronutrient supplementation trials administered prenatally through the school-age years on neurodevelopmental performance have yielded mixed findings. Although at least 1 trial has found an impact of prenatal iron–folic acid supplementation on school-age neurodevelopmental performance,³² others have reported reductions in anemia, with no changes in neurodevelopmental performance. However, few intervention trials have followed children into school-age years. Two reviews of multiple micronutrient supplementation³³ or fortification³⁴ introduced during school-age years have reported

inconsistent findings related to cognition and academic performance.

In addition to adequate nutrition, neurodevelopment is dependent on environmental opportunities for responsive caregiving and early learning. Integrated interventions that combine nutrients with opportunities for responsive caregiving and early learning have been recommended,³⁵ and evidence suggests that trials that include both early education and nutrition are more likely to result in cognitive benefits than single-intervention trials,^{36,37} but relatively few integrated trials or programs have been evaluated systematically.³⁸ Table 1 provides a summary of key gaps in knowledge related to nutrition and neurodevelopment in early to middle child development.

THE IMPACT OF INFLAMMATION ON NEURODEVELOPMENT DURING EARLY TO MIDDLE CHILDHOOD

Fetal exposure to maternal infection and inflammation may have long-term consequences well into early to middle childhood, although these effects are more often studied during the infancy and adolescent periods. This study during infancy and adolescent periods may be because the critical period of

brain development from 0 to 3 years and the interactive effects of hormonal changes in adolescence focus attention on these age groups. Yet, the available evidence suggests a potential for a profound effect of maternal infection on neurodevelopment in early to middle childhood.

Symptomatic congenital cytomegalovirus infection is among the best described congenital infections associated with cognitive impairment in children, starting in infancy and affecting children of all ages.³⁹ In contrast, asymptomatic cytomegalovirus infection is far more common than symptomatic infection, but is associated with cognitive impairment only in school-age,^{40,41} and not older,⁴⁰ children. Limited data from children exposed in utero to maternal HIV infection and antiretroviral drugs, but uninfected with HIV themselves,^{42,43} and from experimental models of malaria exposure in utero⁴⁴ suggest that children in early to middle childhood are affected by maternal infection or inflammation. Numerous studies document the effects of vertically acquired HIV infection in multiple areas of neurodevelopment in school-age children, including overall cognitive ability,^{45,46} memory, perception,⁴⁷ motor function,^{48,49}

executive function,⁵⁰ and language skills.^{49,51} Maternal helminth infection shows effects on children at 1 year of age that may well extend into early to middle childhood,⁵² but these effects are so far unstudied. Findings like this highlight the need for longitudinal studies of the effects of maternal or congenital infections on child neurodevelopment across the age span. The most striking recent example of congenital infection affecting the CNS is Zika virus infection,⁵³ and the full effects of this newly emerging infection on child neurodevelopment remain to be described.⁵⁴

Many infections that occur in children can occur across ages, from the neonatal period to infancy and early to middle childhood. The most common of these infections worldwide, such as helminth infection, diarrheal disease, and malaria, are typically most frequent in children <3 years of age, but the effects of infection in the younger years often lasts into early or middle childhood, and infection does continue, albeit with less frequency and severity, through the latter period.

Diarrheal disease in the first 2 years of life has also been associated with cognitive impairment in early to middle childhood in a number of studies, notably from Brazil^{55–57} and Bangladesh.⁵⁸ Other infections with well-documented effects on neurodevelopment in school-age children include intestinal helminths,^{59–62} schistosomiasis,^{59,63} and malaria.^{64–68} Other than HIV, none of these infections directly infects the CNS; they all act through indirect mechanisms that are still poorly described, yet likely involve inflammation. Because of their high incidence, these infections likely affect neurodevelopment in many more children than infections that directly infect the CNS, such as HIV, bacterial meningitis, and viral encephalitis.

The studies cited above are examples of the substantial body of data showing the effects of infection and inflammation, at various stages, on neurodevelopment during early to middle childhood. However, a recent review noted that, for most infections that affect the CNS, reliable estimates of the incidence of resulting neurodevelopmental impairment (NDI) are not currently available.⁶⁹ Without this information, it is difficult to determine the burden of infection-related NDI in early to middle childhood or implement appropriate interventions. To obtain accurate estimates, better diagnostics (ideally low-cost and point-of-care), more in-depth surveillance, and epidemiologic studies are required.

In addition, studies of the pathogenesis of how infection leads to NDI are urgently needed, because these mechanisms remain largely unknown or poorly characterized, and are not limited solely to inflammation. Infection may lead to NDI through direct CNS injury by the infectious pathogen or, for example, through pathways, such as sequestration and endothelial activation in cerebral malaria,⁷⁰ that may involve inflammation as 1 component but may not be traditionally defined as inflammation. Thus, an understanding of pathogen mechanisms of direct injury and the full range of the host response, including pathways not related to inflammation or of which inflammation is only a component, is another important area for future research. Such research will require better markers of inflammation, tissue injury, and host response, including noninvasive surrogate markers of CNS infection, inflammation, and injury. Current systemic assessments have clear limitations, because they may not reflect CNS findings, and, indeed, in studies in which CNS markers have been assessed through cerebrospinal fluid or brain biopsy, systemic

findings often differ from CNS findings.

Inflammation may also affect child neurodevelopment through noninfectious causes. There is an emerging literature on environmental causes that have been associated with impaired child neurodevelopment, including maternal or child tobacco, pollutant, and pesticide exposure.^{71–74} A few studies have documented an association of environmental pollutants with systemic inflammation^{71,72} and impaired neurodevelopment, suggesting that inflammation could be a mechanism by which these factors affect neurodevelopment. However, there are no studies to date that actually define the pathways by which environmental exposures cause impaired neurodevelopment, highlighting another key area for future research. Table 2 summarizes key gaps in knowledge related to inflammation and neurodevelopment during early to middle childhood.

THE INTERACTION OF NUTRITION, INFLAMMATION, NEURODEVELOPMENT, AND OTHER INFLUENCING FACTORS DURING EARLY TO MIDDLE CHILDHOOD

Interactions between nutrition, infection, inflammation, and environmental factors are clearly important in NDI in early to middle childhood. Micronutrient deficiency, infection, and inflammation interact in complex ways. That is, micronutrient deficiency may predispose to or protect from infection and increased or decreased inflammation, but, conversely, inflammation and infection may lead to micronutrient deficiency. For example, iron deficiency appears to provide fairly strong protection against clinical malaria,⁷⁵ yet malaria-associated inflammation and upregulation of hepcidin likely leads to decreased iron bioavailability and functional iron deficiency. Both

TABLE 2 Gaps in Knowledge Related to Inflammation and Neurodevelopment During Early to Middle Childhood

Problem or Question	Studies Needed
What is the burden of NDI from specific infections or inflammation in children in early to middle childhood?	<ol style="list-style-type: none"> 1. Low-cost, point-of-care, easy-to-use, accurate diagnostics for infection and inflammation. 2. Define the epidemiology of infection- and inflammation-related NDI.
How does infection or inflammation during pregnancy or in early infancy affect a child's neurodevelopment in early to middle childhood and beyond?	<ol style="list-style-type: none"> 1. Longitudinal studies (including birth cohort studies) of infection- and inflammation-related NDI that follow children up through early to middle childhood and beyond. 2. Define correspondence validity of tests that measure specific areas of neurodevelopment in age groups that span infancy to adolescence.
How does infection cause NDI? What are the roles of direct injury, inflammation, or other pathways in infection-related NDI?	<ol style="list-style-type: none"> 1. Define pathogenesis of infection-related NDI. 2. Determine the contributions of direct pathogen injury, infection-related inflammation, and other infection-related pathways (eg, endothelial activation) to NDI. 3. Define noninvasive surrogate markers of brain infection, inflammation, and injury. 4. Assess interaction of these pathways in causing NDI.
What are the noninfectious causes of inflammation in early to middle childhood?	<ol style="list-style-type: none"> 1. Assess inflammation longitudinally in children. 2. Determine environmental and other causes of inflammation in early to middle childhood.
How do environmental factors lead to impaired NDI?	<ol style="list-style-type: none"> 1. Define how environmental factors (eg, pollutants, tobacco use, toxins) relate to NDI. 2. Assess environmental contributions to inflammation-related and noninflammation-related NDI. 3. Define the specific areas of the brain/cognition/behavior affected by particular environmental factors.
Can prevention of environmental, infectious, or inflammatory factors decrease NDI?	<ol style="list-style-type: none"> 1. Well-designed clinical trials to assess whether interventions that reduce known environmental, infectious, or inflammatory risk factors for NDI can reduce NDI in early to middle childhood. 2. Inclusion of NDI as an endpoint in studies aiming to reduce infection, environmental pollution, or toxins.
Can adjunctive treatments decrease infection-related NDI?	Well-designed clinical trials of interventions that appear to block infection-related pathways leading to NDI.

malaria and iron deficiency can lead to neurodevelopmental and behavioral impairment, and children in malaria-endemic areas may also have inflammation due to other causes (eg, helminth infection) that can also lead to NDI. With these complicated relationships, untangling how each factor relates to the other and how each contributes to NDI can be difficult, but is critical if interventions in endemic areas are to successfully prevent NDI without increasing the risk of infection.

Anemia of inflammation, often due to infectious causes, is another understudied but likely frequent cause of NDI in children in low-resource settings. A study in the Philippines showed that children with anemia of inflammation, who did not have iron deficiency by standard biomarkers, had lower cognitive scores than those without

anemia. The authors hypothesized that the worsened outcomes were due to decreased delivery of iron to end organs, including the brain.⁷⁶ This study illustrates complex interactions and the importance of understanding mechanisms of disease, because the children with anemia of inflammation were not iron deficient by standard biomarker measurements, but were likely functionally iron deficient in key end organs, including the brain. Research that contributes to a better definition of the infection-and inflammation-associated pathways that lead to NDI, and how these pathways affect micronutrients, provides the best opportunity to determine the various contributions of these factors to NDI. In addition, such research is needed to help plan for interventions that improve neurodevelopment without disturbing the delicate homeostasis of these relationships in

such a way that the risk of infection, inflammation, or micronutrient deficiency is increased.

Two emerging areas of interest that highlight interactive effects throughout childhood are the microbiome and environmental enteropathy. Inflammation and malnutrition can alter the microbiome, and, conversely, changes in the microbiome can affect both nutrition and systemic inflammation.⁷⁷ Changes in the microbiome have been hypothesized to potentially influence changes in child neurodevelopment and behavior through the “microbiome–gut–brain axis.”⁷⁸ Similarly, environmental enteropathy, involving intestinal inflammation without overt diarrhea, also seems to affect the risk of both malnutrition and impaired child neurodevelopment.⁷⁹ Alterations in the microbiome are

TABLE 3 Gaps in Knowledge Related to Interactions of Nutrition, Inflammation, Neurodevelopment, and Other Influencing Factors During Early to Middle Childhood

Problem or Question	Studies Needed
How do interactions of micronutrient deficiency, infection, and inflammation in a pregnant woman affect neurodevelopment in her child in early to middle childhood?	<ol style="list-style-type: none"> 1. Assess micronutrient deficiency, infection, and inflammation in pregnant women at all stages of pregnancy. 2. Determine the effects of individual factors and interactions between these factors on fetal growth, placental pathology, and long-term neurodevelopment in the affected child in early to middle childhood. 3. Example of a specific study: how do malaria and helminth infection affect development of iron deficiency; what is the role of inflammation in this interaction; how does iron deficiency affect risk of malaria and helminth infection; and how do the 4 factors together affect risk of NDI in endemic areas?
Why and how does environmental enteropathy develop?	Determine the factors and mechanisms that lead to environmental enteropathy.
How does environmental enteropathy contribute to NDI?	Define the pathways by which environmental enteropathy leads to NDI.
How do changes in the microbiome relate to environmental enteropathy and vice versa?	Determine the effects of changes in the microbiome on environmental enteropathy and of development of environmental enteropathy on the microbiome.
What are the interactions between micronutrient deficiency, the microbiome and malnutrition, and how do these relate to NDI?	Comprehensively assess how micronutrient deficiency, the microbiome, and malnutrition interact during pregnancy in the mother, and during infancy and early to middle childhood in the child, lead to early to middle childhood NDI.
How do interactions between micronutrient deficiency, infection, and inflammation in early to middle childhood affect neurodevelopment?	<ol style="list-style-type: none"> 1. Determine how micronutrient deficiency, infection, and inflammation interact from infancy through early to middle childhood. 2. Assess how interactions between these 3 factors in infancy or early to middle childhood contribute to NDI.
How does impaired neurodevelopment affect risk of infection, inflammation, and environmental toxins or pollutants?	Study the risks of infection and infectious or inflammatory processes and the risk of environmental toxin or pollutant exposure in children across the spectrum of neurodevelopment.
What is the relation between structural and functional brain development during middle childhood?	<ol style="list-style-type: none"> 1. Better methods to measure both structural and functional brain development among preschool and early school-age children. 2. Examine the impact of nutrition and inflammation on brain structure and function during middle childhood and the concordance between brain structure and function.

likely to occur in environmental enteropathy and indeed might be part of the pathogenesis of this process. Future research studies should assess how both factors may affect child neurodevelopment in early to middle childhood, how each condition affects the other, and interactions with inflammation, malnutrition, and micronutrient deficiency. Water, sanitation, and hygiene programs may have a role in decreasing environmental enteropathy and have been proposed as a component of early childhood development programs.⁸⁰ Given the complex interactions of microbes, nutritional factors, and inflammation, the components of these programs will need careful consideration and study for optimal effectiveness. Finally, children with disabilities and neurodevelopmental and neurocognitive issues experience

numerous challenges in early to middle childhood. Their access to educational opportunities is often limited, nutritional problems may occur if they have swallowing, motor, or cognitive problems that affect their ability or desire to eat nutritious foods, and they may be neglected in favor of children without impairment. These problems may become particularly acute in settings where resources are limited. These challenges can make children with NDI more susceptible to nutritional deficiencies and infectious/inflammatory exposures than children without NDI. Thus, although the primary focus of this article has been on the effects of infection, inflammation, and nutrition on neurodevelopment, neurodevelopment may also tie back and affect these factors. Table 3 summarizes current gaps in

knowledge related to interactions between nutrition, inflammation, neurodevelopment, and other influencing factors during early to middle childhood.

IMPLICATIONS FOR RESEARCH, PROGRAM, AND POLICY DEVELOPMENT

Primary prevention of infectious and noninfectious inflammation and nutritional deficiencies beginning before conception and extending through childhood would protect children from these threats to early brain development. Children who have avoided early threats or who have emerged with relatively few negative consequences have often been protected by responsive caregiving.⁸¹ In a striking example, at age 12, children raised in Romanian orphanages and randomized at 18 to 24 months

of age to high-quality foster care showed stress responses (marked by cortisol and parasympathetic nervous system reactivity) that differed from children randomized to remain in the orphanages and approached responses expected of noninstitutionalized children.⁸² The foster care group also continued to show better brain electrical activity, measured by EEG, as long as they remained in high-quality foster care.¹¹ This neuroscientific evidence provides support for early intervention programs that focus on responsive caregiving.

Furthermore, preprimary educational programs that include well-qualified personnel working with both parents and children can have beneficial effects on children's cognitive performance⁸³ and prepare children for primary school.⁸⁴ Recent evidence has illustrated the beneficial effects of incorporating neurodevelopmentally-based teaching methods into preprimary classes, particularly among children from low-income families.⁸⁵ Skills often categorized as self-regulation (eg, the ability to attend, to regulate emotions and behavior, and to participate in goal-directed activities) are positively related not only to school-age learning and academic performance, but also to adult well-being.^{86,87} Self-regulation is influenced both by underlying neural and physiologic systems and by environmental expectations, suggesting that it is potentially responsive to interventions. In a trial in which self-regulation activities were systematically incorporated into kindergarten classes (eg, cooperative activities that promote socioemotional and cognitive development, reflective "talk," make-believe play), children experienced improvements in measures of self-regulation, in neuroendocrine functioning, and in academic measures of reading and mathematics.⁸⁸ The academic

benefits were retained through first grade, suggesting that exposure to self-regulation-promoting activities during the middle childhood years may promote subsequent neurodevelopment by enhancing self-regulation and possibly mitigate some of the negative effects of early nutritional deficiencies and inflammation. School feeding programs have been implemented in many resource-limited settings to reduce hunger, promote attendance, and enhance academic performance. A recent review found that school feeding programs have a positive impact on energy intake, micronutrient status, school enrollment, and attendance, with inconsistent effects on growth, cognition, and academic achievement.⁸⁹ The authors postulate that these mixed findings may be attributable to a multitude of factors, including differences in the objectives and methodologies used, quality and quantity of food served, duration of the interventions, degree of malnourishment, and severity of comorbid conditions, such as helminth infection. These findings demonstrate the complexity of assessing neurodevelopmental outcomes in the context of multiple cooccurring and variable risk factors across the life course.

Most studies of NDI are association studies that cannot prove causation. There are multiple reasons why there are few longitudinal studies of children, a key barrier being the high cost of following the large cohorts required. For example, disease states, such as cerebral malaria, are not common in any given cohort, even though, across malaria-endemic areas, hundreds of thousands of children are affected annually. However, for nutritional, infectious/inflammatory, or environmental factors that occur with sufficient frequency, birth cohort studies provide an opportunity to better define the causes of NDI, and the

interactions between factors leading to this impairment, than can be achieved with cross-sectional studies. For example, despite multiple studies showing an association between uncomplicated or severe malaria and cognitive impairment, a large cluster randomized study of intermittent screening and treatment of malaria in school-age children showed that this intervention did not improve educational achievement and was associated with a negative effect in spelling and arithmetic scores.⁹⁰ This study suggests that earlier intervention may be required, or that malaria may be a proxy for other causes of NDI. The failure of randomized controlled trials to detect a treatment effect highlights the need for additional research on the mechanisms of NDI and the importance of interventions for optimal neurodevelopment in early and middle childhood.

Responsive caregiving and early learning may mitigate some of the neuropsychological effects of adversity, emphasizing the importance of interventions to children's health and well-being. In spite of the positive evaluations on the impact of early learning and responsive caregiving interventions,^{37,38,91,92} few intervention options are available for children and families. Future recommendations include strategies to integrate, monitor, and sustain effective interventions for young children.

ABBREVIATIONS

CNS: central nervous system
LMIC: low- and middle-income countries
NDI: neurodevelopmental impairment

REFERENCES

1. Pongcharoen T, Ramakrishnan U, DiGirolamo AM, et al. Influence of prenatal and postnatal growth on

- intellectual functioning in school-aged children. *Arch Pediatr Adolesc Med*. 2012;166(5):411–416
2. Shonkoff JP, Garner AS; Committee on Psychosocial Aspects of Child and Family Health; Committee on Early Childhood, Adoption, and Dependent Care; Section on Developmental and Behavioral Pediatrics. The lifelong effects of early childhood adversity and toxic stress. *Pediatrics*. 2012;129(1). Available at: www.pediatrics.org/cgi/content/full/129/1/e232
3. Grantham-McGregor S, Cheung YB, Cueto S, Glewwe P, Richter L, Strupp B; International Child Development Steering Group. Developmental potential in the first 5 years for children in developing countries. *Lancet*. 2007;369(9555):60–70
4. The United Nations Educational, Scientific and Cultural Organization (UNESCO). Education for all global monitoring report 2000-2015: achievements and challenges. Available at: <http://unesdoc.unesco.org/images/0023/002322/232205e.pdf>. Accessed June 6, 2016
5. The World Bank. World development report 2016: digital dividends. Available at: <http://documents.worldbank.org/curated/en/896971468194972881/pdf/102725-PUB-Replacement-PUBLIC.pdf>. Accessed June 6, 2016
6. Nowakowsk RS, Hayes NL. Overview of early brain development: linking genetics to brain structure. In: Baker W, Benasich A, Fitch RH, eds. *Developmental Dyslexia: Early Precursors, Neurobehavioral Markers, and Biological Substrates*. Baltimore, MD: Brookes Publishing Co; 2012
7. Sowell ER, Trauner DA, Gamst A, Jernigan TL. Development of cortical and subcortical brain structures in childhood and adolescence: a structural MRI study. *Dev Med Child Neurol*. 2002;44(1):4–16
8. Lenroot RK, Gogtay N, Greenstein DK, et al. Sexual dimorphism of brain developmental trajectories during childhood and adolescence. *Neuroimage*. 2007;36(4):1065–1073
9. Sowell ER, Peterson BS, Thompson PM, Welcome SE, Henkenius AL, Toga AW. Mapping cortical change across the human life span. *Nat Neurosci*. 2003;6(3):309–315
10. Nelson CA III, McCleery JP. Use of event-related potentials in the study of typical and atypical development. *J Am Acad Child Adolesc Psychiatry*. 2008;47(11):1252–1261
11. Vanderwert RE, Zeanah CH, Fox NA, Nelson CA III. Normalization of EEG activity among previously institutionalized children placed into foster care: A 12-year follow-up of the Bucharest Early Intervention Project. *Dev Cogn Neurosci*. 2016;17:68–75
12. Johnson SB, Riis JL, Noble KG. State of the art review: poverty and the developing brain. *Pediatrics*. 2016;137(4):e20153075
13. Sudfeld CR, McCoy DC, Danaei G, et al. Linear growth and child development in low- and middle-income countries: a meta-analysis. *Pediatrics*. 2015;135(5). Available at: www.pediatrics.org/cgi/content/full/135/5/e1266
14. Hoddinott J, Behrman JR, Maluccio JA, et al. Adult consequences of growth failure in early childhood. *Am J Clin Nutr*. 2013;98(5):1170–1178
15. Victora CG, Adair L, Fall C, et al; Maternal and Child Undernutrition Study Group. Maternal and child undernutrition: consequences for adult health and human capital. *Lancet*. 2008;371(9609):340–357
16. Walker SP, Chang SM, Wright A, Osmond C, Grantham-McGregor SM. Early childhood stunting is associated with lower developmental levels in the subsequent generation of children. *J Nutr*. 2015;145(4):823–828
17. Wang C, Chan JS, Ren L, Yan JH. Obesity reduces cognitive and motor functions across the lifespan. *Neural Plast*. 2016;2016:2473081
18. Bougma K, Aboud FE, Harding KB, Marquis GS. Iodine and mental development of children 5 years old and under: a systematic review and meta-analysis. *Nutrients*. 2013;5(4):1384–1416
19. Prado EL, Dewey KG. Nutrition and brain development in early life. *Nutr Rev*. 2014;72(4):267–284
20. Anjos T, Altmäe S, Emmett P, et al; NUTRIMENTHE Research Group. Nutrition and neurodevelopment in children: focus on NUTRIMENTHE project. *Eur J Nutr*. 2013;52(8):1825–1842
21. Colombo J, Zavaleta N, Kannass KN, et al. Zinc supplementation sustained normative neurodevelopment in a randomized, controlled trial of Peruvian infants aged 6-18 months. *J Nutr*. 2014;144(8):1298–1305
22. Sandstead HH, Penland JG, Alcock NW, et al. Effects of repletion with zinc and other micronutrients on neuropsychologic performance and growth of Chinese children. *Am J Clin Nutr*. 1998;68(suppl 2):470S–475S
23. Black MM. Zinc deficiency and cognitive development. In: Benton D, ed. *Lifetime nutritional influences on cognition, behaviour and psychiatric illness*. Cambridge, UK: Woodhead Publishing; 2011:79–93
24. Pepper MR, Black MM. B12 in fetal development. *Semin Cell Dev Biol*. 2011;22(6):619–623
25. Bhate V, Deshpande S, Bhat D, et al. Vitamin B12 status of pregnant Indian women and cognitive function in their 9-year-old children. *Food Nutr Bull*. 2008;29(4):249–254
26. Black MM, Baqui AH, Zaman K, et al. Depressive symptoms among rural Bangladeshi mothers: implications for infant development. *J Child Psychol Psychiatry*. 2007;48(8):764–772
27. Wachs TD, Georgieff M, Cusick S, McEwen BS. Issues in the timing of integrated early interventions: contributions from nutrition, neuroscience, and psychological research. *Ann N Y Acad Sci*. 2014;1308:89–106
28. Lozoff B, Georgieff MK. Iron deficiency and brain development. *Semin Pediatr Neurol*. 2006;13(3):158–165
29. Bhutta ZA, Das JK, Rizvi A, et al; Lancet Nutrition Interventions Review Group; Maternal and Child Nutrition Study Group. Evidence-based interventions for improvement of maternal and child nutrition: what can be done and at what cost? *Lancet*. 2013;382(9890):452–477
30. Pasricha SR, Hayes E, Kalumba K, Biggs BA. Effect of daily iron supplementation on health in children aged 4-23 months: a systematic review

- and meta-analysis of randomised controlled trials. *Lancet Glob Health*. 2013;1(2):e77–e86
31. Allen LH. Multiple micronutrients in pregnancy and lactation: an overview. *Am J Clin Nutr*. 2005;81(5):1206S–1212S
32. Christian P, Murray-Kolb LE, Khatry SK, et al. Prenatal micronutrient supplementation and intellectual and motor function in early school-aged children in Nepal. *JAMA*. 2010;304(24):2716–2723
33. Eilander A, Gera T, Sachdev HS, et al. Multiple micronutrient supplementation for improving cognitive performance in children: systematic review of randomized controlled trials. *Am J Clin Nutr*. 2010;91(1):115–130
34. Best C, Neufingerl N, Del Rosso JM, Transler C, van den Briel T, Osendarp S. Can multi-micronutrient food fortification improve the micronutrient status, growth, health, and cognition of schoolchildren? A systematic review. *Nutr Rev*. 2011;69(4):186–204
35. Black MM, Dewey KG. Promoting equity through integrated early child development and nutrition interventions. *Ann N Y Acad Sci*. 2014;1308(1):1–10
36. Nores M, Barnett WS. Benefits of early childhood interventions across the world: (under) Investing in the very young. *Econ Educ Rev*. 2010;29(2):271–282
37. Rao N, Sun J, Pearson V, et al. Is something better than nothing? An evaluation of early childhood programs in Cambodia. *Child Dev*. 2012;83(3):864–876
38. Grantham-McGregor SM, Fernald LC, Kagawa RM, Walker S. Effects of integrated child development and nutrition interventions on child development and nutritional status. *Ann N Y Acad Sci*. 2014;1308:11–32
39. Noyola DE, Demmler GJ, Williamson WD, et al; Congenital CMV Longitudinal Study Group. Cytomegalovirus urinary excretion and long term outcome in children with congenital cytomegalovirus infection. *Pediatr Infect Dis J*. 2000;19(6):505–510
40. Temple RO, Pass RF, Boll TJ. Neuropsychological functioning in patients with asymptomatic congenital cytomegalovirus infection. *J Dev Behav Pediatr*. 2000;21(6):417–422
41. Zhang XW, Li F, Yu XW, Shi XW, Shi J, Zhang JP. Physical and intellectual development in children with asymptomatic congenital cytomegalovirus infection: a longitudinal cohort study in Qinba mountain area, China. *J Clin Virol*. 2007;40(3):180–185
42. Alimenti A, Forbes JC, Oberlander TF, et al. A prospective controlled study of neurodevelopment in HIV-uninfected children exposed to combination antiretroviral drugs in pregnancy. *Pediatrics*. 2006;118(4). Available at: www.pediatrics.org/cgi/content/full/118/4/e1139
43. Le Doaré K, Bland R, Newell ML. Neurodevelopment in children born to HIV-infected mothers by infection and treatment status. *Pediatrics*. 2012;130(5). Available at: www.pediatrics.org/cgi/content/full/130/5/e1326
44. McDonald CR, Cahill LS, Ho KT, et al. Experimental malaria in pregnancy induces neurocognitive injury in uninfected offspring via a C5a-C5a receptor dependent pathway. *PLoS Pathog*. 2015;11(9):e1005140
45. Crowell CS, Huo Y, Tassiopoulos K, et al; PACTG 219C Study Team and the Pediatric HIVAIDS Cohort Study (PHACS). Early viral suppression improves neurocognitive outcomes in HIV-infected children. *AIDS*. 2015;29(3):295–304
46. Puthanakit T, Aupibul L, Louthrenoo O, et al. Poor cognitive functioning of school-aged children in thailand with perinatally acquired HIV infection taking antiretroviral therapy. *AIDS Patient Care STDS*. 2010;24(3):141–146
47. Fundarò C, Miccinesi N, Baldieri NF, Genovese O, Rendeli C, Segni G. Cognitive impairment in school-age children with asymptomatic HIV infection. *AIDS Patient Care STDS*. 1998;12(2):135–140
48. Blanchette N, Smith ML, King S, Fernandes-Penney A, Read S. Cognitive development in school-age children with vertically transmitted HIV infection. *Dev Neuropsychol*. 2002;21(3):223–241
49. Whitehead N, Potterton J, Coovadia A. The neurodevelopment of HIV-infected infants on HAART compared to HIV-exposed but uninfected infants. *AIDS Care*. 2014;26(4):497–504
50. Koekkoek S, de Sonnevile LM, Wolfs TF, Licht R, Geelen SP. Neurocognitive function profile in HIV-infected school-age children. *Eur J Paediatr Neurol*. 2008;12(4):290–297
51. Tardieu M, Mayaux MJ, Seibel N, et al. Cognitive assessment of school-age children infected with maternally transmitted human immunodeficiency virus type 1. *J Pediatr*. 1995;126(3):375–379
52. Mireku MO, Boivin MJ, Davidson LL, et al. Impact of helminth infection during pregnancy on cognitive and motor functions of one-year-old children. *PLoS Negl Trop Dis*. 2015;9(3):e0003463
53. Panchaud A, Stojanov M, Ammerdorffer A, Vouga M, Baud D. Emerging role of zika virus in adverse fetal and neonatal outcomes. *Clin Microbiol Rev*. 2016;29(3):659–694
54. França GV, Schuler-Faccini L, Oliveira WK, et al. Congenital Zika virus syndrome in Brazil: a case series of the first 1501 livebirths with complete investigation. *Lancet*. 2016;388(10047):891–897
55. Guerrant DL, Moore SR, Lima AA, Patrick PD, Schorling JB, Guerrant RL. Association of early childhood diarrhea and cryptosporidiosis with impaired physical fitness and cognitive function four-seven years later in a poor urban community in northeast Brazil. *Am J Trop Med Hyg*. 1999;61(5):707–713
56. Lorntz B, Soares AM, Moore SR, et al. Early childhood diarrhea predicts impaired school performance. *Pediatr Infect Dis J*. 2006;25(6):513–520
57. Niehaus MD, Moore SR, Patrick PD, et al. Early childhood diarrhea is associated with diminished cognitive function 4 to 7 years later in children in a northeast Brazilian shantytown. *Am J Trop Med Hyg*. 2002;66(5):590–593
58. Tarleton JL, Haque R, Mondal D, Shu J, Farr BM, Petri WA Jr. Cognitive effects of diarrhea, malnutrition, and Entamoeba histolytica infection on school age children in Dhaka,

- Bangladesh. *Am J Trop Med Hyg.* 2006;74(3):475–481
59. Ezeamama AE, Friedman JF, Acosta LP, et al. Helminth infection and cognitive impairment among Filipino children. *Am J Trop Med Hyg.* 2005;72(5):540–548
 60. Kvalsvig JD, Cooppan RM, Connolly KJ. The effects of parasite infections on cognitive processes in children. *Ann Trop Med Parasitol.* 1991;85(5):551–568
 61. Nokes C, Grantham-McGregor SM, Sawyer AW, Cooper ES, Bundy DA. Parasitic helminth infection and cognitive function in school children. *Proc Biol Sci.* 1992;247(1319):77–81
 62. Sakti H, Nokes C, Hertanto WS, et al. Evidence for an association between hookworm infection and cognitive function in Indonesian school children. *Trop Med Int Health.* 1999;4(5):322–334
 63. Nokes C, McGarvey ST, Shiue L, et al. Evidence for an improvement in cognitive function following treatment of *Schistosoma japonicum* infection in Chinese primary schoolchildren. *Am J Trop Med Hyg.* 1999;60(4):556–565
 64. Bangirana P, Opoka RO, Boivin MJ, et al. Severe malarial anemia is associated with long-term neurocognitive impairment. *Clin Infect Dis.* 2014;59(3):336–344
 65. Carter JA, Ross AJ, Neville BG, et al. Developmental impairments following severe falciparum malaria in children. *Trop Med Int Health.* 2005;10(1):3–10
 66. Holding PA, Stevenson J, Peshu N, Marsh K. Cognitive sequelae of severe malaria with impaired consciousness. *Trans R Soc Trop Med Hyg.* 1999;93(5):529–534
 67. John CC, Bangirana P, Byarugaba J, et al. Cerebral malaria in children is associated with long-term cognitive impairment. *Pediatrics.* 2008;122(1):e92–e99
 68. Nankabirwa J, Wandera B, Kiwanuka N, Staedke SG, Kanya MR, Brooker SJ. Asymptomatic *Plasmodium* infection and cognition among primary schoolchildren in a high malaria transmission setting in Uganda. *Am J Trop Med Hyg.* 2013;88(6):1102–1108
 69. John CC, Carabin H, Montano SM, Bangirana P, Zunt JR, Peterson PK. Global research priorities for infections that affect the nervous system. *Nature.* 2015;527(7578):S178–S186
 70. Idro R, Marsh K, John CC, Newton CR. Cerebral malaria: mechanisms of brain injury and strategies for improved neurocognitive outcome. *Pediatr Res.* 2010;68(4):267–274
 71. Calderón-Garcidueñas L, Engle R, Mora-Tiscareño A, et al. Exposure to severe urban air pollution influences cognitive outcomes, brain volume and systemic inflammation in clinically healthy children. *Brain Cogn.* 2011;77(3):345–355
 72. Calderón-Garcidueñas L, Macías-Parra M, Hoffmann HJ, et al. Immunotoxicity and environment: immunodysregulation and systemic inflammation in children. *Toxicol Pathol.* 2009;37(2):161–169
 73. Perera FP, Rauh V, Whyatt RM, et al. A summary of recent findings on birth outcomes and developmental effects of prenatal ETS, PAH, and pesticide exposures. *Neurotoxicology.* 2005;26(4):573–587
 74. Rauh VA, Whyatt RM, Garfinkel R, et al. Developmental effects of exposure to environmental tobacco smoke and material hardship among inner-city children. *Neurotoxicol Teratol.* 2004;26(3):373–385
 75. Gwamaka M, Kurtis JD, Sorensen BE, et al. Iron deficiency protects against severe *Plasmodium falciparum* malaria and death in young children. *Clin Infect Dis.* 2012;54(8):1137–1144
 76. Olson CL, Acosta LP, Hochberg NS, et al. Anemia of inflammation is related to cognitive impairment among children in Leyte, the Philippines. *PLoS Negl Trop Dis.* 2009;3(10):e533
 77. Ahmed T, Auble D, Berkley JA, et al. An evolving perspective about the origins of childhood undernutrition and nutritional interventions that includes the gut microbiome. *Ann N Y Acad Sci.* 2014;1332:22–38
 78. Keunen K, van Elburg RM, van Bel F, Benders MJ. Impact of nutrition on brain development and its neuroprotective implications following preterm birth. *Pediatr Res.* 2015;77(1-2):148–155
 79. Gilmartin AA, Petri WA Jr. Exploring the role of environmental enteropathy in malnutrition, infant development and oral vaccine response. *Philos Trans R Soc Lond B Biol Sci.* 2015;370(1671):20140143
 80. Ngure FM, Reid BM, Humphrey JH, Mbuya MN, Peltó G, Stoltzfus RJ. Water, sanitation, and hygiene (WASH), environmental enteropathy, nutrition, and early child development: making the links. *Ann N Y Acad Sci.* 2014;1308:118–128
 81. Luby J, Belden A, Botteron K, et al. The effects of poverty on childhood brain development: the mediating effect of caregiving and stressful life events. *JAMA Pediatr.* 2013;167(12):1135–1142
 82. McLaughlin KA, Sheridan MA, Tibu F, Fox NA, Zeanah CH, Nelson CA III. Causal effects of the early caregiving environment on development of stress response systems in children. *Proc Natl Acad Sci USA.* 2015;112(18):5637–5642
 83. Rao N, Sun J, Wong J, et al. Education Rigorous Literature Review: Early childhood development and cognitive development in developing countries. Available at: <http://eppi.ioe.ac.uk/cms/Portals/0/PDF%20reviews%20and%20summaries/ECD%202014%20Rao%20report.pdf?ver=2014-10-02-145634-017>. Accessed June 6, 2016
 84. Berlinski S, Schady N, eds; Inter-American Development Bank. The early years: Child well-being and the role of public policy. Available at: www.iadb.org/en/research-and-data/dia-2015-the-early-years-child-well-being-and-the-role-of-public-policy,18093.html. Accessed June 6, 2016
 85. Blair C, Raver CC, Berry DJ; Family Life Project Investigators. Two approaches to estimating the effect of parenting on the development of executive function in early childhood. *Dev Psychol.* 2014;50(2):554–565
 86. Moffitt TE, Arseneault L, Belsky D, et al. A gradient of childhood self-control predicts health, wealth, and public safety. *Proc Natl Acad Sci USA.* 2011;108(7):2693–2698
 87. Reynolds AJ, Temple JA. Cost-effective early childhood development programs

- from preschool to third grade. *Annu Rev Clin Psychol.* 2008;4: 109–139
88. Blair C, Raver CC. Closing the achievement gap through modification of neurocognitive and neuroendocrine function: results from a cluster randomized controlled trial of an innovative approach to the education of children in kindergarten. *PLoS One.* 2014;9(11):e112393
 89. Jomaa LH, McDonnell E, Probart C. School feeding programs in developing countries: impacts on children's health and educational outcomes. *Nutr Rev.* 2011;69(2):83–98
 90. Halliday KE, Okello G, Turner EL, et al. Impact of intermittent screening and treatment for malaria among school children in Kenya: a cluster randomised trial. *PLoS Med.* 2014;11(1):e1001594
 91. Engle PL, Black MM, Behrman JR, et al; International Child Development Steering Group. Strategies to avoid the loss of developmental potential in more than 200 million children in the developing world. *Lancet.* 2007;369(9557):229–242
 92. Engle PL, Pelto GH. Responsive feeding: implications for policy and program implementation. *J Nutr.* 2011;141(3):508–511