Dermoscopy of pseudoxanthoma elasticum

Aashim Singh,¹ Neetu Bhari,¹ Anju Bhari²

¹Department of Dermatology and Venereology, All India Institute of Medical Sciences, New Delhi, India

²Dr Rajendra Prasad Centre for Ophthalmic Sciences, All India Institute of Medical Sciences, New Delhi, India

Correspondence to
Dr Neetu Bhari,
drntbhari@gmail.com

Accepted 20 October 2017

DESCRIPTION

A 22-year-old woman presented with complaints of gradually progressive yellowish asymptomatic papules coalescing to form plaques over the lateral aspects of the neck since last 3 years. There were no associated systemic complaints. No similar complaints were noted in any of the family members. Systemic examination was normal while mucocutaneous examination revealed the presence of symmetrically distributed yellowish monomorphic papules arranged in a linear and reticulate manner on both sides of the neck, axillae and periumbilical area ([figure 1](#)). These plaques were confluent at most of the places, occasionally studded with telangiectasias and had a pebble-like feel on palpation with poor elastic recoil. Polarised light dermoscopy at $\times 10$ magnification of these plaques showed multiple irregular yellowish areas alternating with multiple linear vessels. These yellowish plaques coalesced to form parallel strands ([figure 2A](#)). The fundus examination was within normal limits except diffuse pigmentary degeneration ([figure 2B](#)). The histological evaluation of the plaque showed multiple fragmented, clumped, short, curled, basophilic elastic fibres in the mid-dermis in a ravelled ball appearance ([figure 2C,D](#)).¹ A final diagnosis of pseudoxanthoma elasticum (PXE) was made. Similar dermoscopic findings have been

Figure 2 (A) Dermoscopic view ($\times 10$) of pseudoxanthoma elasticum showing linearly arranged yellowish strands intermingled with linear vessels. (B) Fundoscopic view showing absence of angioid streaks. (C, D) Histological photomicrograph (H&E; $\times 10$, $\times 40$) showing multiple short, curled, basophilic elastic fibres in the mid-dermis.

observed by Lacarrubba *et al.*² PXE is a rare hereditary disease affecting connective tissue and causing mineralisation of elastic fibres in the skin, eyes, cardiovascular system and digestive system with an incidence of 1 in 50 000 people.³ We propose that dermoscopy could be a useful tool for a non-invasive prompt diagnosis of PXE for early detection and intervention.

Contributors All authors have contributed substantially and sufficiently to the data collection and completion of this article. AS: data collection and compilation (patient management and dermoscopy), writing and reporting the paper. NB: conception and conduct of the report, supervising and writing the paper. AB: data collection and writing the paper (ophthalmoscopy).

Competing interests None declared.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

© BMJ Publishing Group Ltd (unless otherwise stated in the text of the article) 2017. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

REFERENCES

- Madke B, Doshi B, Khopkar U, *et al.* Appearances in dermatopathology: the diagnostic and the deceptive. *Indian J Dermatol Venereol Leprol* 2013;79:338–48.
- Lacarrubba F, Verzi AE, Caltabiano R, *et al.* Dermoscopy of pseudoxanthoma elasticum. *J Am Acad Dermatol* 2017;76:S69–S70.
- Li Q, Jiang Q, Pfendner E, *et al.* Pseudoxanthoma elasticum: clinical phenotypes, molecular genetics and putative pathomechanisms. *Exp Dermatol* 2009;18:1–11.

Figure 1 Clinical examination showing yellow papules coalescing to form plaques over both sides of the neck, bilateral axillae and periumbilical area.

CrossMark

To cite: Singh A, Bhari N, Bhari A. *BMJ Case Rep* Published Online First: [please include Day Month Year]. doi:10.1136/bcr-2017-221365

Copyright 2017 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow