Published in final edited form as:

Brain Res. 2017 November 15; 1675: 8–19. doi:10.1016/j.brainres.2017.08.026.

Alterations in Histone Acetylases and Acetylated Histone Expression in Aging, Ethanol-Fed Rats

Abhilasha Khurana, M.A.^a and Cynthia A. Dlugos, Ph.D.^b [Associate Research Professor]

^aDepartment of Pathology and Anatomical Sciences, Jacobs School of Medicine and Biomedical Sciences, Stat University of NY at Buffalo

^bJacobs School of Medicine and Biomedical Sciences, State University of NY at Buffalo, Department of Pathology and Anatomical Sciences, 206 Farber Hall, 3435 Main Street, Buffalo, NY 14214, USA, Phone: 1- 716-829-2545, FAX: 1-716-829-2911

Abstract

Ethanol and age-induced pathologies of the Purkinje neuron (PN) may result from histone deacetylases (HDACs), enzymes which repress transcription through coiling of the DNA. The purposes of this study were to investigate expression patterns of Class 1 and IIa HDACs in PN and the effects of aging and alcohol on the density of HDACs and histone acetylation in PN. Ninety, eight month old rats (30/diet) were fed a liquid ethanol, liquid control, or rat chow diet for 10, 20, or 40 weeks (30/ treatment duration). Double immunocytochemical labeling on tissue sections from these rats used antibodies against HDAC isoforms or acetylated histones, and calbindin, a marker for PN. Fluorescent intensities were also measured. Results showed a significant age but not an alcohol-related decrease in the densities of HDACs 2, 3, and 7. In contrast, there were age related-increases in the densities of phosphorylated form of HDAC (4,5,7) PN and in PN nuclei expressing HDAC 7. There were also a trend towards ethanol-induced inhibition of acetylation as the density of AH2b PN nuclei and AH3 and AH2b fluorescent intensity was significantly lower in the EF compared to the PF rats. This study presents unique data concerning which HDACs are commonly expressed in PN and indicates that aging rather than lengthy alcohol expression alters expression of the HDACs studied here. These results also suggest that lengthy ethanol consumption may inhibit histone deacetylation in PN.

Keywords

histone deacetylases; Purkinje neuron; chronic ethanol; aging; cerebellum:histone acetylation

Correspondence to: Cynthia A. Dlugos.

⁶EF= Ethanol-Fed

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

1. Introduction

Ethanol-induced alterations of the cerebellum, the center of motor coordination, have important connotations for elderly humans in the United States who drink alcohol (45%), binge drink (12.2%), and drink heavily (3.25%) (The National Survey on Drugs and Health, 2002). Elderly drinkers suffer more readily from falls than their non-drinking cohorts (Rigler, 2000) with the added concern that abstinent alcoholics retain permanent deficits in cerebellar motor coordination that become more pronounced with age (Sullivan et al., 2000; Deshmukh et al., 2002.) Data from this laboratory show that chronic alcohol exposure for lengthy periods in aging adult rats alters the PN¹, the sole output neuron of the cerebellar cortex. These alterations include regression of the PN's extensive dendritic arbor (Pentney, 1995), decreased synapses on PN dendrites (Dlugos and Pentney, 1997), dilation of the PN smooth endoplasmic reticulum (Dlugos, 2006 a, b), increases in degenerating bodies in PN dendrites (Dlugos, 2008), decreases in levels of the sarco/endoplasmic reticulum calcium adenosine triphosphate pump and balancing ability (Cassidy et al., 2013), and increased accumulation of caspase 12 (Dlugos, 2014).

The mechanisms behind ethanol-induced changes to the rat PN are unknown although epigenetics, specifically removal of acetyl groups by HDACs², is a candidate. Histone deacetylation by HDACs results in removal of acetyl groups from lysines on histone tails resulting in DNA coiling, and repression of transcription (Shahbazian et al., 2007). HDACs deacetylate many other targets within the cell besides histones (Yang and Gregoire, 2005) and are divided into zinc-dependent and nicotinamide adenine dinucleotide dependent groups. The zinc-dependent HDACs are subdivided into four families (Classes I, IIa, IIb and IV) and members of each family are found in the brain (Broide et al., 2007; Volmar and Wahlestedt, 2015). The class I family consists of HDACs 1, 2, 3, and 8 (Yang and Seto, 2008; Haberland et al., 2009) and the class IIa family consists of HDACs 4, 5, 7, and 9 (Yang and Gregoire, 2005; Yang and Seto, 2008; Yao and Yang, 2011). The class IIb family consists of HDACs 6 and 10 whereas HDAC 11 is the sole member of the class IV family.

Members of the of the Class 1 and IIa HDACs have been extensively studied as they are essential to life (Lagger et al., 2002; Vega et al., 2004; Montgomery et al., 2007; Bhaskara et al., 2008; Knutson et al., 2008; Montgomery et al., 2009). In the class I family, HDACs 1 and 2 share similar sequences and nuclear localization but are functionally different as HDAC³ 1 is expressed in glia and HDAC 2 in neurons (MacDonald et al., 2008). HDAC 2 appears to be essential in neuronal function as overexpression reduced dendritic spine density and resulted in poor performance on memory tests (Guan et al., 2009). In contrast to other Class 1 HDACS, HDAC 3 is expressed in the nucleus but may translocate to the cytoplasm with substrates which include p300/CREB binding protein -associated factor (Blanco-Garcia et al., 2009) or its own cleavage products during apoptosis (Escaffit et al., 2007).

¹PN= Purkinje neuron

²HDACs=histone deacetylases

³HDAC= histone deacetylases

Class IIa HDACs normally shuttle between the nucleus and cytoplasm, providing multiple functions for the cell. For example, the export of HDACs 4, 5, and 7 from the nucleus is facilitated by a C-terminal nuclear export sequences and phosphorylation of highly conserved serine residues (Li et al., 2004). Cytoplasmic retention is maintained by anchoring the phosphorylated serines to 14-3-3 chaperone proteins (Annemieke et al., 2003; Li et al., 2004; Parra and Verdin, 2010; Yao and Yang, 2011). HDAC 4 and HDAC 7 appear to have a neuroprotective role that is demonstrated directly in PN as mice lacking HDAC 4 lost PN (Majdzadeh et al., 2008). Class IIa HDACs also appear to have an anti-apoptotic effect on other neurons as forced expression of HDAC 4 (Majdzadeh et al., 2008) and HDAC 7 (Ma and D'Mello, 2011) reduced low potassium apoptosis in cerebellar granule neurons.

An early study using *in situ* hybridization revealed that, of all the 11 zinc-dependent HDACS, only mRNA for HDAC 11 was found in PN (Broide et al., 2007). Later studies with antibody staining showed nuclear localization of HDAC 2 in developing PN (Yoo et al., 2013). HDAC 4 was also reported in the PN cytoplasm of control mice and in the PN nucleus of mice with ataxia telangiectasia (Li et al., 2012). Another study demonstrated a more variable localization of HDAC 4 in normal PN ranging from nuclear to cytoplasmic and including some PN with HDAC 4 in both the nucleus and cytoplasm (Darcy et al., 2010). HDAC7 has not been identified specifically in PN rather in the cytoplasm and nucleus of neurons of mouse cerebellar cortex, cerebral cortex, and striatum (Benn et al., 2009). There is some evidence for the presence of HDAC 3 in PN as PN specific HDAC3 null mice displayed a loss of PN, a thin molecular layer, and decreased rotorod performance (Venkatraman et al., 2014). Similarly, the knockout of HDAC 3 in neuronal progenitor cells resulted in underdeveloped cerebellar folia, lower cell density, and lower staining intensity for a calbindin antibody in PN (Norwood et al., 2014).

There is some precedent for alcohol-induced alterations to the histone deacetylases but no consistent pattern in the results. Some studies indicate that HDACs are upregulated with ethanol. For example, HDAC3 was upregulated in the liver following alcohol exposure. (Kirpich et al., 2013) and acute treatment with alcohol resulted in increased expression of HDAC 2 (Agudelo et al., 2011) and HDACs 1 and 3 (Agudelo et al., 2012) in a human neuroblastoma cell line. Ethanol-related upregulation of HDACS would explain the reported reduction in acetylated histones 3 and 4 in the perinatal rat (third trimester equivalent) cerebellar cortex (Guo et al., 2011). HDAC downregulation with alcohol treatment has also been reported. For example, acute alcohol treatment resulted in downregulation of HDACs 1, 2, and 3 in the caudate and putamen of wild-type mice (Caputi et al., 2015) and also in decreases in m- RNA for HDACs 1 and 4 in hippocampal slices (Zou and Crews, 2014). Ethanol-induced decreases in HDACs might be responsible for increased acetylation of histone 3 in rat liver cultures (Park et al., 2003) or global hyperacetylation of liver proteins such as P53 tumor suppressor protein and α -tubulin (Shepard and Tuma, 2009). There are several reports that ethanol withdrawal alters HDAC expression. In the amygdala, withdrawal resulted in increased HDAC activity and decreased histone acetylation (Pandey et al., 2008) and an HDAC inhibitor was reported to modify increases in HDAC expression resulting in reduced anxiety and increased spine density (You et al., 2014). In contrast, in the prefrontal cortex, withdrawal from ethanol would appear to downregulate HDAC expression

as increased acetylation of histones 3 and 4 have been reported in adolescent (Pascual et al., 2012) and young adult rats (Jung and Metzger, 2015).

Aging is another factor that may influence HDAC activity enhancing or reducing alcohol-induced effects. For example, decreased HDAC expression of Class I HDACs was shown in oligodendrocytes from the corpus callosum of 8 month old rats compared to cells from 3 and 8 week old rats (Siming et al., 2008). Similarly, the level and activity of HDAC 1 decreased in cultured senescent WI-38 cells compared to cells in younger cultures (Place et al., 2005). In contrast, the mean fluorescent intensity of HDAC2 expression in several hippocampal regions increased in mice between 12 and 24 months of age suggesting an age-related increase in HDAC expression (Chouliaras et al., 2013). In addition, HDAC overexpression has been observed in the CNS in aging animals during ischemia (Baltan, 2012).

The following study had three aims. First, to determine the presence and subcellular localization of the Class 1 and IIa HDACs in PN in aging, ethanol-fed rats. Second to assess ethanol and age-induced alterations in the density and/or fluorescent intensity of the predominant Class 1 and IIa HDACs in PN. Third, to determine whether there are age or ethanol-induced changes in histone acetylation in PN at 40 weeks, the maximum duration of ethanol treatment and the duration during which ethanol-induced changes in PN function and structure have been demonstrated in this laboratory.

2. Results

2.1 Blood Alcohol Levels and Body Weight

The mean blood alcohol level ($\pm$ SD) was 125.6 ± 44 mg/dl. Two-way ANOVA showed that with diet and duration of treatment as factors there was an overall effect of weight [F_{overall} (8,81) = 6.427, $P < .001$] that was partially due to dietary treatment [F_{diet} (2,81) = 5.470, $P = .006$]. CF⁴ rats had the lowest weights ($\pm$ S.E.M) overall ($458\text{g} \pm 11.8$) whereas the PF⁵ rats ($491\text{g} \pm 7.4$) had the highest weights. Weights of the PF and CF rats were significantly different from each other ($P < .005$) but EF rat weight (mean = $468.4\text{g} \pm 5.8$) did not differ from any other group ($P > 0.09$). Duration of treatment [F_{duration} (2,81) = 19.58, $P < .001$] was also significant and showed that rats treated for 10 ($P < .001$) and 20 ($P = .003$) weeks weighed less than at 40 weeks of treatment and that, all rats, as expected, gained weight throughout the study.

2.1 Histology

Immunostaining and confocal microscopy (Fig. 1) confirmed the localization of HDACs 2, 3, 4, 7 and phosphorylated HDACs 4,5, and 7 within PN. HDAC2 (Fig. 1 A) and HDAC3 (Fig. 1 B) were localized exclusively in the PN nucleus but was also found within nuclei of other neurons including the large population of cerebellar granule cells. HDAC4 (Fig. 1 C) was specific to PN in the cerebellar cortex and was cytoplasmic in location. Phosphorylated HDAC 4, 5, and 7 (Fig. 1 D) stained a small subset of PN and was located mainly in the nucleus becoming more particulated and dispersed within the cytoplasm. HDAC 7 (Fig. 2 A)

⁴CF=Chow-Fed

⁵PF= Pair-Fed

was present within the nucleus and the cytoplasm although sole nuclear localization was more pronounced with age. The other HDACs tested were not good candidates for further investigation because HDAC 1, HDAC5, and HDAC 6 were absent in PN. As expected, HDAC 1 was contained in Bergmann and other glial cells surrounding PN somas and in the molecular layer. HDAC 5 was not found in the cerebellar cortex whereas HDAC 6 stained Golgi neurons of the granule layer. Acetylated histones 2b and 3 (Fig. 2C) were all localized in the PN nucleus with acetylated histone 3 being most prevalent in PN.

2.2 HDAC and Histone Densities

Two-way ANOVA with age and diet as factors was carried out on HDAC+ PN densities / mm of PN line. The overall effect on the linear density of HDAC2+ PN [$F_{\text{overall}}(8,81)=12.889$, $P<.001$] was due to age [$F_{\text{age}}(2,81)=50.78$, $P<.001$] but not to diet [$F_{\text{diet}}(2,81)=.683$, $P=.508$] (Fig. 3). HDAC 2 density increased in the EF and PF rats at 14 months compared to their 11 month old counterparts ($P<.05$) whereas, at 19 months, HDAC 2 + PN decreased in all dietary groups compared to values in the 14 month old rats ($P<.001$). There was also an overall significant effect on HDAC3+ PN density [$F_{\text{overall}}(8,81)=3.495$, $P=.002$] that was also due to age [$F_{\text{age}}(2,81)=11.457$, $P<.001$] but not to diet [$F_{\text{diet}}(2,81)=1.666$, $P=.195$] (Fig. 4). In this case, there was a decrease of HDAC 3+ PN in the 14 compared to the 11 month old CF and PF rats ($P=.03$) and in the 19 month old PF and EF rats compared to their 14 month old counterparts ($P=.02$). Similarly, a significant decline in HDAC 7 expression [$F_{\text{overall}}(8,81)=9.042$], was due to age [$F_{\text{age}}(2,81)=35.541$, $P<.001$] but not diet [$F_{\text{diet}}(2,81)=.057$, $P=.945$] (Fig 5). When the density of HDAC 7 + PN/10mm was analyzed, there was a decrease at 19 months compared to 11 months in CF, PF, and EF rats ($P=.002$). Additional age-related declines in HDAC 7 + PN densities were seen in CF rats at 19 compared to 14 months of age ($P=.014$) and in PF rats at 14 compared to 11 months of age ($P=.019$). Despite the fact that the total number of HDAC 7+ PN decreased with age, the density of HDAC7 + PN with staining solely in the nucleus increased [$F_{\text{overall}}(8,81)=4.052$, $P<.001$] with age [$F_{\text{age}}(2,81)=15.823$, $P<.001$] but not with dietary treatment [$F_{\text{diet}}(2,81)=.157$, $P<.855$]. Increases in sole nuclear expression of HDAC 7 occurred in all dietary groups at 19 compared to 11 months of age ($P=.02$).

The density of phosphorylated HDAC4, 5, and 7+ PN [$F_{\text{overall}}(8,81)=11.398$, $P<.001$] was again due to age [$F_{\text{age}}(2,81)=44.015$, $P<.001$] but not to diet [$F_{\text{diet}}(2,81)=.668$, $P=.516$] (Fig. 6). Rather than the decrease shown with the other HDACS, the density of phosphorylated HDAC 4,5,7 increased at 19 months of age compared to the 11($P<.001$) and 14($P<.001$) month old CF, PF, and EF rats. There was no overall effect on the linear density of HDAC4+ PN /mm [$F_{\text{overall}}(8,81)=1.616$, $P=.133$] (Fig. 7). Two way ANOVA with treatment and histone as factors, showed an effect on the density of acetylated histones on PN/ mm [$F(5,54)=3.701$, $P=.006$] due to decreases in the EF compared to the PF rats ($P=.004$) (Fig 8) that was localized in AH2b⁷ ($P=.009$) but not in AH3⁸ ($P=.246$).

⁷AH2b= Acetylated histone 2b

⁸AH3=Acetylated histone 3

2.3 Corrected Total Cell Fluorescence

Two-way ANOVA on CTCF⁹ values from HDACs 2,3, and 7 at the 40 week treatment duration, using the factors of HDAC and diet, showed no overall effect [$F_{\text{overall}}(8,81)=.705$, $P=.686$] (Fig.9). The area of the particle measured for CTCF was also tested for HDACs 2,3, and 7 at 40 weeks and showed an overall difference [$F_{\text{area}}(8,81)=6.437$, $P<.001$] among the HDACs [$F_{\text{HDAC}}=25.207$, $P<.001$]. This difference was due to smaller areas for HDAC 2 ($P<.001$) and HDAC 3($P<.001$) compared to HDAC 7, reflecting the nuclear location of HDACs 2 and 3 and the most common cytoplasmic location of HDAC 7. Diet had no effect on particle size. [$F_{\text{diet}}(8,81)=.145$, $P=.866$]. Two-way ANOVA on CTCF for AH2b and AH3 showed an overall effect on CTCF [$F_{\text{overall CTCF}}(5,54)=6.526$, $P<.001$] that was due to the ethanol diet [$F_{\text{diet}}(2,54)=10.973$, $P<.001$] as the CTCF for AH2b ($P=.017$) and AH3 ($P=.001$) were lowest in the EF compared to the PF rats. AH3 was also highest in the PF relative to the CF rats ($P=.002$). There was also an effect of the acetylated histone used [$F_{\text{histone}}(1,54)=8.291$, $P=.006$] that was due to a higher CTCF for the AH3 compared to the AH2b antibody. There was no difference in the areas of the particle measured for the acetylated histones [$F_{\text{overall}}(5,54)=1.764$, $P=.136$].

2.4 Volume measures of PN and Molecular Layers

Two-Way ANOVA on volume measures of the molecular layer (including PN line) showed no alterations in the volumes of the molecular and Purkinje neuron layers [$F_{\text{overall}}(8,81)=1.528$, $P=.157$] (Table1). This lack of change in molecular layer volume with age and treatment suggests that alterations in densities are reliable indicators of increases or decreases in HDACs or acetylated histones.

3. Discussion

The results of this study showed a distinct expression pattern for specific Class I and Class IIa HDACs in PN. There were age-induced decreases in densities of PN + HDAC 2, HDAC3, and in PN/10 mm for HDAC 7. In contrast, there were also age-related increases in the densities of phosphorylated HDAC 4, 5, 7+ PN and in PN expressing HDAC 7 solely in the nucleus. There was no effect of ethanol treatment on HDAC 4 density in PN within any age group nor was the fluorescent intensity for HDACs 2, 3, or 7 in PN different after 40-weeks of dietary treatment, the time when ethanol-induced effects commonly occur. (Cassidy et al., 2013; Dlugos and Pentney, 1997; Dlugos, 2006a, b; Dlugos, 2014). The density of AH 2b+ PN/mm at 40 weeks, however, was significantly lower in the EF compared to the PF rats. There was also a significant decrease in fluorescence intensity in the EF relative to the PF rats indicative of lower levels of AH3 and AH2b after 40 weeks of ethanol treatment.

These findings were consistent with characterization of Class I and Class IIa HDACs as HDACs 2 and 3 were localized within PN nuclei and HDAC 4 was mainly cytoplasmic. HDAC 3 localization within the cytoplasm of PN was not observed although cytoplasmic substrates of HDAC 3 such as nuclear factor- light-chain-enhancer (Li et al., 2010) and

⁹CTCF=Corrected total cell fluorescence

signal transducer activator of transcription 1 expression (Yao and Yan, 2011) has been described. Further investigation into the cytoplasmic substrates of HDAC3 suggests that they may be minimally expressed or absent in PN. For example, signal transducer activator of transcription 1 appears not to be expressed in PN (<http://www.proteinatlas.org/ENSG00000115415-STAT1/tissue/primary+data>). Another substrate, transcription factor p65, a member of the nuclear factor- light-chain-enhancer of activated B cells family, is expressed in PN during development and capable of cytoplasmic localization (Li et al., 2010). Its expression in PN during adulthood has not been verified, however. We were also unable to detect cleaved portions of HDAC3 in the cytoplasm in PN such as were described during Fas ligand-induced apoptosis in Jurkat cells (Escaffit et al., 2007). Cleaved HDAC 3 may not have been detected in the present study because an antibody against the whole HDAC 3 molecule rather than the cleaved fragment was used. It is also possible that Fas ligand-induced apoptosis did not occur in PN as age-related apoptosis or cell death in PN has not been observed in this particular animal model (Dlugos and Pentney, 1994),.

The present findings showing expression of HDACs 2, 3, 4, 7 in rat PN contradicted the findings of another study that showed only HDAC11 expression in PN (Broide et al., 2007). A possible reason for this discrepancy is that the Broide et al. study (2007) used neuron-specific nuclear protein to recognize PN, although PN do not stain with this antibody (Kim et al., 2009). The present study used calbindin D-28 to recognize PN as it is a calcium binding protein that is specific for PN within the cerebellar cortex (Bastinelli, 2003). We did not find HDAC 6 + PN in our study although it has been reported by others in PN axons, not visualized to any extent here as the focus was on the cerebellar cortex rather than the white matter of the cerebellum (Southwood et al., 2007). Support for the observed lack of HDAC 6 in PN is provided by a study in which transfection of HDAC 6 was required to visualize it in PN (Ohkawa et al., 2008). HDAC 5 expression in PN was not observed here and its expression in PN has only been reported in disease states, such as Huntington's chorea (Yeh et al., 2013).

The present data showing that the densities of PN which express HDACs 2 (Fig 3) and 3 (Fig 4) decrease with age corroborate similar findings in which levels of class I HDACs were decreased in oligodendrocytes of the corpus callosum of aged mice (Siming et al., 2008) and in aging WI-38 cells (Place et al., 2005). Several studies, however, support age-induced increases in HDAC staining. In these studies, HDAC 2 staining intensity was increased in the hippocampus of 12 and 24 month old mice (Chouliaras et al., 2013), in the hippocampus of 3 and 6 month old mice and in human Alzheimer's patients (Graff et al., 2012). Comparison of the age-related decreases in HDACs 2 and 3 reported in the present study and the age-related increases in HDAC 2 shown in the hippocampal studies (Graff et al., 2012; Chouliaras et al., 2013) indicate that aging effects on the hippocampus and cerebellum are different and that cell specificity, cell function, and brain region may play a role in these differences. The age and species of the animals studied may also be a factor despite the brain region studied. For example, a study in the rat hippocampus showed significant increases in mRNA levels in HDAC 2 between 2 and 15 months of age (Suri et al., 2014). In the present study, the density of HDAC2+ PN showed a similar significant increase from 11-14 months of age but then significantly decreased between 14-19 months of age (Fig. 3). Should the age of the animals in the study of Suri et al. (2014) been extended

to 19 months of age, it is possible that a decline in HDAC 2 may have occurred in hippocampal neurons making age a more critical factor than neuronal type.

Age-related changes in PN expressing HDACs, 2 (Fig 3), 3 (Fig 4) and 7 (Fig 5) could be strongly suggestive of age-related decline in PN if it were not for the fact that HDAC 4, (Fig 7), present in every PN along the PN line, did not decrease with age. Age-related decreases in PN have been demonstrated in some studies (Rogers et al., 1984; Woodruff-Pak et al., 2006) whereas other studies showed no age-related changes in PN number (Druge et al., 1986; Bakalian et al., 1991; Dlugos and Pentney, 1994; Larsen et al., 2000). There are many factors involved that explain the differing results, including the counting method employed but external factors, such as the simple paradigm of exercise, can make a difference in age-related PN loss. For example, one study shows that exercise prevented the loss of PN in old rats compared to cohorts that were not exercised (Larsen et al., 2000) while in another, rats, of ages as high as 27 months, did not show age-related PN loss compared to their 3, 9, and 18 month old cohorts that were exercised (Dlugos and Pentney, 1994). Other factors in the present study argue against substantial age-related PN. For example, Table 1 shows that there was no age-related difference in the volume of the PN line and molecular layer among the 11, 14 and 19 month old rats used here that would be indicative of PN loss. Gliosis, an indicator of cell loss, was also not apparent in the present study.

In contrast to HDAC density, the density of the phosphorylated form of HDAC4, 5 and 7 increased with age (Fig 6), a finding that may reflect an age-induced increase in phosphorylation to support cytoplasmic class IIa family members in their neuroprotective role (Li et al., 2004; Wang et al., 2014). Our finding of localization of the phosphorylated 4, 5, and 7 antibody mainly in the nucleus with particulate matter in the cytoplasm (Fig. 1.) is supportive of the model of phosphorylation within the nucleus presumably by calmodulin dependent protein kinases. Phosphorylation enhances nuclear export of the HDACs and cytoplasmic association with the 14-3-3 proteins (Kao et al., 2001; Annemieke et al., 2003; Li et al., 2004). Current results with the phosphorylated antibody cannot predict whether there is a greater difference in the phosphorylated form of HDAC 4 or 7 with age but HDAC 5 can be ruled out as it does not appear to be localized within PN. HDAC dephosphorylation in the cytoplasm results in solely nuclear expression (Parra and Verdin, 2010), a finding that was observed in our older rats with HDAC 7 but not with HDAC 4. More needs to be known about the specific functions of HDACs 4 and 7 in PN in order to decipher the differences in the nuclear versus the cytoplasmic expression of these isoforms and what these differences mean with respect to function in the cell.

HDAC 4, highly expressed in PN in this study, never became localized in the nucleus although nuclear localization of HDAC4 in PN has been reported with the neuronal degeneration that accompanies ataxia telangiectasia (Li et al., 2012). Aging was, therefore, not sufficient to induce nuclear localization of HDAC 4 in PN although we observed small populations of PN showing HDAC7 nuclear localization with age. There are reported differences between HDACs 4 and 7 that are suggestive of different functions in PN. For example, hypoxia inducible factor 1 alpha, a common substrate between HDACs 4 and 7, has a cytoplasmic location for HDAC 4 but a nuclear in location for HDAC 7 (Yao and Yang, 2011). There are also structural differences between HDAC 4 and 7 that may result in

functional dichotomy in PN. For example, HDAC 7 has the most divergent sequence in its amino terminus of the IIa HDACS (Kao et al., 2001) and is more similar to HDAC 5 than to HDAC4 (Annemieke et al., 2003).

The present study shows no alcohol-induced effects on the Class I and Class IIa HDACs studied. Several studies with different alcohol paradigms showed upregulation of one or more members of the HDAC 1 family in the liver (Kirpich et al., 2013), a human neuroblastoma cell line (Aguedelo et al., 2011, 2012), and the amygdala (Pandey et al., 2008). Others showed ethanol-induced downregulation of the HDAC I family in the caudate and putamen (Caputi et al., 2015)a and decreased mRNA content for HDAC I and 4 in the hippocampus (Zou and Crews, 2014). These studies are so diverse in their ethanol treatments and the cell types used that it is difficult to relate them to the current study showing that chronic ethanol treatment in PN does not alter expression of the HDACs studied. This is the sole study specifically on HDACs in PN in aging, adult rats exposed to lengthy alcohol exposure.

The alcohol effect in this study was not on HDACs but on acetylation of histones. Studies using antibodies against AH2b and AH3 showed a significant decrease in the density of AH2b in the EF rats following 40 weeks of ethanol treatment and nonsignificant trend towards a decrease in the density of AH3 (Fig 8). Fluorescent intensity measures showed significant decreases in both as AH2b and AH3 with respect to the PF rats (Fig 9), the true dietary controls in this study. Significant changes in fluorescent intensity measures were unique to the acetylated histones as there was not similar decrease in fluorescent intensity in the HDACs after 40 weeks. These ethanol-induced decreases in fluorescent intensity and density corroborate data showing decreased levels of histone acetylation in the cerebellar cortex following prenatal alcohol exposure although this study was not specific to PN (Guo et al., 2011). Decreased histone acetylation has also been linked to alcohol exposure in the cerebral cortex and it has been proposing that this action is through multiple histone modifying enzymes (Finegersh and Homanics, 2014),

The appropriate histone modified enzyme or enzymes related to ethanol-induced decreases in acetylation was not detected in this study, despite the focus on the class I HDACs which are predominantly nuclear in location and likely candidates to modifying histone coiling. There are other HDACs that were not tested, such as HDACs 9-11 or any member of the nicotinamide adenine dinucleotide dependent sirtuin family which may show ethanol-induced increases and be responsible for the decrease in H2b and H3 acetylation shown here. Sirtuin 1, has been found in the cerebellum and other central nervous system neurons (Zakhary et al., 2010) and is worthy of further investigation. An alcohol-related decrease in histone acetylation may also be due to a decrease in one member of the five families of histone acetyl transferases (Schneider et al., 2013). CREB binding protein is an excellent candidate as it appears to be decreased in the cerebellum following perinatal treatment with ethanol in a study in which acetylation of H3 and H4 were also decreased (Guo et al., 2011), although the specific localization of CREB binding protein in PN was not shown. Future studies will, therefore, be aimed at investigation of the sirtuin and histone acetyl transferases families in aging rats exposed to chronic alcohol.

In summary, this study confirmed localization of Class I and Class IIa HDACs in PN of aging animals. Densities of two members of the Class I HDACs, (HDAC2 and HDAC3) and one member of the Class II a family decreased with age (HDAC 7). Densities of HDAC 4 remained stable with age. There was an age-induced increase in the density of phosphorylated HDAC 4,5, and 7 as well as an age-induced increase in HDAC 7+ PN with solely nuclear localization. There was also an ethanol-induced decrease in the density of A H2b + PN and in the fluorescent intensity of AH2b and AH3 reflecting decreases in gene expression. This change may reflect increased expression of HDACs not tested in this study, decreased expression of HATs and/or a change in the balance between HDACs and HATs with lengthy chronic alcohol exposure. There was no alteration in the volume of the Purkinje and molecular layer with increasing age or alcohol treatment that would confound interpretation of the densities determined in this study.

4. Experimental Procedure

4.1 Animal model and diet

Ninety male 8 month old Fischer 344 rats and three male 3 month old Fischer 344 rats were purchased from the National Institute on Aging colony. The three month old rats were used to test the antibodies in preliminary studies and for confocal microscopy. The 8 month old rats were sorted into one of three dietary treatment groups (30 rats/diet) for durations of 10, 20 or 40 weeks (30 rats/treatment duration). CF rats received Teklad Global 18% Protein Rodent Diet (Teklad) and water *ad libitum*. Liquid diets were purchased from Dyets Inc. (Bethlehem PA). EF rats received the AIN¹⁰-93M liquid diet *ad libitum* modified so that 35% of the daily caloric intake was ethanol. PF rats received the same volume of the AIN-93M liquid control, isocaloric to the ethanol diet with the addition of maltose dextrans. PF rats received the same volume of diet that their EF partners consumed on the previous day. Blood alcohol levels were determined enzymatically one hour into the twelve hour dark cycle with the Analox Alcohol Analyzer (Analox, Instruments, USA, Inc.) in a subset of ten EF rats after 4 weeks of alcohol treatment. Dietary intakes were recorded daily and animals were weighed every week during treatment. All procedures were performed with approval of the Institutional Animal Care and Use Committee of the State University of New York at Buffalo (Institutional Animal Welfare Assurance A3354-01).

4.2 Tissue preparation

At the end of the study when rats were 11, 14 and 19 months of age, they were euthanized by an overdose of Fatal Plus® (100 mg/kg) and perfused through the aorta with 4% paraformaldehyde in 0.1 M phosphate buffer (pH 7.4). The cerebellum was separated from the cerebrum and divided into the vermis and two cerebellar hemispheres. All cerebella were coded to prevent investigator bias. Following cryoprotection in 30% sucrose in 0.1M phosphate buffer, the cerebellum in each animal was completely sectioned at 40 µm on a rotary microtome and the total number of sections recorded. All sections were collected in each rat. Sixteen sections from each rat, collected at a set interval throughout the cerebellum, were affixed on slides and stained immediately with 0.5% cresyl violet in 1% sodium borate.

¹⁰AIN= American Institute of Nutrition

These sections were used for determination of molecular layer volume. As all of the immunocytochemical staining could not be accomplished at one time, all the other sections were stored in a 30% ethylene glycol, 30% glycerol mix in 0.1M phosphate buffer at -20° C until staining could be accomplished (Alshammari et al., 2016).

4.3 HDAC Immunocytochemistry

As little is known about HDAC expression in PN and to select HDAC antibodies for the study, a preliminary study using monoclonal mouse antibodies against HDAC1 (e.g. RRID: AB_10612242), HDAC2 (e.g. RRID: AB_10624871) and HDAC3 (e.g. RRID: AB_2118371), monoclonal rabbit antibodies against HDAC4 (e.g. RRID: AB_1086255), HDAC6 (e.g. RRID: AB_10891804), phosphorylated HDAC 4,5,7 (e.g. RRID: AB_2118725), and polyclonal antibodies against HDAC5 (e.g. RRID: AB_10828358), HDAC 6 (e.g. RRID: AB_10696889) and HDAC7 (e.g. RRID: AB_2540658) was undertaken on 3 month old male control Fischer 344 rats (n=3). Sections labelled for the monoclonal mouse HDACs were co-labelled with polyclonal (e.g. RRID: AB_258818) antibodies against calbindin, a marker specific for PN in the cerebellar cortex. Sections labelled with the monoclonal rabbit or the polyclonal antibodies were co-labeled with monoclonal (e.g. RRID: AB_476894) calbindin antibody. HDAC antibodies were diluted to 1:250 and calbindin diluted to 1:500. Prior to immunocytochemistry of quantitation, a subset of sections, stained with each of the above antibody concentrations was examined with the Zeiss LSM-520 Meta NLO laser scanning microscope to confirm co-localization of the HDACs within PN.

Immunocytochemistry was undertaken with standard techniques. Briefly, free floating sections selected systematically throughout the cerebellum, were washed thoroughly in PBS (3×), blocked for 1 hour in a solution of phosphate buffer to which 1% BSA and 0.4% Triton-X100 were added, and incubated in the antibody solution overnight at 4° C. Two sections in which the primary antibody was eliminated were included in each staining and served as negative controls. Sections were then washed 3× with PBS (5 min/wash) and exposed to a mix of 2 secondary antibodies for one hour at 25 °C. Sections were then washed in PBS (3×), mounted on slides, dried on a slide warmer, and cover-slipped with Prolong Gold (ThermoFisher Scientific, Waltham, MA). Mouse monoclonal HDACs were stained with a mix (1:100 dilution) of Alexa Fluor 488 (Ex 495/Em 519) goat-anti mouse (e.g. RRID: AB_2534069) and Alexa Fluor 568 (Ex 578/Em 603) (e.g. RRID: AB_143157). Rabbit monoclonal and the polyclonal antibodies were stained with a mix (1:100 dilution) of Alexa Fluor 488 (Ex 495/Em 519) goat anti-rabbit (e.g. RRID: AB_143165) and Alexa Fluor 568 (Ex 578/Em 603) anti-mouse (e.g. RRID: 2534072). Localization of HDACs within the PN nucleus, cytoplasm, or nucleus and cytoplasm was the criteria for selection of HDAC antibodies for quantitation.

Following the preliminary selection HDACs 2,3,4, phosphorylated HDAC4, 5,7, and HDAC 7 were chosen for staining and quantitation. The number of sections stained/ animal varied by antibody and were systematically selected at regular intervals throughout the cerebellum. For HDAC2 and 3, localized within the nucleus, 6 and 8 sections/ rat were stained and quantitated, respectively. For HDACs 4, phosphorylated HDAC 4,5, and 7, and HDAC 7,

three sections / rat were stained and quantitated for each antibody. To insure equal staining conditions sections for each triplet of CF, PF and EF rats were stained together. Identical immunocytochemical procedures as in the preliminary study was used.

4.4 Histone staining

A preliminary study to determine which acetylated histones are most highly expressed in PN using an acetylated histone staining kit (e.g. RRID: AB_10699455). From these results 3 sections/ rat in the 40 week group were labelled with rabbit monoclonal antibodies against AH3 (e.g. RRID: AB_823528) and AH2b (e.g. RRID: AB_2636805) and the mouse calbindin (e.g. RRID: AB_476894) used in HDAC staining. Due to the paucity of sections and the repeated finding in this laboratory that chronic ethanol effects take lengthy periods to occur (Dlugos and Pentney, 1997; Dlugos 2006a, Dlugos 2006b, Dlugos, 2008; Cassidy et al., 2011; Dlugos, 2014), only the 40 week group was stained. Dilutions of 1:250 for the acetylated histone antibodies and 1:500 for the calbindin antibody were used and the secondary antibodies were Alexa Fluor 488 goat anti-rabbit (e.g. RRID: AB_143165) and Alexa Fluor 568 anti-mouse (e.g. RRID: 2534072) diluted to 1:100. Localization of AH2b and AH3 within PN nuclei was confirmed with confocal microscopy. Immunostaining was performed as described above.

4.5 Quantitative Analyses

A Zeiss Axioimager fluorescent microscope with rhodamine and fluorescein filters was used to determine the densities of HDACs and histones. Ten fields/ slide were photographed with both filters at 40 $\times$ (1600 $\times$ = total magnification). HDAC+, AH3+, and AH2b+ PN were counted in each field by comparing HDAC and acetylated histone+ cells with calbindin D-28+ PN. This was done by side-to-side comparison of the fluorescein images stained with HDACs or histone antibodies with the rhodamine images stained with calbindin D-28. HDAC or histone + neurons were only counted if the cell was positive for calbindin D-28+, the marker for PN. The standard rules for inclusion and exclusion of profiles was used in that PN touching two sides of the image were excluded; cells touching the opposite two sides were included (Gundersen, 1977). The length of the PN line in each field was determined using the UTHSCSA Image Tool Program (<http://compdent.uthscsa.edu/dig/itdesc.html>). Mean ($\pm$ S.E.M.) cell counts and PN line measurements were found for each slide and for each animal and a mean density of HDAC + or histone + PN calculated per rat by dividing the mean number of HDAC+ or histone + PN by the mean length of the PN line. For HDAC7, the number of HDAC 7 PN in which localization was solely nuclear also were counted.

CTCF was also determined using the Image J program (<https://imagej.nih.gov/ij/>) for HDAC2, HDAC3, HDAC 7, AH3 and AH2b (Turano et al., 2015). Ten PN/rat were used in these determinations. The area (nucleus or nucleus and cytoplasm) of the fluorescent particle was also measured as significant increases in the area of the particle measures might be due to significant increases in fluorescence between dietary treatment groups. CTCF for anti-HDAC 4 was not measured because the program did not allow for isolation of the fluorescent cytoplasm from the non-fluorescent nucleus, making the results difficult to

interpret. CTCF of anti-phosphorylated 4,5, and 7 was not conducted because of the difficulty in finding fluorescent cells in many age groups.

The sixteen sections stained with cresyl violet were digitized and area of the molecular layer + PN line / slide determined using the UTHSCSA Image Tool program. This area in each rat was the average from the values of sixteen sections. Section thickness/ animal was determined by taking three measures on three slides with the DG-3601 microcator (Ohno Sokki, Addison, IL). A mean section thickness / animal (t) was determined by averaging all values / slide and then the values from the three slides. The interval, k, between tissue sections was recorded. These values and the total number of sections / cerebellum were used to determine the reference volume (V ref) of the molecular layer including the PN line with a previously reported modification of Cavalieri's principle ($V_{ref} = \Sigma P. ap. t. k$) (Sterio, 1984; Dlugos and Pentney, 2001). In this modification, molecular layer area is determined with the UTHSCSA Image Tool program instead of point counting ($\Sigma P. ap$) on each slice. This was done because changes in the volume in which the PN reside (Purkinje and molecular layer) would confound interpretation of density changes.

4.6 Statistical Analyses

Means ($\pm$ S.E.M.) for all variables were determined within each rat, treatment group, and duration of treatment. ANOVA analysis was used for immunocytochemical data using dietary treatment and duration of treatment as fixed variables. One-way ANOVA with the post-hoc Tukey test was used to localize differences. An alpha value of .05 was selected as significant.

Acknowledgments

Many thanks to Dr. Thaddeus Szczesny for help with the illustrations and for Frederick Dlugos for rat care. This work was supported by grant AA017195 from the National Institute on Alcohol Abuse and Alcoholism, and Office of Extramural Research, National Institutes of Health.

References

- Agudelo M, Gandhi N, Saiyed Z, Pichili V, Thangavel S, Khatavkar P, Yndart-Arias A, Nair M. Effects of alcohol on histone deacetylase 2 (HDAC2) and the neuroprotective role of trichostatin A (TSA). *Alc Clin Exp Res*. 2011; 35:1550–1556.
- Agudelo M, Yoo C, Nair MP. Alcohol-induced serotonergic modulation: the role of histone deacetylases. *Alcohol*. 2012; 46:635–642. [PubMed: 22796363]
- Alshammari, MA., Alshammari, TK., Laezza, F. Improved methods for fluorescence microscopy detection of macromolecules at the axon initial segment. *Front Cell Neurosci*. 2016. [//dx.doi.org/10.3389/fncel.2016.00005](https://doi.org/10.3389/fncel.2016.00005)
- Annemieke JM, de Ruijter AH, van Gennip HN, Caron SK, van Kuilenburg AB. Histone deacetylases (HDACs): Characterization of the classical HDAC family. *Biochem J*. 2003; 370:737–749. [PubMed: 12429021]
- Bakalian A, Corman B, Delhay-Bouchaud N, Mariani J. Quantitative analysis of the Purkinje cell population during extreme ageing in the cerebellum of the Wistar/Louvain rat. *Neurobiol Aging*. 1991; 12:425–230. [PubMed: 1770976]
- Baltan S. Histone deacetylase inhibitors preserve function in aging axons. *J Neurochem*. 2012; 123:108–115. [PubMed: 23050648]
- Bastinelli E. Distribution of calcium-binding proteins in the cerebellum. *Cerebellum*. 2003; 2:242–262. [PubMed: 14964684]

- Benn, CL., Butler, R., Mariner, L., Nixon, J., Moffitt, H., Mielcarek, M., Woodman, B., Bates, GP. Genetic knock-down of HDAC7 does not ameliorate disease pathogenesis in the R6/2 mouse model of Huntington's disease. *PLoS ONE*. 2009. <http://dx.doi.org/doi:10.1371/journal.pone.0005747>
- Blanco-Garcia N, Asensio-Juan E, de la Cruz X, Martínez-Balbás MA. Autoacetylation regulates P/CAF Nuclear Localization. 2009; 284:1343–1352.
- Bhaskara S, Chyla BJ, Amann JM, Knutson SK, Cortez D, Sun ZW, Hiebert SW. Deletion of histone deacetylase 3 reveals critical role in S phase progression and DNA damage control. *Mol Cell*. 2008; 30:61–72. [PubMed: 18406327]
- Broide RS, Redwine JM, Aftahi N, Young W, Bloom FE, Winrow CJ. Distribution of histone deacetylases 1-11 in the rat brain. *J Mol Neurosci*. 2007; 31:47–58. [PubMed: 17416969]
- Caputi FF, Palmisano M, D'Addario C, Candeletti S, Romualdi P. Effects of acute ethanol exposure on class I HDACs family enzymes in wild-type and BDNF+/- mice. *Drug Alcohol Dep*. 2015; 155:68–75.
- Cassidy LL, Dlugos FF, Dlugos CA. Time course of SERCA 2b and calreticulin expression in Purkinje neurons of ethanol-fed rats with behavioral correlates. *Alcohol Alcohol*. 2013; 48:667–678. [PubMed: 23884168]
- Chouliaras L, van den Hove DLA, Kenis G, van Draanen M, Hof PR, van Os J, Steinbusch HWM, Schmitz C, Rutten BPF. Histone deacetylase 2 in the mouse hippocampus: Attenuation of age-related increase by caloric restriction. *Curr Alzheimer Res*. 2013; 10:868–876. [PubMed: 24093534]
- Darcy MJ, Calvin K, Cavnar K, Ouimet CC. Regional and subcellular distribution of HDAC4 in mouse brain. *J Comp Neurol*. 2010; 518:722–740. [PubMed: 20034059]
- Deshmukh A, Rosenbloom MJ, Pfefferbaum A, Sullivan EV. Clinical signs of cerebellar dysfunction in schizophrenia, alcoholism, and their comorbidity. *Schizophrenia Res*. 2002; 57:281–291.
- Dlugos CA, Pentney RJ. Morphometric analyses of Purkinje and granule cells in aging F344 rats. *Neurobiol Aging*. 1994; 15:435–440. [PubMed: 7969720]
- Dlugos CA, Pentney RJ. Morphometric evidence that the total number of synapses on Purkinje neurons of old F344 rats is reduced after long-term ethanol treatment and restored to control levels after recovery. *Alcohol Alcohol*. 1997; 32:161–72. [PubMed: 9105510]
- Dlugos CA, Pentney RP. Quantitative immunocytochemistry of glia in the cerebellar cortex of old ethanol-fed rats. *Alcohol*. 2001; 23:63–69. [PubMed: 11331103]
- Dlugos CA. Ethanol-related smooth endoplasmic reticulum dilation in Purkinje dendrites of aging rats. *Alc Clin Exp Res*. 2006a; 30:883–891.
- Dlugos CA. Smooth endoplasmic reticulum dilation and degeneration in Purkinje neuron dendrites of aging ethanol-fed female rats. *Cerebellum*. 2006b; 5:155–162. [PubMed: 16818390]
- Dlugos CA. Ethanol-related increases in degenerating bodies in the Purkinje neurons dendrites of aging rats. *Brain Res*. 2008; 1221:98–107. [PubMed: 18559274]
- Dlugos CA. ATF6 and caspase 12 expression in Purkinje neurons in acute slices from adult, ethanol-fed rats. *Brain Res*. 2014; 1577:11–20. [PubMed: 24976582]
- Dlugos CA, Pentney RJ. Morphometric evidence that the total number of synapses on Purkinje neurons of old F344 rats is reduced after long-term ethanol treatment and restored to control levels after recovery. *Alcohol Alcohol*. 1997; 32:161–172. [PubMed: 9105510]
- Druge H, Heinsen H, Heinsen YL. Quantitative studies in ageing Chbb:THOM (Wistar) rats II. Neuron numbers in lobules I, VI_{b+c}, and X. *Bib Anat*. 1986; 28:121–137.
- Escaffit F, Vaute O, Chevillard-Briet M, Segui B, Takami Y, Nakayama T, Trouche D. Cleavage and cytoplasmic delocalization of histone deacetylase 3 are important for apoptosis progression. *Mol Cell Biol*. 2007; 27:554–567. [PubMed: 17101790]
- Finegersh A, Homanics GE. Acute ethanol alters multiple histone modifications at model gene promoters in the cerebral cortex. *Alcohol Clin Exp Res*. 2014; 38:1865–1873. [PubMed: 24942484]
- Graff G, Rei D, Guan JS, Wang WY, Seo J, Hennig KM, Nieland TJF, Fass DM, Kao PF, Kahn M, Su SC, Samiei A, Joseph N, Haggart ST, Delalle I, Tsai LH. *Nature*. 2012; 484:222–228.

- Guan JS, Haggarty SJ, Giacometti E, Dannenberg JH, Joseph N, Gao J, Nieland TJ, Zhou Y, Wang X', Mazitschek R, Bradner JE, Depinho RA, Jaenisch R, Tsai LH. HDAC2 Negatively Regulates Memory Formation and Synaptic Plasticity. *Nature*. 2009; 459:55–60. [PubMed: 19424149]
- Guo, W., Erossey, EL., Zhang, L., Zucca, S., George, OL., Balenzuela, G., Zhao, X. Alcohol exposure decreases CREB binding protein expression and histone acetylation in the developing cerebellum. *PLoS One*. 2011. <http://dx.doi.org/doi:10.1371/journal.pone.0019351>
- Gundersen HJG. Notes on the estimation of the numerical density of arbitrary profiles: the edge effect. *J Microsc*. 1977; 11:219–223.
- Haberland, M., Montgomery, R., Olson, E. The many roles of histone deacetylases in development and physiology: implications for disease and therapy. *Nature Rev Genet*. 2009. <http://dx.doi.org/10.1038/nrg2485>.2008
- Jung ME, Metzger DB. Aberrant histone acetylation promotes mitochondrial respiratory suppression in the brain of alcoholic rats. *J Pharmacol Exp Ther*. 2015; 352:258–266. [PubMed: 25406171]
- Kao HY, Verdel A, Tsai CC, Juguilon CS, Khochbin S. Mechanism for nucleocytoplasmic shuttling of histone deacetylase 7. *J Biol Chem*. 2001; 276:47496–47507. [PubMed: 11585834]
- Kim KK, Adelstein RS, Kawamoto S. Identification of neuronal nuclei (NeuN) as fox 3, a new member of the fox-1 gene family of splicing factors. *J Biol Chem*. 2009; 284:31052–31061. [PubMed: 19713214]
- Kirpich I, Zhang J, Gobejishvili L, Kharebava G, Barker D, Ghare S, Joshi-Barve S, McClain C, Barve S. Binge ethanol-induced HDAC3 down-regulates Cpt1alpha expression leading to hepatic steatosis and injury. *Alc Clin Exp Res*. 2013; 11:1920–1929.
- Knutson SK, Chyla BJ, Amann JM, Bhaskara S, Huppert SS, Hiebert SW. Liver specific deletion of histone deacetylase 3 disrupts metabolic transcriptional networks. *EMBO J*. 2008; 27:1017–1028. [PubMed: 18354499]
- Lagger G, O'Carroll D, Rembold M, Khier H, Tischler J, Weitzer G, Schuettengruber B, Hauser C, Brunmeir R, Jenuwein T, Seiser C. Essential function of histone deacetylase I in proliferation control and CDK inhibitor repression. *EMBO J*. 2002; 21:2672–2681. [PubMed: 12032080]
- Larsen JO, Skalicky M, Viidik A. Does long-term physical exercise counteract age-related Purkinje cell loss? A stereological study of rat cerebellum. *J Comp Neurol*. 2000; 428:213–222. [PubMed: 11064362]
- Li J, Gu X, Ma Y, Calicchio ML, Kong D, Teng YD, Yu L, Crain AM, Vartanian TK, Pasqualini R, Arap W, Libermann TA, Snyder EY, Sidman RL. Nn1 mediates Purkinje cell dendritic development via lysyl oxidase propeptide and NF- κ B signaling. *Neuron*. 2010; 68:45–60. [PubMed: 20920790]
- Li J, Chen J, Ricupero CL, Hart RP, Schwarz MS, Kusnecov A, Herrup K. Nuclear accumulation of HDAC4 in ATM deficiency promotes neurodegeneration in ataxia telangiectasia. *Nat Med*. 2012; 18:783–791. [PubMed: 22466704]
- Li X, Song S, Yu L, Sung-Hwan K, Hung-Ying K. Phosphorylation of the histone deacetylase 7 modulates its stability and association with 14-3-3 proteins. *J Biol Chem*. 2004; 279:34201–34208. [PubMed: 15166223]
- Ma C, D'Mello S. Neuroprotection by histone deacetylase-7 (HDAC7) occurs by inhibition of c-jun expression through a deacetylase-independent mechanism. *J Biol Chem*. 2011; 286:4819–4828. [PubMed: 21118817]
- MacDonald JL, Roskams AJ. Histone deacetylases 1 and 2 are expressed at distinct stages of neuroglial development. *Dev Dyn*. 2008; 237:2256–2267. [PubMed: 18651664]
- Majdzadeh N, Wang L, Morrison BE, Bassel-Duby R, Olson EN, D'Mello SR. HDAC4 inhibits cell cycle progression and protects neurons from cell death. *Dev Neurobiol*. 2008; 68:1076–1092. [PubMed: 18498087]
- Montgomery RL, Davis CA, Potthoff MJ, Haberland M, Fielita J, Qi X, Hill JA, Richardson JA, Olson EN. Histone deacetylases 1 and 2 redundantly regulate cardiac morphogenesis, growth, and contractility. *Genes Dev*. 2007; 21:1790–1802. [PubMed: 17639084]
- Montgomery RL, Hsieh J, Barbosa AC, Richardson JA, Olson EN. Histone deacetylases 1 and 2 control the progression of neural precursors to neurons during brain development, 2009. *PNAS*. 2009; 106:7876–7881. , [PubMed: 19380719]

- Norwood J, Franklin JM, Sharma D, D'Mello SR. Histone deacetylase 3 is necessary for proper brain development. *J Biol Chem*. 2014; 289:34569–34582. [PubMed: 25339172]
- Ohkawa N, Sugisaki S, Tokunaga E, Fujitani K, Hayasaka T, Setou M, Inokuchi K. N-acetyltransferase ARD 1-NAT1 regulates neuronal dendritic development. *Genes Cells*. 2008; 13:1171–1183. [PubMed: 19090811]
- Park PH, Miller R, Shukla SD. Acetylation of histone H3 at lysine 9 by ethanol in rat hepatocytes. *Biochem Biophys Res Comm*. 2003; 306:501–504. [PubMed: 12804592]
- Pandey SC, Ugale R, Zhang H, Tang L, Prakash A. Brain chromatin remodeling: A novel mechanism of alcoholism. *J Neurosci*. 2008; 28:3729–3737. [PubMed: 18385331]
- Parra M, Verdin E. Regulatory signal transduction pathways for class IIa histone deacetylases. *Curr Opin Pharmacol*. 2010; 10:454–460. [PubMed: 20447866]
- Pascual M, Do Couta BR, Alfonso-Loeches S, Aguilar MA, Rodriguez-Arias M, Guerri C. Changes in histone acetylation in the prefrontal cortex of ethanol-exposed adolescent rats are associated with ethanol-induced place conditioning. *Neuropharmacology*. 2012; 62:2309–2319. [PubMed: 22349397]
- Pentney RJ. Measurements of dendritic path lengths provide evidence that ethanol-induced lengthening of the terminal dendritic segments may result from dendritic regression. *Alcohol Alcohol*. 1995; 30:87–96. [PubMed: 7748280]
- Place RF, Noonan EJ, Giardina C. HDACs and the senescent phenotype of WI-38 cells. *BMC Cell Biol*. 2005; 6:37. www.biomedcentral.com/. doi: 10.1186/1471-2121-6/37 [PubMed: 16250917]
- Rigler SK. Alcoholism in the Elderly. *Am Fam Physician*. 2000; 15:1710–1716.
- Rogers J, Zornetzer SF, Bloom FE, Mervis RE. Senescent microstructural changes in rat cerebellum. *Brain Res*. 1984; 292:23–32. [PubMed: 6199084]
- Schneider A, Chatterjee S, Bousiges O, Selvi BR, Swaminathan A, Cassel R, Blanc F, Kundu TK, Boutillier AL. Acetyltransferases (HATS) as targets for neurological therapeutics. *Neurotherapeutics*. 2014; 10:568–588.
- Shahbazian MD, Grunstein M. Functions of site-specific histone acetylation and deacetylation. *Annu Rev Biochem*. 2007; 76:75–100. [PubMed: 17362198]
- Shepard B, Tuma P. Alcohol-induced protein hyperacetylation: Mechanisms and consequences. *World J Gastroenterol*. 2009; 15:1219–1230. [PubMed: 19291822]
- Siming S, Aixiao L, Jiadong L, Candy W, Patrizia CB. Epigenetic memory loss in aging oligodendrocytes in the corpus callosum. *Neurobiol Aging*. 2008; 29:452–463. [PubMed: 17182153]
- Southwood CM, Peppi M, Dryden S, Tainsky MA, Gow A. Microtubule deacetylases, SirT2 and HDAC6, in the nervous system. *Neurochem Res*. 2007; 32:187–195. [PubMed: 16933150]
- Sterio DC. The unbiased estimation of number and sizes of arbitrary particles using the disector. *J Microsc*. 1984; 134:127–136. [PubMed: 6737468]
- Sullivan EV, Rosenbloom MJ, Pfefferbaum A. Pattern of motor and cognitive deficits in detoxified alcoholic men. *Alc Clin Exp Res*. 2000; 24:611–621. .
- Suri D, Bhattacharya A, Vaidya VA. Early stress evokes temporally distinct consequences on the hippocampal transcriptome, anxiety and cognitive behavior. *Int J Neuropsychopharmacology*. 2014; 17:289–301.
- The National Survey on Drug Use and Health. Substance Use among Older Adults, 2002 and 2003 Update. Apr 22. 2005 <http://www.oas.samhsa.gov/2k5/olderadults/olderadults.htm>
- Turano E, Busetto G, Marconi S, Guzzo F, Farinazzo A, Comisso M, Bistaffa E, Angiari S, Musumeci S, Sotgiu S, Boneti B. Neurotoxicity and synaptic plasticity impairment of N-acetylglucosamine polymers: implications for Alzheimer's disease. *Neurobiol Aging*. 2015; 36:1780–1791. [PubMed: 25735590]
- Venkatraman A, Hu YS, Didonna A, Cvetanovic M, Krbanjevic A, Bilesimo P, Opal P. The histone deacetylase HDAC3 is essential for Purkinje cell function, potentially complicating the use of HDAC inhibitors in SCA1. *Human Mol Gen*. 2014; 23:3733–3745.
- Vega RB, Matsuda K, Oh J, Barbosa AC, Yang X, Meadows E, McAnally J, Pomajzi C, Shelton JM, Richardson JA, Karsenty G, Olson EN. Histone deacetylase 4 controls chondrocyte hypertrophy during skeletogenesis. *Cell*. 2004; 119:555–566. [PubMed: 15537544]

- Volmar CH, Wahlestedt C. Histone deacetylases (HDACs) and brain function. *Neuroepigenetics*. 2015; 1:20–27.
- Wang Z, Qin G, Zhao TC. Histone deacetylase 4 (HDAC 4): mechanism of regulations and biological functions. *Epigenomics*. 2014; 6:139–150. <http://dx.doi.org/10.2217/epi13.73>. [PubMed: 24579951]
- Woodruff-Pak DS. Stereological estimation of Purkinje neuron number in C57BL/6 mice and its relation to associative learning. *Neuroscience*. 2006; 141:233–243. [PubMed: 16815479]
- Yao YL, Yang WM. Beyond Histone and deacetylases: an overview of cytoplasmic histone deacetylases and their nonhistone substrates. *J Biomed Biotech*. 2011; 2011:1–15. <http://dx.doi.org/10.1155/2011/146493>.
- Yang XJ, Grégoire S. Class II Histone Deacetylases : from sequence to function, regulation, and clinical implication. *Mol Cell Biol*. 2005; 25:2873–2884. <http://dx.doi.org/10.1128/MCB.25.8.2873-2874.2005>. [PubMed: 15798178]
- Yang XJ, Seto E. The Rpd3/Hda1 family of lysine deacetylases: from bacteria and yeast to mice and men. *Nature Rev Mol Cell Biol*. 2008; 9:206–218. [PubMed: 18292778]
- Yeh HH, Young D, Gelovani JG, Robinson A, Davidson Y, Herholz K, Mann DMA. Histone deacetylase class II and acetylated core histone immunohistochemistry in human brains with Huntington's disease. *Brain Res*. 2013; 1504:16–24. [PubMed: 23419892]
- Yoo JYJ, Larouche M, Goldowitz D. The expression of HDAC1 and HDAC2 during cerebellar cortical development. *Cerebellum*. 2013; 12:534–546. [PubMed: 23436026]
- You C, Zhang H, Sakharkar AJ, Teppen T, Pandey SC. Reversal of deficits in dendritic spines, BDNF and Arc expression in the amygdala during alcohol dependence by HDAC inhibitor treatment. *Int J Neuropsychopharmacol*. 2014; 17:313–322. [PubMed: 24103311]
- Zakhary SM, Ayubcha D, Dileo JN, Jose R, Leheste JR, Horowitz JM, Torres G. Distribution analysis of deacetylase SIRT1 in rodent and human nervous system. *Anat Rec*. 2010; 293:1024–1032.
- Zou, J., Crews, F. Release of neuronal HMGB1 by ethanol through decreased HDAC activity activates brain neuroimmune signaling. *PloS ONE*. 2014. <http://dx.doi.org/10.1371/journal.pone.0087915>

- Purkinje neurons contain Class I histone deacetylases (HDACs) 2,3 and Class IIb HDACs 4 and 7.
- Densities of HDACs 2,3, and 7+ Purkinje neurons decrease significantly with age in adult Fischer 344 rats.
- Density of phosphorylated HDAC4,5, and 7 significantly increases with age as does nuclear localization of HDAC 7.
- Chronic ethanol consumption in aging rats results in decreases in the density and intensity of acetylated histone 2b and 3.

Fig. 1. Slices from confocal stacks of Purkinje neurons of the cerebellar cortex double labelled with antibodies against A) HDAC 2 and calbindin D-28 B) HDAC 3 and calbindin D-28 C) HDAC4 and calbindin D-28, and D) phosphorylated HDAC 4,5,7 and calbindin D-28. A, B, D, Colocalization of HDACs 2, 3, and the phosphorylated form of HDAC 4,5,7 with calbindin D-28 can be visualized within the nucleus (arrows). Colocalization of HDAC4 and calbindin D-28 can be visualized within the cytoplasm (asterisks) of the Purkinje neuron. The dendritic arbor is calbindin positive (white arrowheads) whereas stellate (A) and granule neuron nuclei (B) are HDAC 2(A) and HDAC 3(B) positive (blue arrowheads). Bar on lower right for all images.

Fig. 2. Side by side frames of HDAC 7 and calbindin D-28 + and acetylated histone 3 and calbindin D-28 +3 fields viewed with fluorescein (left) and rhodamine (right) filters. (A) The same HDAC 7+ and calbindin D-28 (arrowheads) Purkinje neuron compared with a different HDAC7- Purkinje neuron (arrow). (B) Acetylated histone 3 + and calbindin D-28 Purkinje neurons (arrowhead) compared to a different acetylated histone 3- Purkinje neuron (arrow). Bar on lower right for all images.

Fig. 3. Mean linear density (+ S.E.M) of HDAC 2+ PN nuclei/ mm in Chow-Fed, Pair-Fed, and EtOH-Fed Rats after 10 (n=30), 20 (n=30) and 40 (n=30) weeks of dietary treatment. The overall significant effect with ANOVA [$F_{\text{overall}}(8,81)=12.889$, $P<.001$] was attributed to decreased HDAC 2 expression with age [$F_{\text{age}}(2,81)=50.78$, $P<.001$] but not to diet [$F_{\text{diet}}(2, 81)=.683$, $P=.508$]. The age changes were due to an overall rise in HDAC 2 + PF and EF PN at 14 compared to 11 months of age (+, $P<.05$) followed by a subsequent decrease in HDAC 2 + PN densities in the CF, PF, and EF rats at 19 compared to 14 months of age (**, $P<.001$).

Fig. 4.

Mean linear density (+ S.E.M) of HDAC 3+ nuclei/mm in Chow-Fed, Pair-Fed, and EtOH-Fed Rats after 10 (n=30), 20 (n=30) and 40 (n=30) weeks of dietary treatment. The overall significant effect with ANOVA [$F_{\text{overall}}(8,81)=3.495$, $P=.002$] was attributed to decreased HDAC 3 expression with age [$F_{\text{age}}(2,81)=11.457$, $P<.001$] but not with diet [$F_{\text{diet}}(2, 81)=1.666$, $P=.195$]. Localization of the effect showed that it was due to decreased densities of HDAC 3+PN in the 11 versus 19 month old CF and PF rats (+, $P=.03$) and in the PF and EF rats of 19 versus 14 months of age. (**, $P=.02$).

Fig. 5.

Mean linear density (+S.E.M.) of HDAC7+ PN /10mm in Chow-Fed, Pair-Fed and EtOH-Fed Rats after 10 (n=30), 20(n=30) and 40 (n=30) weeks of dietary treatment. The overall density of HDAC 7 expressing PN was significant [$F_{\text{overall}}(8,81)=9.042$, $P<.001$] due to age [$F_{\text{age}}(2,81)=35.541$, $P<.001$] but not diet [$F_{\text{diet}}(2,81)=.057$, $P=.945$]. There was a significant decrease in HDAC 7 densities in PN from all treatment groups at 19 compared to 11 months of age (+++, $P=.002$). At 14 months, there was also a significant decline in HDAC 7+ PN in PF rats compared to their 11 month old counterparts (++, $P=.019$). In the CF rats, there was a decrease at 19 months compared to 14 month old rats (+, $P=.014$). In contrast, the density of HDAC7 + PN with staining solely in the nucleus increased [$F_{\text{overall}}(8,81)=4.052$, $P<.001$] with age [$F_{\text{age}}(2,81)=15.823$, $P<.001$] but not with dietary treatment [$F_{\text{diet}}(2,81)=.157$, $P<.855$]. The major increase in sole nuclear expression of HDAC 7 occurred at 19 versus 11 months of age in all dietary groups (*, $P=.02$).

Fig. 6. Mean linear density (+ S.E.M) of phosphorylated HDAC (4,5, 7) + PN/ mm in Chow-Fed, Pair-Fed, and EtOH-Fed rats after 10 (n=30), 20 (n=30) and 40 (n=30) weeks of dietary treatment. The overall significant effect with ANOVA [$F_{\text{overall}}(8,81)=11.398$, $P<.001$] was attributed to increased expression of the phosphorylated form of HDAC 4,5,7 with age [$F_{\text{age}}(2,81)=44.015$, $P<.001$] but not to diet [$F_{\text{diet}}(2,81)=.668$, $P=.516$]. Localization of the effect showed that it was due to increased densities in the phosphorylated form of HDAC 4,5,7 between 11 and 19 mo (**, $P<.001$) and 14 and 19 mo (*, $P<.001$) of age in all dietary groups.

Fig. 7. Mean linear density (+ S.E.M) density of HDAC 4+ Purkinje neurons/ mm in Chow-Fed, Pair-Fed, and EtOH-Fed Rats after 10 (n=30), 20 (n=30) and 40 (n=30) weeks of dietary treatment. There was no significant effect of ANOVA testing on HDAC density [F_{overall} (8.81)=1.616, $P=.133$].

Fig. 8. Mean linear density (+S.E.M.) of acetylated histone 2b and 3+ Purkinje neuron nuclei/mm in Chow-Fed, Pair-Fed, and EtOH-Fed rats after 40 (n=30) weeks of dietary treatment. Two-way ANOVA with treatment and histone as factors, showed an effect of dietary treatment on PN density [$F(5,54)=3.701$, $P=.006$]. The ethanol induced effect was due to decreased density of acetylated histones in the 40 week old EF compared to PF rats of the same age ($P=.004$) and was localized in AH2b ($++$, $P=.009$).

Fig. 9. Mean (+S.E.M) Corrected Total Cell Fluorescence (CTCF) for HDACs 2,3,7 and Acetylated Histone 2b and 3 in Chow-Fed, Pair-Fed, and EtOH-Fed rats after 40 weeks of dietary treatment. There were no overall effects on CTCF [$F_{\text{overall CTCF}}(8,81)=.705$, $P=.686$] for the HDACs. There was, however a main effect on CTCF for the acetylated histones [$F_{\text{overall CTCF}}(5,54)=6,526$, $P=.001$] that was due to treatment [$F_{\text{diet}}(2,54)=10.973$, $P<.001$] and acetylated histone [$F_{\text{histone}}(1,54)=8.291$, $P=.006$]. Localization of the treatment effect showed that AH2b (+, $P=.017$) and AH3 (**, $P=.001$) was lowest in the EF compared to the PF rats. The CTCF for AH3 was also higher in the PF relative to the CF rats (*, $P=.002$).

Table 1
Mean ($\pm$ S.E.M.) Volumes (mm^3) of Molecular Layer Area in 11,14 and 19 mo Chow-Fed, Pair-Fed, and Ethanol-Fed Rats

Age	Chow-Fed Volume (mm^3)	Pair-Fed Volume (mm^3)	Ethanol-Fed Volume (mm^3)
11 months	57.2 $\pm$ 2.0	52.1 $\pm$ 4.4	51.4 $\pm$ 3.5
14 months	61.3 $\pm$ 4.0	57.7 $\pm$ 4.2	58.9 $\pm$ 3.5
19 months	69.9 $\pm$ 2.15	60.1 $\pm$ 2.5	57.4 $\pm$ 1.9