Published in final edited form as:

J Intellect Disabil Res. 2017 December ; 61(12): 1083–1093. doi:10.1111/jir.12424.

Comparing the Broad Socio-cognitive Profile of Youth with Williams Syndrome and 22q11.2 Deletion Syndrome

Omri Weisman^{a,b}, Ruth Feldman^c, Merav Burg-Malki^{a,d}, Miri Keren^{d,e}, Ronny Geva^c, Gil Diesendruck^c, and Doron Gothelf^{a,b,d}

^aThe Behavioral Neurogenetics Center, The Edmond and Lily Safra Children's Hospital, Sheba Medical Center, Tel Hashomer 52621, Israel

^bSagol School of Neuroscience, Tel Aviv University, Tel Aviv 69978, Israel

^cDepartment of Psychology and Gonda Brain Research Center, Bar-Ilan University, Ramat Gan 52900, Israel

^dSackler School of Medicine, Tel Aviv University, Tel Aviv 69978, Israel

^eChild and Adolescent Division, Geha Mental Health Center, Petah Tikva, Israel

Abstract

Background—Numerous studies have assessed the socio-cognitive profile in Williams syndrome (WS) and, independently, in 22q11.2 deletion syndrome (22q11.2DS). Yet, a cross-syndrome comparison of these abilities between individuals with these two syndromes with known social deficits has not been conducted.

Methods—Eighty-two children participated in four study groups: WS (n=18), 22q11.2DS (n=24), age-matched individuals with idiopathic developmental disability (IDD; n=20), and typically developing (TD) controls (n=20). Participants completed four socio-cognitive tests; facial emotion recognition, mental state attribution, differentiating real from apparent emotions, and trait inference based on motives and actions-outcomes.

Results—The current findings demonstrate that children with WS were better in labeling happy faces compared to children with 22q11.2DS, partially reflecting their exaggerated social drive. In the false belief task, however, the WS and IDD groups performed poorly compared to the 22q11.2DS group, possibly due to their difficulty to interpret subtle social cues. When asked to identify the gap between real-negative vs. apparent-positive emotions, the 22q11.2DS group performed similarly to TD children but better than the WS group, possibly due to their anxious personality and their innate bias towards negatively-valence cues. Finally, individuals with WS were more willing to become friends with a story-character even when the character's motives were negative, reflecting their difficulty to avoid potentially harmful real-life situations.

Corresponding Author(s): Prof. Doron Gothelf, M.D., and Dr. Omri Weisman, Ph.D., Sagol School of Neuroscience, Tel Aviv University, and The Behavioral Neurogenetics Center, The Edmond and Lily Safra Children's Hospital, Sheba Medical Center, Tel Hashomer 52621, Israel. Tel: 972-3-530-2663; Fax: 972-77-349-8317; gothelf@post.tau.ac.il, omweisman@gmail.com.

Conflict of Interest: The authors declare having no conflict of interest.

Conclusions—Overall, our multi-facet socio-cognitive battery uncovered strengths and weaknesses in social cognition that are syndrome-specific, shared among the genetic syndromes, or common to the three clinical groups compared to healthy controls. Our findings underscore the need to devise age- and condition-specific assessment tools and intervention programs toward improving these children's socio-cognitive deficits.

Keywords

Velocardiofacial syndrome; DiGeorge; Social cognition; Theory of mind (ToM); Neurogenetic disorders; Children

Introduction

Williams syndrome (WS) and chromosome 22q11.2 deletion syndrome (22q11.2DS) are two neurogenetic disorders with prevalence rates of 1:7,500 and 1:4,000 live births, respectively (Botto et al., 2003; Grati et al., 2015; Shprintzen, 2008; Strømme, Bjørnstad, & Ramstad, 2002). Individuals with WS usually present mild to moderate intellectual disability (with an average full-scale IQ score [FSIQ] of 60 points), coupled with weakness in executive and visuospatial abilities (Bellugi, Wang, & Jernigan, 1994; Zarchi et al., 2014). However, their language and short term verbal memory abilities are considered to be relatively intact (Mervis, 2003).

In terms of their social phenotype, individuals with WS are hyper-social, demonstrate excessive desire to approach people, exhibit indiscriminate trust toward others (Martens, Wilson, & Reutens, 2008), and are extremely attracted to human faces (Pober, 2010; Vivanti, Hocking, Fanning, & Dissanayake, 2017a). As such, their performance on tasks that require the processing of human facial expressions or the categorization of affective facial stimuli is relatively preserved (Dodd, Porter, Peters, & Rapee, 2010; Santos, Rosset, & Deruelle, 2009), and it appears to be commensurate with mental age, rather than chronological age.

Conversely, their understanding of others' mental states (such as others' false beliefs) is impaired and they often have difficulties to understand intricate or ironic social circumstances (Deutsch, Rosse, & Schwartz, 2007). As a result, they usually stumble upon their naïve and over-trusting personality when confronted with social situations that involve disguise, conflicting interests and social complexity (Porter, Coltheart, & Langdon, 2008; Santos & Deruelle, 2009). Other studies have reported that WS individuals show a specific impairment in recognizing angry or negative faces (Plesa-Skwerer, Faja, Schofield, Verbalis, & Tager-Flusberg, 2006; Porter, Shaw, & Marsh, 2010).

The cognitive profile of individuals with 22q11.2DS is characterized by an average FSIQ score of 75 points (at the borderline range; Swillen et al., 1999), alongside adequate visuospatial and numerical processing but impaired verbal processing (Bearden et al., 2001; Gothelf, 2007; Simon, Bearden, Mc-Ginn, & Zackai, 2005). The social profile in 22q11.2DS is characterized by extreme shyness, avoidance, coupled with restricted affect (Schonherz et al., 2014; Shprintzen, 2008; Swillen & McDonald-McGinn, 2015), and deficient ability to categorize facial stimuli (Andersson et al., 2008; Campbell et al., 2010). Other studies point for impairments in false belief and emotion attribution abilities in 22q11.2DS compared to

typically-developing (TD) controls (Campbell, McCabe, Melville, Strutt, & Schall, 2015). A recent review of 16 studies further confirmed that social cognition is indeed impaired in 22q11.2DS, mostly abilities such as emotion processing and complex theory of mind (Norkett, Lincoln, Gonzalez-Heydrich, & D'Angelo, 2017).

In addition, several studies have compared neuropsychiatric and neurocognitive profiles between the two syndromes (Weinberger et al., 2016; Zarchi et al., 2014). For example, Zarchi and colleagues (2014) showed that individuals with WS are characterized by more severe impairments in both executive and visuospatial functions compared to 22q11.2DS, but that the two syndromes exhibit similarly deficient performance-IQ and verbal-IQ, when compared to individuals with idiopathic developmental disability (IDD) and TD controls.

Yet, research that looks into and compares between the socio-cognitive profiles of individuals harboring these microdeletions is missing. Moreover, contrasting the syndromes against individuals with IDD and TD controls is likely to be informative, as it may uncover patterns that are unique to the syndromes but differ from that seen in IDD, or that are shared between the three clinical groups versus TD controls. This comparison is further expected to promote our understanding of nonverbal deficits in these neurogenetic conditions, potentially paving the way for the development of adequate interventions. While some studies compared between the socio-cognitive abilities of WS, 22q11.2DS, and those with autism spectrum disorder (ASD), findings highlight fundamental differences between the groups which merit that such an investigation will be conducted independently. As such, the current study focused on the two syndromes while referring to the IDD and TD groups as controls.

Considering the above, the current study sought to compare the socio-cognitive performance of children with WS, 22q11.2DS, IDD, and TD, across a range of tasks, including: (1) facial emotion recognition task – testing the ability to infer emotional states from facial stimuli; (2) first-order false belief task – testing the ability to represent others' thoughts and perspectives; (3) a paradigm that assesses the ability to differentiate between real (i.e. concealed) vs. apparent emotions; and, finally, (4) a paradigm that tests one's capacity to infer others' traits based on their motives and actions-outcomes.

We hypothesized that as compared to the 22q11.2DS and IDD groups, individuals with WS will demonstrate better ability to recognize emotions from facial stimuli, especially when stimuli are coupled with verbal descriptors. This superior ability would be emotion-specific, given prior findings of impaired recognition of angry faces in WS. However, individuals with WS are expected to perform worse than the 22q11.2DS and IDD groups on tasks that require to grasp others' mental states (also referred to as mental attribution). Finally, individuals with 22q11.2DS are likely to recognize others' negative emotions even if they are concealed due to their anxious and alert personality and their inclination towards negative social cues (Shprintzen, 2008).

Method

Participants

The study sample included 82 children: 18 participants with WS, 24 participants with 22q11.2DS, 20 participants with IDD, and 20 TD participants. The three clinical groups were between 7 – 15 years of age. The chronological age (CA) of TD controls served as a proxy for their mental age (MA), and as the basis for matching them with the three clinical groups (see Table 1 for additional data). The diagnosis of 22q11.2DS and WS was confirmed in all affected individuals by fluorescent in situ hybridization (FISH) and by multiplex ligation probe amplification (MLPA) (Michaelovsky et al., 2012). The WS and 22q11.2DS participants were recruited from the Behavioral Neurogenetics Center at a large tertiary referral center in Israel. Participants with IDD were recruited from schools for special education for children and adolescents with developmental disability. TD controls were recruited through advertisements within the local community. They were all students in mainstream classes and none had a major psychopathology. The study was approved by the local Institutional Review Board (IRB) committee. Written informed consents were obtained from the parents of all children prior to their participation in the study.

Behavioral paradigms

Facial emotion recognition—Participants' ability to correctly *label* and *match* facial stimuli was assessed using a task especially developed for this study. Stimuli consisted of 24 short colored video-clips of male and female actors (Baron-Cohen, Tager-Flusberg, & Lomardo, 2013) presented on a computer screen (23 × 30 cm wide), representing six basic emotions: happiness, sadness, anger, fear, disgust, and surprise; i.e., four clips for each emotion. The average duration of the clips was 5.47 seconds ($sd = 0.76$; range: 3.2 – 7.4 sec). The bank of stimuli was originally developed for learning and practicing purposes for individuals with ASD and later adopted for research. For details about the validity of stimuli, refer to www.jkp.com/mindreading. Moreover, as part of a pilot test, 20 students were presented with 8 video clips for each emotion (48 videos in total). Participants were then asked to specify the emotion shown and its intensity level (on a 1 – 4 likert scale). Half of stimuli (24 videos) were selected for the study based on the likelihood that they will be identified correctly and with equivalent mean intensity between the different emotions.

Participants were asked to answer two questions about each video-clip. In the *labeling* module, the experimenter provided four emotion-labels (e.g., “happy”, “sad”, etc.), and participants were prompted to choose the label that best fits the stimulus they just saw. In the *matching* module, participants watched the same video clip again followed by three videos: two distractor videos and one identical to the target stimulus. The entire task took about 35 – 40 minutes to complete. Correct answer credited participants with 1 point. Since less than half of TD participants gave correct answers on two of the labeling items (representing fear) these items were excluded from further analysis, resulting in a total score of 22 in the labeling module and 24 in the matching module.

False belief—Participants were assessed with two false-belief tasks (Wimmer & Perner, 1983). In the first task, participants were shown a familiar container (with a picture of

cookies on it) that, when opened, revealed an unexpected content (pencils). Children were asked: “*What your mother will think is in the box?*”. This task requires to impute to another person a false belief about the content of the box (which differs from both reality and what the child knows to be true). A correct answer would credit 1 point.

In the second false belief task, participants were presented with four scenarios in which a character places an object in a specific location and then exits the room. In his/her absence, the object is moved to a new location. Children were asked three questions: “*Upon his return, where will the character think the object is?*”; “*Where will the character look for the object?*”; “*Why will the character look for the object there?*”. Two (out of four) scenarios were adapted from an earlier study (Shatz, Diesendruck, Martinez-Beck, & Akar, 2003), while two scenarios were especially created for the current study. Providing correct answers to the first two questions would credit 1 point. Referring to the character’s false belief in the third question would credit 2 points. Scoring in this task ranged between 0 – 4 points for each of the scenarios (16 points in total for 4 scenarios), thus the overall scoring in the two false belief tasks ranged between 0 – 17 points.

Differentiating real from apparent emotions—In this task, adapted from Harris et al. (Harris, Donnelly, Guz, & Pitt-Watson, 1986), children were asked to listen to four stories in which it would be appropriate for the character to feel a positive or negative emotion but to hide that emotion. Participants were then asked: “*How will the character look like (referring to perceived emotion)?*”; “*How will the character really feel (referring to real emotion)?*”; Next, the experimenter assessed whether the child grasped the difference between real and apparent emotion (in the right direction); finally, participants were requested to elaborate (justify) their answers. Each correct answer in the first three questions would credit 1 point. Explaining why the character expresses different emotions than the one felt would credit 2 additional points (but only 1 point for answering correctly without referring to the hidden emotion). Thus, scoring in this task ranged between 0 – 8, reflecting overall task performance.

Traits inference based on motives and outcomes—Children’s ability to infer others’ traits based on their motives or their actions was assessed using a task developed by Heyman and Gelman (Heyman & Gelman, 1998). Participants were presented with four illustrated stories in which a character’s motives were either positive or negative, while the outcome of his/her action was the opposite. For example, in one of the stories, a girl (the character) sprinkles water on her friend. A positive motive in that story would be that the girl wanted to make her friend feel cooler because of the weather, although the outcome was negative (i.e., the friend got angry about getting wet). An opposite scenario would read as negative motive (i.e., to get the other person wet) coupled with a positive outcome (i.e., that person enjoyed getting wet). Children were asked four questions: “*Do you think the character is nice/mean?*”; “*Do you think what the character did was nice/mean?*”; “*Would you like to be friends with the character?*” Possible answers included: “yes”, “maybe”, and “no”. Answering “maybe” would qualify for half a point. The last question was, “*Do you think the character would help another child who fell in the park?*”. Answers that matched the character’s motive were regarded as correct (i.e., worthy of 1 point).

Procedure

The initial meeting with children in the three clinical groups and their parents (usually the mother) were held at the Behavioral Genetics Center and the adjunct clinic. An intelligence test was conducted at the first session; FSIQ scores of individuals in the three clinical groups were assessed using the Wechsler Intelligence Scale for Children-Revised (WISC-R; Wechsler, 1974). The FSIQ scores of control participants are missing because of the IRB restraints on administering IQ tests to TD children. Second meeting was held either at the clinic or at children's home, as preferred (8 of 18 WS subjects asked to conduct the second meeting at home, so thus 13 of 24 22q11.2DS subjects, and 9 of 20 IDD subjects). Children in the TD group were tested at home.

The socio-cognitive battery was employed by one of the authors of the current study (M. B-M.) as part of her doctoral thesis and clinical training. The emotion recognition paradigm took 30–45 minutes to complete, including breaks to accommodate to the child's attention span etc. Next, the three socio-cognitive tests (mental state attribution, differentiating real from apparent emotions, and trait inference) were employed according to a pre-defined randomized order. Occasional breaks between the tests were taken when needed. The parent was not present at the room during the tests.

Planned analyses

Two steps were taken to count for group differences in MA and FSIQ scores. As a first step, statistical analyses were conducted while controlling for MA. As a second step, each analysis was conducted twice - once with a sample of 82 participants which differed in FSIQ (with 22q11.2DS having the highest score, then IDD, and finally WS), and a second analysis with a smaller sample of 70 participants (comprising the three clinical groups) which were matched in terms of FSIQ. Since the findings of the larger sample did not differ from the findings of the smaller sample, we further detail the analysis based on the larger cohort ($n = 82$). Statistical analyses conducted for each of the tasks is detailed at the beginning of each of the below mentioned sections. Post-hoc *Bonferroni* comparisons were computed where appropriate.

Results

Facial emotion recognition

Separate mixed-design analysis of covariance (ANCOVA) were computed for each of the modules in the facial emotion recognition task (i.e., labeling and matching) with valence being the within-subject variable and group being the between-subject variable, respectively. Participants' FSIQ score was treated as covariate in the analysis (as well as in the analyses detailed further below).

In the labeling module, an interaction of group by valence was found, $F_{(3,73)} = 5.01$, $p < .01$. Breaking down this interaction by valence revealed that the WS group was better in labeling positive emotions compared to the 22q11.2DS group, $F_{(3,73)} = 2.77$, $p < .05$ (see Table 2 for additional data). The effect was dominated by WS labeling "happy" faces more accurately than the 22q11.2DS, $F_{(3,73)} = 2.54$, $p < .05$, as revealed by a post-hoc analysis. When asked to

label negative emotions, the TD group showed superior ability compared to the three clinical groups, $F_{(3,73)} = 8.22, p < .001$. The latter finding replicated in the matching module, with TD participants exhibiting higher performance than all three clinical groups when asked to match negative or positive emotions, $p's < .001$. All participants labeled positive stimuli more accurately than negative stimuli, above and beyond group identity, $F_{(1,73)} = 9.86, p < .01$.

False belief

Univariate ANCOVA was employed to uncover group differences in the false belief task. A main effect of Group was found, $F_{(3,73)} = 13.40, p < .001$. Bonferroni post-hoc revealed that TD participants scored higher compared to the three clinical groups, and that the 22q11.2DS group showed superior performance compared to the WS and IDD groups (see Figure 1).

Differentiating real from apparent emotions

Mixed-design ANCOVA with emotion type (apparent/real) and stimuli valence (positive/negative) as the within-subject variables was computed to assess children's ability to differentiate real from apparent emotions. FSIQ was treated as covariate.

A group main effect was found, $F_{(3,69)} = 8.31, p < .001$. Post-hoc comparisons revealed that the TD group had higher scores than the WS and IDD groups, while the 22q11.2DS group performed similarly to the rest of the groups (see Figure 2). However, when asked to identify the shift between the character's real and apparent emotions, the WS and IDD groups showed lower abilities compared to TD controls, whereas the 22q11.2DS and TD groups performed comparably well, $F_{(3,73)} = 5.40, p < .01$. All three clinical groups had difficulty to identify the shift between *real-positive* and *apparent-negative* emotions compared to TD, $F_{(3,73)} = 8.98, p < .001$.

Finally, a main effect for emotion type was found, as participants across all groups identified real emotions more accurately than apparent emotion, $F_{(1,69)} = 5.05, p < .05$ (mean $\pm$ SD; 1.74 ± 0.35 vs. 1.35 ± 0.69). An interaction between emotion type and stimuli valence emerged, with real emotions of negative valence more accurately identified compared to real emotions of positive valence, $F_{(1,69)} = 6.35, p < .05$.

Inferring traits from motives and actions-outcomes

Mixed-design ANCOVA with motive (positive/negative) as the within-subject variable and group identity as the between-subject variable was computed to examine whether participants rely on the character's motives when asked about his/her traits. Next, finer ANCOVA analyses were conducted to evaluate possible differences in the four questions. Scores on each question for each motive valence were entered as dependent variables (scoring ranged between 0–2) and group identity served as the between-subject variable. Analysis was conducted separately for the positive and negative motive scenarios.

Whether the motive was positive or negative did not alter task performance for any of the groups, $p = ns$. In addition, the three clinical groups exhibited significantly lower ability to infer others' traits based on their motives compared to the TD group, $F_{(3,73)} = 7.77, p < .001$ (TD: 3.28 ± 0.79 > 22q11.2DS: 2.56 ± 0.94 , IDD: 2.36 ± 0.97 , WS: 1.96 ± 0.86). When

presented with positive scenarios, the TD group answered the first question (“*Do you think the character is nice/mean?*”) more accurately than the WS and IDD groups, while the 22q11.2DS group performed comparable to the rest (TD: 1.75 ± 0.63 > IDD: 1.05 ± 0.91 , WS: 1.0 ± 0.73 ; 22q11.2DS: 1.52 ± 0.66). This pattern extended to the second question (“*Do you think what the character did was nice/mean?*”), $F_{(3,73)} = 5.66$, $p < .01$ (TD: 1.6 ± 0.75 > 22q11.2DS: 1.08 ± 0.66 , IDD: 0.89 ± 0.65 , WS: 0.75 ± 0.77). No difference was found with respect to the last two questions.

Finally, a group main effect was found when children were asked whether they would like to become friends with the character when the character’s motive was described as negative, $F_{(3,73)} = 7.10$, $p < .001$. Post-hoc comparisons indicated that children with WS performed worse in this test than children from the other groups, i.e., they were more willing to become friends with the character (Figure 3).

Discussion

To the best of our knowledge, this study is the first to directly compare the socio-cognitive profile of children with WS to that of children with 22q11.2DS. As expected, our results uncover a specific socio-cognitive profile in each disorder. In line with our first hypothesis, the WS group was found to be better at labeling happy faces compared to the 22q11.2DS group, possibly due to their disposition towards positive facial stimuli coupled with their intense social appetite (Gagliardi et al., 2003; Plesa-Skwerer et al., 2006). This pattern was not driven by a general bias to recognize other facial stimuli as happy as their error rate (i.e., erroneously labeling different facial stimuli as happy) was similar to that of other groups. Since it is easier to identify “happy face” based on specific parts of the face as compared to other emotions (Calder, Young, Keane, & Dean, 2000), our results are consistent with studies showing that individuals with WS focus on specific parts of the face rather than on it entirely (Tager-Flusberg, Plesa-Skwerer, Faja, & Joseph, 2003). Indeed, findings emerging from brain imaging studies further suggest that the brain activity of individuals with WS vs. TD is altered during face processing tasks, with happy faces inducing greater right amygdala activation and fearful faces inducing lower overall amygdala activation (Haas et al., 2009).

Additionally, the WS group demonstrated intact ability to label facial stimuli but poor ability to match the same stimuli, compared to TD controls. Their performance in the labeling module can be explained by their relative verbal proficiency and their tendency to rely on verbal descriptors for this matter (Brock, Jarrold, Farran, Laws, & Riby, 2007; Vivanti, Hocking, Fanning, & Dissanayake, 2016). Their poor performance in the matching module is likely to be explained by their disrupted visuospatial capacity (Bellugi et al., 1994). In 22q11.2DS, the poor face recognition ability may be sub-served by their difficulty to process facial configurations as a unified, *gestalt*-like, unit (Glaser et al., 2010), coupled with disruption in sub-cortical regions responsible for face processing, including the superior temporal sulcus and the amygdala (Drew et al., 2011; Johnson, 2005; Leleu et al., 2016; Pessoa & Adolphs, 2010).

In the false belief tasks a different pattern emerged; individuals with WS and IDD performed relatively well, but worse than individuals with 22q11.2DS and TD controls (M. A. Porter et

al., 2008). A plausible explanation is that false belief involve several cognitive functions – such as perceptual and auditory processing, working memory, and executive control – all of which are abnormal in WS (Meyer-Lindenberg et al., 2005; Porter, Coltheart, & Langdon, 2007; Zarchi et al., 2015).

When asked to identify situations where positive emotions were concealed by negative ones, the three clinical groups performed below par compared to TD controls. This pattern replicated also when real-negative emotions were concealed by apparent-positive emotions, except that 22q11.2DS performed comparably well to TD in this condition. The poor performance of individuals with WS in both conditions parallels their difficulty to identify others' concealed intentions and thoughts, which often prevents them from developing alarming signals when they face potentially harmful circumstances (Riby, Kirk, Hanley, & Riby, 2014). Children with 22q11.2DS performed relatively well when asked to recognize negative emotions concealed by positive ones. A possible explanation for that may be nested in their anxious temperament and their innate bias towards negative social cues (Azuma et al., 2015).

Limitations

Limitations of the current study may include the relatively small sample size of each group, the fact that 2 stimuli from the emotion recognition task were omitted after TD participants did not recognize them adequately, and, the lack of reliable diagnostic instruments - such as the Read the Mind in the Eyes Test [RMET] (Baron-Cohen, Wheelwright, Hill, Raste, & Plumb, 2001), or the Interpersonal Reactivity Index [IRI] (Davis, 1980) - to assess autistic characteristics among enrolled children.

In addition, the absence of an ASD group to serve as an additional control group for the genetic syndromes is a methodological caveat of the current study that merits future research. The importance of contrasting the syndromes with an ASD group is strengthened given the well-characterized neurodevelopmental deficits in ASD (despite the lack of clear genetic loci), and initial findings showing fundamental discrepancies in some socio-cognitive abilities between ASD, 22q11.2DS, and WS (McCabe et al., 2013; Vivanti, Hocking, Fanning, Uljarevic, et al., 2017; Vivanti, Hocking, Fanning, & Dissanayake, 2017b).

Future research into social cognition in children with Williams and 22q11.2DS is recommended to endorse the points above mentioned when aiming to highlight syndrome-unique, syndrome-shared, and neurodevelopment-shared socio-cognitive abilities of affected individuals. Moreover, the fourth behavioral tasks that was employed (which included social scripts) suggest that stories may be a prolific manner to expose children with WS to different social circumstances and to educate train them how to respond in potentially harmful situations to reduce their risk of being abused.

Conclusions

The multi-task broad battery of abilities employed here enabled us to expose strengths and weaknesses in several socio-cognitive domains that are either specific to one of the

syndromes, shared between the syndromes, or shared between the three clinical (intellectual disabilities) groups compared to controls. Together with the existing literature, our findings support the development of age- and condition- appropriate clinical evaluation practices and intervention approaches that are geared towards alleviating the suffering of affected individuals and their families.

Acknowledgments

The authors thank the Keren Shalem Fund for the Development of Services for the Retarded in the Local Councils, the Gonda Family Foundation, the Binational Science Foundation, grant number 2011378 and the National Institute of Mental Health of the National Institutes of Health under Award Number U01MH101722 for their support of this research. The work conducted by Dr. Merav Burg-Malki was supported by Bar-Ilan University excellence scholarship for doctoral students. The work conducted by Dr. Omri Weisman was supported by a fellowship from the Sagol School of Neuroscience, Tel Aviv University.

References

- Andersson F, Glaser B, Spiridon M, Debbané M, Vuilleumier P, Eliez S. Impaired Activation of Face Processing Networks Revealed by Functional Magnetic Resonance Imaging in 22q11.2 Deletion Syndrome. *Biological Psychiatry*. 2008; 63(1):49–57. <http://doi.org/10.1016/j.biopsych.2007.02.022>. [PubMed: 17651704]
- Azuma R, Deeley Q, Campbell LE, Daly EM, Giampietro V, Brammer MJ, Murphy DG. An fMRI study of facial emotion processing in children and adolescents with 22q11.2 deletion syndrome. *Journal of Neurodevelopmental Disorders*. 2015; 7(1):1. <http://doi.org/10.1186/1866-1955-7-1>. [PubMed: 25972975]
- Baron-Cohen, S., Tager-Flusberg, H., Lomardo, MV. *Understanding Other Minds - Perspectives from Developmental Social Neuroscience*. Oxford University Press; 2013. <http://doi.org/10.1007/s13398-014-0173-7.2>
- Baron-Cohen S, Wheelwright S, Hill J, Raste Y, Plumb I. The “Reading the Mind in the Eyes” Test Revised Version: A Study with Normal Adults, and Adults with Asperger Syndrome or High-functioning Autism. *Journal of Child Psychology and Psychiatry*. 2001; 42(2):241–251. <http://doi.org/10.1111/1469-7610.00715>. [PubMed: 11280420]
- Bearden CE, Woodin MF, Wang PP, Moss E, McDonald-McGinn D, Zackai E, Cannon TD. The neurocognitive phenotype of the 22q11.2 deletion syndrome: selective deficit in visual-spatial memory. *Journal of Clinical and Experimental Neuropsychology*. 2001; 23(4):447–64. <http://doi.org/10.1076/jcen.23.4.447.1228>. [PubMed: 11780945]
- Bellugi, U., Wang, PP., Jernigan, TL. *Williams Syndrome: an Unusual Neuropsychological Profile; Atypical Cognitive Deficits in Developmental Disorders: Implications for Brain Function*. 1994. p. 1-59. Retrieved from http://www.wsfamilyofhope.org/uploads/An_Unusual_Neuropsychological_Profile.pdf
- Botto LD, May K, Fernhoff PM, Correa A, Coleman K, Rasmussen SA, Campbell RM. A Population-Based Study of the 22q11.2 Deletion: Phenotype, Incidence, and Contribution to Major Birth Defects in the Population. *Pediatrics*. 2003; 112(1):101–107. <http://doi.org/10.1542/peds.112.1.101>. [PubMed: 12837874]
- Brock J, Jarrold C, Farran EK, Laws G, Riby DM. Do children with Williams syndrome really have good vocabulary knowledge? Methods for comparing cognitive and linguistic abilities in developmental disorders. *Clinical Linguistics & Phonetics*. 2007; 21(9):673–88. <http://doi.org/10.1080/02699200701541433>. [PubMed: 17701755]
- Calder AJ, Young AW, Keane J, Dean M. Configural information in facial expression perception. *Journal of Experimental Psychology: Human Perception and Performance*. 2000; 26(2):527–551. <http://doi.org/10.1037/0096-1523.26.2.527>. [PubMed: 10811161]
- Campbell LE, McCabe KL, Melville JL, Strutt PA, Schall U. Social cognition dysfunction in adolescents with 22q11.2 deletion syndrome (velo-cardio-facial syndrome): Relationship with

- executive functioning and social competence/functioning. *Journal of Intellectual Disability Research*. 2015; 59(9):845–859. <http://doi.org/10.1111/jir.12183>. [PubMed: 25726953]
- Campbell L, McCabe K, Leadbeater K, Schall U, Loughland C, Rich D. Visual scanning of faces in 22q11.2 deletion syndrome: Attention to the mouth or the eyes? *Psychiatry Research*. 2010; 177(1–2):211–215. <http://doi.org/10.1016/j.psychres.2009.06.007>. [PubMed: 20381171]
- Davis MH. Interpersonal Reactivity Index. *JSAS Catalog of Selected Documents in Psychology*. 1980; 10(1980):14–15. <http://doi.org/10.1037/t01093-000>.
- Deutsch SI, Rosse RB, Schwartz BL. Williams syndrome: a genetic deletion disorder presenting clues to the biology of sociability and clinical challenges of hypersociability. *CNS Spectrums*. 2007; 12(12):903–7. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/18163035>. [PubMed: 18163035]
- Dodd HF, Porter MA, Peters GL, Rapee RM. Social approach in pre-school children with Williams syndrome: The role of the face. *Journal of Intellectual Disability Research*. 2010; 54(3):194–203. <http://doi.org/10.1111/j.1365-2788.2009.01241.x>. [PubMed: 20070473]
- Drew LJ, Crabtree GW, Markx S, Stark KL, Chaverneff F, Xu B, Karayiorgou M. The 22q11.2 microdeletion: Fifteen years of insights into the genetic and neural complexity of psychiatric disorders. *International Journal of Developmental Neuroscience*. 2011; 29(3):259–281. <http://doi.org/10.1016/j.ijdevneu.2010.09.007>. [PubMed: 20920576]
- Gagliardi C, Frigerio E, Burt DM, Cazzaniga I, Perrett DI, Borgatti R. Facial expression recognition in Williams syndrome. *Neuropsychologia*. 2003; 41(6):733–738. [http://doi.org/10.1016/S0028-3932\(02\)00178-1](http://doi.org/10.1016/S0028-3932(02)00178-1). [PubMed: 12591030]
- Glaser B, Debbane M, Ottet MC, Vuilleumier P, Zesiger P, Antonarakis SE, Eliez S. Eye Gaze During Face Processing in Children and Adolescents With 22q11.2 Deletion Syndrome. *Journal of the American Academy of Child and Adolescent Psychiatry*. 2010; 49:665–674. <http://doi.org/DOI10.1016/j.jaac.2010.04.004>. [PubMed: 20610136]
- Gothelf D. Velocardiofacial Syndrome. *Child and Adolescent Psychiatric Clinics of North America*. 2007; 16(3):677–693. <http://doi.org/10.1016/j.chc.2007.03.005>. [PubMed: 17562586]
- Grati FR, Molina Gomes D, Ferreira JCPB, Dupont C, Alesi V, Gouas L, Vialard F. Prevalence of recurrent pathogenic microdeletions and microduplications in over 9500 pregnancies. *Prenatal Diagnosis*. 2015; 35(8):801–9. <http://doi.org/10.1002/pd.4613>. [PubMed: 25962607]
- Haas BW, Mills D, Yam A, Hoeft F, Bellugi U, Reiss A. Genetic influences on sociability: heightened amygdala reactivity and event-related responses to positive social stimuli in Williams syndrome. *The Journal of Neuroscience*: The Official Journal of the Society for Neuroscience. 2009; 29(4):1132–9. <http://doi.org/10.1523/JNEUROSCI.5324-08.2009>. [PubMed: 19176822]
- Harris PL, Donnelly K, Guz GR, Pitt-Watson R. Children's understanding of the distinction between real and apparent emotion. *Child Development*. 1986; 57(4):895–909. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/3757608>. [PubMed: 3757608]
- Heyman GD, Gelman SA. Young Children Use Motive Information to Make Trait Inferences. *Developmental Psychology*. 1998; 34(2):310–321. <http://doi.org/10.1037/0012-1649.34.2.310>. [PubMed: 9541783]
- Johnson MH. Subcortical face processing. *Nature Reviews. Neuroscience*. 2005; 6(10):766–74. <http://doi.org/10.1038/nrn1766>. [PubMed: 16276354]
- Leleu A, Saucourt G, Rigard C, Chesnoy G, Baudouin J-Y, Rossi M, Demily C. Facial emotion perception by intensity in children and adolescents with 22q11.2 deletion syndrome. *European Child & Adolescent Psychiatry*. 2016; 25(3):297–310. <http://doi.org/10.1007/s00787-015-0741-1>. [PubMed: 26149605]
- Martens MA, Wilson SJ, Reutens DC. Research Review: Williams syndrome: a critical review of the cognitive, behavioral, and neuroanatomical phenotype. *Journal of Child Psychology and Psychiatry*. 2008; 49(6):576–608. <http://doi.org/10.1111/j.1469-7610.2008.01887.x>. [PubMed: 18489677]
- M McCabe KL, Melville JL, Rich D, Strutt PA, Cooper G, Loughland CM, Campbell LE. Divergent Patterns of Social Cognition Performance in Autism and 22q11.2 Deletion Syndrome (22q11DS). *Journal of Autism and Developmental Disorders*. 2013; 43(8):1926–1934. <http://doi.org/10.1007/s10803-012-1742-2>. [PubMed: 23292161]

- Mervis CB. Williams syndrome: 15 years of psychological research. *Developmental Neuropsychology*. Special Issue: Williams Syndrome. 2003; 23(1–2):1–12. <http://doi.org/10.1080/87565641.2003.9651884>.
- Meyer-Lindenberg A, Hariri AR, Munoz KE, Mervis CB, Mattay VS, Morris Ca, Berman KF. Neural correlates of genetically abnormal social cognition in Williams syndrome. *Nature Neuroscience*. 2005; 8(8):991–3. <http://doi.org/10.1038/nm1494>. [PubMed: 16007084]
- Michaelovsky E, Frisch A, Carmel M, Patya M, Zarchi O, Green T, Gothelf D. Genotype-phenotype correlation in 22q11.2 deletion syndrome. *BMC Medical Genetics*. 2012; 13(1):122. <http://doi.org/10.1186/1471-2350-13-122>. [PubMed: 23245648]
- Norkett, EM., Lincoln, SH., Gonzalez-Heydrich, J., D'Angelo, EJ. Social cognitive impairment in 22q11 deletion syndrome: A review. *Psychiatry Research*. 2017. <http://doi.org/10.1016/j.psychres.2017.01.103>
- Pessoa L, Adolphs R. Emotion processing and the amygdala: from a “low road” to “many roads” of evaluating biological significance. *Nature Reviews. Neuroscience*. 2010; 11(11):773–83. <http://doi.org/10.1038/nrn2920>. [PubMed: 20959860]
- Plesa-Skwerer D, Faja S, Schofield C, Verbalis A, Tager-Flusberg H. Perceiving Facial and Vocal Expressions of Emotion in Individuals With Williams Syndrome. *American Journal on Mental Retardation*. 2006; 111(1):15. [http://doi.org/10.1352/0895-8017\(2006\)111\[15:PFAVEO\]2.0.CO;2](http://doi.org/10.1352/0895-8017(2006)111[15:PFAVEO]2.0.CO;2). [PubMed: 16332153]
- Pober BR. Williams-Beuren Syndrome. *New England Journal of Medicine*. 2010; 362(3):239–252. <http://doi.org/10.1056/NEJMra0903074>. [PubMed: 20089974]
- Porter MA, Coltheart M. Cognitive heterogeneity in Williams syndrome. *Developmental Neuropsychology*. 2005; 27(2):275–306. <http://doi.org/10.1207/s15326942dn2702>. [PubMed: 15753050]
- Porter MA, Coltheart M, Langdon R. The neuropsychological basis of hypersociability in Williams and Down syndrome. *Neuropsychologia*. 2007; 45(12):2839–2849. <http://doi.org/10.1016/j.neuropsychologia.2007.05.006>. [PubMed: 17597166]
- Porter MA, Coltheart M, Langdon R. Theory of mind in Williams syndrome assessed using a nonverbal task. *Journal of Autism and Developmental Disorders*. 2008; 38(5):806–814. <http://doi.org/10.1007/s10803-007-0447-4>. [PubMed: 17874179]
- Porter, Ma, Shaw, Ta, Marsh, PJ. An unusual attraction to the eyes in Williams-Beuren syndrome: a manipulation of facial affect while measuring face scanpaths. *Cognitive Neuropsychiatry*. 2010; 15(6):505–30. <http://doi.org/10.1080/13546801003644486>. [PubMed: 20432078]
- Riby DM, Kirk H, Hanley M, Riby LM. Stranger danger awareness in Williams syndrome. *Journal of Intellectual Disability Research*. 2014; 58(6):572–582. <http://doi.org/10.1111/jir.12055>. [PubMed: 23718278]
- Santos A, Deruelle C. Verbal Peaks and visual Valleys in theory of mind ability in Williams syndrome. *Journal of Autism and Developmental Disorders*. 2009; 39(4):651–659. <http://doi.org/10.1007/s10803-008-0669-0>. [PubMed: 19039658]
- Santos A, Rosset D, Deruelle C. Human versus non-human face processing: Evidence from Williams syndrome. *Journal of Autism and Developmental Disorders*. 2009; 39(11):1552–1559. <http://doi.org/10.1007/s10803-009-0789-1>. [PubMed: 19557509]
- Schonherz Y, Davidov M, Knafo A, Zilkha H, Shoval G, Zalsman G, Gothelf D. Shyness discriminates between children with 22q11.2 deletion syndrome and Williams syndrome and predicts emergence of psychosis in 22q11.2 deletion syndrome. *Journal of Neurodevelopmental Disorders*. 2014; 6(1):3. <http://doi.org/10.1186/1866-1955-6-3>. [PubMed: 24517288]
- Shatz M, Diesendruck G, Martinez-Beck I, Akar D. The influence of language and socioeconomic status on children's understanding of false belief. *Developmental Psychology*. 2003; 39(4):717–729. <http://doi.org/10.1037/0012-1649.39.4.717>. [PubMed: 12859125]
- Shprintzen RJ. Velo-cardio-facial syndrome: 30 Years of study. *Developmental Disabilities Research Reviews*. 2008; 14(1):3–10. <http://doi.org/10.1002/ddrr.2>. [PubMed: 18636631]
- Simon TJ, Bearden CE, Mc-Ginn DM, Zackai E. Visuospatial and numerical cognitive deficits in children with chromosome 22q11.2 deletion syndrome. *Cortex; a Journal Devoted to the Study of*

- the Nervous System and Behavior. 2005; 41(2):145–155. [http://doi.org/10.1016/S0010-9452\(08\)70889-X](http://doi.org/10.1016/S0010-9452(08)70889-X). [PubMed: 15714897]
- Strømme P, Bjørnstad PG, Ramstad K. Prevalence Estimation of Williams Syndrome. *Journal of Child Neurology*. 2002; 17(4):269–271. <http://doi.org/10.1177/088307380201700406>. [PubMed: 12088082]
- Swillen A, McDonald-McGinn D. Developmental trajectories in 22q11.2 deletion syndrome. *American Journal of Medical Genetics, Part C: Seminars in Medical Genetics*. 2015; 169(2):172–181. <http://doi.org/10.1002/ajmg.c.31435>.
- Swillen A, Vandeputte L, Cracco J, Maes B, Ghesquière P, Devriendt K, Fryns JP. Neuropsychological, learning and psychosocial profile of primary school aged children with the velo-cardio-facial syndrome (22q11 deletion): evidence for a nonverbal learning disability? *Child Neuropsychology*. 1999; 5(4):230–241. [http://doi.org/10.1076/0929-7049\(199912\)05:04;1-R;FT230](http://doi.org/10.1076/0929-7049(199912)05:04;1-R;FT230). [PubMed: 10925707]
- Tager-Flusberg H, Plesa-Skwerer D, Faja S, Joseph RM. People with Williams syndrome process faces holistically. *Cognition*. 2003; 89(1):11–24. [http://doi.org/10.1016/S0010-0277\(03\)00049-0](http://doi.org/10.1016/S0010-0277(03)00049-0). [PubMed: 12893122]
- Vivanti, G., Hocking, DR., Fanning, PAJ., Uljarevic, M., Postorino, V., Mazzone, L., Dissanayake, C. Attention to novelty versus repetition: Contrasting habituation profiles in Autism and Williams syndrome. *Developmental Cognitive Neuroscience*. 2017. <http://doi.org/10.1016/j.dcn.2017.01.006>
- Vivanti G, Hocking DR, Fanning P, Dissanayake C. Verbal labels increase the salience of novel objects for preschoolers with typical development and Williams syndrome, but not in autism. *Journal of Neurodevelopmental Disorders*. 2016; 8(1):46. <http://doi.org/10.1186/s11689-016-9180-7>. [PubMed: 28050217]
- Vivanti G, Hocking DR, Fanning P, Dissanayake C. The social nature of overimitation: Insights from Autism and Williams syndrome. *Cognition*. 2017a; 161:10–18. <http://doi.org/10.1016/j.cognition.2017.01.008>. [PubMed: 28088702]
- Vivanti G, Hocking DR, Fanning P, Dissanayake C. The social nature of overimitation: Insights from Autism and Williams syndrome. *Cognition*. 2017b; 161:10–18. <http://doi.org/10.1016/j.cognition.2017.01.008>. [PubMed: 28088702]
- Wechsler, D. Wechsler Intelligence Scale for Children-Revised (WISC-R). TEA Ediciones. 1974.
- Weinberger R, Yi J, Calkins M, Guri Y, McDonald-McGinn DM, Emanuel BS, Gothelf D. Neurocognitive profile in psychotic versus nonpsychotic individuals with 22q11.2 deletion syndrome. *European Neuropsychopharmacology*. 2016; 26(10):1610–1618. <http://doi.org/10.1016/j.euroneuro.2016.08.003>. [PubMed: 27524298]
- Wimmer H, Perner J. Beliefs about beliefs: Representation and constraining function of wrong beliefs in young children's understanding of deception. *Cognition*. 1983; 13:103–128. [http://doi.org/10.1016/0010-0277\(83\)90004-5](http://doi.org/10.1016/0010-0277(83)90004-5). [PubMed: 6681741]
- Zarchi O, Avni C, Attias J, Frisch A, Carmel M, Michaelovsky E, Gothelf D. Hyperactive auditory processing in Williams syndrome: Evidence from auditory evoked potentials. *Psychophysiology*. 2015; 52(6):782–9. <http://doi.org/10.1111/psyp.12407>. [PubMed: 25603839]
- Zarchi O, Diamond A, Weinberger R, Abbott D, Carmel M, Frisch A, Gothelf D. A comparative study of the neuropsychiatric and neurocognitive phenotype in two microdeletion syndromes: Velocardiofacial (22q11.2 deletion) and Williams (7q11.23 deletion) syndromes. *European Psychiatry*. 2014; 29(4):203–210. <http://doi.org/10.1016/j.eurpsy.2013.07.001>. [PubMed: 24054518]

Figure 1. Performance in a first-order false belief task

Legend: The 22q11.2DS group showed higher performance compared to the WS and IDD groups; Typically-developing (TD) participants demonstrated superior ability compared to all three clinical groups; Error bars depict standard deviations (SD); Mean $\pm$ SD: TD: 13.15 $\pm$ 1.72 > 22q11.2DS: 10.95 $\pm$ 3.03 > IDD: 8.52 $\pm$ 4.0, WS: 6.81 $\pm$ 4.15; * p <.05, *** p <.001.

Figure 2. Distinguishing real from apparent emotions

Legend: The WS and IDD groups performed poorly compared to TD controls; The 22q11.2DS group performed similarly to TD controls; Error bars depict standard deviations; Mean $\pm$ SD: TD: 7.25 $\pm$ 1.3 > WS: 5.56 $\pm$ 1.5, IDD: 5.32 $\pm$ 1.3; 22q11.2DS: 6.35 $\pm$ 1.3; *** p <.001.

Figure 3. Becoming friends with a story-character

Legend: Children with WS were more willing to become friends with a story-character even when the character's motives were negative (i.e., they scored poorly in this question compared to the rest of the groups); Error bars depict standard deviation; Mean $\pm$ SD: WS: 0.90 ± 0.73 < TD: 1.82 ± 0.37 , 22q11.2DS: 1.58 ± 0.57 , IDD: 1.52 ± 0.69 ; *** $p<.001$.

Table 1

Demographic characteristics of participants in the four study groups (N=82)

	WS	22q11.2DS	IDD	TD	ANOVA <i>F</i>	Comparisons
N	18	24	20	20		
Child's age, years	11.96 (2.33)	11.41 (1.96)	10.86 (2.72)	7.82 (2.38)	$F_{(3,78)} = 12.33^{***}$	TD < WS, 22q11.2, IDD
Mental age	7.82 (1.36)	9.5 (2.01)	7.83 (2.00)	7.82 (2.38)	$F_{(3,74)} = 3.75^*$	<i>n.s.</i>
IQ	67.63 (8.78)	83.57 (11.95)	73.84 (9.52)		$F_{(2,75)} = 11.70^{***}$	22q11.2DS > WS, IDD
PIQ	68.19 (9.95)	85.64 (11.42)	75.43 (14.17)		$F_{(2,48)} = 10.52^{***}$	22q11.2DS > WS, IDD
VIQ	67.94 (11.83)	83.27 (12.21)	77.77 (12.54)		$F_{(2,48)} = 7.36^{**}$	22q11.2DS > WS
Gender (M/F)	9/9	17/7 [†]	14/6 [†]	11/9		

Mean (SD); WS – Williams syndrome; 22q11.2DS – 22q11.2 deletion syndrome; IDD – idiopathic developmental disability; TD – typically developing (control group); ANOVA – analysis of variance; Mental age (MA) was calculated as: MA = [IQ * CA]/100.

* $p < .05$,

** $p < .01$,

*** $p < .001$;

[†] Chi-square analysis (X^2), $p < .05$.

Table 2

Group comparisons of performance in the facial emotion recognition paradigm

Labeling	Mean ($\pm$ SD)					ANOVA F	Comparisons
	WS	22q11.2DS	IDD	TD			
Happy	3.62 ($\pm$.61)	2.78 ($\pm$ 1.04)	3.15 ($\pm$.83)	3.10 ($\pm$.96)		$F_{(3,73)} = 2.54^*$	WS > 22q11.2DS
Sad	2.87 ($\pm$ 1.02)	2.78 ($\pm$.99)	3.26 ($\pm$.93)	3.75 ($\pm$.44)		$F_{(3,73)} = 5.38^{***}$	TD > WS, 22q11.2DS, IDD
Scared	1.43 ($\pm$.62)	1.21 ($\pm$.67)	1.10 ($\pm$.87)	1.45 ($\pm$.60)			
Angry	3.37 ($\pm$.61)	3.26 ($\pm$.75)	3.42 ($\pm$.69)	3.70 ($\pm$.65)			
Disgust	2.37 ($\pm$ 1.20)	3.00 ($\pm$ 1.08)	2.73 ($\pm$ 1.14)	3.50 ($\pm$.76)		$F_{(3,73)} = 4.72^{**}$	TD > WS, 22q11.2DS
Surprised	2.18 ($\pm$ 1.04)	2.47 ($\pm$ 1.12)	2.05 ($\pm$ 1.22)	2.70 ($\pm$ 1.03)			
Overall	17.33 ($\pm$.59)	15.87 ($\pm$.52)	17.19 ($\pm$.54)	19.35 ($\pm$.53)		$F_{(3,73)} = 7.13^{***}$	TD > 22q11.2DS, IDD
Matching							
	Mean ($\pm$ SD)						
Happy	2.68 ($\pm$.79)	3.13 ($\pm$.69)	2.73 ($\pm$.73)	3.65 ($\pm$.58)		$F_{(3,73)} = 7.62^{***}$	TD > WS, IDD
Sad	2.68 ($\pm$.60)	2.60 ($\pm$.83)	2.52 ($\pm$.84)	3.30 ($\pm$.65)		$F_{(3,73)} = 4.79^{**}$	TD > 22q11.2DS, IDD
Scared	2.93 ($\pm$.35)	3.34 ($\pm$.88)	3.00 ($\pm$.74)	3.60 ($\pm$.75)			
Angry	3.31 ($\pm$.70)	3.56 ($\pm$.58)	3.36 ($\pm$.76)	3.75 ($\pm$.44)			
Disgust	3.37 ($\pm$.71)	3.47 ($\pm$.66)	3.42 ($\pm$.90)	3.90 ($\pm$.76)		$F_{(3,73)} = 2.84^*$	TD > WS, 22q11.2DS, IDD
Surprised	2.31 ($\pm$.87)	2.56 ($\pm$ 1.23)	2.42 ($\pm$ 1.12)	2.85 ($\pm$.87)			
Overall	17.55 ($\pm$.55)	18.11 ($\pm$.48)	17.71 ($\pm$.51)	21.29 ($\pm$.49)		$F_{(3,73)} = 12.38^{***}$	TD > WS, 22q11.2DS, IDD

WS – Williams syndrome; 22q11.2DS – 22q11.2 deletion syndrome; IDD – idiopathic developmental disability; TD – typically developing (controls); ANOVA – analysis of variance;

* $p < .05$,** $p < .01$,*** $p < .001$