Published in final edited form as:

Mol Cell Neurosci. 2017 December ; 85: 183–189. doi:10.1016/j.mcn.2017.10.003.

Cocaine alters *Homer1* natural antisense transcript in the nucleus accumbens

Gregory C. Sartor^{*}, Samuel K. Powell^{*}, Dmitry Velmeshchev, David Y. Lin, Marco Magistri, Hannah J. Wiedner, Andrea M. Malvezzi, Nadja S. Andrade, Mohammad A. Faghihi, and Claes Wahlestedt

Center for Therapeutic Innovation, Department of Psychiatry and Behavioral Sciences, University of Miami Miller School of Medicine, Miami FL 33136

Abstract

Natural antisense transcripts (NATs) are an abundant class of long noncoding RNAs that have recently been shown to be key regulators of chromatin dynamics and gene expression in nervous system development and neurological disorders. However, it is currently unclear if NAT-based mechanisms also play a role in drug-induced neuroadaptations. Aberrant regulation of gene expression is one critical factor underlying the long-lasting behavioral abnormalities that characterize substance use disorder, and it is possible that some drug-induced transcriptional responses are mediated, in part, by perturbations in NAT activity. To test this hypothesis, we used an automated algorithm that mines the NCBI AceView transcriptomics database to identify NAT overlapping genes linked to addiction. We found that 22% of the genes examined contain NATs and that expression of *Homer1* natural antisense transcript (*Homer1-AS*) was altered in the nucleus accumbens (NAc) of mice 2 hours and 10 days following repeated cocaine administration. In *in vitro* studies, depletion of *Homer1-AS* led to an increase in the corresponding sense gene expression, indicating a potential regulatory mechanism of *Homer1* expression by its corresponding antisense transcript. However, future *in vivo* studies are needed to definitely determine a role for *Homer1-AS* in cocaine-induced behavioral and molecular adaptations.

Keywords

long noncoding RNA; natural antisense transcripts; cocaine; nucleus accumbens; epigenetics

Introduction

Recent high-throughput sequencing studies have uncovered tens of thousands of RNA transcripts with little or no protein-coding potential (termed noncoding RNAs) (Carninci *et*

Correspondence to: Claes Wahlestedt, M.D., Ph.D., Department of Psychiatry and Behavioral Sciences, University of Miami Miller School of Medicine, 1501 NW 10th Avenue, BRB-407 (M-860), Miami, FL 33136, CWahlestedt@med.miami.edu, Telephone: 305-243-1367, Fax: 305-243-2523.

^{*}Contributed equally to this work.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

et al., 2005; Katayama *et al.*, 2005a; Birney *et al.*, 2007). Initially thought of as transcriptional noise, research has now shown that noncoding RNAs have fundamental roles in multiple aspects of cellular physiology (Magistri *et al.*, 2012; Qureshi & Mehler, 2012). Long noncoding RNAs (lncRNAs), defined as noncoding transcripts greater than 200 nucleotide bases in length, are pervasively expressed across mammalian genomes and have been implicated in several aspects of transcriptional and posttranscriptional regulation (Feng *et al.*, 2006; Rinn *et al.*, 2007; Nowacki *et al.*, 2008; Huarte *et al.*, 2010). lncRNAs are generally classified based on their anatomical properties and their gene loci. For example, natural antisense transcripts (NATs) overlap other genes on the opposite strand of the DNA while intergenic lncRNAs (lincRNAs) are located between genes. Interestingly, many of these lncRNAs show specific spatial and temporal expression patterns within tissues and cells (Clark *et al.*, 2011). In the brain, for example, numerous lncRNAs are selectively expressed in neurons (Mercer *et al.*, 2008; Belgard *et al.*, 2011), indicating that some lncRNAs may be specifically involved in neuronal function (Qureshi & Mehler, 2012).

Natural Antisense Transcripts (NATs) are transcribed from the opposite DNA strand of other RNA transcripts with which they share sequence complementarities (Magistri *et al.*, 2012). Transcriptome-wide analyses by the FANTOM and ENCODE consortia have revealed that up to 66% of protein-coding genes contain NATs (Katayama *et al.*, 2005b; Djebali *et al.*, 2012), indicating their vast prevalence and potential importance in biological mechanisms. Of the proposed functional mechanisms of NATs, regulation of chromatin architecture and epigenetic memory have received much attention, as some antisense transcripts may provide a scaffold by which proteins can interact with DNA and chromatin in a locus-specific manner (Saxena & Carninci, 2011; Magistri *et al.*, 2012). Moreover, many NATs are expressed in the CNS where they have been shown to play fundamental roles in a variety of biological processes. For example, our previous work demonstrates that NAT dysregulation is associated with Alzheimer's disease pathophysiology (Modarresi *et al.*, 2011; Magistri *et al.*, 2016), and in separate reports we have also revealed that NATs contribute to Parkinson's disease, Autism, Fragile X mental retardation, seizures and metabolism (Scheele *et al.*, 2007; Faghihi *et al.*, 2010a; Pastori *et al.*, 2014; Hsiao *et al.*, 2016; Silva *et al.*, 2016). Mechanistically, we and others have shown that NATs can mediate epigenetic modification of genes in *cis*, making them unexplored targets to achieve specific up-regulation of gene expression (Modarresi *et al.*, 2012). Additionally, NATs have been found to affect disease-related gene expression by modifying mRNA stability and microRNA function (Faghihi *et al.*, 2010b; Modarresi *et al.*, 2011). Thus, these findings suggest that NATs are dysregulated in multiple neuropathologies and that endogenous gene expression can be specifically regulated by targeting NATs.

With such integral roles in transcriptional and posttranscriptional regulation, it is not surprising that dysregulation of NATs have been linked to multiple diseases (Scheele *et al.*, 2007; Faghihi *et al.*, 2008; Han *et al.*, 2012; Yang *et al.*, 2012), but little is known about their involvement in substance use disorder. Two recent reports, however, indicate that lncRNAs, in general, may play a role in cocaine-induced adaptations, as expression of several lncRNAs was altered in the nucleus accumbens (NAc) and midbrain of post-mortem human cocaine abusers (Michelhaugh *et al.*, 2011; Bannon *et al.*, 2015) and in mice following cocaine conditioned place preference (Bu *et al.*, 2012). To better understand a role for NATs

in response to repeated cocaine use, we took advantage of a recently published automated algorithm that mines the AceView transcriptomic database to identify biologically relevant sense-antisense pairs (Velmeshev *et al.*, 2013) and revealed that numerous genes that have been previously associated with cocaine-induced neuroadaptations contain NATs. In *in vivo* studies, we found that expression of *Homer1-AS* was significantly altered in the NAc of mice following repeated cocaine administration, and that siRNA-mediated depletion of *Homer1-AS in vitro* altered corresponding sense gene expression. Together, these data indicate NAT-based mechanisms may contribute to cocaine-mediated molecular adaptations.

Methods and materials

Animals

Male C57BL/6 mice (8–10 weeks old, Charles River Laboratories) were housed 4 animals per cage, under a regular 12h/12h light/dark cycle with *ad libitum* access to food and water. Mice were housed in a humidity and temperature-controlled, AAALAC-accredited, animal facility at the University of Miami Miller School of Medicine. All experiments were approved by the Institutional Animal Care and Use Committee (IACUC) and conducted according to specifications of the NIH as outlined in the Guide for the Care and Use of Laboratory Animals.

Drug Treatments

Cocaine HCl (NIDA, Research Triangle Park, NC) was dissolved in 0.9% sterile saline. Animals received one intraperitoneal (i.p.) injection of saline or cocaine (20 mg/kg) once a day for 10 consecutive days for repeated exposure experiments and a single injection of saline or cocaine (20 mg/kg) for acute studies. Two hours or 10 days after the last injection, bilateral tissue punches of the nucleus accumbens (NAc) were collected on ice and flash frozen in liquid nitrogen and then processed for RNA analysis as described below. Doses for all drugs were based on their salt form.

Bioinformatics analysis

We utilized a previously published bioinformatics algorithm to identify Natural Antisense Transcripts (NATs) from AceView transcriptome database (Velmeshev *et al.*, 2013). The open-access algorithm and the manual are available at <https://github.com/DmitryVel/NATs>. Briefly, the algorithm takes an input list of genes and first identifies all transcripts expressed from the antisense strand of the target locus or the gene promoter (1kb downstream from the transcription start site). Next, transcripts containing a significant open reading frame according to the annotation on AceView are filtered out to retain noncoding NATs. We utilized this pipeline to investigate the expression of antisense transcripts overlapping with 545 genes. This list was based on previously published genes that have been implicated in cocaine-induced neuroadaptations (Renthall *et al.*, 2009; Maze *et al.*, 2010; Ferguson *et al.*, 2013; Feng *et al.*, 2014) and from the top gene searches on Allan Brain Atlas related to neuroplasticity. The pipeline output yielded the existence of an antisense overlap, the position of the overlap, the NAT Aceview name, the NAT length and the overlap start and stop sites (Table S1). PANTHER (<http://www.pantherdb.org/>) was used to categorize NAT-containing genes based on protein class.

Cell culture

Neuro-2a (N2a) cell lines (ATCC) were maintained in Advanced DMEM supplemented with 10% FBS and 1% primocin. In a 6 well plate, ~400,000 cells were plated and transfected 24 h later with scrambled control (Qiagen SI03650318) or *Homer1-AS* siRNA (Forward: 5' GUACAAAUGCAAUGUGUUAUU 3', Reverse: 5' UAACACAUUGCAUUUGUACUU 3') in lipofectamine RNAiMax (Invitrogen) according to the manufacturer's instructions (20 or 40 nM for 48 h). A blank group treated with transfection reagent only was also included as an additional control. After treatment, N2a cells (6 biological replicates with the same passage number and treatment time) were harvested for RNA analysis as described below.

Quantitative Real-Time PCR

Following treatment, RNA was isolated using a Trizol and column RNA extraction kit as described by the manufacturer (RNeasy kit, Qiagen). Samples were run in triplicate replicates, normalized to beta-actin and analyzed using the 2^{-CT} method. NAT primers (Table S1) were preferentially designed to span a spliced junction of the antisense transcript that does not overlap any spliced junction of the sense gene to avoid sense transcript amplification. In cases where no junctional primers could be designed, we designed a primer pair targeting an exon of the antisense transcript and utilized a strand-specific qRT-PCR to amplify only the antisense transcript. To perform strand-specific measurement of antisense transcript expression, we designed primers for a region of antisense transcript that overlaps with an intron or the promoter of the sense gene. Using the one-step RNA-to-Ct SYBR Green Kit (Life Technologies, 4389986), the reverse transcription (RT) step was performed in a 384-well optical plate using reverse primers to specifically reverse-transcribe antisense RNA and to exclude the possibility of measuring the expression of the sense pre-mRNA. Samples were then incubated at 95°C for 5 minutes to inactivate the reverse transcriptase enzyme. Forward primers were then added to the reaction and quantitative PCR was performed on the same plate. We included no-RT control and no-template controls for each set of primers to control for non-specific binding and melting curves were analyzed to verify PCR specificity. *Homer1* and *beta-actin* expression were measured with Taqman assays as described by the manufacturer (ThermoFisher, Mm00516275_m1 and Mm02619580_g1). A map of the sense/antisense overlap of *Homer1/Homer1-AS* may be found on the Aceview mouse database (<https://www.ncbi.nlm.nih.gov/iebr/research/acembly/>) by searching for *mahemo*. Using SnapGene software, an additional map of *Homer1* sense and antisense is shown in Figure 4. In Figures 2, 3 and 5, the spliced form of *Homer1-AS* was measured.

Data Analysis

Graphpad Prism software was used for graph preparation and statistical analysis. Rq values were normalized to saline or vehicle-treated samples and compared between groups using one- or two-way ANOVA. Newman-Keuls, Bonferroni or Benjamini and Hochberg false discovery rate test was performed for *post-hoc* comparisons. Data are mean $\pm$ SEM, and the level of significance was set to 0.05.

Results

To explore a potential role for NATs in response to repeated cocaine exposure, we compiled a list of 545 genes that have been associated with neuroplasticity and/or addiction, and used our previously described algorithm to mine the AceView database to identify overlapping sense-antisense pairs (Velmeshov *et al.*, 2013) (Table S1). Using this pipeline (Figure 1A), we found that 22% (121/545) of the genes examined contained NATs, several of which (23/121) contained two or more NATs per gene (Figure 1B). Gene ontologies were diverse, with the largest proportion of the NAT-containing genes classified under receptor and signaling molecules (Figure 1C). The vast majority of these antisense RNAs were classified as intronic overlap with respect to the sense gene, though some were found to overlap with the promoter and exonic regions (Table S1).

To determine if the expression of these transcripts was altered in response to cocaine, we selected 25 antisense RNAs, many of which overlap with genes that have been implicated in cocaine-induced adaptations (Table 1), and quantified their expression in the nucleus accumbens (NAc) 2 hours following the last cocaine or saline injection. A two-way ANOVA showed significant interaction between antisense transcript and treatment ($F_{(24, 394)} = 3.0$, $P < 0.0001$), a main effect for antisense transcript ($F_{(24, 394)} = 3.0$, $P < 0.0001$, $n = 7-9$) but no main effect for treatment ($F_{(1, 394)} = 0.6$, $P > 0.05$) (Figure 2). Benjamini and Hochberg *post hoc* test revealed that *Homer1-AS* was significantly altered following repeated cocaine treatment ($P < 0.0005$, Figure 2, Table S2). Although *Homer1-AS* was increased in the NAc following repeated cocaine administration, no change in *Homer1* expression was observed at the same time point following repeated cocaine injections (Figure 3A). *Homer1-AS* remained significantly elevated 10 days following repeated cocaine injections ($P < 0.005$, Figure 3B). In additional experiments, an acute injection of cocaine increased *Homer1*, but not *Homer1-AS*, expression in the NAc ($P < 0.0005$, Figure 3C). A schematic of the sense-antisense overlap between *Homer1/Homer1-AS* is shown in Figure 4.

NATs typically employ *cis*-regulatory mechanisms to positively (concordant) or negatively (discordant) regulate the expression of their overlapping sense mRNA (Magistri *et al.*, 2012). Though *Homer1* is known to be altered in the NAc following cocaine administration (Szumlinski *et al.*, 2008), it is unclear if *Homer1-AS* modulates its expression. To identify the relationship between *Homer1/Homer1-AS* expression, we utilized siRNA-mediated knockdown to significantly deplete *Homer1-AS* in N2a cells ($F_{(3, 20)} = 8.1$, $P < 0.005$, $n = 6$) (Figure 5A). *Homer1-AS* knockdown resulted in a discordant regulation of *Homer1* ($F_{(3, 20)} = 60.7$, $P < 0.0001$, $n = 6$) (Figure 5B), indicating a potential regulatory mechanisms between the protein-coding gene by its corresponding antisense transcript.

Discussion

Recent transcriptomic studies by FANTOM, GENCODE and ENCODE consortia have revealed that RNA plays a variety of roles in nuclear architecture, X chromosome inactivation, imprinting, DNA looping, transcription, translation and epigenetic processes (Magistri *et al.*, 2012). While small noncoding RNAs such as microRNAs have been shown to be important in regulating addiction-relevant neuroplasticity in the brain and cocaine-

seeking behaviors (Bali & Kenny, 2013), little is known about the involvement of long noncoding RNAs, particularly natural antisense transcripts (NATs), in cocaine-induced neuroadaptations and behavior. Our results show that numerous genes contain NATs and reveal that *Homer1-AS* was altered in the nucleus accumbens following repeated cocaine administration. Thus, these data, along with previous reports, indicate that NAT-based mechanisms may regulate transcriptional and/or post-transcriptional responses to cocaine and contribute to cocaine-induced neuroadaptations.

Although cocaine-mediated regulation of *Homer1* is an important adaptation underlying drug-seeking behaviors, the effects of cocaine on *Homer1* expression are quite complex and dependent upon the duration of cocaine administration, length of withdrawal, and the brain region/isoform examined (Szumlinski *et al.*, 2008). For example, acute cocaine administration transiently increases mRNA and protein of the immediate early gene, *Homer1a*, and the constitutively expressed isoform, *Homer1b/c*, in the NAc (Brakeman *et al.*, 1997; Fourgeaud *et al.*, 2004; Zhang *et al.*, 2007). Following 3 weeks of withdrawal from repeated cocaine injections, *Homer1b/c* protein is decreased in the NAc, which also coincides with reductions in mGluR5 protein and manifestation of cocaine-induced behavioral sensitization (Swanson *et al.*, 2001). In functional studies, knockout or oligo-induced depletion of *Homer1* in the NAc enhances locomotor responses to cocaine, whereas over-expression of the constitutive isoform prevents cocaine sensitization (Ghasemzadeh *et al.*, 2003; Szumlinski *et al.*, 2004). The data presented here add another layer of complexity to the regulation of *Homer1* in response to cocaine. We identified a noncoding antisense RNA that overlaps with *Homer1* and whose expression is increased in the NAc in response to repeated, but not acute, cocaine injections. Although we revealed that acute cocaine transiently increases *Homer1* in the NAc, we did not observe a change in *Homer1* mRNA expression 2 hours after repeated cocaine administration. These data are consistent with previous studies that have shown a temporary increase in *Homer1* mRNA and protein following an acute cocaine injection (Brakeman *et al.*, 1997; Fourgeaud *et al.*, 2004; Zhang *et al.*, 2007) and a delayed decrease in *Homer1* protein following repeated cocaine injections (Swanson *et al.*, 2001). Thus, based on the differences in temporal regulation of *Homer1* and *Homer1-AS*, it is conceivable that *Homer1-AS* affects *Homer1* protein expression in the NAc via post-transcriptional processes. In *in vitro* studies, we revealed that siRNA-mediated knockdown of *Homer1-AS* increased *Homer1* mRNA expression, indicating a discordant regulation mechanism between *Homer1* sense and antisense expression. Although we have previously shown that NAT knockdown produced a similar upregulation of sense gene expression in both *in vivo* and *in vitro* experiments (Modarresi *et al.*, 2012), it is possible that *Homer1/Homer-AS* interact differently in specific cells types and under *in vitro* vs. *in vivo* conditions. Thus, future *in vivo* studies are needed to determine a mechanistic role for *Homer1-AS*. Nonetheless, given that cocaine increases *Homer1-AS* and that changes in *Homer1* contribute to the long-lasting neurochemical and behavioral abnormalities that characterize addiction (Szumlinski *et al.*, 2008), antisense transcripts that overlap with *Homer1* may play an important role in cocaine-induced neuroadaptations.

Our earlier findings indicate that the expression of *Bdnf*, a gene importantly involved in the development of cocaine-induced neuroplasticity and behaviors (Li & Wolf, 2014), is regulated by a *Bdnf* antisense transcript (Modarresi *et al.*, 2012). Interestingly, in our current

studies, *Bdnf-AS* was reduced 38% following repeated cocaine exposure, though this reduction was not statistically significant when adjusted for multiple comparisons. Due to the stringency of the Benjamini and Hochberg false discovery rate test, Type II errors (false negatives) are possible and future studies are needed to address the functional role of *Bdnf-AS* in response to cocaine. However, given that *Bdnf* is known to increase in the NAc following cocaine exposure and self-administration (Graham *et al.*, 2007), it is possible that *Bdnf* and *Bdnf-AS* also interact in a discordant manner to regulate behavioral and molecular responses to drugs of abuse. Indeed, a recent study revealed a correlation between polymorphisms in *BDNF-AS* and heroin abuse in the Han Chinese population (Jin *et al.*, 2016), indicating the potential clinical importance of *BDNF-AS* in substance use disorders. Likewise, the antisense transcript of *Kcnq1* (referred to as *Kcnq1ot1*), which was reduced by 35% in the NAc by cocaine, has been studied extensively and is known to silence multiple genes in the *Kcnq1* domain by establishing a repressive higher order chromatin structure (Mohammad *et al.*, 2009) and, similar to our results, is altered in the mouse NAc by methamphetamine (Zhu *et al.*, 2015). Thus, the epigenetic mechanisms by which *Kcnq1-AS* and other NATs regulate transcriptional and behavioral responses to psychostimulants merit further investigation.

Utilizing microarray analysis, previous studies have also identified several long noncoding RNAs that were altered in the NAc and midbrain dopamine neurons of human cocaine and heroin abusers (Michelhaugh *et al.*, 2011; Bannon *et al.*, 2015) and in the NAc of mice following cocaine conditioned place preference (Bu *et al.*, 2012). Although NATs were not the primary focus of these previous studies, several of the long noncoding RNAs that were altered by cocaine were antisense transcripts (52/603 in the mouse NAc and 16/32 in the human midbrain) (Bu *et al.*, 2012; Bannon *et al.*, 2015). To our knowledge, *Homer1-AS* and other NATs examined in the current study were not investigated in previous cocaine studies. However, our current results and previous studies are far from exhaustive. Subsequent RNA-seq studies utilizing the latest lncRNA annotations are needed in order to have comprehensive understanding of the NATs involved in substance use disorder. Another interesting observation in the current study is that few NATs were significantly altered in the NAc by repeated cocaine injections. The stringency of the Benjamini and Hochberg false discovery rate test likely accounts for why only one of the NATs examined met statistical significance. Although only a portion of lncRNAs that were significantly altered by drugs of abuse were NATs in the above studies, it is important to note that many NATs are expressed at low levels and in a cell type-specific manner (Cabili *et al.*, 2011; Derrien *et al.*, 2012; Li *et al.*, 2013; Dong *et al.*, 2015). Thus, small changes in expression that are observed in tissue homogenate may have functional importance in specific cell types. In fact, recent FANTOM5 analyses indicate the potential functionality of 69% of lncRNAs (Hon *et al.*, 2017). Future studies examining brain region-, cell type- and temporal-specific expression profiles of noncoding antisense RNAs following cocaine use, along with functional experiments examining the behavioral consequences of NAT manipulation on drug-seeking behaviors, will lead to a better understanding of NATs in drug-induced neurobehavioral adaptations.

In summary, imbalanced expression of NATs may result in the silencing or activation of partner protein-coding genes, thus providing a potentially interesting and novel mechanism

to explain the aberrant gene regulation following repeated drug use. Based on these initial findings and previously published data, we hypothesize that repeated cocaine administration leads to a dysregulation of natural antisense transcripts that may act in concert with transcriptional and post-transcriptional machinery to pathologically alter gene expression and behavior. However, additional studies are needed to better understand the anatomical and temporal dynamics of NAT expression following acute and repeated cocaine exposure and to reveal the functional importance of specific NATs during drug-seeking behaviors. Such experiments may lead to the incorporation of NATs into epigenetic models of drug-induced plasticity and open the door for critically-needed new therapeutic avenues for substance use disorder.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

In remembrance of David Lin, for his skill and dedication were essential to this work. This research was supported by National Institute on Drug Abuse Grant: R21DA035592. S.K.P. was supported by Howard Hughes Medical Institute (HHMI) Undergraduate Research Program. The cocaine used in these studies was obtained from the NIDA Drug Supply Program. G.C.S., S.K.P., D.V., M.M. and C.W. designed the experiments. G.C.S., S.K.P., D.Y.L., H.J.W., A.M. and D.V. conducted the experiments. G.C.S. and D.V. wrote the paper with input from S.K.P., M.M., H.J.W., A.M., M.A.F. and C.W.

References

- Bali P, Kenny PJ. MicroRNAs and Drug Addiction. *Front Genet.* 2013; 4:43. [PubMed: 23717324]
- Bannon MJ, Savonen CL, Jia H, Dacht F, Halter SD, Schmidt CJ, Lipovich L, Kapatos G. Identification of long noncoding RNAs dysregulated in the midbrain of human cocaine abusers. *J Neurochem.* 2015; 135:50–59. [PubMed: 26222413]
- Belgard TG, Marques AC, Oliver PL, Abaan HO, Sirey TM, Hoerder-Suabedissen A, Garcia-Moreno F, Molnar Z, Margulies EH, Ponting CP. A transcriptomic atlas of mouse neocortical layers. *Neuron.* 2011; 71:605–616. [PubMed: 21867878]
- Birney E, Stamatoiyannopoulos JA, Dutta A, Guigo R, Gingeras TR, Margulies EH, Weng Z, Snyder M, Dermitzakis ET, Thurman RE, Kuehn MS, Taylor CM, Neph S, Koch CM, Asthana S, Malhotra A, Adzhubei I, Greenbaum JA, Andrews RM, Flicek P, Boyle PJ, Cao H, Carter NP, Clelland GK, Davis S, Day N, Dhami P, Dillon SC, Dorschner MO, Fiegler H, Giresi PG, Goldy J, Hawrylycz M, Haydock A, Humbert R, James KD, Johnson BE, Johnson EM, Frum TT, Rosenzweig ER, Karnani N, Lee K, Lefebvre GC, Navas PA, Neri F, Parker SC, Sabo PJ, Sandstrom R, Shafer A, Vetrie D, Weaver M, Wilcox S, Yu M, Collins FS, Dekker J, Lieb JD, Tullius TD, Crawford GE, Sunyaev S, Noble WS, Dunham I, Denoeud F, Reymond A, Kapranov P, Rozowsky J, Zheng D, Castelo R, Frankish A, Harrow J, Ghosh S, Sandelin A, Hofacker IL, Baertsch R, Keefe D, Dike S, Cheng J, Hirsch HA, Sekinger EA, Lagarde J, Abril JF, Shahab A, Flamm C, Fried C, Hackermuller J, Hertel J, Lindemeyer M, Missal K, Tanzer A, Washietl S, Korb J, Emanuelsson O, Pedersen JS, Holroyd N, Taylor R, Swarbreck D, Matthews N, Dickson MC, Thomas DJ, Weirauch MT, Gilbert J, Drenkow J, Bell I, Zhao X, Srinivasan KG, Sung WK, Ooi HS, Chiu KP, Foissac S, Alioto T, Brent M, Pachter L, Tress ML, Valencia A, Choo SW, Choo CY, Ucla C, Manzano C, Wyss C, Cheung E, Clark TG, Brown JB, Ganesh M, Patel S, Tammanna H, Chrast J, Henrichsen CN, Kai C, Kawai J, Nagalakshmi U, Wu J, Lian Z, Lian J, Newburger P, Zhang X, Bickel P, Mattick JS, Carninci P, Hayashizaki Y, Weissman S, Hubbard T, Myers RM, Rogers J, Stadler PF, Lowe TM, Wei CL, Ruan Y, Struhl K, Gerstein M, Antonarakis SE, Fu Y, Green ED, Karaoz U, Siepel A, Taylor J, Liefer LA, Wetterstrand KA, Good PJ, Feingold EA, Guyer MS, Cooper GM, Asimenos G, Dewey CN, Hou M, Nikolaev S, Montoya-Burgos JI, Loytynoja A, Whelan S, Pardi F, Massingham T, Huang H, Zhang NR, Holmes I, Mullikin JC, Ureta-Vidal A, Paten B, Serinhaus M, Church D, Rosenbloom

K, Kent WJ, Stone EA, Batzoglou S, Goldman N, Hardison RC, Haussler D, Miller W, Sidow A, Trinklein ND, Zhang ZD, Barrera L, Stuart R, King DC, Ameur A, Enroth S, Bieda MC, Kim J, Bhinge AA, Jiang N, Liu J, Yao F, Vega VB, Lee CW, Ng P, Yang A, Moqtaderi Z, Zhu Z, Xu X, Squazzo S, Oberley MJ, Inman D, Singer MA, Richmond TA, Munn KJ, Rada-Iglesias A, Wallerman O, Komorowski J, Fowler JC, Couttet P, Bruce AW, Dovey OM, Ellis PD, Langford CF, Nix DA, Euskirchen G, Hartman S, Urban AE, Kraus P, Van Calcar S, Heintzman N, Kim TH, Wang K, Qu C, Hon G, Luna R, Glass CK, Rosenfeld MG, Aldred SF, Cooper SJ, Halees A, Lin JM, Shulha HP, Xu M, Haidar JN, Yu Y, Iyer VR, Green RD, Wadelius C, Farnham PJ, Ren B, Harte RA, Hinrichs AS, Trumbower H, Clawson H, Hillman-Jackson J, Zweig AS, Smith K, Thakapallayil A, Barber G, Kuhn RM, Karolchik D, Armengol L, Bird CP, de Bakker PI, Kern AD, Lopez-Bigas N, Martin JD, Stranger BE, Woodroffe A, Davydov E, Dimas A, Eyraes E, Hallgrimsdottir IB, Huppert J, Zody MC, Abecasis GR, Estivill X, Bouffard GG, Guan X, Hansen NF, Idol JR, Maduro VV, Maskeri B, McDowell JC, Park M, Thomas PJ, Young AC, Blakesley RW, Muzny DM, Sodergren E, Wheeler DA, Worley KC, Jiang H, Weinstock GM, Gibbs RA, Graves T, Fulton R, Mardis ER, Wilson RK, Clamp M, Cuff J, Gnerre S, Jaffe DB, Chang JL, Lindblad-Toh K, Lander ES, Koriabine M, Nefedov M, Osoegawa K, Yoshinaga Y, Zhu B, de Jong PJ. Identification and analysis of functional elements in 1% of the human genome by the ENCODE pilot project. *Nature*. 2007; 447:799–816. [PubMed: 17571346]

Brakeman PR, Lanahan AA, O'Brien R, Roche K, Barnes CA, Haganir RL, Worley PF. Homer: a protein that selectively binds metabotropic glutamate receptors. *Nature*. 1997; 386:284–288. [PubMed: 9069287]

Bu Q, Hu Z, Chen F, Zhu R, Deng Y, Shao X, Li Y, Zhao J, Li H, Zhang B, Lv L, Yan G, Zhao Y, Cen X. Transcriptome analysis of long non-coding RNAs of the nucleus accumbens in cocaine-conditioned mice. *J Neurochem*. 2012; 123:790–799. [PubMed: 22957495]

Cabili MN, Trapnell C, Goff L, Koziol M, Tazon-Vega B, Regev A, Rinn JL. Integrative annotation of human large intergenic noncoding RNAs reveals global properties and specific subclasses. *Genes Dev*. 2011; 25:1915–1927. [PubMed: 21890647]

Carninci P, Kasukawa T, Katayama S, Gough J, Frith MC, Maeda N, Oyama R, Ravasi T, Lenhard B, Wells C, Kodzius R, Shimokawa K, Bajic VB, Brenner SE, Batalov S, Forrest AR, Zavolan M, Davis MJ, Wilming LG, Aidinis V, Allen JE, Ambesi-Impimbato A, Apweiler R, Aturaliya RN, Bailey TL, Bansal M, Baxter L, Beisel KW, Bersano T, Bono H, Chalk AM, Chiu KP, Choudhary V, Christoffels A, Clutterbuck DR, Crowe ML, Dalla E, Dalrymple BP, de Bono B, Della Gatta G, di Bernardo D, Down T, Engstrom P, Fagiolini M, Faulkner G, Fletcher CF, Fukushima T, Furuno M, Futaki S, Gariboldi M, Georgii-Hemming P, Gingeras TR, Gojobori T, Green RE, Gustincich S, Harbers M, Hayashi Y, Hensch TK, Hirokawa N, Hill D, Huminieccki L, Iacono M, Ikeo K, Iwama A, Ishikawa T, Jakt M, Kanapin A, Katoh M, Kawasawa Y, Kelso J, Kitamura H, Kitano H, Kollias G, Krishnan SP, Kruger A, Kummerfeld SK, Kurochkin IV, Lareau LF, Lazarevic D, Lipovich L, Liu J, Liuni S, McWilliam S, Madan Babu M, Madera M, Marchionni L, Matsuda H, Matsuzawa S, Miki H, Mignone F, Miyake S, Morris K, Mottagui-Tabar S, Mulder N, Nakano N, Nakauchi H, Ng P, Nilsson R, Nishiguchi S, Nishikawa S, Nori F, Ohara O, Okazaki Y, Orlando V, Pang KC, Pavan WJ, Pavesi G, Pesole G, Petrowsky N, Piazza S, Reed J, Reid JF, Ring BZ, Ringwald M, Rost B, Ruan Y, Salzberg SL, Sandelin A, Schneider C, Schonbach C, Sekiguchi K, Semple CA, Seno S, Sessa L, Sheng Y, Shibata Y, Shimada H, Shimada K, Silva D, Sinclair B, Sperling S, Stupka E, Sugiura K, Sultana R, Takenaka Y, Taki K, Tammoja K, Tan SL, Tang S, Taylor MS, Tegner J, Teichmann SA, Ueda HR, van Nimwegen E, Verardo R, Wei CL, Yagi K, Yamanishi H, Zabarovsky E, Zhu S, Zimmer A, Hide W, Bult C, Grimmond SM, Teasdale RD, Liu ET, Brusic V, Quackenbush J, Wahlestedt C, Mattick JS, Hume DA, Kai C, Sasaki D, Tomaru Y, Fukuda S, Kanamori-Katayama M, Suzuki M, Aoki J, Arakawa T, Iida J, Imamura K, Itoh M, Kato T, Kawaji H, Kawagashira N, Kawashima T, Kojima M, Kondo S, Konno H, Nakano K, Ninomiya N, Nishio T, Okada M, Plessy C, Shibata K, Shiraki T, Suzuki S, Tagami M, Waki K, Watahiki A, Okamura-Oho Y, Suzuki H, Kawai J, Hayashizaki Y. The transcriptional landscape of the mammalian genome. *Science*. 2005; 309:1559–1563. [PubMed: 16141072]

Clark MB, Amaral PP, Schlesinger FJ, Dinger ME, Taft RJ, Rinn JL, Ponting CP, Stadler PF, Morris KV, Morillon A, Rozowsky JS, Gerstein MB, Wahlestedt C, Hayashizaki Y, Carninci P, Gingeras TR, Mattick JS. The reality of pervasive transcription. *PLoS Biol*. 2011; 9:e1000625. discussion e1001102. [PubMed: 21765801]

- Derrien T, Johnson R, Bussotti G, Tanzer A, Djebali S, Tilgner H, Guernec G, Martin D, Merkel A, Knowles DG, Lagarde J, Veeravalli L, Ruan X, Ruan Y, Lassmann T, Carninci P, Brown JB, Lipovich L, Gonzalez JM, Thomas M, Davis CA, Shiekhata R, Gingeras TR, Hubbard TJ, Notredame C, Harrow J, Guigo R. The GENCODE v7 catalog of human long noncoding RNAs: analysis of their gene structure, evolution, and expression. *Genome Res.* 2012; 22:1775–1789. [PubMed: 22955988]
- Djebali S, Davis CA, Merkel A, Dobin A, Lassmann T, Mortazavi A, Tanzer A, Lagarde J, Lin W, Schlesinger F, Xue C, Marinov GK, Khatun J, Williams BA, Zaleski C, Rozowsky J, Roder M, Kokocinski F, Abdelhamid RF, Alioto T, Antoshechkin I, Baer MT, Bar NS, Batut P, Bell K, Bell I, Chakraborty S, Chen X, Chrast J, Curado J, Derrien T, Drenkow J, Dumais E, Dumais J, Dutttagupta R, Falconnet E, Fastuca M, Fejes-Toth K, Ferreira P, Foissac S, Fullwood MJ, Gao H, Gonzalez D, Gordon A, Gunawardena H, Howald C, Jha S, Johnson R, Kapranov P, King B, Kingswood C, Luo OJ, Park E, Persaud K, Preall JB, Ribeca P, Risk B, Robyr D, Sammeth M, Schaffer L, See LH, Shahab A, Skancke J, Suzuki AM, Takahashi H, Tilgner H, Trout D, Walters N, Wang H, Wrobel J, Yu Y, Ruan X, Hayashizaki Y, Harrow J, Gerstein M, Hubbard T, Reymond A, Antonarakis SE, Hannon G, Giddings MC, Ruan Y, Wold B, Carninci P, Guigo R, Gingeras TR. Landscape of transcription in human cells. *Nature.* 2012; 489:101–108. [PubMed: 22955620]
- Dong X, Chen K, Cuevas-Diaz Duran R, You Y, Sloan SA, Zhang Y, Zong S, Cao Q, Barres BA, Wu JQ. Comprehensive Identification of Long Non-coding RNAs in Purified Cell Types from the Brain Reveals Functional LncRNA in OPC Fate Determination. *PLoS Genet.* 2015; 11:e1005669. [PubMed: 26683846]
- Faghihi MA, Kocerha J, Modarresi F, Engstrom PG, Chalk AM, Brothers SP, Koesema E, St Laurent G, Wahlestedt C. RNAi screen indicates widespread biological function for human natural antisense transcripts. *PLoS One.* 2010a;5.
- Faghihi MA, Modarresi F, Khalil AM, Wood DE, Sahagan BG, Morgan TE, Finch CE, St Laurent G 3rd, Kenny PJ, Wahlestedt C. Expression of a noncoding RNA is elevated in Alzheimer's disease and drives rapid feed-forward regulation of beta-secretase. *Nat Med.* 2008; 14:723–730. [PubMed: 18587408]
- Faghihi MA, Zhang M, Huang J, Modarresi F, Van der Brug MP, Nalls MA, Cookson MR, St-Laurent G 3rd, Wahlestedt C. Evidence for natural antisense transcript-mediated inhibition of microRNA function. *Genome Biol.* 2010b; 11:R56. [PubMed: 20507594]
- Feng J, Bi C, Clark BS, Mady R, Shah P, Kohtz JD. The Evf-2 noncoding RNA is transcribed from the Dlx-5/6 ultraconserved region and functions as a Dlx-2 transcriptional coactivator. *Genes Dev.* 2006; 20:1470–1484. [PubMed: 16705037]
- Feng J, Wilkinson M, Liu X, Purushothaman I, Ferguson D, Vialou V, Maze I, Shao N, Kennedy P, Koo J, Dias C, Laitman B, Stockman V, LaPlant Q, Cahill ME, Nestler EJ, Shen L. Chronic cocaine-regulated epigenomic changes in mouse nucleus accumbens. *Genome Biol.* 2014; 15:R65. [PubMed: 24758366]
- Ferguson D, Koo JW, Feng J, Heller E, Rabkin J, Heshmati M, Renthall W, Neve R, Liu X, Shao N, Sartorelli V, Shen L, Nestler EJ. Essential role of SIRT1 signaling in the nucleus accumbens in cocaine and morphine action. *J Neurosci.* 2013; 33:16088–16098. [PubMed: 24107942]
- Fourgeaud L, Mato S, Bouchet D, Hemar A, Worley PF, Manzoni OJ. A single in vivo exposure to cocaine abolishes endocannabinoid-mediated long-term depression in the nucleus accumbens. *J Neurosci.* 2004; 24:6939–6945. [PubMed: 15295029]
- Ghasemzadeh MB, Permenter LK, Lake R, Worley PF, Kalivas PW. Homer1 proteins and AMPA receptors modulate cocaine-induced behavioural plasticity. *Eur J Neurosci.* 2003; 18:1645–1651. [PubMed: 14511343]
- Graham DL, Edwards S, Bachtell RK, DiLeone RJ, Rios M, Self DW. Dynamic BDNF activity in nucleus accumbens with cocaine use increases self-administration and relapse. *Nat Neurosci.* 2007; 10:1029–1037. [PubMed: 17618281]
- Han L, Zhang K, Shi Z, Zhang J, Zhu J, Zhu S, Zhang A, Jia Z, Wang G, Yu S, Pu P, Dong L, Kang C. LncRNA profile of glioblastoma reveals the potential role of lncRNAs in contributing to glioblastoma pathogenesis. *Int J Oncol.* 2012; 40:2004–2012. [PubMed: 22446686]
- Hon CC, Ramilowski JA, Harshbarger J, Bertin N, Rackham OJ, Gough J, Denisenko E, Schmeier S, Poulsen TM, Severin J, Lizio M, Kawaji H, Kasukawa T, Itoh M, Burroughs AM, Noma S, Djebali

- S, Alam T, Medvedeva YA, Testa AC, Lipovich L, Yip CW, Abugessaisa I, Mendez M, Hasegawa A, Tang D, Lassmann T, Heutink P, Babina M, Wells CA, Kojima S, Nakamura Y, Suzuki H, Daub CO, de Hoon MJ, Arner E, Hayashizaki Y, Carninci P, Forrest AR. An atlas of human long non-coding RNAs with accurate 5' ends. *Nature*. 2017
- Hsiao J, Yuan TY, Tsai MS, Lu CY, Lin YC, Lee ML, Lin SW, Chang FC, Liu Pimentel H, Olive C, Coito C, Shen G, Young M, Thorne T, Lawrence M, Magistri M, Faghihi MA, Khorkova O, Wahlestedt C. Upregulation of Haploinsufficient Gene Expression in the Brain by Targeting a Long Non-coding RNA Improves Seizure Phenotype in a Model of Dravet Syndrome. *EBioMedicine*. 2016
- Huarte M, Guttman M, Feldser D, Garber M, Koziol MJ, Kenzelmann-Broz D, Khalil AM, Zuk O, Amit I, Rabani M, Attardi LD, Regev A, Lander ES, Jacks T, Rinn JL. A large intergenic noncoding RNA induced by p53 mediates global gene repression in the p53 response. *Cell*. 2010; 142:409–419. [PubMed: 20673990]
- Jin T, Zhang H, Yang Q, Li L, Ouyang Y, Yang M, Wang F, Wang Z, Zhang J, Yuan D. The relationship between polymorphisms of BDNFOS and BDNF genes and Heroin addiction in the Han Chinese population. *J Gene Med*. 2016
- Katayama S, Tomaru Y, Kasukawa T, Waki K, Nakanishi M, Nakamura M, Nishida H, Yap CC, Suzuki M, Kawai J, Suzuki H, Carninci P, Hayashizaki Y, Wells C, Frith M, Ravasi T, Pang KC, Hallinan J, Mattick J, Hume DA, Lipovich L, Batalov S, Engstrom PG, Mizuno Y, Faghihi MA, Sandelin A, Chalk AM, Mottagui-Tabar S, Liang Z, Lenhard B, Wahlestedt C. Antisense transcription in the mammalian transcriptome. *Science*. 2005a; 309:1564–1566. [PubMed: 16141073]
- Katayama S, Tomaru Y, Kasukawa T, Waki K, Nakanishi M, Nakamura M, Nishida H, Yap CC, Suzuki M, Kawai J, Suzuki H, Carninci P, Hayashizaki Y, Wells C, Frith M, Ravasi T, Pang KC, Hallinan J, Mattick J, Hume DA, Lipovich L, Batalov S, Engstrom PG, Mizuno Y, Faghihi MA, Sandelin A, Chalk AM, Mottagui-Tabar S, Liang Z, Lenhard B, Wahlestedt C, Group RGER. Antisense transcription in the mammalian transcriptome. *Science*. 2005b; 309:1564–1566. [PubMed: 16141073]
- Li S, Liberman LM, Mukherjee N, Benfey PN, Ohler U. Integrated detection of natural antisense transcripts using strand-specific RNA sequencing data. *Genome Res*. 2013; 23:1730–1739. [PubMed: 23816784]
- Li X, Wolf ME. Multiple faces of BDNF in cocaine addiction. *Behav Brain Res*. 2014; 279C:240–254.
- Magistri M, Faghihi MA, St Laurent G 3rd, Wahlestedt C. Regulation of chromatin structure by long noncoding RNAs: focus on natural antisense transcripts. *Trends Genet*. 2012; 28:389–396. [PubMed: 22541732]
- Magistri M, Khoury N, Mazza EM, Velmeshev D, Lee JK, Bicciato S, Tsoulfas P, Ali Faghihi M. A comparative transcriptomic analysis of astrocytes differentiation from human neural progenitor cells. *Eur J Neurosci*. 2016
- Maze I, Covington HE, Dietz DM, LaPlant Q, Renthall W, Russo SJ, Mechanic M, Mouzon E, Neve RL, Haggarty SJ, Ren YH, Sampath SC, Hurd YL, Greengard P, Tarakhovsky A, Schaefer A, Nestler EJ. Essential Role of the Histone Methyltransferase G9a in Cocaine-Induced Plasticity. *Science*. 2010; 327:213–216. [PubMed: 20056891]
- Mercer TR, Dinger ME, Sunkin SM, Mehler MF, Mattick JS. Specific expression of long noncoding RNAs in the mouse brain. *Proc Natl Acad Sci U S A*. 2008; 105:716–721. [PubMed: 18184812]
- Michelhaugh SK, Lipovich L, Blythe J, Jia H, Kapatos G, Bannon MJ. Mining Affymetrix microarray data for long non-coding RNAs: altered expression in the nucleus accumbens of heroin abusers. *J Neurochem*. 2011; 116:459–466. [PubMed: 21128942]
- Modarresi F, Faghihi MA, Lopez-Toledano MA, Fatemi RP, Magistri M, Brothers SP, van der Brug MP, Wahlestedt C. Inhibition of natural antisense transcripts in vivo results in gene-specific transcriptional upregulation. *Nat Biotechnol*. 2012; 30:453–459. [PubMed: 22446693]
- Modarresi F, Faghihi MA, Patel NS, Sahagan BG, Wahlestedt C, Lopez-Toledano MA. Knockdown of BACE1-AS Nonprotein-Coding Transcript Modulates Beta-Amyloid-Related Hippocampal Neurogenesis. *Int J Alzheimers Dis*. 2011; 2011:929042. [PubMed: 21785702]
- Mohammad F, Mondal T, Kanduri C. Epigenetics of imprinted long noncoding RNAs. *Epigenetics*. 2009; 4:277–286.

- Nowacki M, Vijayan V, Zhou Y, Schotanus K, Doak TG, Landweber LF. RNA-mediated epigenetic programming of a genome-rearrangement pathway. *Nature*. 2008; 451:153–158. [PubMed: 18046331]
- Pastori C, Peschansky VJ, Barbouth D, Mehta A, Silva JP, Wahlestedt C. Comprehensive analysis of the transcriptional landscape of the human FMR1 gene reveals two new long noncoding RNAs differentially expressed in Fragile X syndrome and Fragile X-associated tremor/ataxia syndrome. *Hum Genet*. 2014; 133:59–67. [PubMed: 24005575]
- Qureshi IA, Mehler MF. Emerging roles of non-coding RNAs in brain evolution, development, plasticity and disease. *Nat Rev Neurosci*. 2012; 13:528–541. [PubMed: 22814587]
- Renthal W, Kumar A, Xiao G, Wilkinson M, Covington HE 3rd, Maze I, Sikder D, Robison AJ, LaPlant Q, Dietz DM, Russo SJ, Vialou V, Chakravarty S, Kodadek TJ, Stack A, Kabbaj M, Nestler EJ. Genome-wide analysis of chromatin regulation by cocaine reveals a role for sirtuins. *Neuron*. 2009; 62:335–348. [PubMed: 19447090]
- Rinn JL, Kertesz M, Wang JK, Squazzo SL, Xu X, Bruggmann SA, Goodnough LH, Helms JA, Farnham PJ, Segal E, Chang HY. Functional demarcation of active and silent chromatin domains in human HOX loci by noncoding RNAs. *Cell*. 2007; 129:1311–1323. [PubMed: 17604720]
- Saxena A, Carninci P. Long non-coding RNA modifies chromatin: epigenetic silencing by long non-coding RNAs. *Bioessays*. 2011; 33:830–839. [PubMed: 21915889]
- Scheele C, Petrovic N, Faghihi MA, Lassmann T, Fredriksson K, Rooyackers O, Wahlestedt C, Good L, Timmons JA. The human PINK1 locus is regulated in vivo by a non-coding natural antisense RNA during modulation of mitochondrial function. *BMC Genomics*. 2007; 8:74. [PubMed: 17362513]
- Silva JP, Lambert G, van Booven D, Wahlestedt C. Epigenomic and metabolic responses of hypothalamic POMC neurons to gestational nicotine exposure in adult offspring. *Genome Med*. 2016; 8:93. [PubMed: 27609221]
- Swanson CJ, Baker DA, Carson D, Worley PF, Kalivas PW. Repeated cocaine administration attenuates group I metabotropic glutamate receptor-mediated glutamate release and behavioral activation: a potential role for Homer. *J Neurosci*. 2001; 21:9043–9052. [PubMed: 11698615]
- Szumliński KK, Ary AW, Lominac KD. Homers regulate drug-induced neuroplasticity: implications for addiction. *Biochem Pharmacol*. 2008; 75:112–133. [PubMed: 17765204]
- Szumliński KK, Dehoff MH, Kang SH, Frys KA, Lominac KD, Klugmann M, Rohrer J, Griffin W 3rd, Toda S, Champiaux NP, Berry T, Tu JC, Shealy SE, During MJ, Middaugh LD, Worley PF, Kalivas PW. Homer proteins regulate sensitivity to cocaine. *Neuron*. 2004; 43:401–413. [PubMed: 15294147]
- Velmeshev D, Magistri M, Faghihi MA. Expression of non-protein-coding antisense RNAs in genomic regions related to autism spectrum disorders. *Mol Autism*. 2013; 4:32. [PubMed: 24007600]
- Yang F, Bi J, Xue X, Zheng L, Zhi K, Hua J, Fang G. Up-regulated long non-coding RNA H19 contributes to proliferation of gastric cancer cells. *FEBS J*. 2012
- Zhang GC, Mao LM, Liu XY, Parelkar NK, Arora A, Yang L, Hains M, Fibuch EE, Wang JQ. In vivo regulation of Homer1a expression in the striatum by cocaine. *Mol Pharmacol*. 2007; 71:1148–1158. [PubMed: 17234898]
- Zhu L, Zhu J, Liu Y, Chen Y, Li Y, Huang L, Chen S, Li T, Dang Y, Chen T. Methamphetamine induces alterations in the long non-coding RNAs expression profile in the nucleus accumbens of the mouse. *BMC Neurosci*. 2015; 16:18. [PubMed: 25884509]

Highlights

- Many genes that have been implicated in addiction overlap with natural antisense transcripts.
- Homer1 antisense transcript is significantly increased in the nucleus accumbens by cocaine.
- Knockdown of Homer1 antisense transcript increases expression of Homer1 *in vitro*.

Figure 1. Bioinformatics pipeline and classification of noncoding antisense RNAs

(A) Schematic of bioinformatics pipeline used to identify natural antisense transcripts. (B) One hundred and twenty-one noncoding antisense (AS) RNAs were identified from 545 genes examined. (C) Gene ontology analysis of NAT-containing genes based on protein class.

Figure 2. Expression of noncoding antisense RNAs in the NAc

Following repeated cocaine or saline injections, NAc was collected for qPCR analysis.

Significant change in *Homer1-AS* expression was revealed in cocaine-treated mice. * $P < 0.0001$ compared to saline via Benjamini and Hochberg false discovery rate test, $n = 7-9$.

Figure 3. *Homer1* and *Homer1-AS* expression following repeated or acute cocaine injections
Repeated cocaine injections increased *Homer1-AS*, but not *Homer1*, in the NAc 2 hours (A) and 10 days (B) after the last injection. C) Increased *Homer1*, but not *Homer1-AS*, expression in the NAc 2 hours following a single injection of cocaine. * $P < 0.005$ and ** $P < 0.0005$ compared to saline via Bonferroni post-hoc test, $n = 6-9$.

Figure 4. Graphical representation of *Homer1* sense and antisense map
Genomic location of mouse antisense (blue) and sense (green) RNA pairs for *Homer1*.

Figure 5. Sense and antisense expression following *Homer1-AS* knockdown

A) Reduction of *Homer1-AS* following siRNA treatment in N2a cells. B) Increase in *Homer1* expression following *Homer1-AS* knockdown in N2a cells. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.0001$ compared to si-Control via Newman-Keuls *post-hoc* test.

Table 1

Antisense RNAs examined in the mouse Nac

Gene name	NAT name	Interaction type	Overlap with sense
adenylate cyclase 2	<i>Adcy2-AS</i>	intronic	69137671-69138140
beta arrestin 1	<i>Arrb1-AS</i>	intronic	106751848-106751978
Brain-derived neurotrophic factor	<i>Bdnf-AS</i>	promoter	109513857-109514010
Chromodomain helicase DNA binding protein 4	<i>Chd4-AS</i>	intronic	125078861-125078948
cholinergic receptor, muscarinic 1B, CNS	<i>Chrm1-AS</i>	promoter	8737279-8737826
catechol-O-methyltransferase	<i>Comt-AS</i>	intronic	18421262-18421605
DNA (Cytosine-5-)-Methyltransferase 3 Alpha	<i>Dnmt3a-AS</i>	intronic	3884675-3884792
dopamine receptor 5	<i>Drd5-AS</i>	intronic	38712870-38713757
GABA receptor 1	<i>Gabbr1-AS</i>	intronic	37169668-37169798
glutamate receptor, AMPA 4, ionotropic	<i>Gria4-AS</i>	intronic	4514665-4514849
glutamate receptor, AMPA 4, ionotropic	<i>Gria3-AS</i>	intronic	39016690-39017360
glutamate kainate receptor 2	<i>Grik2-AS</i>	intronic	48863035-48865786
Glutamate Metabotropic Receptor 7	<i>Grm7-AS</i>	exonic	110595506-110595915
glycogen synthase kinase 3	<i>Gsk3-AS</i>	intronic	38089077-38089347
Homer homolog 1	<i>Homer1-AS</i>	intronic	94073408-94073537
5-HT1a receptor	<i>Htr1a-AS</i>	intronic	106236748-106238201
5-HT receptor 1B	<i>Htr1b-AS</i>	intronic	81525537-81525886
5-HT receptor 1D	<i>Htr1d-AS</i>	intronic	135990295-135990924
5-HT2b receptor	<i>Htr2b-AS</i>	promoter	88009163-88009546
potassium inwardly-rectifying channel, subfamily J, member 6	<i>Kcnj6-AS</i>	exonic	94970290-94970498
potassium channel, subfamily K, member 1	<i>Kcnk1-AS</i>	exonic	128553148-128553570
K large conductance Ca-activated channel, subfamily M, a1	<i>Kcnma1-AS</i>	intronic	24824213-24825059
K large conductance Ca-activated channel, subfamily M, b4	<i>Kcnmb4-AS</i>	exonic	115911266-115911408
potassium voltage-gated channel, subfamily Q, member 1.	<i>Kcnq1-AS</i>	intronic	150399016-150403528
sirtuin 2	<i>Sirt2-AS</i>	intronic	29566257-29566534